

## Terrestrial Small Mammals as Reservoirs of *Mycobacterium ulcerans* in Benin<sup>∇</sup>

Lies Durnez,<sup>1,2\*</sup> Patrick Suykerbuyk,<sup>2</sup> Violaine Nicolas,<sup>3</sup> Patrick Barrière,<sup>4,5</sup> Erik Verheyen,<sup>6</sup>  
Christian R. Johnson,<sup>7</sup> Herwig Leirs,<sup>1,8†</sup> and Françoise Portaels<sup>2‡</sup>

*Evolutionary Ecology, Department of Biology, University of Antwerp, Antwerp, Belgium*<sup>1</sup>; *Mycobacteriology Unit, Department of Microbiology, Institute of Tropical Medicine, Antwerp, Belgium*<sup>2</sup>; *Laboratoire Mammifères et Oiseaux, Département de Systématique et Evolution, Muséum National d'Histoire Naturelle, Paris, France*<sup>3</sup>; *CREG-AICA, Nessadiou BP37, 98870 Bourail, New Caledonia*<sup>4</sup>; *Laboratoire Ecobio UMR6553, Université de Rennes I, 35380 Paimpont, France*<sup>5</sup>; *Vertebrate Department, Royal Belgian Institute of Natural Sciences, Brussels, Belgium*<sup>6</sup>; *Programme National de Lutte contre l'Ulcère de Buruli, Cotonou, Bénin*<sup>7</sup>; and *Danish Pest Infestation Laboratory, University of Aarhus, Kongens Lyngby, Denmark*<sup>8</sup>

Received 26 January 2010/Accepted 23 April 2010

***Mycobacterium ulcerans*, the causative agent of Buruli ulcer (BU), is considered an environmental pathogen. Different mycobacteria were detected in 68 (12%) out of 565 small mammals collected in areas in Benin where BU is endemic. Although *M. ulcerans* was not found, we suggest that more research on *M. ulcerans* in African (small) mammals is needed.**

*Mycobacterium ulcerans* is the causative agent of Buruli ulcer (BU), a serious skin disease (7, 29). Epidemiological evidence strongly associates BU with aquatic ecosystems (29). *M. ulcerans* DNA has been identified in water, fish, aquatic insects, detritus, leeches, crustaceans, mollusks, and mosquitoes (13, 18, 20, 25, 36). However, the difficulty in culturing the bacillus from environmental specimens and the low bacillary concentration shown by PCR (28) strongly suggest that *M. ulcerans* does not multiply in these specimens. Recent findings in Australia show high concentrations of *M. ulcerans* DNA in possum feces in sites where BU is endemic (C. O'Brien et al., presented at the WHO Annual Meeting on Buruli Ulcer, Geneva, Switzerland, 2008; C. O'Brien et al., presented at the WHO Annual Meeting on Buruli Ulcer, Geneva, Switzerland, 2009; J. Fyfe et al., presented at the WHO Annual Meeting on Buruli Ulcer, Geneva, Switzerland, 2010). *M. ulcerans* DNA also has been found in mosquitoes trapped in the same sites of endemicity where the possum feces were collected (18) and in feces of the black rat *Rattus rattus* (Linnaeus, 1758) (O'Brien et al., presented at the WHO Annual Meeting on Buruli Ulcer, Geneva, Switzerland, 2009; Fyfe et al., presented at the WHO Annual Meeting on Buruli Ulcer, Geneva, Switzerland, 2010). Similarly, in West Africa, mammals present in watery environments, such as rodents and insectivores (17), could be a reservoir of *M. ulcerans*. African rodents and insectivores (shrews) can carry pathogenic mycobacteria (9) and are sensitive to experimental infection with *M. ulcerans* (1, 6, 14, 24, 34). Moreover, emergence of BU has been

associated with environmental disturbances (29), which could also alter the transmission of rodent-borne diseases (26). To date, only one study has attempted to systematically culture *M. ulcerans* from rodents in an area of Africa where BU is endemic (Uganda) (31), but since the development of PCR assays, no such study has been carried out. In this study we hypothesize that small terrestrial mammals are part of the reservoir of *M. ulcerans* in which the bacteria can multiply and from which the environment can be contaminated.

By using Sherman live traps and box traps (9), 326 rodents and 222 shrews were caught around bodies of water and in the houses of three villages with high BU endemicity and three villages with low BU endemicity (Table 1) in the dry (January and February) and wet (October and November) seasons of 2006. Animal species identifications (Table 2) were based on external and/or cranio-dental analysis and were confirmed by molecular analysis. Cytochrome *b* gene sequences were compared to those presented by several researchers (8, 21, 23, 27, 32, 35). From each animal, a piece of liver, the spleen, a lung, the mesenteric lymph nodes, and external lesions, if present, were kept in semisolid transport medium (12) at  $-20^{\circ}\text{C}$  until further analysis. The organs of each animal were pooled for analysis by culture and PCR or analyzed individually when the animal presented external or internal lesions. Culture and identification of mycobacteria were performed as described earlier (9), but with inoculation at 30 to 32°C (22) and additional use of charcoal medium (30). DNA was extracted using the modified Boom method (2, 10) and amplified in a nested PCR specific for all mycobacteria (9) and specific for *M. ulcerans* (33). From 49 (8.7%) animals, nontuberculous mycobacteria were cultured, but no *M. ulcerans* was isolated. Most of the mycobacterial isolates in this study can cause disease in humans (Table 3). Twenty-six animals (4.6%) were positive for mycobacteria by PCR, but no *M. ulcerans* was detected. Com-

\* Corresponding author. Mailing address: Mycobacteriology Unit, Department of Microbiology, Institute of Tropical Medicine, Nationalestraat 155, B-2000 Antwerp, Belgium. Phone: 3 2476311. Fax: 3 2476333. E-mail: ldurnez@itg.be.

† H.L. and F.P. contributed equally to this study.

∇ Published ahead of print on 30 April 2010.

TABLE 1. Location of the study villages

CDTUB <sup>a</sup>	Village	No. of BU cases in 2005-2006	Level of BU endemicity	Geographical coordinate	
				Latitude	Longitude
Lalo	Tandji	25	High	6.94122098	1.97169973
	Adjassagon	2	Low	7.00175107	1.95172347
Zagnanado	Houedja	6	High	7.13986237	2.44355033
	Agonvè	2	Low	7.25431818	2.45778188
Allada	Sedje Houégoudo	11	High	6.74590738	2.37206443
	Ahonzonoude	0	Low	6.8136599	2.37536363

<sup>a</sup> CDTUB, Centres de Dépistage et Traitement d'Ulçère de Buruli.

binning culture and PCR, a total of 68 animals (12.0%) were carriers of mycobacteria. In 14 (15.5%) out of 90 fecal samples collected from a subset of the trapped animals, mycobacteria were detected by PCR, but *M. ulcerans* was not detected. Whether rodents and shrews can, indeed, transmit mycobacteria, e.g., by excretion in their feces, should be investigated further experimentally.

Although a slightly higher presence of mycobacteria was found in the animals trapped in the villages with high BU endemicity (13.8%) than in villages with low BU endemicity (9.8%), the difference was not statistically significant ( $P = 0.162$ ). Eddyani et al. (11) did find more mycobacteria in amoebae in areas of high BU endemicity than in areas of low BU endemicity.

Similar to findings of a previous study carried out in Tanzania (9), the presence of mycobacteria in shrews (21.2%) was significantly higher than in rodents (6.1%) ( $P < 0.001$ ). For shrews, a significantly higher presence of myco-

bacteria was found in the wet season (33.3%) than in the dry season (10.3%) ( $P < 0.001$ ). On the other hand, for rodents we found that in the dry season relatively more mycobacteria were present (8.5%) than in the wet season (2.7%) ( $P = 0.025$ ). These findings could be due to a difference in behavior or feeding habits between shrews and rodents. Shrews mainly forage on invertebrates from the ground surface and among leaf litter (4, 5). Several mycobacteria have already been found in several invertebrates (15, 16, 18, 20, 28). It is possible that the seasonal distribution of mycobacteria in shrews observed in this study is a consequence of a seasonal distribution of mycobacteria in the invertebrates on which the shrews forage although no information is available on seasonality of mycobacteria in invertebrates. In other studies on environmental mycobacteria, more mycobacteria were found in the environment (soil and water) in the dry season than in the wet season (3), which could be a possible explanation for the seasonality of mycobacteria in rodents.

The fact that *M. ulcerans* was not found in the animals collected in the present study could be due to several factors. The size and type of the traps favor certain species of rodents and shrews. Some animal species are too large to enter the traps or too small to trigger them. Additionally, several animal species were caught in low numbers only. The prevalence of BU in humans varies between 0 and 5.61% in villages in the district of Lalo (Benin) (19). In order to have 95% probability of trapping at least one positive individual, assuming a prevalence of between 5 and 10%, we would need to test between 30 and 80 animals per species in a certain area, which is more than the numbers we have trapped for most species.

The fact that we did not find *M. ulcerans* DNA in the feces of *R. rattus* trapped in Benin although it has been detected in the same species in Australia could be due to a lower sensitivity of our methods (gel-based PCR in the present study versus real-time PCR in the study of C. O'Brien et al. (presented at the WHO Annual Meeting on Buruli Ulcer, Geneva, Switzerland, 2009). However, it is also possible that in Australia *R. rattus* obtained *M. ulcerans* only from eating contaminated possum feces while a similar source of *M. ulcerans* is absent in Benin.

*M. ulcerans* disease in wild and domestic animals has never been described in the literature from any of the West and Central African countries, probably because of the lack of attention to diseases in wild (and domestic) animals in this region. Taking all the above into consideration, we do

TABLE 2. Number of animals collected and number of animals positive for mycobacteria per animal species

Animal species	No. of animals collected	No. of animals positive for mycobacteria
<b>Rodents</b>		
<i>Rattus rattus</i> (Linnaeus, 1758)	78	4
<i>Lemniscomys striatus</i> (Linnaeus, 1758)	66	5
<i>Mastomys natalensis</i> (Smith, 1834)	64	3
<i>Praomys misonnei</i> Van der Straeten & Dieterlen, 1987	52	5
<i>Praomys cf. derooi</i> Vanderstraeten & Verheyen, 1978	38	1
<i>Praomys</i> sp. n. 1	7	0
<i>Uranomys ruddi</i> Dollman, 1909	7	0
<i>Mus</i> ( <i>Nannomys</i> ) spp.	6	1
<i>Dasymys bentleyae</i> (Thomas, 1892)	3	0
<i>Hybomys cf. trivirgatus</i>	1	0
<i>Hylomyscus</i> sp.	1	0
<i>Mastomys erythroleucis</i> (Temminck, 1853)	1	1
<i>Mus</i> sp.	1	0
<i>Taterillus gracilis</i> (Thomas, 1892)	1	0
<i>Thryonomys swinderianus</i> (Temminck, 1827)	14	1
<i>Xerus erythropus</i> Desmarest, 1817	3	0
<b>Insectivores</b>		
<i>Crocidura cf. foxi</i> Dollman, 1915	146	31
<i>Crocidura olivieri</i> (Lesson, 1827)	56	13
<i>Crocidura</i> spp.	20	3

TABLE 3. Mycobacteria isolated from small mammals in areas of high and low BU endemicity assigned to risk groups<sup>a</sup>

Risk group and isolated mycobacterium <sup>b</sup>	Small mammal (n) <sup>c</sup>	Field code	Location (BU endemicity level)	Trapping site	Season	Comment <sup>d</sup>	
<b>Risk group 2</b>							
<i>Mycobacterium scrofulaceum</i>	<i>Crocidura cf. foxi</i> (3)	BN899, BN902, BN934	Zagnanado (high)	Near water body	Wet		
	<i>Crocidura olivieri</i> (2)	BN906, BN961	Zagnanado (high)	Near water body	Wet		
	<i>Crocidura cf. foxi</i> (2)	BN968, BN974	Zagnanado (low)	Near water body	Wet		
	<i>Crocidura cf. foxi</i> (1)	BN986	Allada (high)	Near water body	Wet		
	<i>Crocidura olivieri</i> (1)	BN997	Allada (high)	Near water body	Wet		
	<i>Crocidura olivieri</i> (1)	BN998	Allada (high)	House	Wet		
	<i>Crocidura olivieri</i> (1)	BN1062	Allada (low)	Near water body	Wet	Feces PCR+	
	<i>Mycobacterium simiae</i>	<i>Crocidura cf. foxi</i> (1)	BN321	Lalo (high)	Near water body	Dry	
		<i>Lemniscomys striatus</i> (1)	BN1007	Lalo (high)	Near water body	Dry	
		<i>Crocidura olivieri</i> (1)	BN825	Lalo (low)	Near water body	Wet	
		<i>Crocidura cf. foxi</i> (2)	BN912, BN959	Zagnanado (high)	Near water body	Wet	
		<i>Crocidura sp.</i> (1)	BN947	Zagnanado (high)	Near water body	Wet	
		<i>Crocidura cf. foxi</i> (1)	BN341	Zagnanado (high)	Near water body	Dry	
		<i>Mastomys erythroleucus</i> (1)	BN410	Zagnanado (high)	Near water body	Dry	Spleen, ML
<i>Mus (Nannomys) sp.</i> (1)		BN980	Zagnanado (low)	Near water body	Wet		
<i>Crocidura cf. foxi</i> (1)		BN987	Allada (high)	Near water body	Wet		
<i>Crocidura olivieri</i> (1)		BN1017	Allada (high)	House	Wet		
<i>Mycobacterium avium</i> complex	<i>Crocidura cf. foxi</i> (1)	BN1020	Allada (high)	Near water body	Wet		
	<i>Praomys misonnei</i> (1)	BN308	Lalo (high)	Near water body	Dry	Lung	
	<i>Crocidura sp.</i> (1)	BN875	Lalo (high)	House	Dry		
<i>Mycobacterium intracellulare</i>	<i>Praomys misonnei</i> (1)	BN346	Zagnanado (high)	Near water body	Dry	Wound on back	
	<i>Crocidura olivieri</i> (1)	BN300	Lalo (high)	House	Dry		
<i>Mycobacterium asiaticum</i> (-like)	<i>Crocidura cf. foxi</i> (1)	BN969	Zagnanado (low)	House	Dry		
	<i>Crocidura sp.</i> (1)	BN1061	Allada (low)	House	Wet		
<i>Mycobacterium shimoidei</i> -like	<i>Praomys misonnei</i> (1)	BN308	Lalo (high)	Near water body	Dry	ML	
	<i>Mastomys natalensis</i> (1)	BN474	Zagnanado (low)	House	Dry	Tail	
<b>Risk group 1</b>							
<i>Mycobacterium interjectum</i>	<i>Crocidura olivieri</i> (1)	BN900	Zagnanado (high)	Near water body	Wet		
	<i>Crocidura cf. foxi</i> (1)	BN1002	Allada (high)	Near water body	Wet		
<i>Mycobacterium lentiflavum</i>	<i>Mus (Nannomys) sp.</i> (1)	BN980	Zagnanado (low)	Near water body	Wet		
	<i>Crocidura cf. foxi</i> (1)	BN337	Zagnanado (high)	Near water body	Dry	Spleen	
<i>Mycobacterium triplex</i>	<i>Crocidura cf. foxi</i> (1)	BN478	Zagnanado (low)	Near water body	Dry		
<b>Not assigned to a risk group</b>							
<i>Mycobacterium paraffinicum</i> (-like)	<i>Crocidura cf. foxi</i> (1)	BN456	Zagnanado (low)	Near water body	Dry	Spleen, lung	
	<i>Mastomys natalensis</i> (1)	BN517	Zagnanado (low)	Near water body	Dry	Lung	
<i>Mycobacterium saskatchewanense</i>	<i>Crocidura cf. foxi</i> (1)	BN206	Lalo (high)	Near water body	Dry		
<i>Mycobacterium sherrisii</i>	<i>Crocidura cf. foxi</i> (1)	BN967	Zagnanado (low)	Near water body	Wet		
	<i>Crocidura cf. foxi</i> (2)	BN1015, BN1027	Allada (high)	Near water body	Wet	Feces PCR+	
	<i>Mastomys natalensis</i> (1)	BN958	Zagnanado (high)	House	Wet		
<i>Mycobacterium colombiense</i>	<i>Crocidura olivieri</i> (1)	BN1001	Allada (high)	Near water body	Wet		
	<i>Crocidura cf. foxi</i> (2)	BN1002, BN1030	Allada (high)	Near water body	Wet	Feces PCR+	
<i>Mycobacterium angelicum</i>	<i>Crocidura olivieri</i> (1)	BN1043	Allada (low)	Near water body	Wet		
<i>Mycobacterium barombii</i>	<i>Crocidura cf. foxi</i> (1)	BN990	Allada (high)	Near water body	Wet		
<i>Mycobacterium spp.</i>	<i>Crocidura cf. foxi</i> (2)	BN970, BN982	Zagnanado (low)	Near water body	Wet		
	<i>Crocidura olivieri</i> (1)	BN291	Lalo (high)	House	Dry		

<sup>a</sup> Leão et al. (22).<sup>b</sup> Risk group 2 contains pathogens that pose a moderate individual risk and of which disease with average severity exists in the community. Risk group 1 contains pathogens that pose a low risk of infection for both the human individual and the community. Diseases are never or rarely described in normal adults (22).<sup>c</sup> n, number of animals.<sup>d</sup> The body site from which the mycobacterium is isolated (ML, mesenteric lymph nodes) is mentioned. If no body site is mentioned, the mycobacterium was isolated from the pooled organs. Feces PCR+, the feces sample was also positive by PCR.

not reject our initial hypothesis that rodents or shrews are part of the reservoir; instead, we broaden it to other mammals.

This research was supported by a Ph.D. grant of the Flemish Interuniversity Council, the Directorate-General for Development Cooperation (Brussels, Belgium), the Damien Foundation, and the European Union (project INCOCT-2005-051476-BURULICO).

We thank our laboratory staff for their excellent technical assistance and all field staff for their support during the field work.

#### REFERENCES

1. Addo, P., B. Adu-Addai, M. Quartey, M. Abbas, I. Okang, E. Owusu, D. Ofori-Adjei, and B. Awumbila. 2007. Clinical and histopathological presentation of Buruli ulcer in experimentally infected grasscutters (*Thryonomys swinderianus*). *Internet J. Trop. Med.* 3:e2.
2. Boom, R., C. J. Sol, M. M. Salimans, C. L. Jansen, P. M. Wertheim-van Dillen, and J. van der Noorda. 1990. Rapid and simple method for purification of nucleic acids. *J. Clin. Microbiol.* 28:495–503.
3. Chilima, B. Z., I. M. Clark, S. Floyd, P. E. Fine, and P. R. Hirsch. 2006. Distribution of environmental mycobacteria in Karonga District, northern Malawi. *Appl. Environ. Microbiol.* 72:2343–2350.
4. Churchfield, S. 1990. The natural history of shrews. Christopher Helm/A. & C. Black, London, United Kingdom.
5. Churchfield, S., P. Barrière, R. Hutterer, and M. Colyn. 2004. First results on the feeding ecology of sympatric shrews (Insectivora: Soricidae) in the Tai National Park, Ivory Coast. *Acta Theriol.* 49:1–15.
6. Clancey, J. K. 1964. Mycobacterial skin ulcers in Uganda: description of a new *Mycobacterium* (*Mycobacterium buruli*). *J. Pathol. Bacteriol.* 88:175–187.
7. Debacker, M., J. Aguiar, C. Steunou, C. Zinsou, W. M. Meyers, A. Guédénon, J. T. Scott, M. Dramaix, and F. Portaels. 2004. *Mycobacterium ulcerans* disease (Buruli ulcer) in rural hospital, Southern Benin, 1997–2001. *Emerg. Infect. Dis.* 10:1391–1398.
8. Dobigny, G., E. Lecompte, C. Tatard, P. Gauthier, K. Ba, C. Denys, J. M. Duplantier, and L. Granjon. 2008. An update on the taxonomy and geographic distribution of the cryptic species *Mastomys kollmannspergeri* (Muridae, Murinae) using combined cytogenetic and molecular data. *J. Zool.* 276:368–374.
9. Durnez, L., M. Eddyani, G. F. Mgone, A. Katakweba, C. R. Katholi, R. R. Machang'u, R. R. Kazwala, F. Portaels, and H. Leirs. 2008. First detection of mycobacteria in African rodents and insectivores, using stratified pool screening. *Appl. Environ. Microbiol.* 74:768–773.
10. Durnez, L., P. Stragier, K. Roebben, A. Ablordey, H. Leirs, and F. Portaels. 2009. A comparison of DNA extraction procedures for the detection of *Mycobacterium ulcerans*, the causative agent of Buruli ulcer, in clinical and environmental specimens. *J. Microbiol. Methods* 76:152–158.
11. Eddyani, M., J. F. De Jonckheere, L. Durnez, P. Suykerbuyk, H. Leirs, and F. Portaels. 2008. Occurrence of free-living amoebae in communities of low and high endemicity for Buruli ulcer in southern Benin. *Appl. Environ. Microbiol.* 74:6547–6553.
12. Eddyani, M., M. Debacker, A. Martin, J. Aguiar, C. R. Johnson, C. Uwizeye, K. Fissette, and F. Portaels. 2008. Primary culture of *Mycobacterium ulcerans* from human tissue specimens after storage in semisolid transport medium. *J. Clin. Microbiol.* 46:69–72.
13. Eddyani, M., D. Ofori-Adjei, G. Teugels, D. De Weirtd, D. Boakye, W. M. Meyers, and F. Portaels. 2004. Potential role for fish in transmission of *Mycobacterium ulcerans* disease (Buruli ulcer): an environmental study. *Appl. Environ. Microbiol.* 70:5679–5681.
14. Fenner, F. 1956. The pathogenic behavior of *Mycobacterium ulcerans* and *Mycobacterium balnei* in the mouse and the developing chick embryo. *Am. Rev. Tuberc.* 73:650–673.
15. Fischer, O., L. Matlova, L. Dvorska, P. Svastova, J. Bartl, I. Melicharek, R. T. Weston, and I. Pavlík. 2001. Diptera as vectors of mycobacterial infections in cattle and pigs. *Med. Vet. Entomol.* 15:208–211.
16. Fischer, O. A., L. Matlova, J. Bartl, L. Dvorska, P. Svastova, R. du Maine, I. Melicharek, M. Bartos, and I. Pavlík. 2003. Earthworms (Oligochaeta, Lumbricidae) and mycobacteria. *Vet. Microbiol.* 91:325–338.
17. Hanney, P. W. 1975. Rodents: their lives and habits. Taplinger, New York, NY.
18. Johnson, P. D., J. Azuolas, C. J. Lavender, E. Wishart, T. P. Stinear, J. A. Hayman, L. Brown, G. A. Jenkin, and J. A. Fyfe. 2007. *Mycobacterium ulcerans* in mosquitoes captured during outbreak of Buruli ulcer, southeastern Australia. *Emerg. Infect. Dis.* 13:1653–1660.
19. Johnson, R. C., M. Makoutode, G. E. Sopoh, P. Elsen, J. Gbovi, L. H. Pouteau, W. M. Meyers, M. Boko, and F. Portaels. 2005. Buruli ulcer distribution in Benin. *Emerg. Infect. Dis.* 11:500–501.
20. Kotłowski, R., A. Martin, A. Ablordey, K. Chemlal, P. A. Fonteyne, and F. Portaels. 2004. One-tube cell lysis and DNA extraction procedure for PCR-based detection of *Mycobacterium ulcerans* in aquatic insects, molluscs and fish. *J. Med. Microbiol.* 53:927–933.
21. Kouassi, S. K., V. Nicolas, V. Aniskine, A. Lalis, C. Cruaud, A. Couloux, M. Colyn, M. Dosso, L. Koivogui, E. Verheyen, C. Akoua-Koffi, and C. Denys. 2008. Taxonomy and biogeography of the African pygmy mice, subgenus *Nannomys* (Rodentia, Murinae, *Mus*) in Ivory Coast and Guinea (West Africa). *Mammalia* 72:237–252.
22. Leão, S. C., A. Martin, G. I. Mejia, J. Palomino, J. Robledo, M. A. da Silva Telles, and F. Portaels. 2004. Practical handbook for the phenotypic and genotypic identification of mycobacteria. Vanden Broele, Bruges, Belgium.
23. Lecompte, E., K. Aplin, C. Denys, F. Catzeflis, M. Chades, and P. Chrevet. 2008. Phylogeny and biogeography of African Murinae based on mitochondrial and nuclear gene sequences, with a new tribal classification of the subfamily. *BMC Evol. Biol.* 8:199.
24. MacCallum, P., G. Buckle, and H. A. Sissons. 1948. A new mycobacterial infection in man. *J. Pathol. Bacteriol.* 60:93–122.
25. Marsollier, L., R. Robert, J. Aubry, J. P. Saint Andre, H. Kouakou, P. Legras, A. L. Manceau, C. Mahaza, and B. Carboneille. 2002. Aquatic insects as a vector for *Mycobacterium ulcerans*. *Appl. Environ. Microbiol.* 68:4623–4628.
26. Mills, J. N. 2005. Regulation of rodent-borne viruses in the natural host: implications for human disease. *Arch. Virol. Suppl.* 19:45–57.
27. Nicolas, V., B. Akpatou, W. Wendelen, J. Kerbis Peterhans, A. Olayemi, J. Decher, A. D. Missou, C. Denys, P. Barrière, C. Cruaud, and M. Colyn. Molecular and morphometric variation in two sibling species of the genus *Praomys* (Rodentia: Muridae): implications for biogeography. *Zool. J. Linn. Soc.*, in press.
28. Portaels, F., W. M. Meyers, A. Ablordey, A. G. Castro, K. Chemlal, P. de Rijk, P. Elsen, K. Fissette, A. G. Fraga, R. Lee, E. Mahrous, P. L. Small, P. Stragier, E. Torrado, A. Van Aerde, M. T. Silva, and J. Pedrosa. 2008. First cultivation and characterization of *Mycobacterium ulcerans* from the environment. *PLoS Negl. Trop. Dis.* 2:e178.
29. Portaels, F., M. T. Silva, and W. M. Meyers. 2009. Buruli ulcer. *Clin. Dermatol.* 27:291–305.
30. Realini, L., K. De Ridder, B. Hirschel, and F. Portaels. 1999. Blood and charcoal added to acidified agar media promote the growth of *Mycobacterium genavense*. *Diagn. Microbiol. Infect. Dis.* 34:45–50.
31. Reville, W. D. L., R. H. J. Morrow, W. Parson, and J. W. M. Kiryabwire. 1972. *Mycobacterium ulcerans* infection (Buruli ulcer), p. 19–31. In A. G. Shaper, J. W. Kibukamusoke, and M. S. R. Hutt (ed.), *Medicine in a tropical environment*. British Medical Association, London, United Kingdom.
32. Robins, J. H., M. Hingston, E. Matisoo-Smith, and H. A. Ross. 2007. Identifying *Rattus* species using mitochondrial DNA. *Mol. Ecol. Notes* 7:717–729.
33. Ross, B. C., L. Marino, F. Oppedisano, R. Edwards, R. M. Robins-Browne, and P. D. Johnson. 1997. Development of a PCR assay for rapid diagnosis of *Mycobacterium ulcerans* infection. *J. Clin. Microbiol.* 35:1696–1700.
34. Singh, N. B., A. Srivastava, V. K. Verma, A. Kumar, and S. K. Gupta. 1984. *Mastomys natalensis*: a new animal model for *Mycobacterium ulcerans* research. *Indian J. Exp. Biol.* 22:393–394.
35. Verheyen, W. N., J. L. J. Hulselmans, T. Dierckx, M. Colyn, H. Leirs, and E. Verheyen. 2003. A craniometric and genetic approach to the systematics of the genus *Dasymys*, Peters, 1875, selection of a neotype and description of three new taxa (Rodentia, Muridae, Africa). *Bull. Inst. R. Sci. Nat. Belg. Biol.* 73:27–71.
36. Williamson, H. R., M. E. Benbow, K. D. Nguyen, D. C. Beachboard, R. K. Kimbirauskas, M. D. McIntosh, C. Quayle, E. O. Ampadu, D. Boakye, R. W. Merritt, and P. L. Small. 2008. Distribution of *Mycobacterium ulcerans* in buruli ulcer endemic and non-endemic aquatic sites in Ghana. *PLoS Negl. Trop. Dis.* 2:e205.