



Published in final edited form as:

*Transplant Proc.* 1987 October ; 19(5 Suppl 6): 57–61.

## Canine Kidney Transplantation With FK-506 Along or in Combination With Cyclosporine and Steroids

S. Todo, A.J. Demetris, Y. Ueda, O. Imventarza, K. Okuda, A. Casavilla, S. Cemaj, A. Ghalab, V. Mazzaferro, B.S. Rhoe, Y. Tonghua, L. Makowka, and T.E. Starzl

From the Department of Surgery, University Health Center of Pittsburgh, and the Veterans Administration Medical Center, Pittsburgh

### Abstract

In this study, the new drug FK-506 (FK) was tested in dogs for its ability to prevent rejection of kidneys when given alone, with small doses of cyclosporine or steroids, or both.

### MATERIALS AND METHODS

Mongrel donors weighing 13 to 15 kg provided kidneys for beagle recipients of the same size under 25 mg/kg pentobarbital anesthesia intravenously. When needed, supplementary pentobarbital or ketamine, 2 mg/kg, or both, were given intravenously. The renal grafts were placed intraabdominally by connecting the renal artery and vein to the recipient iliac vessels and using uretero-neocystotomies. The recipient native kidneys were removed at the same operation. One gram of cephalosporin was given intravenously or intramuscularly during the operation and for three days afterwards. A diet was started the next day.

All immunosuppressive drugs were given orally starting on the morning after transplantation and continuing daily. To prevent vomiting caused by FK or other postoperative factors, 2 mg atropine sulphate was given intramuscularly twice a day for the first week and once a day for the second week. During the third week, the atropine dose was reduced to 1 mg/d.

The FK was provided in powder form by the Fujisawa Pharmaceutical Co, Ltd, Osaka, Japan, and placed in commercial capsules. The dogs were docile well-trained beagles who allowed deep digital placement of the capsules in the oropharynx. Prednisone (Pred) was given as a 5 mg tablet. Cyclosporine (CyA) in the commercial oil carrier used clinically was given orally by syringe.

### FK Administration Alone

FK was the sole immunosuppressive treatment. The animals were placed in four groups: I, untreated control, n = 6; II, FK (0.5 mg/kg/d), n = 6; III, FK (1.0 mg/kg/d), n = 6; and IV, FK (1.5 mg/kg/d), n = 6.

### FK in Combination

The smallest dose in the foregoing studies (group II, 0.5 mg/kg/d) or half this dose was combined with small doses of other agents as follows: group V, FK (0.5 mg/kg/d), CyA (5.0

mg/kg/d), Pred (5.0 mg/d), n = 6; group VI, FK (0.25 mg/kg/d), CyA (2.50 mg/kg/d), Pred (2.5 mg/d), n = 6; and group VII, CyA (5.0 mg/kg/d), Pred (5.0 mg/d), n = 6.

The results in group II were considered to be a single-agent control for those obtained with the triple-drug combination of group V. The animals of group VII who received CyA and Pred were considered controls for the triple-drug, FK-containing regimen of group V.

## Biochemical and Pathological Studies

Blood samples were taken every three mornings for the measurement of levels of blood urea nitrogen, creatinine, SGOT, and total bilirubin. Complete postmortem examination was performed immediately after the animals died. Tissues were fixed with formalin and stained with hematoxylin and eosin. Histopathologic changes were scored blindly according to a subjective scale from 1 to 4.

## Statistics

The Wilcoxon rank sum test and Student's *t* test were applied for the statistical analysis.

## RESULTS

The outcome in each individual experiment is summarized in Table 1. The untreated animals died of rejection in eight to 19 days. Survival was significantly prolonged with all doses of FK, the best results being with 1.5 mg/kg. With this high dose, rejection as judged histopathologically was absent or mild in the two animals who provided tissue for histopathologic analysis. The other four are alive after 39 to 71 days. Increases in serum creatinine levels caused by rejection were minimized or avoided in animals receiving 1.0 and 1.5 mg/kg/d (Fig 1). With doses of 0.5 mg/kg/d, rejection was always found histopathologically (Table 1), and elevated creatinine levels were the rule (Fig 1).

However, animals treated with the higher doses of FK had significant evidence on histopathologic examination of toxicity involving the liver, heart, and pancreas (Table 2). The findings were similar to those reported elsewhere in this symposium in dogs not subjected to transplantation.

Vomiting and weight loss were common in all of the treatment groups. Severe emaciation with loss of more than 30% of the body weight was seen in four of the 18 FK-treated animals. Two animals each in groups II and III died with severe rejection plus lethal emaciation.

## FK in Combination

Five of six dogs in group V that were treated with 0.5 mg/kg FK, 5 mg/kg CyA, and 5 mg Pred are alive with normal renal function after 51 to 71 days. The survival results (Table 3) were better than those in group II in which 0.5 mg/kg FK was used alone ( $P < .01$ ) and better than those in group VII in which 5 mg/kg CyA and Pred were given ( $P < .01$ ). None of the animals in group V lost body weight.

The superiority of the triple-drug combination of group V was also evident from the serum creatinine levels (Fig 2). However, the three drugs when given at half dosage (group VI) were no longer effective (Table 3 and Fig 2).

## DISCUSSION

The principal studies of FK-506 in dogs by Ochiai et al<sup>1</sup> and those by us<sup>2</sup> have not yet been published. Ochiai's Chiba University group had almost universal survival after canine renal

transplantation without much evidence of toxicity.<sup>1</sup> Our findings were less dramatic or favorable. Rejection often was not controlled, even at doses that led to profound illness or sometimes fatal emaciation. In addition, there was histopathologic evidence of toxicity in multiple extrarenal organs.

The toxicity of dogs may be a species-specific phenomenon. Rats have tolerated high-dose FK therapy well.<sup>3,4</sup> The effect of FK on subhuman primates is under study at several institutions, but no results are available at this time. However, even if FK proves to have substantial dose-related toxicity in all species including humans, it could still be a valuable agent for clinical use. The synergism of small doses of FK with suboptimal doses of CyA and steroids was striking in our canine experiments herein reported. The results were consonant with those using the same drug combination in rats submitted to heterotopic heart transplantation.<sup>5</sup> With *in vitro* studies of human lymphocytes, Zeevi et al<sup>6</sup> also have obtained convincing evidence for a true synergism as opposed to an additive effect of FK and CyA.

More and more, polypharmaceutical therapy has been used to allow potent immunosuppressive agents to be used in doses small enough so that side effects inherent with high doses of single agents can be avoided.

## SUMMARY


The immunosuppressive agent FK permitted increased kidney transplant survival in dogs over a wide dose range, but with weight loss and manifold evidence of toxicity. The best use of FK at low doses was in combination with CyA and Pred.

## Acknowledgments


Supported by research grants from the Veterans Administration and Project Grant No. AM-29961 from the National Institutes of Health, Bethesda, MD.

## REFERENCES

1. Ochini T, Nagata M, Nakajima K, et al. Transplantation. 1987 (in press).
2. Todo S, Makowka L, Starzl TE. Surg Res Comm. 1987 (in press).
3. Ochiai T, Nakajima K, Nagata M, et al. Transplant Proc 1987;19:1284. [PubMed: 2484094]
4. Lee P, Murase N, Todo S, et al. Surg Res Comm. 1987 (in press).
5. Murase N, Lee P, Lei H, et al. Transplant Proc. (this issue).
6. Zeevi A, Duquesnoy R, Eiras G, et al. Transplant Proc. (this issue).



**Fig. 1.** Influence of FK dose on serum creatinine level. The rises in the serum creatinine concentration in animals of all groups were usually caused by rejection.



**Fig. 2.** Serum creatinine levels in dogs treated with drug combinations. Note the superior results in group V.

**Table 1**

Effect of FK Dose on Results of Canine Kidney Transplantation

Groups	Oral Dosage (mg/kg/d)	Animal No	Survival (d)	Cause of Death	Lastest Value		Histological severity of Rejection		
					BUN (mg/dl)	Cr (mg/dl)			
I	0	3	19	Rejection	359	14.1	Severe		
		4	9	Rejection	154	8.2	Severe		
		5	12	Rejection	142	16.0	Severe		
		6	11	Rejection	271	13.5	Severe		
		7	8	Rejection	193	13.0	Severe		
		8	11	Rejection	135	8.7	Severe		
		II	0.5	102	16	Rejection	172	13.5	Moderate
				104	24	Intussusception	44	1.1	Mild
109	>70			—	77	4.3	—		
112	44			Rejection	195	5.1	Moderate		
121	13			Rejection	336	7.7	Moderate		
122	28			Rejection	140	14.4	Severe		
III	1.0	101	37	Unknown	154	2.5	Moderate		
		105	63	Unknown	24	1.0	Severe		
		108	16	Rejection	219	6.3	Severe		
		114	21	Unknown	59	2.9	Severe		
		123	14	Unknown	39	2.0	None		
		124	35	Unknown	41	1.5	Moderate		
		IV	1.5	103	>71	—	20	1.0	—
				107	28	Pancreatitis (?)	108	8.8	Mild
113	>67			—	20	1.3	—		
125	17			Unknown	42	1.2	None		
127	>44			—	19	0.9	—		
133	>39	—	95	4.0	—				

Abbreviation: Cr, creatinine.

**Table 2**  
 Histological Changes in Extrarenal Organs After Kidney Transplantation Under FK

Group	Dosage (mg/kg/d)	Liver		Heart		Pancreas: Acinar cell Degeneration (Focal Necrosis)
		Swelling of Hepatocytes	Fatty Change of Hepatocytes	Arteritis	Infarction (Focal Large)	
I	0	0/6 (0%)	1/6 (16%)	0/6 (0%)	0/6 (0%)	0/6 (0%)
II	0.5	1/5 (20%)	1/5 (20%)	0/5 (0%)	0/5 (0%)	0/5 (0%)
III	1.0	1/6 (16%)	0/6 (0%)	2/6 (33%)	2/6 (33%)	3/6 (50%)
IV	1.5	1/2 (50%)	0/2 (0%)	1/2 (50%)	1/2 (50%)	2/2 (100%)

**Table 3**  
 Combination Therapy With Low Doses of FK, CyA, and Pred After Canine Kidney Transplantation

Groups	Treatment Dose (Oral)	Animal No.	Survival (d)	Cause of Death	Latest Value	
					BUN (mg/dL)	Cr (mg/dL)
V	FK, 0.5 mg/kg/d CyA, 5/0 mg/kg/d Pred, 5.0 mg/d	106	>71	—	16	1.0
		115	23	Unknown	159	3.2
		116	>60	—	24	1.3
		118	>53	—	16	1.2
		119	>51	—	29	1.0
VI	FK, 0.25 mg/kg/d CyA, 2.5 mg/kg/d Pred, 2.5 mg/d	120	>51	—	24	1.2
		129	13	Rejection	94	6.6
		130	19	Rejection	113	22.4
		132	31	Rejection	107	6.2
		134	30	Rejection	176	11.6
VII	CyA, 5.0 mg/kg/d Pred, 5.0 mg/d	135	19	Rejection	123	6.5
		136	26	Rejection	105	9.3
		137	>24	—	30	1.9
		138	12	Rejection	288	13.3
		139	12	Rejection	181	8.9
140	12	Rejection	147	15.5		
141	>12	—	120	12.5		
142	>12	—	52	2.8		