

Breast cancer care in the Canada and the United States: Ecological comparisons of extremely impoverished and affluent urban neighborhoods

Kevin M. Gorey^{a,*}, Isaac N. Luginaah^b, Caroline Hamm^c, Karen Y. Fung^d, and Eric J. Holowaty^e

^a School of Social Work, University of Windsor, 401 Sunset Avenue, Windsor, Ontario, Canada N9B 3P4

^b Department of Geography, University of Western Ontario, London, ON, Canada

^c Clinical Trials and Research, Windsor Regional Cancer Center, Windsor, ON, Canada

^d Department of Mathematics and Statistics, University of Windsor, Windsor, ON, Canada

^e Population Studies and Surveillance, Cancer Care Ontario, Toronto, ON, Canada

Abstract

This study examined the differential effect of extreme impoverishment on breast cancer care in urban Canada and the United States. Ontario and California registry-based samples diagnosed between 1998 and 2000 were followed until 2006. Extremely poor and affluent neighborhoods were compared. Poverty was associated with non-localized disease, surgical and radiation therapy (RT) waits, nonreceipt of breast conserving surgery, RT and hormonal therapy, and shorter survival in California, but not in Ontario. Extremely poor Ontario women were consistently advantaged on care indices over their California counterparts. More inclusive health insurance coverage in Canada seems the most plausible explanation for such Canadian breast cancer care advantages.

Keywords

Breast cancer care; Poverty; Affluence; Survival; United States; Canada

1. Introduction

Social, political and economic forces converged in twentieth century America to produce extreme socioeconomic segregation in and around many urban places. Extremely poor neighborhoods tended to concentrate in inner-cities at the same time that extremely affluent neighborhoods were developing in suburban to exurban areas that tended to sprawl away from cities. Extremes of impoverishment and affluence and so relative socioeconomic inequities have fluctuated over recent generations, but distinct very low-income ghettos and well-to-do enclaves clearly persist in twenty-first century urban America. Such

*Corresponding author. Tel.: +1519 253 3000x3085; fax: +1519 9737036.

socioeconomic extremes are not unknown in Canada (Duncan et al., 1993; Gorey, 1998), but perhaps because of their greater prevalence and apparent virulence as well as their stronger association with race in America, they have been studied much more there. In particular, substantially increased risks of diverse population health problems in extremely poor neighborhoods have been well described in America, but not in Canada. One exemplary sentinel indicator of population health—breast cancer care—has been consistently observed to be of much lower quality in low-income neighborhoods and communities in the United States and of relatively higher quality in similar Canadian places. Though probably similar on many risks and vulnerabilities, low-income Canadian women with breast cancer, indeed all low-income Canadians are relatively less deprived than their American counterparts on at least one potentially critical characteristic. Their access to medically necessary health care is guaranteed. Such is clearly not the case for Americans. This between-country health insurance difference, therefore, is at the heart of this study's theoretical context. The health insurance theory predicts that breast cancer care will be much more equitable in Canada and that Canadian patients who reside in extremely poor neighborhoods will receive much higher quality health care than do their counterparts in America.

William Julius Wilson's (1987) germinal work in the high poverty neighborhoods of 1960s Chicago began the description and analysis of so-called underclass neighborhoods where 30% or more of the households had annual incomes below the US Census Bureau's poverty criterion. Modestly advancing the predictive validity of such high poverty areas while greatly extending this field's external validity, Paul Jargowsky (1997); Jargowsky and Mary Jo Bane (1991) studied census tract-based areas of extreme impoverishment where 40% or more of the households were poor in 239 US metropolitan areas during the generational time frame of the 1970s through the 1990s. Together they described high to extreme poverty areas as places of prevalent demographic vulnerability, where all of the following people tended to be more concentrated: racial/ethnic minority group members, young adults without a high school diploma, single mothers, the unemployed and those who had withdrawn from the labor market altogether, and welfare recipients. Perhaps not surprisingly, analysts have since observed consistent and generally strong associations between extreme impoverishment and diverse indicators of familial, social and personal illness in America: child neglect and abuse, teen pregnancy, violent crime, low birth weight, obesity, hypertension, diabetes, heart disease, cancer, AIDS, depression and suicide (Drake and Pandey, 1996; Geronimus et al., 2006; Harding, 2003; Krieger et al., 2003; Krivo and Peterson, 1996; Pearl et al., 2001; Rehkopf and Buka, 2006; Robbins and Webb, 2004; Zierler et al., 2000). Similar poverty–illness associations have been observed in Canada, though Canadian analysts have tended to use less extreme poverty criteria (e.g., 20% or more poor) or to study the linear health affects of relatively low-income areas that are characterized by their median incomes (Dupere et al., 2009; Gorey et al., 1998; Hou and Chen, 2003; Lemstra et al., 2006; Mustard et al., 1999).

1.1. North American health care policy laboratory

Sharing a 5000km border and having many social, cultural, lifestyle and physical environmental similarities, it seems that the myriad risks associated with extremely poor neighborhoods probably operate similarly to cause diverse diseases in the United States and Canada. The factors that are ultimately causally related to disease occurrences, however, are

not necessarily the same as the factors that are related to their effective care and outcomes. For instance, though common coronary heart disease and cancer morbidities are well known to be strongly associated with poverty in both the US and Canada, their mortalities and survival rates remain strongly associated with poverty in the US, while such associations seem null to nil in Canada (Gorey et al., 1998; Pilote et al., 2007). This pattern may be most parsimoniously explained by between-country health insurance differences. Their social-cultural-lifestyle-environmental similarities notwithstanding, all Canadians, be they extremely poor or affluent, employed, unemployed or having withdrawn from the labor market are distinctly advantaged as compared with their American counterparts. They universally enjoy access to a single payer system of health care. Low-income Americans are essentially much more prevalently exposed to various under- or uninsured statuses that greatly increase their risk of experiencing substandard health care or no health care at all (DeNavas-Walt et al., 2006; Gorey, 1999).

1.1.1. Breast cancer care in Canada and the US—Breast cancer care is one sentinel indicator of a health care system's performance. The most common type of cancer among North American women, its prognosis is typically excellent with early diagnosis and timely access to the best available treatments (Canadian Cancer Society, 2006; Ries et al., 2008). Moreover, for a number of reasons breast cancer seems particularly instructive for Canada-US cancer care comparisons. First, though the US and Canada, respectively, rank number one and two at the top of the world's breast cancer survival distribution, the overall difference between them is miniscule (RR=1.02; Coleman et al., 2008). Second, Canada-US comparative studies of breast cancer survival that accounted for socioeconomic factors consistently observed income by country interactions (Gorey, in press; Gorey et al., 1997, 2000a, 2000b, 2003b, 2009c; Zhang-Salomons et al., 2006). Moderate to strong inverse income-survival associations were consistently observed among US cohorts, but not among Canadian cohorts. Within-country social forces then seemed to operate so that low-income Canadian women experienced moderate to large survival advantages compared with their counterparts in the US, but between-country differences among, respective, middle- and high-income groups were consistently null. All of these studies used census tract-based US poverty measures and analogous low-income measures in Canada, but these did not measure the construct of extreme poverty areas as defined by either Wilson (1987) or Jargowsky and Bane (1991). Most of the low-income area comparisons, for example, were of lowest income third to fifths that typically only approached prevalence estimates of 20% poor. Third and finally, breast cancer diagnosis and treatments (screening, stage at diagnosis, waits for care, access to surgery, chemotherapy and radiation therapy) seem very sensitive to poverty in the US and have demonstrated similar poverty by country interactions in Canada-US comparisons that have been observed for breast cancer survival (Gold et al., 2008; Gorey et al., in press, 2009d; Polednak, 2002, 2004; Schootman et al., 2009). But again, their lowest income areas typically only ranged from 10% to 20% poor.

1.1.2. Hypotheses—We are unaware of any previous study that compared cancer care in high poverty urban areas of the United States and Canada. Focusing on breast cancer, this one will do so. Placing a greater emphasis on the “haves and have nots” than previous of this field's studies have, its findings could perhaps be of incrementally greater practical-policy

significance. Consistent with health insurance theoretical explanations we hypothesized the following. Within-country comparisons: (1) extremely poor urban neighborhoods will be significantly disadvantaged as compared with extremely affluent urban neighborhoods on breast cancer stage at diagnosis, waits for surgical and adjuvant treatments, receipt of surgical and adjuvant treatments and survival in the US, but not in Canada. Between-country comparisons: (2) women with breast cancer in extremely poor urban Canadian neighborhoods will be significantly advantaged on all of the cancer care and outcome measures as compared with their American counterparts. A hypothetical addendum predicts such Canadian advantages among the extremely poor to be qualitatively larger than those previously observed among the poor. (3) In contrast, extremely affluent Canadian and American urban neighborhoods are not expected to differ significantly on any measure of cancer care or outcome.

2. Methods

2.1. Samples

This historical cohort study is one of a series of analyses of cancer care in diverse urban and rural places in Ontario and California. For the present urban analysis, the Ontario Cancer Registry (OCR) and the California Cancer Registry (CCR), respectively, provided 624 and 660 primary, invasive, adult (25 or older) female breast cancer cases diagnosed between January 1, 1998 and December 31, 2000 in comparable urban areas. The OCR and CCR comprehensively surveil the most populace Canadian province and state in America with demonstrated validity. They have both been estimated to ascertain nearly all breast cancer cases (greater than 98%) with nearly perfect rates of microscopic confirmation and nearly nil rates of autopsy or death certificate only identification (Hall et al., 2006; North American Association of Central Cancer Registries, 2009; Walter et al., 1994; Zippin et al., 1995). The CCR incorporated additional hospital and physician follow back procedures to more completely capture breast cancer stage and treatment data than is typical of most other US cancer registries (Wright, 1996). The OCR abstracted the same stage and treatment variables from health records as it did not routinely collect them. Agreements were extremely high across study variables among three chart abstractors who were trained by an experienced cancer registrar. An inter-rater reliability assessment of 150 randomly sampled health records found that kappa (κ) coefficients ranged from 0.88 to 0.96 across the additional abstracted study variables.

Provincial and state samples (660 each), stratified by place, were randomly selected from megalopolises with more than 5 million residents (greater metropolitan Toronto [GMT] and the San Francisco bay area [SFBA]) and small cities with populations between 300,000 and 400,000 (Windsor-Essex county and Modesto-Stanislaus county) (Statistics Canada, 2002; US Bureau of the Census, 2002). Only 36 of the Ontario patient health records were unavailable for retrospective review. In a statistical sense this represented a significant between-country difference as all of the California records were available; $\chi^2(1, N=1320)=37.01, p<.05$. However, such losses to retrospective chart abstraction and database enhancement did not differ significantly on key study independent, dependent or co-variables that were routinely collected by the OCR (age, place, neighborhood income and 5-

year survival). Therefore, it seems very unlikely that this meager sample loss confounds any of this study's hypothesized income–breast cancer care relationships.

2.1.1. Extremely poor and extremely affluent neighborhoods—Statistics Canada and the US Bureau of the Census use conceptually similar indices of economic deprivation—respectively, “low-income” and “poverty” thresholds. Both are based on annual household income from all sources adjusted for household size. The Canadian low-income cutoff is a more liberal criterion though, approximately equal to 140% of the US poverty threshold (Osberg, 2000). Though not a poverty measure, per se, the Canadian low-income measure seems a close conceptual match for “near poverty” status that is sometimes used in US contexts (Gorey and Vena, 1995). Our previous analytic experience also suggested that though these two measures were not compositionally identical, they would allow for the valid contextual definition of relatively low- to high-income neighborhoods in both countries. In constructing extremely poor and extremely affluent urban neighborhoods, the following procedural goals were balanced. First, it was deemed most important that such neighborhood definitions be face validly aligned with this field's germinal measures. Such was balanced against a second important goal that planned comparisons have adequate statistical power to detect modest to large rate differences of 15% or more. Samples of 50 each, in extremely poor and extremely affluent, Canadian and American neighborhoods were thus required ($\alpha=0.05$ [2-tailed] and power $[1-\beta]=0.80$; Fleiss, 1981). Finally, because previous research observed similar income–breast cancer care gradients in large and small urban areas, they were represented equally (Gorey et al., 2009c, 2009d, in press).

Breast cancer cases in Ontario and California were first, respectively, joined to the 2001 Canadian and 2000 US censuses based on each patient's residential census tract (CT) at the time of diagnosis (Statistics Canada, 2002; US Bureau of the Census, 2002). Then to maximize the predictive validity of extremely poor to affluent groups, low- to high-income deciles were defined by their prevalence of, respective, low-income or poor households (Krieger et al., 2002). Finally, to maximize between-country construct validity as well as to satisfy this study's power demands, the 50 most extremely low or high CT residences on median annual household income within the lowest and highest deciles ultimately defined each country's extremely poor and extremely affluent neighborhoods. For comparison, less extremely poor and affluent areas were defined similarly, but they were based on low- to high-income quintiles. These merely poor and affluent areas each included 100 breast cancer patients. Poverty/low-income and median-income distributions of this study's CT or neighborhood-based, aggregated extremely poor to extremely affluent areas are displayed in Table 1. This study's procedures seem to have selected extremely poor urban neighborhoods in California that most closely converge with Wilson's (1987) high poverty neighborhood definition (median=29% poor [65% of them were between 30% and 39% poor]). Moreover, household incomes typically differed by less than \$3000 in such, respective, extremely poor urban neighborhoods of America and Canada. Perhaps not surprisingly, very extreme affluence was a bit more prevalent in the California sample, but on both sides of the border, this study's definition seemed to converge quite closely with contemporary definitions of extreme affluence (e.g., less than 5% poor or typical incomes of \$75,000 to \$100,000 or more; Barrett et al., 2008; Brookfield et al., 2009; Henry et al., 2009; Lee and Marlay, 2007).

2.1.2. Breast cancer care—Key study variables that had been routinely coded by the CCR were retrospectively abstracted in the same manner from hospital and physician office-based patient health records for the OCR sample: summary stage (localized, regional or metastasized), receipt of initial cancer directed surgery, type of surgery (breast conserving lumpectomy or mastectomy), receipt of adjuvant radiation therapy, chemotherapy or hormonal therapy, wait-times from diagnosis to surgery and radiation therapy (California Cancer Registry, 2003; Young et al., 2001). All of these cancer care variables had less than 5% missing data. Because such missing statuses were not significantly associated with either poverty-affluence or 5-year survival, they probably were not potent confounds. Chemotherapy and hormonal therapy initiation dates were prevalently missing from both the Ontario and California samples (12–20%), so their wait-times were not validly calculable. Long wait criteria of 2 months for surgery and 3 months post-lumpectomy for radiation therapy were used. Previous research suggested that such waits may be associated with disease recurrences, metastases and shorter survival (Chen et al., 2008b; Hershman et al., 2006; Lund et al., 2008). All breast cancer cases were followed until January 1, 2006 allowing, minimally, for the analysis of 5-year survival.

2.2. Analyses

All of the rates were directly age-adjusted, using this study's combined California–Ontario population of cases as the standard. So all of the rates (e.g., of lumpectomy or radiation therapy) within any table can be directly compared. Within- and between-country comparisons used rate ratios (RR) with 95% confidence intervals (CI) that were based on the χ^2 -test (Mantel and Haenszel, 1959; Miettinen, 1976). Ninety percent CIs were reported for findings that approached statistical significance at $p < .10$. Both the Ontario and California wait distributions were distinctly skewed. That is, many more patients experienced relatively short waits than long ones. Therefore, median wait-times in days were compared within- and between-country with the non-parametric Mann–Whitney *U*-test (Hollander and Wolfe, 1999). Maximum likelihood logistic regression models were used to estimate the associations of breast cancer care (stage, waits, receipt of surgery and adjuvant treatments) with 5-year all-cause survival (Hosmer and Lemeshow, 2000). Odds ratios (OR) and CIs were estimated from regression statistics.

3. Results

3.1. Extreme poverty and breast cancer care and survival

All except one of the breast cancer care and outcome indices were significantly associated with extreme impoverishment among the urban California sample (Table 2). Extremely poor urban Californian women with breast cancer were much less likely than their extremely affluent counterparts to be diagnosed with localized disease (RR=0.71), to receive breast conserving surgery (RR=0.74), to receive adjuvant radiation (RR=0.46) or hormonal therapy (RR=0.60) or to survive for 5 years after their diagnosis (RR=0.82). No such associations were observed among the urban Ontario sample. Consequently, extremely poor patients in Ontario were extremely advantaged as compared with similarly poor American patients: receipt of lumpectomy (RR=1.51), radiation therapy (RR=2.21) or hormonal therapy (RR=1.77) and survival (RR=1.21). A similar pattern was observed for within-country

treatment waits (Table 3). Relatively long initial surgical waits (RR=12.08) and long post-surgical waits for radiation therapy (RR=2.09) were very strongly associated with extreme poverty in California, but not in Ontario. The between-country pattern was notably different. For such waits it seems that affluent to extremely affluent Americans were advantaged, while among the poor to extremely poor, Canadians may have been.

3.2. Breast cancer care mediation by poverty-affluence

Age-, but not income-adjusted regression models observed that disease stage, long waits for care (radiation therapy in California and surgery in Ontario) and receipt of radiation therapy similarly predicted 5-year survival in the Canadian and American samples (Table 4, left column). After adjustment for income extremes, localized disease and radiation therapy waits were no longer predictive, and the radiation therapy receipt–survival association diminished significantly in California. Whereas, the income-adjusted model in Ontario was essentially identical to the unadjusted model (Table 4, right column). It seems that income extremes significantly mediate breast cancer care–survival relationships in the US, but not in Canada.

4. Discussion

This study, the first we are aware of that compared sentinel health processes and outcomes—breast cancer care and survival—in extremely poor urban neighborhoods in Canada and the United States, found near unequivocal support for its hypotheses. Extreme impoverishment was strongly associated with nonlocalized disease at diagnosis, long waits for initial surgery and adjuvant RT, non-receipt of breast conserving surgery, RT and hormonal therapy, and shorter survival in California. All such associations were null in Ontario. Consequently, extremely poor Ontario women were largely advantaged on most breast cancer care indices and 5-year survival over their counterparts in California. Moreover, such Canadian advantages were larger among extremely poor women than they were among less extremely poor women. The qualitative hypothesis that such Canadian advantages among the extremely poor would be larger than those previously observed among the poor was also supported. For example, among otherwise similar, but less impoverished samples of women with node negative breast cancer in Ontario and California, Canadian women were significantly advantaged on their receipt of post-surgical adjuvant RT (RRs of 1.39 and 1.23 with and without chemotherapy; Gorey et al., 2009d), but not nearly as advantaged as this study's sample of extremely poor Canadian women were (RR=2.21). Relatedly, a synthesis of 10 previous Canada–US comparisons of 5-year breast cancer survival observed a smaller aggregate Canadian advantage in near poor to poor areas (RR=1.14; Gorey, in press) than this one did in extremely poor areas (RR=1.21). These findings, along with the regression-based suggestion that breast cancer care–survival relationships are largely mediated by income extremes in the United States, but not in Canada seem to most parsimoniously indict inadequate health insurance coverage among America's extremely poor. In fact, this study's findings seem extraordinarily convergent with US studies that have consistently observed strong relationships between low-income, various under- and uninsured statuses, and relatively later stage at breast cancer diagnosis, lack of treatment access and early death (Anderson and Eamon, 2005; Chen et al., 2008a; Coburn et al., 2008; Griggs et al., 2007;

Hahn et al., 2007; Purc-Stephenson and Gorey, 2008; Studts et al., 2006; Ward et al., 2008). It is likely that the differential outcomes this study observed will persist until medical coverage and the best available care is made available to all Americans with cancer (Freedman, 2004).

It was also hypothesized that extremely affluent Canadian and American urban neighborhoods were not expected to differ significantly on any measure of cancer care or outcome. Indeed, such was the case for breast cancer stage at diagnosis, receipt of lumpectomy and all adjuvant treatments as well as 5-year survival. However, non-significant trends to significant differences indicative of American advantages in affluent to extremely affluent neighborhoods were observed on waits for initial surgery and on waits for post-lumpectomy RT. Affluent to extremely affluent Americans enjoyed shorter waits, but typically their waits were only one (surgery) to four (RT) weeks shorter than those of their Canadian counterparts. This difference is probably clinically insignificant for most patients as waits of that magnitude were not significantly related to survival in this study nor in previous ones (Chen et al., 2008a, 2008b; Hershman et al., 2006). Moreover, the implementation of federal and provincial government wait time guarantees for many services including cancer treatments suggests that cancer survival in Canada is likely to become even more equitable in the future (Cancer Care Ontario, 2009).

4.1. Potential study limitations

A number of potential alternative explanations related to this study's use of ecological measures, particularly of extremely impoverished neighborhoods, could be advanced by alternative theorists. One might legitimately wonder, for instance, if the racial/ethnic composition of such neighborhoods, rather than their prevalent representation of extremely low incomes, could account for this study's observed United States–Canada breast cancer care and survival differences. First, it should be noted that this study's sample of 50 women with breast cancer who resided in extremely poor urban Californian neighborhoods were predominantly non-Hispanic white women ($n=42$). Three each were African and Asian American, and two were Hispanic. We were not able to adjust for this factor directly as the OCR does not code race/ethnicity. We were able, however, to replicate key findings with the following conservative comparison: non-Hispanic white women in California versus the entire racial/ethnically diverse sample of women in Ontario. Among them, extremely poor Ontario women remained advantaged on early stage at diagnosis (RR=2.09, 95% CI 1.22, 3.58), shorter waits for surgery (RR=0.31, 90% CI 0.11, 0.84), receipt of lumpectomy (RR=1.99, 95% CI 1.25, 3.16), RT (RR=1.52, 95% CI 1.00, 2.30) and hormonal therapy (RR=1.96, 95% CI 1.06, 3.62), and on 5-year survival (RR=1.30, 90% CI 1.01, 1.67). One might also wonder if the, respective, neighborhood-level measures of extreme impoverishment were actually measuring the same contextual construct in California and Ontario. They were, after all, not compositionally identical; one being based in the US Census Bureau's "poverty" threshold, the other on Statistics Canada's "low-income" criterion. We cannot know for sure because no previous study has directly compared the construct or predictive validities of such ecological measures in Canada and the US. Some analytic comfort was provided though by the fact that both national censuses provide estimates of median CT or neighborhood-level income in urban areas. Using this fact, we

observed that household incomes typically differed by less than \$3000 in the, respective, extremely poor urban neighborhoods of California (\$27,400) and Ontario (\$30,275) that we studied. This suggests their similar aggregate lack of purchasing power, which is probably also the best contextual definition of this study's central ecological measure. Though probably similarly challenged to purchase life's necessities, residents of such neighborhoods clearly differ contextually in one important way. Canadians, even in such extremely poor neighborhoods, seem to be able to "purchase" much higher quality health care than can similarly poor Americans.

One might also wonder if other systemic health care factors, beyond payer and health insurance differences, potentially confound this study's key between-country findings. Though this study seemed to most parsimoniously indict inadequate health insurance coverage as well as its corollaries of inaccessible primary care and cancer care among America's poor (Gorey, in press; Starfield, in press), one might plausibly argue that between-country primary care differences themselves could independently explain our findings. We think probably not for the following reasons. Analyses of the same Ontario-California database observed significant primary care physician-cancer care effects that were not moderated by income in Ontario, but no such independent effects of primary care in California (Gorey et al., 2009a, 2009b). Moreover, consistent with other analysts (McGrail et al., 2009), we found that after income or insurance status are accounted for, primary care itself seems to explain very little of the health inequalities in Canada or the United States. Between-country screening differences could be advanced as another plausible alternative explanation. Again, we think it improbable because overall screening mammography rates as well as income-mammography associations seemed roughly similar in Ontario and California between 1995 and 2005, and apparently effective publicly funded screening programs had been instituted in both Ontario and California during the 1990s (Meersman et al., in press; Minore et al., 2001; Purc-Stephenson and Gorey, 2008; Schueler et al., 2008; Shields and Wilkins, 2009). But even if screening differences did influence this study's findings, they would probably have only affected its findings related to breast cancer stage at the time of diagnosis. This and other studies' between-country comparisons of treatments and survival are unlikely to have been influenced at all by any screening differences.

This study's samples of women with breast cancer may not be generalizable to all such women in Ontario and California or to other provinces and states. Samples were drawn from purposively diverse and potentially policy-important places in Ontario and California: very large and small cities. Admittedly, our findings are most generalizable to such places. It should be noted, however, that after accounting for demographic, socioeconomic and clinical factors, place, per se, did not seem to matter in any of this study's analytic models nor in those of other of this field's studies (e.g., Gorey, in press; Gorey et al., 2009c, 2009d). Furthermore, though not as well controlled, this study's general pattern of findings had previously been observed in Manitoba and in a number of other states: Michigan, Washington, Connecticut, Iowa and Hawaii (Gorey et al., 1997, 2000a, 2000b, 2003). Its inferences, therefore, can probably be confidently generalized to urban areas, small cities to megalopolises, across Canada and the United States.

5. Conclusions

Extremely poor Ontario women were advantaged on breast cancer care and survival over their counterparts in California. Breast cancer care–survival relationships seem to be largely mediated by income extremes in the United States, but not in Canada. More inclusive health care insurance coverage in Canada versus America, particularly among each country's extremely poor people, seems the most plausible explanation for such observed Canadian breast cancer care and survival advantages.

Acknowledgments

This work was supported by the Canadian Breast Cancer Research Alliance (Canadian Institutes of Health Research [CIHR] Grant no. 67161), the Canadian Cancer Society (National Cancer Institute of Canada Grant no. 016160), the Social Sciences and Humanities Research Council of Canada (Grant no. 410-2002-0173) as well as a CIHR investigator award and an Assumption University research chair. The authors gratefully acknowledge the administrative and logistical assistance of William E. Wright, chief, Cancer Surveillance Section of the California Department of Health Services at the time this study was initiated. We also gratefully acknowledge the research and technical assistance of Carole Herbert (Cancer Care Ontario), Madhan Balagurusamy, Leah Archambault, Natalie Herbert, Dylan Herbert, Nancy Richter (University of Windsor) and Mark Allen (California Cancer Registry). This study was reviewed and cleared by the University of Windsor's Research Ethics Committee as well as the Ontario Cancer Research Network's Research Ethics Board.

References

- Anderson SG, Eamon MK. Stability of health care coverage among low-income working women. *Health & Social Work*. 2005; 30:7–17. [PubMed: 15847233]
- Barrett RE, Cho YI, Weaver KE, Ryu K, Campbell RT, Dolecek TA, Warnecke RB. Neighborhood change and distant metastasis at diagnosis of breast cancer. *Annals of Epidemiology*. 2008; 18:43–47. [PubMed: 17890103]
- Brookfield KF, Cheung MC, Lucci J, Fleming LE, Koniaris LG. Disparities in survival among women with invasive cervical cancer: a problem of access to care. *Cancer*. 2009; 115:166–178. [PubMed: 19097209]
- California Cancer Registry. Cancer reporting in California: abstracting and coding procedures for hospitals. California Cancer Reporting System standards. 7. Vol. I. Department of Health Services, Cancer Surveillance Section; Sacramento: 2003.
- Canadian Cancer Society. Canadian cancer statistics. National Cancer Institute of Canada; Toronto: 2006.
- Cancer Care Ontario. 2009 Wait times. Accessed on June 2, 2009 Available at: <<http://www.cancercare.on.ca/ocs/wait-times>>
- Chen AY, Halpern MI, Schrag NM, Stewart A, Leitch M, Ward E. Disparities and trends in sentinel lymph node biopsy among early-stage breast cancer patients (1998–2005). *Journal of the National Cancer Institute*. 2008a; 100:462–474. [PubMed: 18364506]
- Chen Z, King W, Pearcey R, Kerba M, Mackillop WJ. The relationship between waiting time for radiotherapy and clinical outcomes: a systematic review of the literature. *Radiotherapy and Oncology*. 2008b; 87:3–16. [PubMed: 18160158]
- Coburn N, Fulton J, Pearlman DN, Law C, DiPaolo B, Cady B. Treatment variation by insurance status for breast cancer patients. *The Breast Journal*. 2008; 14:128–134. [PubMed: 18315690]
- Coleman MP, Quaresma M, Berrino F, Lutz JM, Capocaccia R, Baili P, Rachet B, Gatta G, Hakulinen T, Micheli A, Sant M, Weir HK, Elwood JM, Tsukuma H, Koifman SE, Silva GA, Francisci S, Santaquilani M, Verdecchia A, Storm HH, Young JL. CONCORD Working Group. Cancer survival in five continents: a worldwide population-based study (CONCORD). *The Lancet Oncology*. 2008; 9:730–756. [PubMed: 18639491]

- DeNavas-Walt, C., Proctor, BD., Lee, CH. US Bureau of the Census, Current Population Reports (P60-231). Income, Poverty, and Health Insurance Coverage in the United States, 2005. US Government Printing Office; Washington, DC: 2006.
- Drake B, Pandey S. Understanding the relationship between neighborhood poverty and specific types of child maltreatment. *Child Abuse & Neglect*. 1996; 20:1003–1018. [PubMed: 8958452]
- Duncan GJ, Gustafsson B, Hauser R, Schmauss G, Messinger H, Muffels R, Nolan B, Ray JC. Poverty dynamics in eight countries. *Journal of Population Economics*. 1993; 6:215–234.
- Dupere V, Leventhal T, Lacourse E. Neighborhood poverty and suicidal thoughts and attempts in late adolescence. *Psychological Medicine*. 2009; 39:1295–1306. [PubMed: 18845013]
- Flleiss, JL. *Statistical Methods for Rates and Proportions*. 2. John Wiley & Sons; New York: 1981.
- Freedman HP. Poverty, culture and social injustice: determinants of cancer disparities. *CA Cancer Journal for Clinicians*. 2004; 54:72–77.
- Geronimus AT, Colen CG, Shochet T, Ingler LB, James SA. Urban–rural differences in excess mortality among high-poverty populations: evidence from the Harlem Household Survey and the Pitt County, North Carolina Study of African American Health. *Journal of Health Care for the Poor and Underserved*. 2006; 17:532–558. [PubMed: 16960321]
- Gold HT, Do HT, Dick AW. Correlates and effect of suboptimal radiotherapy in women with ductal carcinoma in situ or early invasive breast cancer. *Cancer*. 2008; 113:3108–3115. [PubMed: 18932243]
- Gorey KM. Prevalent low income status in Canadian and United States metropolitan areas, 1980 and 1990. *International Journal of Comparative Sociology*. 1998; 39:378–383.
- Gorey KM. What is wrong with the US health care system? It does not effectively exist for one of every eight Americans. *The Milbank Quarterly*. 1999; 77:401–407. [PubMed: 10526551]
- Gorey KM. Breast cancer survival in Canada and the United States: Meta-analytic evidence of a Canadian advantage in low-income areas. *International Journal of Epidemiology*. in press.
- Gorey KM, Fung KY, Luginaah IN, Holowaty EJ, Hamm C. Income and long-term breast cancer survival: comparisons of vulnerable urban places in Ontario and California *The Breast Journal*. in press.
- Gorey KM, Holowaty EJ, Fehringer G, Laukkanen E, Moskowitz A, Webster DJ, Richter NL. An international comparison of cancer survival: Toronto, Ontario, and Detroit, Michigan, metropolitan areas. *American Journal of Public Health*. 1997; 87:1156–1163. [PubMed: 9240106]
- Gorey KM, Holowaty EJ, Laukkanen E, Fehringer G, Richter NL. Association between socioeconomic status and cancer incidence in Toronto, Ontario: possible confounding of cancer mortality by incidence and survival. *Cancer Prevention & Control*. 1998; 2:236–241. [PubMed: 10093638]
- Gorey KM, Holowaty EJ, Fehringer G, Laukkanen E, Richter NL, Meyer CM. An international comparison of cancer survival: metropolitan Toronto, Ontario and Honolulu, Hawaii. *American Journal of Public Health*. 2000a; 90:1866–1872. [PubMed: 11111258]
- Gorey KM, Holowaty EJ, Fehringer G, Laukkanen E, Richter NL, Meyer CM. An international comparison of cancer survival: Toronto, Ontario and three resourceful US metropolitan areas. *Journal of Public Health Medicine*. 2000b; 22:343–348. [PubMed: 11077908]
- Gorey KM, Kliever E, Holowaty EJ, Laukkanen E, Ng EY. An international comparison of breast cancer survival: Winnipeg, Manitoba and Des Moines, Iowa, metropolitan areas. *Annals of Epidemiology*. 2003; 13:32–41. [PubMed: 12547483]
- Gorey KM, Luginaah IN, Bartfay E, Fung KY, Holowaty EJ, Wright FC, Hamm C, Kanjeekal SM, Balagurusamy M. Associations of physician supplies with colon cancer care in Ontario and California, 1996 to 2006. Unpublished results. 2009a
- Gorey KM, Luginaah IN, Hamm C, Zou GY, Balagurusamy M, Holowaty EJ. Associations of primary care and specialist physician supplies with breast cancer care in Ontario and California, 1996 to 2006. Unpublished results. 2009b
- Gorey KM, Luginaah IN, Holowaty EJ, Fung KY, Hamm C. Breast cancer survival in Ontario and California, 1998–2006: socioeconomic inequity remains much greater in the United States. *Annals of Epidemiology*. 2009c; 19:121–124. [PubMed: 19185806]
- Gorey KM, Luginaah IN, Holowaty EJ, Fung KY, Hamm C. Wait times for initial surgical and adjuvant treatment of breast cancer in Canada and the United States: evidence of greater

- socioeconomic inequity in America. *Clinical and Investigative Medicine*. 2009d; 32:E239–E249. [PubMed: 19480739]
- Gorey KM, Vena JE. The association of near poverty status with cancer incidence among black and white adults. *Journal of Community Health*. 1995; 20:359–366. [PubMed: 7593741]
- Griggs JJ, Culakova E, Sorbero MES, Poniewierski MS, Wolff DA, Crawford J, Dale DC, Lyman GH. Social and racial differences in selection of breast cancer adjuvant chemotherapy regimens. *Journal of Clinical Oncology*. 2007; 25:2522–2527. [PubMed: 17577029]
- Hahn KM, Bondy ML, Selvan M, Lund MJ, Liff JM, Flagg EW, Brinton LA, Porter P, Eley JW, Coates RJ. Factors associated with advanced disease stage at diagnosis in a population-based study of patients with newly diagnosed breast cancer. *American Journal of Epidemiology*. 2007; 166:1035–1044. [PubMed: 17690220]
- Hall S, Schulze K, Groome P, Mackillop W, Holowaty E. Using cancer registry data for survival studies: the example of the Ontario Cancer Registry. *Journal of Clinical Epidemiology*. 2006; 59:67–76. [PubMed: 16360563]
- Harding DJ. Counterfactual models of neighborhood effects: the effect of neighborhood poverty on dropping out and teenage pregnancy. *American Journal of Sociology*. 2003; 109:676–719.
- Henry KA, Sherman R, Roche LM. Colorectal cancer stage at diagnosis and area socioeconomic characteristics in New Jersey. *Health & Place*. 2009; 15:505–513. [PubMed: 19028134]
- Hershman DL, Wang X, McBride R, Jacobson JS, Grann VR, Neugut AI. Delay in initiating adjuvant radiotherapy following breast conservation surgery and its impact on survival. *International Journal of Radiation Oncology Biology Physics*. 2006; 65:1353–1360.
- Hollander, M., Wolfe, DA. *Nonparametric Statistical Methods*. 2. John Wiley & Sons; New York: 1999.
- Hosmer, DW., Lemeshow, S. *Applied Logistic Regression*. 2. John Wiley & Sons; New York: 2000.
- Hou F, Chen J. Neighbourhood low income, income inequality and health in Toronto. *Health Reports*. 2003; 14:21–34.
- Jargowsky, PA. *Poverty and Place: Ghettos, Barrios, and the American City*. Russell Sage Foundation; New York: 1997.
- Jargowsky, PA., Bane, MJ. Ghetto poverty in the United States, 1970–1980. In: Jencks, C., Peterson, PE., editors. *The Urban Underclass*. The Brookings Institution; Washington, DC: 1991. p. 235–273.
- Krieger N, Chen JT, Waterman PD, Rehkopf DH, Subramanian SV. Race/ethnicity, gender, and monitoring socioeconomic gradients in health: a comparison of area-based socioeconomic measures—The Public Health Disparities Geocoding Project. *American Journal of Public Health*. 2003; 93:1655–1671. [PubMed: 14534218]
- Krieger N, Chen JT, Waterman PD, Soobader M, Subramanian SV, Carson R. Geocoding and monitoring of US socioeconomic inequalities in mortality and cancer incidence: does the choice of area-based measure and geographic level matter? The Public Health Disparities Geocoding Project. *American Journal of Epidemiology*. 2002; 156:471–482. [PubMed: 12196317]
- Krivo LJ, Peterson RD. Extremely disadvantaged neighborhoods and urban crime. *Social Forces*. 1996; 75:619–650.
- Lee BA, Marlay M. The right side of the tracks: affluent neighborhoods in the metropolitan United States. *Social Science Quarterly*. 2007; 88:766–789.
- Lemstra M, Neudorf C, Opondo J. Health disparity by neighbourhood income. *Canadian Journal of Public Health*. 2006; 97:435–439. [PubMed: 17203720]
- Lund MJ, Brawley OP, Ward KC, Young JL, Gabram SSG, Eley JW. Parity and disparity in first course treatment of invasive breast cancer. *Breast Cancer Research and Treatment*. 2008; 109:545–557. [PubMed: 17659438]
- Mantel N, Haenszel W. Statistical aspects of the analysis of data from retrospective studies of disease. *Journal of the National Cancer Institute*. 1959; 22:719–748. [PubMed: 13655060]
- McGrail KM, van Doorslaer E, Ross NA, Sanmartin C. Income-related health inequalities in Canada and the United States: a decomposition analysis. *American Journal of Public Health*. 2009; 99:1856–1863. [PubMed: 19150915]

- Meersman SC, Breen N, Pickle LW, Meissner HI, Simon P. Access to mammography screening in a large urban population: a multi-level analysis. *Cancer Causes and Control*. in press.
- Miettinen OS. Estimability and estimation in case-referent studies. *American Journal of Epidemiology*. 1976; 103:226–235. [PubMed: 1251836]
- Minore B, Hill ME, Kurm MJ, Vergidis D. Knowledgeable, consistent, competent care: meeting the challenges of delivering quality care in remote northern communities. *International Journal of Circumpolar Health*. 2001; 60:196–204. [PubMed: 11507969]
- Mustard CA, Derksen S, Berthelot JM, Wolfson M. Assessing ecologic proxies for household income: a comparison of household and neighbourhood level income measures in the study of population health status. *Health & Place*. 1999; 5:157–171. [PubMed: 10670997]
- North American Association of Central Cancer Registries. 2009Data quality assessments. Accessed on March 17, 2009Available at: <<http://www.naacr.org>>
- Osberg L. Poverty in Canada and the United States: measurement, trends, and implications. *Canadian Journal of Economics*. 2000; 33:847–877.
- Pearl M, Braveman P, Abrams B. The relationship of neighborhood socioeconomic characteristics to birthweight among 5 ethnic groups in California. *American Journal of Public Health*. 2001; 91:1808–1814. [PubMed: 11684609]
- Pilote L, Tu JV, Humphries K, Behouli H, Belisle P, Austin PC, Joseph L. Socioeconomic status, access to health care, and outcomes after acute myocardial infarction in Canada's universal health care system. *Medical Care*. 2007; 45:638–646. [PubMed: 17571012]
- Polednak AP. Survival of breast cancer patients in Connecticut in relation to socioeconomic and health care access indicators. *Journal of Urban Health*. 2002; 79:211–218. [PubMed: 12023496]
- Polednak AP. Chemotherapy of nonelderly breast cancer patients by poverty-rate of area of residence in Connecticut. *Breast Cancer Research and Treatment*. 2004; 83:245–248. [PubMed: 14758094]
- Purc-Stephenson RJ, Gorey KM. Lower adherence to screening mammography guidelines among ethnic minority women in America: a meta-analytic review. *Preventive Medicine*. 2008; 46:479–488. [PubMed: 18295872]
- Rehkopf DH, Buka SL. The association between suicide and the socioeconomic characteristics of geographical areas: a systematic review. *Psychological Medicine*. 2006; 36:145–157. [PubMed: 16420711]
- Ries, LAG., Eisner, MP., Kosary, CL., Hankey, BF., Miller, BA., Clegg, L., Edwards, BK. SEER Cancer Statistics Review, 1975–2005. National Cancer Institute; Bethesda, MD: 2008.
- Robbins JM, Webb DA. Neighborhood poverty, mortality rates, and excess deaths among African Americans: Philadelphia, 1999–2001. *Journal of Health Care for the Poor and Underserved*. 2004; 15:530–537. [PubMed: 15531812]
- Schootman M, Jeffe DB, Lian M, Gillanders WE, Aft R. The role of poverty rate and racial distribution in the geographic clustering of breast cancer survival among older women: a geographic and multilevel analysis. *American Journal of Epidemiology*. 2009; 169:554–561. [PubMed: 19103608]
- Schueler KM, Chu PW, Smith-Bindman R. Factors associated with mammography utilization: a systematic quantitative review of the literature. *Journal of Women's Health*. 2008; 17:1477–1498.
- Shields M, Wilkins KM. An update on mammography use in Canada. *Health Reports*. 2009; 20:1–13.
- Starfield B. Commentary: how does 'insurance' improve equity in health? *International Journal of Epidemiology*. in press.
- Statistics Canada. Profiles of Census Tracts, 2001 (Ontario). Author; Ottawa, ON: 2002.
- Studs CR, Stone RF, Barber GM. Predictors of access to health care services among groups of TANF recipients in Kentucky. *Social Service Review*. 2006; 80:527–548.
- US Bureau of the Census. 2000 Census of Population and Housing in California: Summary tape file 3 on CD-ROM. US Department of Commerce; Washington, DC: 2002.
- Walter SD, Birnie SE, Marrett LD, Taylor SM, Reynolds D, Davies J, Drake JJ, Hayes M. The geographic variation of cancer incidence in Ontario. *American Journal of Public Health*. 1994; 84:367–376. [PubMed: 8129051]

- Ward E, Halpern M, Schrag N, Cokkinides V, DeSantis C, Bandi P, Siegel R, Stewart A, Jemal A. Association of insurance with cancer care utilization and outcomes. *CA: A Cancer Journal for Clinicians*. 2008; 58:9–31. [PubMed: 18096863]
- Wilson, WJ. *The Truly Disadvantaged: the Inner City, the Underclass, and Public Policy*. The University of Chicago Press; 1987.
- Wright, WE. *California Cancer Registry Enhancement for Breast Cancer Research*. California Public Health Foundation; Berkeley, CA: 1996.
- Young, JL., Jr, Roffers, SD., Ries, LAG., Fritz, AG., Hurlbut, AA. *SEER Summary Staging Manual—2000: Codes and Coding Instructions* (NIH pub. no. 01-4969). National Cancer Institute; Bethesda, MD: 2001.
- Zhang-Salomons J, Qian H, Holowaty E, Mackillop WJ. Associations between socioeconomic status and cancer survival: choice of SES indicator may affect results. *Annals of Epidemiology*. 2006; 16:521–528. [PubMed: 16386924]
- Zierler S, Krieger N, Tang Y, Coady W, Siegfried E, DeMaria A, Auerbach J. Economic deprivation and AIDS incidence in Massachusetts. *American Journal of Public Health*. 2000; 90:1064–1073. [PubMed: 10897184]
- Zippin C, Lum D, Hankey BF. Completeness of hospital case reporting from the SEER program of the National Cancer Institute. *Cancer*. 1995; 76:2343–2350. [PubMed: 8635041]

Descriptive profiles of census-tract based income areas in urban California (San Francisco and Modesto, 2000) and Ontario (Toronto and Windsor, 2001).

Table 1

Income areas	California		Ontario	
	Range	Median	Range	Median
Extremely affluent	0.79–3.99	2.14	1.30–4.50	2.90
Affluent	0.79–4.99	2.62	1.30–5.99	3.65
Poor	11.00–59.32	21.95	24.50–52.80	30.45
Extremely poor	21.30–59.32	29.35	30.00–52.80	37.10
				Income ^b
				88,400
				82,900
				35,125
				30,275

^aCensus tract median annual household income in US dollars.

^bCensus tract median annual household income in Canadian dollars.

Associations of country and place with breast cancer care and survival: relatively poor compared with affluent neighborhoods, and extremely poor compared with extremely affluent neighborhoods.

Table 2

Cancer care indicator	California			Ontario			Ontario/California					
	Neighborhood	n	Rate ^a	RR ^b	95% CI ^c	n	Rate ^a	RR ^b	95% CI ^c	RR	95% CI ^c	
<i>Localized disease at diagnosis</i>												
Extremely affluent	50	0.720	1.00	50	0.680	1.00	...	0.94	0.71, 1.24	
Affluent	99	0.677	1.00	100	0.641	1.00	...	0.95	0.80, 1.13	
Poor	100	0.600	0.89	0.73, 1.09	...	100	0.648	1.01	0.89, 1.15	1.08	0.88, 1.33	
Extremely poor	50	0.512	0.71	0.52, 0.97	...	50	0.640	0.94	0.71, 1.25	1.25	0.90, 1.74	
<i>Received lumpectomy</i>												
Extremely affluent	50	0.644	1.00	50	0.646	1.00	...	1.00	0.94, 1.06	
Affluent	99	0.596	1.00	100	0.630	1.00	...	1.06	0.84, 1.34	
Poor	99	0.545	0.91	0.70, 1.18	...	99	0.708	1.12	0.92, 1.36	1.30	1.05, 1.62	
Extremely poor	50	0.477	0.74	0.55, 0.99^d	...	49	0.718	1.11	0.85, 1.45	1.51	1.09, 2.09	
<i>Received radiation therapy</i>												
Extremely affluent	50	0.620	1.00	50	0.564	1.00	...	0.91	0.67, 1.23	
Affluent	99	0.566	1.00	100	0.520	1.00	...	0.92	0.72, 1.18	
Poor	100	0.480	0.85	0.65, 1.10	...	100	0.572	1.10	0.85, 1.43	1.19	0.91, 1.56	
Extremely poor	50	0.286	0.46	0.29, 0.73	...	50	0.631	1.12	0.85, 1.47	2.21	1.42, 3.45	
<i>Received chemotherapy</i>												
Extremely affluent	49	0.426	1.00	50	0.406	1.00	...	0.95	0.67, 1.34	
Affluent	98	0.425	1.00	100	0.371	1.00	...	0.87	0.64, 1.19	
Poor	99	0.414	0.97	0.72, 1.30	...	100	0.386	1.04	0.80, 1.35	0.93	0.70, 1.24	
Extremely poor	50	0.411	0.96	0.60, 1.52	...	49	0.423	1.04	0.56, 1.95	1.03	0.74, 1.44	
<i>Received hormonal therapy</i>												
Extremely affluent	48	0.520	1.00	50	0.498	1.00	...	0.96	0.64, 1.43	
Affluent	98	0.455	1.00	100	0.541	1.00	...	1.19	0.91, 1.55	
Poor	98	0.287	0.63	0.43, 0.92	...	100	0.547	1.01	0.88, 1.16	1.91	1.36, 2.67	
Extremely poor	49	0.312	0.60	0.36, 0.99	...	49	0.551	1.11	0.75, 1.65	1.77	1.12, 2.79	
<i>Five-year survival</i>												

Cancer care indicator	California				Ontario				Ontario/California			
	<i>n</i>	Rate ^a	RR ^b	95% CI ^c	<i>n</i>	Rate ^a	RR ^b	95% CI ^c	RR	95% CI ^c	RR	95% CI ^c
Extremely affluent	50	0.812	1.00	...	50	0.883	1.00	...	1.09	0.89, 1.33	1.10	0.98, 1.23
Affluent	99	0.806	1.00	...	100	0.887	1.00	...	1.10	0.98, 1.23	1.20	1.03, 1.40
Poor	100	0.690	0.86	0.74, 1.00	100	0.828	0.94	0.85, 1.04	1.20	1.03, 1.40	1.21	1.01, 1.45^d
Extremely poor	50	0.663	0.82	0.68, 0.99^d	50	0.803	0.91	0.77, 1.08	1.21	1.01, 1.45^d	1.21	1.01, 1.45^d

Notes: Within-country comparisons were of the poor versus the affluent and of the extremely poor versus the extremely affluent. *n*=number of incident breast cancer cases, RR=standardized rate ratio, CI=confidence interval. Bolded RRs and CIs are statistically significant.

^aAll rates were directly age-adjusted using this study's combined Ontario-California population of cases as the standard (age strata: 25–44, 45–54, 55–64, 65–74 and 75 years or older), so all of the rates for each cancer care indicator are directly comparable.

^bA survival rate ratio of 1.00 is the within-country baseline.

^cConfidence intervals are based on the Mantel-Haenszel χ^2 -test.

^d90% confidence interval.

Table 3

Associations of country and place with breast cancer care delays: relatively poor compared with affluent neighborhoods, and extremely poor compared with extremely affluent neighborhoods.

Cancer care indicator	California			Ontario			Ontario/California		
	n	Rate ^a (Mdn)	RR ^b 95% CI ^c	n	Rate ^a (Mdn)	RR ^b 95% CI ^c	MDD	RR	95% CI ^c
<i>Waited 60 days or longer for initial surgery</i>									
Extremely affluent	47	0.013 (3)	1.00 ...	50	0.072 (8)	1.00 ...	5	5.53	1.00, 30.58^d
Affluent	90	0.045 (5)	1.00 ...	100	0.101 (13)	1.00 ...	8*	2.24	0.75, 6.71
Poor	94	0.100 (10)	2.22 0.77, 6.41	99	0.081 (13)	0.80 0.33, 1.95	3	0.81	0.40, 1.63
Extremely poor	47	0.157 (11)*	12.08 1.65, 88.66	49	0.065 (9)	0.90 0.53, 1.52	32	0.41	0.17, 0.99^d
<i>Received lumpectomy and waited 90 days or longer for radiation therapy</i>									
Extremely affluent	24	0.375 (68)	1.00 ...	20	0.700 (97)	1.00 ...	29	1.87	1.05, 3.33
Affluent	42	0.405 (89)	1.00 ...	41	0.699 (110)	1.00 ...	21*	1.73	1.15, 2.60
Poor	35	0.584 (105)	1.44 1.01, 2.05^d	56	0.573 (110)	0.82 0.62, 1.09	5	0.98	0.41, 2.34
Extremely poor	17	0.785 (111)*	2.09 1.17, 3.74	27	0.556 (95)	0.79 0.50, 1.24	316	0.71	0.51, 0.99^d

Notes: Within-country comparisons were of the poor versus the affluent and of the extremely poor versus the extremely affluent. *n*=number of incident breast cancer cases, Mdn=median days waited, RR=standardized rate ratio, CI=confidence interval, MDD=median days difference. Bolded RRs and CIs are statistically significant.

^a All rates were directly age-adjusted using this study's combined Ontario-California population of cases as the standard (age strata: 25–44, 45–54, 55–64, 65–74 and 75 years or older), so all of the rates for each cancer care indicator are directly comparable.

^b A survival rate ratio of 1.00 is the within-country baseline.

^c Confidence intervals are based on the Mantel-Haenszel χ^2 -test.

^d 90% confidence interval.

* Median wait differed significantly from baseline (within-country) or between-countries: Mann-Whitney *U*-test, *P*<0.05.

Table 4

Associations of stage of disease at diagnosis and treatments with 5-year breast cancer survival: entire urban sample analyses within countries.

Cancer care indicator	Logistic regression models			
	Age-adjusted ^a		Age & income-adjusted ^b	
	OR	95% CI	OR	95% CI
<i>California (N=656)</i>				
Localized disease at diagnosis	2.17	1.39, 3.39	1.53	0.54, 4.33
Waited 90 days or longer for RT	0.35	0.16, 0.74	0.83	0.50, 1.38
Receipt of radiation therapy (RT)	3.95	2.07, 7.53	2.33	1.55, 3.51
<i>Ontario (N=624)</i>				
Localized disease at diagnosis	2.19	1.38, 3.47	2.14	1.35, 3.42
Waited 60 days or longer for surgery	0.43	0.22, 0.85	0.42	0.21, 0.83
Receipt of radiation therapy (RT)	1.56	1.05, 2.31^c	1.61	1.01, 2.56

Notes: All breast cancer care indices were entered into regression models in temporal order.

OR=odds ratio, CI=confidence interval. Bolded ORs and CIs are statistically significant.

^a Adjusted across the following age strata: 25–44, 45–54, 55–64, 65–74 and 75 years or older.

^b Adjusted across the following categorical income areas: extremely poor, poor, middle-income areas, affluent and extremely affluent.

^c 90% confidence interval.