

# Bradykinin B<sub>2</sub> Receptor Interacts with Integrin $\alpha 5\beta 1$ to Transactivate Epidermal Growth Factor Receptor in Kidney Cells

Inga I. Kramarenko, Marlene A. Bunni, John R. Raymond, and Maria N. Garnovskaya

Medical and Research Services, Ralph H. Johnson Veterans Affairs Medical Center, and Nephrology Division, Department of Medicine, Medical University of South Carolina, Charleston, South Carolina

Received March 18, 2010; accepted April 12, 2010

## ABSTRACT

We have shown previously that the vasoactive peptide bradykinin (BK) stimulates proliferation of a cultured murine cell model of the inner medullary collecting duct (mIMCD-3 cells) via transactivation of epidermal growth factor receptor (EGFR) by a mechanism that involves matrix metalloproteinases (collagenase-2 and -3). Because collagenases lack an integral membrane domain, we hypothesized that receptors for extracellular matrix proteins, integrins, may play a role in BK-induced signaling by targeting collagenases to the membrane, thus forming a functional signaling complex. BK-induced phosphorylation of extracellular signal-regulated protein kinase (ERK) in mIMCD-3 cells was reduced by ~65% by synthetic peptides containing an Arg-Gly-Asp sequence, supporting roles for integrins in BK-induced signaling. Neutralizing antibody against  $\alpha 5\beta 1$  integrin partially (~60%) blocked BK-induced ERK acti-

vation but did not affect EGF-induced ERK activation. Silencing of  $\alpha 5$  and  $\beta 1$  expression by transfecting cells with small interfering RNAs (siRNA) significantly decreased BK-induced ERK activation (~80%) and EGFR phosphorylation (~50%). This effect was even more pronounced in cells that were cotransfected with siRNAs directed against both collagenases and  $\alpha 5\beta 1$  integrin. On the basis of our results, we suggested that integrin  $\alpha 5\beta 1$  is involved in BK-induced signaling in mIMCD-3 cells. Using immunoprecipitation/Western blotting, we demonstrated association of BK B<sub>2</sub> receptor with  $\alpha 5\beta 1$  integrin upon BK treatment. Furthermore, BK induced association of  $\alpha 5\beta 1$  integrin with EGFR. These data provide the first evidence that specific integrins are involved in BK B<sub>2</sub> receptor-induced signaling in kidney cells, and ultimately might lead to development of new strategies for treatment of renal tubulointerstitial fibrosis.

The vasoactive nonapeptide bradykinin (BK) plays important roles in the regulation of kidney functions, such as electrolyte and water excretion (Mukai et al., 1996). Specifically, a role of BK in the control of absorptive function in the kidney collecting ducts is well established (Tomita et al., 1985; Zeidel et al., 1990). BK also acts directly as a potent

cellular growth factor for multiple cell types, including kidney cells. We established previously that the BK B<sub>2</sub> receptor stimulates early mitogenic signals associated with activation of extracellular signal-regulated protein kinase (ERK) in a murine epithelial cells derived from the inner medullary collecting duct (mIMCD-3 cells), and demonstrated that BK-induced cell proliferation depends on transactivation of the epidermal growth factor receptor (EGFR) (Mukhin et al., 2003). Furthermore, we demonstrated that BK B<sub>2</sub> receptor-induced EGFR transactivation involves activation of matrix metalloproteinases (MMPs), namely collagenase-2 and -3 (Mukhin et al., 2006). Because collagenases lack an integral membrane domain, we hypothesized that integrins may play a role in BK-induced signaling by targeting collagenases to the membrane, thus forming a functional signaling complex. Integrins are heterodimeric receptors for cell-surface adhesion molecules and extracellular matrix proteins, which are composed of two subunits,  $\alpha$  and  $\beta$ . Each  $\alpha\beta$  combination has

This work was supported by the National Institutes of Health National Institute of Diabetes and Digestive and Kidney Diseases [Grant DK52448], the National Institutes of Health National Institute of General Medical Sciences [Grant GM3909], the Department of Veterans Affairs Merit and Research Enhancement Award Program (to J.R.R. and M.N.G.), the American Heart Association [Grant-in-Aid 0655445U], and a laboratory endowment jointly supported by the Medical University of South Carolina Division of Nephrology and Dialysis Clinics, Incorporated.

Parts of this work were presented previously in abstract form: Kramarenko I, Bunni M, Raymond JR, and Garnovskaya MN (2008) Bradykinin B<sub>2</sub> receptor interacts with integrin  $\alpha 5/\beta 1$  to transactivate epidermal growth factor receptor in kidney cells [abstract]. *FASEB J* 22:829.5.

Article, publication date, and citation information can be found at <http://molpharm.aspetjournals.org>.  
doi:10.1124/mol.110.064840.

**ABBREVIATIONS:** BK, bradykinin; ERK, extracellular signal-regulated protein kinase; EGFR, epidermal growth factor receptor; MMP, matrix metalloproteinase; GPCR, G protein-coupled receptor; FAK, focal adhesion kinase; EGF, epidermal growth factor; GAPDH, glyceraldehyde-3-phosphate dehydrogenase; PCR, polymerase chain reaction; RIPA, radioimmunoprecipitation assay; ANOVA, analysis of variance; siRNA, small interfering RNA.

specific signaling properties (for review, see Juliano, 2002). To date, 18  $\alpha$  and 8  $\beta$  subunits that form at least 24 different  $\alpha\beta$  integrins have been identified (Humphries et al., 2006).

The first interaction between integrins and MMPs was identified in melanoma cells in which it was demonstrated that the C-terminal domain of gelatinase-A [matrix metalloproteinase (MMP)-2] binds directly to integrin  $\alpha V\beta 3$ , which localizes MMP-2 in a proteolytically active form on the surface of invasive cells (Brooks et al., 1996). Furthermore, the involvement of the  $\alpha V\beta 3$ /MMP-2 complex in tumor growth and angiogenesis has been demonstrated in vivo (Brooks et al., 1998). Integrin  $\alpha V\beta 3$  also cooperates with gelatinase-B (MMP-9) to regulate migration of breast cancer cells (Rolli et al., 2003). Purified  $\beta 2$  integrin has been shown to bind to the catalytic domain in pro-MMP-9 gelatinase to form complexes of pro-MMP-9 with both the  $\alpha M\beta 2$  and  $\alpha L\beta 2$  integrins in leukemic cell lines; these associations probably control the activation of the proenzyme (Stefanidakis et al., 2003). Cell-surface interactions between  $\beta 2$  integrins and gelatinases play roles in normal leukocyte migration and in cancer progression (Stefanidakis and Koivunen, 2006). Interactions with integrins also have been demonstrated for collagenase-1 (MMP-1). Thus, pro-MMP-1 specifically binds to  $\alpha 2\beta 1$  integrin on keratinocytes, facilitating the cleavage of type I collagen and keratinocyte migration (Dumin et al., 2001). This binding occurs via the I-domain of the  $\alpha 2$  integrin subunit and requires both the linker domain and the hemopexin-like domain of MMP-1 (Stricker et al., 2001). MMP-1 also interacts with  $\alpha 2\beta 1$  integrin in human neurons (Conant et al., 2004) and with  $\alpha 1\beta 1$  integrin in monocytes (Stricker et al., 2001). However, no interactions between integrins and collagenases 2 or 3 have been described.

Interesting connections between integrins and G protein-coupled receptors (GPCRs) have been described recently. Integrin-mediated cell anchorage affects GPCR signaling to the ERK/mitogen-activated protein kinase (Della Rocca et al., 1999; Short et al., 2000). Muscarinic receptors stimulate tyrosine phosphorylation of focal adhesion kinase via an integrin-dependent mechanism (Slack, 1998). Focal adhesion kinase phosphorylation in response to bombesin and muscarinic signaling depends on the integrity of the cytoskeleton (Hunger-Glaser et al., 2003). Activation of histamine receptors results in translocation of the nonreceptor tyrosine kinase PYK2 to focal adhesions and enhances PYK2 tyrosine phosphorylation leading to ERK activation in HeLa cells (Litvak et al., 2000).

The precise molecular mechanism underlying integrin-mediated GPCR signaling to ERK remains to be defined, but one possibility involves integrin-mediated recruitment of cytoskeletal components to form a scaffold that facilitates efficient assembly of the components of the signaling pathway. Alternatively, some GPCRs interact directly with integrins: P2Y<sub>2</sub> nucleotide receptor contains an RGD motif in the first extracellular loop that enables it to interact with  $\alpha V\beta 3$  and  $\alpha V\beta 5$  integrins, leading to increased astrocyte migration (Bagchi et al., 2005; Wang et al., 2005). In addition,  $\beta 1$  and  $\beta 3$  integrins colocalize with the  $\mu$ -opioid receptor in sensory neurons and regulate receptor signaling, probably by altering its coupling to either G $\alpha_i$  or G $\alpha_s$  proteins (Berg et al., 2007).

Possible interactions between the BK B<sub>2</sub> receptor and integrins have not been studied. Herein, we provide multiple lines of evidence that suggest the involvement of integrin

$\alpha 5\beta 1$  in BK-induced signaling, and we describe physical and functional connections among BK B<sub>2</sub> receptor,  $\alpha 5\beta 1$  integrin, and collagenase-2 and -3 in mIMCD-3 cells.

## Materials and Methods

**Materials.** Cell culture supplies were obtained from Invitrogen (Carlsbad, CA), or Corning Life Sciences (Lowell, MA). Bradykinin, EGF, and other reagent-grade chemicals were from Sigma-Aldrich (St. Louis, MO). Cyclic RGD/control RGD peptides and MMP inhibitors were from Calbiochem (La Jolla, CA). Phospho-ERK kit was obtained from Cell Signaling Technology (Danvers, MA). Monoclonal anti-BK B<sub>2</sub> receptor antibody was from BD Biosciences Transduction Laboratories (Lexington, KY). EGFR polyclonal antibody; anti-phospho-EGFR (Tyr 1173) monoclonal antibody; monoclonal anti-phosphotyrosine antibody; MMP-8 monoclonal antibody; MMP-13 monoclonal antibody; anti- $\alpha 5\beta 1$  integrin polyclonal antibody; monoclonal blocking antibodies against  $\alpha 2\beta 1$ ,  $\alpha 5\beta 1$ ,  $\alpha V\beta 6$ , and  $\alpha 6$  integrins; antibodies against  $\alpha 2$ ,  $\alpha 3$ ,  $\alpha V$ , and  $\beta 1$  integrin subunits; and mouse monoclonal glyceraldehyde-3-phosphate dehydrogenase (GAPDH) antibody were from Millipore (Billerica, MA). Antibodies against integrin  $\alpha 2$ ,  $\alpha 5$ ,  $\alpha 6$ ,  $\beta 6$ , and  $\beta 1$  subunits, blocking antibodies against  $\alpha 3$ ,  $\alpha V$ , and  $\beta 3$  integrins, MMP-8 siRNA, MMP-13 siRNA, integrin  $\alpha 2$  siRNA, integrin  $\alpha 5$  siRNA, integrin  $\beta 1$  siRNA, and control siRNA were from Santa Cruz Biotechnology (Santa Cruz, CA). The RNA Stat-60 reagent was from Tel-Test, Inc., (Friendswood, TX). The RT<sup>2</sup> Profiler PCR Array System for mouse extracellular matrix and adhesion molecules and the Oligo GEArray kit for mouse extracellular matrix and adhesion molecules were from SuperArray Bioscience Corporation (Frederick, MD).

**Cell Culture.** mIMCD-3 cells were obtained from the American Type Culture Collection (Manassas, VA). mIMCD-3 cells were grown in equal mixtures of Dulbecco's modified Eagle's medium and Ham's F12 medium, supplemented with 10% fetal bovine serum, 100 U/ml penicillin, and 100  $\mu$ g/ml streptomycin at 37°C in a humidified atmosphere of 95% air and 5% CO<sub>2</sub>.

**Study of Expression Levels of Integrin Subunits in mIMCD-3 Cells.** To monitor the expression profile of integrins in mIMCD-3 cells, we employed the Oligo GEArray kit for mouse extracellular matrix and adhesion molecules. Using the RNA Stat-60 reagent, we extracted total RNA from mIMCD-3 cells that were grown on plastic plates and converted RNA into biotin-labeled cRNA target probes for Microarray hybridization by using the TrueLabeling-AMP linear RNA amplification kit (SuperArray Bioscience Corporation). The cRNA targets (6  $\mu$ g of cRNA) next were hybridized with oligonucleotide probes, representing different genes, printed on a nylon membrane. The resulting products on arrayed membranes were detected by a chemiluminescent detection kit, and analyzed by GEArray Analyzer data analysis software. We also used an RT<sup>2</sup> Profiler PCR Array System for mouse extracellular matrix and adhesion molecules that validates the expression of 84 relevant genes for cell-cell and cell-matrix interactions, including nine  $\alpha$  and four  $\beta$  integrin subunits. Total RNA from mIMCD-3 cells was extracted as above and converted to cDNA using the RT<sup>2</sup> Profiler PCR Array first strand kit. Quantitative real-time PCR was performed according to the manufacturer's protocol. The presence of various integrin subunits in lysates from mIMCD-3 cells was supported by Western blots using commercially available antibodies.

**Transfections of mIMCD-3 Cells: Integrin and MMP Silencing.** Transfections of mIMCD-3 cells were achieved by nucleofection with an Amaxa Biosystems instrument (Giessen, Germany). Cells (10<sup>6</sup>) were resuspended in 100  $\mu$ l of Nucleofector solution R and nucleofected with either 100 nM siRNA for MMP-8, MMP-13, or integrin  $\alpha 5\beta 1$  siRNA, or with the same amount of control siRNA using manufacturer's protocol T-16. Forty-eight hours after nucleofection, cells were stimulated with BK, EGF, or vehicle; lysed; and analyzed for integrin and MMP expression by Western blotting with anti-MMP-8,

anti-MMP-13, or anti-integrin  $\alpha 5$  and  $\beta 1$  rabbit polyclonal antibodies and for EGFR phosphorylation and ERK activation. Blots were re-probed with a mouse monoclonal GAPDH antibody to control for protein loading and for silencing specificity and efficiency.

**ERK Assay.** ERK phosphorylation was assessed by Western blot using a phosphorylation-state specific antibody that specifically recognizes Thr<sup>202</sup>- and Tyr<sup>204</sup>-phosphorylated ERK-1 and ERK-2 as described previously (Mukhin et al., 2003) in mIMCD-3 cells treated for 5 min with BK, EGF, or vehicle. The membranes were stripped using Re-Blot Plus antibody stripping solution (Millipore) and re-probed with the control ERK antibody, which recognizes phosphorylated and nonphosphorylated ERK equally well. Results are presented as intensities of phospho-ERK bands relative to total ERK bands and are expressed as fold of basal phosphorylation (nontreated cells).

**EGF Receptor Phosphorylation Assay.** The phosphorylation state of EGFR was assessed by immunoprecipitation/Western blotting studies as described previously (Mukhin et al., 2003). Quiescent mIMCD-3 cells, grown in 100-mm dishes, were pretreated with vehicle or inhibitors for 30 min. Subsequently, cells were treated with 100 nM BK, 1 ng/ml EGF, or vehicle for 5 min and lysed in RIPA buffer (50 mM Tris-HCl, pH 7.4, 1% Nonidet P-40, 0.25% sodium deoxycholate, 150 mM NaCl, 1 mM EDTA, 1 mM sodium fluoride, 1 mM Na<sub>3</sub>VO<sub>4</sub>, 1  $\mu$ g/ml each aprotinin, leupeptin, and pepstatin). Cell lysates were precleared by incubating with protein A-agarose bead slurries for 30 min at 4°C. Precleared lysates (1  $\mu$ g/ $\mu$ l total cell protein) were incubated with 4  $\mu$ g of anti-EGFR polyclonal IgG overnight at 4°C. The immunocomplexes were captured by the addition of protein A-agarose bead slurries, with incubation for a further 2 h at 4°C. The agarose beads were collected by centrifugation, washed three times with RIPA buffer, resuspended in 2 $\times$  Laemmli sample buffer, boiled for 5 min, and subjected to SDS-polyacrylamide gel electrophoresis. After wet transfer to polyvinylidene difluoride membranes, the membranes were probed with monoclonal anti-phospho-EGFR (Tyr<sup>1173</sup>) antibody to assess the phosphorylation state of EGFR, or with EGFR antibody.

**Studies of Possible Complex Formation between Integrin  $\alpha 5\beta 1$ , BK B<sub>2</sub> Receptor, and Collagenases.** Quiescent mIMCD-3 cells, grown in 100-mm dishes, were treated with 100 nM BK, 1 ng/ml EGF, or with vehicle for different time periods and lysed in RIPA buffer. Cell lysates were precleared by incubating with protein A-agarose bead slurries for 30 min at 4°C. Precleared lysates (1  $\mu$ g/ $\mu$ l total cell protein) were incubated with 4  $\mu$ g of anti- $\alpha 5\beta 1$  integrin polyclonal IgG overnight at 4°C. The immunocomplexes were captured by addition of protein A-agarose bead slurry and incubation for a further 2 h at 4°C. The agarose beads were collected by centrifugation, washed three times with RIPA buffer, resuspended in 2 $\times$  Laemmli sample buffer, boiled for 5 min, and subjected to SDS-polyacrylamide gel electrophoresis. After wet transfer to polyvinylidene difluoride membranes, the membranes were probed with monoclonal anti-BK B<sub>2</sub> receptor, with monoclonal anti-MMP-8 or anti-MMP-13 antibody, and with anti-EGFR antibody to study possible physical associations of these proteins.

**Data Analysis.** Data are presented as mean + S.E.M. and were analyzed for repeated measures by Student's *t* test for unpaired two-tailed analysis and by ANOVA. Differences were considered significant at *p* < 0.05.

## Results

### Integrins are Involved in BK-Induced ERK1/2 Activation and EGFR Transactivation in mIMCD-3 Cells.

In the first set of experiments, we tested the involvement of integrins in B<sub>2</sub> receptor-induced EGFR transactivation and ERK phosphorylation using Arg-Gly-Asp (RGD) peptides. RGD was originally identified as the sequence in fibronectin that is a recognition site for  $\alpha 5\beta 1$  integrin, but it also serves as a recognition motif for other integrins, including  $\alpha 3\beta 1$ ,

$\alpha 8\beta 1$ ,  $\alpha V\beta 1$ ,  $\alpha V\beta 3$ ,  $\alpha V\beta 5$ ,  $\alpha V\beta 6$ , and  $\alpha 2\beta 3$  (Plow et al., 2000). Quiescent mIMCD-3 cells were pretreated for 1 h with 28  $\mu$ M concentrations of either control or cyclic RGD peptides before stimulation with 100 nM BK or with 1 ng/ml EGF for 5 min. Control RGD peptides were without effect, whereas cyclic RGD peptides reduced the activation of ERK by BK by ~65%, suggesting that integrins with RGD recognition specificity may be involved in BK-induced ERK phosphorylation (Fig. 1A). At the same time, cyclic RGD peptides did not affect EGF-induced ERK activation (Fig. 1C).

### Expression Profile of Integrin Subunits in mIMCD-3 Cells.


Having established a probable role for integrins in BK-induced signaling in mIMCD-3 cells, we attempted to examine specific integrins that could mediate this response. To plan more specific experiments, we needed to determine which integrins are present in mIMCD-3 cells, because at least 18  $\alpha$  and 8  $\beta$  integrin subunits have been identified. First, we employed the Oligo GEArray kit for mouse extracellular matrix and adhesion molecules, which represents 113 genes encoding proteins important for the attachment of cells to their surroundings including various types of cell adhesion molecules (such as the integrins, IgG superfamily members, cadherins and catenins, and selectins) as well as extracellular matrix proteins, proteases (such as the matrix metalloproteinases and the serine and cysteine proteinases), and their inhibitors. This array allowed us to determine simultaneously the expression profile of 13  $\alpha$  and 7  $\beta$  integrin subunits in mIMCD-3 cells. Table 1 summarizes mRNA expression of integrins in mIMCD-3. Integrin subunits  $\alpha 2b$ ,  $\alpha 3$ ,  $\alpha V$ , and  $\beta 1$  seemed to be the most abundant in mIMCD-3 cells. The messages for  $\alpha 2$ ,  $\alpha 6$ ,  $\alpha 7$ ,  $\alpha X$ ,  $\beta 6$ , and  $\beta 7$  integrin subunits also were detectable. To our surprise, we did not detect message for  $\alpha 5$  integrin subunit, although we expected to find  $\alpha 5\beta 1$  integrin in mIMCD-3 cells based on the ability of RGD peptides to affect BK-induced signaling (Fig. 1A).

To verify the microarray data, we employed an RT<sup>2</sup> Profiler PCR Array System for mouse extracellular matrix and adhesion molecules that measures the expression of 84 relevant genes for cell-cell and cell-matrix interactions, including nine  $\alpha$  ( $\alpha 2$ ,  $\alpha 3$ ,  $\alpha 4$ ,  $\alpha 5$ ,  $\alpha E$ ,  $\alpha L$ ,  $\alpha M$ ,  $\alpha V$ , and  $\alpha X$ ) and four  $\beta$  ( $\beta 1$ ,  $\beta 2$ ,  $\beta 3$ , and  $\beta 4$ ) integrin subunits. The expression of the *GAPDH* gene was used as control. Reverse transcription-PCR results confirmed the expression of  $\alpha 2$ ,  $\alpha 3$ ,  $\alpha X$ ,  $\alpha V$  and  $\beta 1$  genes identified by the Oligo microarray (not shown). In addition, an RT<sup>2</sup> Profiler PCR Array System allowed us to detect the message for  $\alpha 5$  integrin subunit. Our inability to detect the message for  $\alpha 5$  integrin subunit by the Oligo GEArray could be caused by a technical issue with this particular Oligo GEArray layout, because the message for  $\alpha 5$  integrin (spot 48) is probably masked by a strong message for fibronectin-1 (spot 40), which is highly expressed in mIMCD-3 cells. Furthermore, to confirm the mRNA data and to support the expression of integrins on a protein level, we performed Western blotting on mIMCD-3 lysates using commercially available antibodies against integrins. Results presented in Fig. 2 support the presence of integrins  $\alpha 2$ ,  $\alpha 3$ ,  $\alpha 5$ ,  $\alpha 6$ ,  $\alpha V$ ,  $\beta 1$ , and  $\beta 6$  in mIMCD-3 cells.

### Roles of Specific Integrins in BK-Induced ERK1/2 Activation in mIMCD-3 Cells.

To examine specific integrins that are involved in BK-induced ERK phosphorylation, we employed commercially available neutralizing antibodies against mouse integrin subunits. Quiescent mIMCD-3 cells





**Fig. 1.** Involvement of integrins in BK-induced ERK phosphorylation. Quiescent mIMCD-3 cells were pretreated for 1 h either with 28  $\mu$ M control or with cyclic RGD peptides before stimulation with 100 nM BK (A) or 1 ng/ml EGF (C) for 5 min. Quiescent mIMCD-3 cells were pretreated for 2 h with 0.1 mg/ml neutralizing anti-integrin antibodies before stimulation with 100 nM BK (B) or 1 ng/ml EGF (D) for 5 min. Control samples were preincubated with 0.1 mg/ml normal rabbit IgG. ERK phosphorylation was measured as described under *Materials and Methods*. Bars represent intensities of phospho-ERK bands relative to total ERK expressed as fold of basal (cells treated with vehicle). Experiments were performed three times in duplicate. Data are presented as mean + S.E.M. \*\*\*,  $p < 0.001$ .

were pretreated for 2 h with anti-integrin antibodies or normal rabbit IgG before stimulation with 100 nM BK or with 1 ng/ml EGF for 5 min. Neutralizing antibody against  $\alpha 5\beta 1$  integrin, when used in a concentration of 100 ng/ml, significantly ( $\sim 60\%$ ) blocked BK-induced phosphorylation of ERK in mIMCD-3 cells (Fig. 1B;  $n = 3$ ,  $p < 0.001$ ), but did not affect EGF-induced ERK activation (Fig. 1D). Neutralizing antibody against  $\alpha 2\beta 1$  integrin in some experiments also partially ( $\sim 25\%$ ) decreased BK-induced phosphorylation of ERK in mIMCD-3 cells, although these results were not statistically significant (not shown). At the same time, neutralizing antibodies against  $\alpha 3$ ,  $\alpha V$ ,  $\beta 6$ , and  $\alpha V\beta 6$  were without effect (not shown), suggesting specificity in the involvement of  $\alpha 5\beta 1$  integrin in BK-induced ERK phosphorylation.

#### Transfection of mIMCD-3 Cells with Integrin and MMP siRNAs Decreases BK-Induced ERK Activation.


To further support the involvement of integrins in BK-induced ERK activation, we employed RNA-mediated interference to knock down the expression of  $\alpha 5\beta 1$  integrin. mIMCD-3 cells were nucleofected with either 100 nM siRNA for integrin  $\alpha 5\beta 1$  or with the same amount of control siRNA. Forty-eight hours after nucleofection, cells were stimulated either with vehicle or 100 nM BK or with 1 ng/ml EGF for 5 min, lysed, and analyzed for ERK phosphorylation. Cells

TABLE 1

mIMCD-3 cells express mRNA for  $\alpha$  and  $\beta$  integrin subunits

The expression of mRNA was assessed by the Oligo GEArray as described under *Materials and Methods*. Proteins for which we were able to detect message are shown in bold

	Gene Symbol	Position	Signal Intensity
1	<b><math>\alpha 2</math></b>	44	+
2	<b><math>\alpha 2b</math></b>	45	++++
3	<b><math>\alpha 3</math></b>	46	+++
4	$\alpha 4$	47	-
5	$\alpha 5$	48	-
6	<b><math>\alpha 6</math></b>	49	+
7	<b><math>\alpha 7</math></b>	50	+
8	$\alpha 8$	51	-
9	$\alpha E$	52	-
10	$\alpha L$	53	-
11	$\alpha M$	54	-
12	<b><math>\alpha V</math></b>	55	+++
13	<b><math>\alpha X</math></b>	56	++
14	<b><math>\beta 1</math></b>	57	++++
15	$\beta 2$	58	-
16	$\beta 3$	59	-
17	$\beta 4$	60	-
18	$\beta 5$	61	-
19	<b><math>\beta 6</math></b>	62	++
20	<b><math>\beta 7</math></b>	63	++



**Fig. 2.** Western blot analysis of mIMCD-3 lysates. Western blot analyses of mIMCD-3 lysates (40  $\mu$ g of total protein) were performed with commercially available anti-integrin antibodies to demonstrate the expression of these integrins on a protein level. The bands of  $\alpha 2$  (155 kDa),  $\alpha 3$  (150 and 130 kDa),  $\alpha 5$  (150 kDa),  $\alpha 6$  (140 kDa),  $\alpha V$  (150 kDa),  $\beta 1$  (130 kDa), and  $\beta 6$  (97 kDa) are indicated.

transfected with  $\alpha 5\beta 1$  integrin siRNAs demonstrated  $\sim 80\%$  less BK-induced ERK activation than cells transfected with control siRNA (Fig. 3A). EGF-induced ERK activation was not affected by  $\alpha 5\beta 1$  integrin down-regulation (Fig. 3B). Silencing of  $\alpha 2\beta 1$  integrin did not affect either BK-induced or EGF-induced phosphorylation of ERK in mIMCD-3 cells (not shown), suggesting specificity in the involvement of  $\alpha 5\beta 1$  integrin in BK-induced ERK phosphorylation.

We demonstrated previously the involvement of MMP-8 and MMP-13 in BK-induced ERK activation (Mukhin et al., 2006); therefore, we performed the combined inhibition of MMP-8 or MMP-13 and  $\alpha 5\beta 1$  integrins by cotransfecting cells with MMP-8 or MMP-13 siRNA in addition to integrin siRNAs. Cells transfected with a combination of MMP-8 and  $\alpha 5\beta 1$  integrin siRNAs demonstrated  $\sim 75\%$  decrease in BK-induced ERK phosphorylation (Fig. 3A). Cells transfected with a combination of MMP-13 and  $\alpha 5\beta 1$  integrin siRNAs demonstrated  $\sim 90\%$  decrease in BK-induced ERK phosphorylation. Because in our previous study, we proposed roles for both collagenases in BK-induced ERK phosphorylation (Mukhin et al., 2006), we blocked both MMP-8 and MMP-13 by transfecting cells with a combination of MMP-8 and MMP-13 siRNAs in addition to silencing  $\alpha 5\beta 1$  integrin. Each set of transfected cells demonstrated significant decreases in BK-induced ERK phosphorylation compared with the set transfected with scrambled siRNA. Although differences between cells transfected only with  $\alpha 5\beta 1$  integrin siRNA and cells cotransfected with MMP-8 and MMP-13 siRNAs in addition to  $\alpha 5\beta 1$  integrin siRNA were not statistically significant; in all experiments, there was a trend that down-regulation of both collagenases in addition to  $\alpha 5\beta 1$  integrin resulted in a complete inhibition of the signal. Down-regulation of  $\alpha 5\beta 1$  integrin either alone or in combination with MMP-8 and/or MMP-13 did not affect EGF-induced ERK activation (Fig. 3B). Effective silencing of integrin and MMP expression in mIMCD-3 cells transfected with siRNAs was supported by immunoblotting (Fig. 4C). These results support the hypothesis that integrin  $\alpha 5\beta 1$  very probably mediates BK-induced signaling in mIMCD-3 cells.

**Transfection of mIMCD-3 Cells with Integrin  $\alpha 5\beta 1$  and MMP-13 and MMP-8 siRNAs Decreases BK-Induced EGFR Phosphorylation.** Because BK activates ERK in mIMCD-3 cells via EGFR transactivation (Mukhin et al., 2003), in the next series of experiments, we assessed the involvement of  $\alpha 5\beta 1$  integrin in BK-induced EGFR phosphorylation. To test the involvement of integrin  $\alpha 5\beta 1$ , we transfected mIMCD-3 cells with either 100 nM integrin  $\alpha 5\beta 1$  siRNA or with control (scrambled) siRNA. To block the activity of collagenases, we silenced MMP-8 and/or MMP-13 expression by nucleofection with MMP-8 siRNA, MMP-13 siRNA, or with both MMP-8 and MMP-13 siRNAs. Forty-eight hours after nucleofection, cells were stimulated either with vehicle or 100 nM BK or with 1 ng/ml EGF for 5 min, lysed, and analyzed for EGFR phosphorylation as described under *Materials and Methods*. Silencing of integrin  $\alpha 5\beta 1$  significantly ( $p < 0.01$ ) decreased BK-induced EGFR phosphorylation (Fig. 4A). At the same time, the combined inhibition of MMP-8 and  $\alpha 5\beta 1$  integrin significantly blocked BK-induced EGFR phosphorylation by  $\sim 70\%$ , and the combined inhibition of MMP-13 and  $\alpha 5\beta 1$  integrin completely abolished the effect of BK (Fig. 4). Similar results were obtained when we blocked the activity of collagenases by pretreatment


of mIMCD-3 cells transfected with integrin  $\alpha 5\beta 1$  siRNA with chemical inhibitors of MMP-8 and MMP-13 (not shown), suggesting that MMP-8, MMP-13, and  $\alpha 5\beta 1$  integrin are involved in BK-induced phosphorylation of EGFR. However, differences in BK-induced EGFR phosphorylation between cells transfected only with  $\alpha 5\beta 1$  integrin siRNA and cells cotransfected with MMP-8 and MMP-13 siRNAs in addition to  $\alpha 5\beta 1$  integrin siRNA were not statistically significant. At the same time, down-regulation of  $\alpha 5\beta 1$  integrin either alone or in combination with MMP-8 and/or MMP-13 did not affect EGF-induced EGFR phosphorylation (Fig. 4B).

**BK Induces Complex Formation between  $\alpha 5\beta 1$  Integrin, EGFR, MMP-13, MMP-8, and BK B<sub>2</sub> Receptor.** We next proposed that BK stimulates the assembly of a signal transduction complex, which includes molecules involved in EGFR phosphorylation, and examined whether BK could induce a physical interaction between  $\alpha 5\beta 1$  integrin, EGFR, and collagenases. To explore this possibility, we used immunoprecipitation of lysates from cells treated with vehicle, BK, or EGF with anti- $\alpha 5\beta 1$  integrin antibody, followed by Western blotting with antibody against EGFR, MMP-8, MMP-13, and BK B<sub>2</sub> receptor. Figure 5A shows that  $\alpha 5\beta 1$  integrin and EGFR (175 kDa) coimmunoprecipitate and that their association can be increased by stimulation of mIMCD-3 cells with BK. Stimulation with EGF did not significantly increase association between  $\alpha 5\beta 1$  integrin and EGFR, supporting the idea that  $\alpha 5\beta 1$  integrin is not essential for EGF-induced EGFR activation. Figure 5B shows that immunoprecipitation of  $\alpha 5\beta 1$  integrin from mIMCD-3 cells treated with BK, but not with EGF, resulted in coprecipitation of the BK B<sub>2</sub> recep-


tor (42 kDa). The BK B<sub>2</sub> receptor antibody that we employed for these studies has been used previously by our group (Kramarenko et al., 2009) and others (Golser et al., 2000; Xie et al., 2000) to demonstrate the presence of the endogenous and/or transfected BK B<sub>2</sub> receptor in different cell lines by Western blotting, and also to immunoprecipitate BK B<sub>2</sub> receptor (Golser et al., 2000). The bands corresponding to EGFR or BK B<sub>2</sub> receptor were not detectable in precipitates obtained from identical samples precipitated in the presence of normal rat immunoglobulins instead of  $\alpha 5\beta 1$  integrin antibody (data not shown). These results suggest that BK induces coprecipitation of  $\alpha 5\beta 1$  integrin with EGFR and BK B<sub>2</sub> receptor. Furthermore, we found that MMP-8 and MMP-13 also are present in BK-induced signaling complex (not shown).

## Discussion


The current work describes a novel mechanism of EGFR transactivation by the G<sub>q</sub>-coupled BK B<sub>2</sub> receptor in mIMCD-3 cells that involves integrin  $\alpha 5\beta 1$ . What is new about this work is that we have (1) characterized the repertoire of integrins in mIMCD-3 cells using reverse transcription-PCR, microarray detection, and Western blotting; (2) implicated integrins as key mediators of BK-induced ERK activation using RGD peptides; (3) provided evidence that integrin  $\alpha 5\beta 1$  is involved in BK-induced EGFR transactivation based on results of experiments using neutralizing anti-integrin antibodies and siRNA; and (4) demonstrated for the first time that BK induces formation of a signaling complex among



**Fig. 3.** Transfection of mIMCD-3 cells with integrin and MMP siRNAs decreases BK-induced ERK activation. mIMCD-3 cells were nucleofected either with 100 nM siRNA for integrin  $\alpha 5\beta 1$  alone (- $\alpha 5\beta 1$ ) or with combinations of MMP-8 siRNA (- $\alpha 5\beta 1$ -MMP-8) and/or MMP-13 siRNA (- $\alpha 5\beta 1$ -MMP-13), or with combinations of all siRNAs (- $\alpha 5\beta 1$ -MMP-8-MMP-13), or with the same amount of control siRNA (control). Forty-eight hours after nucleofection, cells were stimulated with vehicle or 100 nM BK (A) or with 1 ng/ml EGF (B) for 5 min, lysed, and analyzed for ERK phosphorylation. ERK phosphorylation was measured as described under *Materials and Methods*. Bars represent intensities of phospho-ERK bands relative to total ERK expressed as fold of basal (cells treated with vehicle). Experiments were performed three times in duplicate. Data are presented as mean  $\pm$  S.E.M. \*\*,  $p < 0.01$  compared with control BK-treated cells. ANOVA - $\alpha 5\beta 1$  compared with - $\alpha 5\beta 1$ -MMP-8, - $\alpha 5\beta 1$ -MMP-13, or with - $\alpha 5\beta 1$ -MMP-8-MMP-13, not significant.



**Fig. 4.** Transfection of mIMCD-3 cells with integrin  $\alpha 5\beta 1$  and MMP siRNAs decreases BK-induced EGFR phosphorylation. Cells were nucleofected with 100 nM  $\alpha 5\beta 1$  siRNA (- $\alpha 5\beta 1$ ) or - $\alpha 5\beta 1$  with a combination of either MMP-8 siRNA (- $\alpha 5\beta 1$ -MMP-8) or MMP-13 siRNA (- $\alpha 5\beta 1$ -MMP-13); with a combination of all siRNAs (- $\alpha 5\beta 1$ -MMP-8-MMP-13); or with the same amount of control siRNA (control), as described under *Materials and Methods*. Forty-eight hours after nucleofection, cells were stimulated with vehicle or 100 nM BK (A) or with 1 ng/ml EGF (B) for 5 min, lysed, and analyzed for EGFR phosphorylation as described under *Materials and Methods*. Experiments were performed at least three times. Data are presented as mean + S.E.M. \*\*,  $p < 0.01$  compared with control BK-treated cells. ANOVA (- $\alpha 5\beta 1$ ) compared with - $\alpha 5\beta 1$ -MMP-8, - $\alpha 5\beta 1$ -MMP-13, or - $\alpha 5\beta 1$ -MMP-8-MMP-13, not significant. C, Western blot analyses of lysates of mIMCD-3 cells transfected with either scrambled siRNA or siRNAs for  $\alpha 5$ ,  $\beta 1$ , MMP-8, and MMP-13 (40  $\mu$ g of total protein) were performed with commercially available antibodies against  $\alpha 5$  and  $\beta 1$  integrin subunits, MMP-8 and MMP-13 to demonstrate down-regulation of these proteins. Blots were stripped and re-probed with antibody against GAPDH to control for the specificity of silencing and protein loading.



**Fig. 5.** BK induces complex formation between EGFR and  $\alpha 5\beta 1$  integrin. Lysates from mIMCD-3 cells treated with vehicle, 100 nM BK, or 1 ng/ml EGF were immunoprecipitated with anti- $\alpha 5\beta 1$  integrin antibody as described under *Materials and Methods*. Immunoblotting was performed with antibodies against EGFR (A) and BK B<sub>2</sub> receptor (B). The blots shown are representative of four experiments. A, coimmunoprecipitation experiments show that  $\alpha 5\beta 1$  integrin and EGFR coimmunoprecipitate and that their association can be increased by stimulation of mIMCD-3 cells with 100 nM BK but not with EGF. Inset, representative Western blot with antibody against EGFR showing immunoreactive band at 175 kDa. Blot was stripped and re-probed with antibody against  $\alpha 5$  integrin to control for immunoprecipitation and protein loading. Immunoreactive band at 150 kDa is shown. B, BK B<sub>2</sub> receptor coimmunoprecipitates with  $\alpha 5\beta 1$  integrin. Inset, representative Western blot with antibody against BK B<sub>2</sub> receptor showing immunoreactive duplet at 42/40 kDa. Blot was stripped and re-probed with antibody against  $\alpha 5$  integrin to control for immunoprecipitation and protein loading. Immunoreactive band at 150 kDa is shown. IP, immunoprecipitation; IB, immunoblot.



$\alpha 5\beta 1$  integrin, EGFR, the BK  $B_2$  receptor, and probably MMP-13 and MMP-8.

In our previous work, we established that the BK  $B_2$  receptor stimulates early mitogenic signals associated with activation of ERK1/2 in mIMCD-3 cells and demonstrated that BK-induced cell proliferation was dependent on activation of EGFR (Mukhin et al., 2003). Furthermore, we described a novel mechanism of EGFR transactivation by the  $G_q$ -coupled BK  $B_2$  receptor that involves activation of MMPs, namely collagenases 2 and 3 (MMP-8 and MMP-13). We demonstrated that collagenases 2 and 3 are activated by the BK  $B_2$  receptor in mIMCD-3 cells, and are involved in cross-talk between the  $B_2$  receptor and EGFR (Mukhin et al., 2006). In the current study, we looked further into this mechanism by testing the hypothesis that in mIMCD-3 cells, integrins may play a role in BK-induced signaling by targeting collagenases to the membrane, thus forming a functional signaling complex. Several GPCRs have been shown to activate ERK in an integrin-dependent manner: thrombin, BK, and lysophosphatidic acid receptors in PC12 rat pheochromocytoma cells (Della Rocca et al., 1999); gonadotropin-releasing hormone receptors expressed in HEK293 cells (Davidson et al., 2004); P2Y receptors in endothelial cells (Short et al., 2000); histamine receptors in HeLa cells (Litvak et al., 2000); and  $\delta$ -opioid receptors transfected into HEK293 cells and endogenously expressed in neuroblastoma x glioma hybrid NG108-15 cells (Eisinger and Ammer, 2008).

In our study, we assessed the involvement of integrins in BK-induced signaling in mIMCD-3 cells, using RGD-containing synthetic peptides that inhibit ligand binding to integrins with RGD recognition specificity (e.g., integrins  $\alpha 3\beta 1$ ,  $\alpha 5\beta 1$ ,  $\alpha 8\beta 1$ ,  $\alpha V\beta 1$ ,  $\alpha V\beta 3$ ,  $\alpha V\beta 5$ ,  $\alpha V\beta 6$ , and  $\alpha 2\beta 3$ ). RGD peptides decreased BK-induced ERK activation by  $\sim 65\%$  (Fig. 1A) without affecting EGF-induced ERK activation (Fig. 1C), supporting the hypothesis that integrins are involved in BK-induced ERK phosphorylation in mIMCD-3 cells. Next, we aimed to determine specific integrins responsible for mediating BK-induced ERK activation. Using two different methods of mRNA analysis and Western blotting, we established that mIMCD-3 cells express the following integrin subunits:  $\alpha 2$ ,  $\alpha 2b$ ,  $\alpha 3$ ,  $\alpha 5$ ,  $\alpha V$ ,  $\alpha X$ ,  $\beta 1$ , and  $\beta 6$  (Table and Fig. 2). These subunits may form four integrins with RGD recognition specificity:  $\alpha 3\beta 1$ ,  $\alpha 5\beta 1$ ,  $\alpha V\beta 1$ , and  $\alpha V\beta 6$ . We employed neutralizing antibodies against these integrins to study their possible involvement in BK-induced ERK activation. Neutralizing antibody against  $\alpha 5\beta 1$  integrin significantly blocked BK-induced phosphorylation of ERK by  $\sim 60\%$  in mIMCD-3 cells without affecting EGF-induced ERK activation (Fig. 1, B and D), suggesting the involvement of  $\alpha 5\beta 1$  integrin in BK-induced ERK phosphorylation. At the same time, neutralizing antibodies against  $\alpha 3$ ,  $\alpha V$ ,  $\beta 6$ , and  $\alpha V\beta 6$  integrins did not change BK-induced ERK phosphorylation, suggesting a lack of involvement of  $\alpha 3\beta 1$ ,  $\alpha V\beta 1$ , and  $\alpha V\beta 6$  integrins. We further supported the involvement of  $\alpha 5\beta 1$  integrin in BK-induced signaling by using mIMCD-3 cells, in which the expression of  $\alpha 5$  and  $\beta 1$  subunits was knocked down by an RNA-mediated interference (Fig. 3A). Because silencing of integrin  $\alpha 5\beta 1$  also decreased BK-induced EGFR phosphorylation (Fig. 4A), the present findings indicate that BK-induced ERK activation is mediated by integrin-stimulated EGFR. At the same time, down-regulation of  $\alpha 5\beta 1$  integrin, either alone or in combination with MMP-8 and/or MMP-13,

did not affect EGF-induced EGFR phosphorylation and ERK activation (Figs. 3B and 4B). Thus, for the first time, we have identified a specific integrin ( $\alpha 5\beta 1$ ) that specifically mediates BK-induced EGFR transactivation and ERK phosphorylation in mIMCD-3 cells.

The ability of integrins to cooperate with receptor tyrosine kinases, including EGFR, to transduce proliferative signals and to regulate cell survival and migration has been discussed previously (Miranti and Brugge, 2002; Schwartz and Ginsberg, 2002). Integrins are able to form physical complexes with EGFR at the cell membrane and to trigger ligand-independent phosphorylation of Tyr<sup>845</sup>, Tyr<sup>1068</sup>, Tyr<sup>1086</sup>, and Tyr<sup>1173</sup> residues in the EGFR molecule (Moro et al., 1998). This integrin-dependent EGFR activation seems to be necessary for full EGFR-dependent transcriptional responses (Cabodi et al., 2004). GPCR-dependent transactivation of EGFR is usually mediated by MMP-dependent processing of membrane EGFR pro-ligands (Prenzel et al., 1999). In our previous study, we demonstrated the involvement of MMP-8 and MMP-13 in BK-induced EGFR transactivation in mIMCD-3 cells; at the same time, we found that this transactivation does not require extracellular release of EGF-like growth factors such as HB-EGF and/or TGF- $\alpha$  (Mukhin et al., 2006). Our present data favor the possibility that BK induces association of EGFR with integrin  $\alpha 5\beta 1$ , thus causing ligand-independent phosphorylation and activation of EGFR. In support of that idea, we showed that  $\alpha 5\beta 1$  integrin and EGFR coimmunoprecipitate upon BK treatment (Fig. 5A).

The concept that the association of MMPs with integrins can modify intracellular signaling has been demonstrated in several cell models (for review, see Stefanidakis and Koivunen, 2006). However, most reports describe interactions between integrins and gelatinases (MMP-2 and MMP-9) either in tumor cells (Brooks et al., 1998; Rolli et al., 2003) or in leukocytes (Stefanidakis et al., 2003). The only collagenase described to interact with integrins, collagenase-1 (MMP-1), was reported to be associated with  $\alpha 2\beta 1$  integrin in keratinocytes (Dumin et al., 2001) and in human neurons (Conant et al., 2004) and with  $\alpha 1\beta 1$  integrin in monocytes (Stricker et al., 2001). In the current work, the involvement of MMP-8 and MMP-13 in BK-induced ERK activation was supported by combined inhibition of MMP-8 and/or MMP-13 and  $\alpha 5\beta 1$  integrin by cotransfecting cells with MMP-8 or MMP-13 siRNA in addition to integrin siRNAs. Although differences between cells transfected only with  $\alpha 5\beta 1$  integrin siRNA and cells cotransfected with MMP-8 and MMP-13 siRNAs in addition to  $\alpha 5\beta 1$  integrin siRNA were not statistically significant, in all experiments, there was a trend that down-regulation of both collagenases in addition to  $\alpha 5\beta 1$  integrin resulted in a complete inhibition of BK-induced EGFR phosphorylation and ERK activation. (Figs. 3A and 4A). Thus, the current work suggests that collagenase-2 and -3 (MMP-8 and MMP-13) may act in concert with integrin  $\alpha 5\beta 1$  to mediate BK-induced phosphorylation of EGFR and ERK in kidney cells. It is noteworthy that the BK  $B_2$  receptor also was coimmunoprecipitated with  $\alpha 5\beta 1$  integrin after BK treatment of mIMCD-3 cells (Fig. 5B).

This work is novel in that there are only a few reports regarding interactions between GPCRs and integrins. Colocalization of  $\beta 1$  and  $\beta 3$  integrins with the  $\mu$ -opioid receptor was detected in trigeminal ganglion neurons by immunocytochemistry and confocal imaging (Berg et al., 2007). These



RGD-binding integrins probably regulate the spatial distribution of G proteins in plasma membrane microdomains containing GPCRs; therefore, the relative amounts of activated integrins ( $\beta 1$  or  $\beta 3$ ) at focal adhesions govern G protein subunit composition ( $G_{\alpha_i}$  versus  $G_{\alpha_s}$ ) coupled to the  $\mu$ -opioid receptor, thus regulating the signaling of this receptor in sensory neurons. The only GPCR that has been shown to interact directly with integrins, the P2Y<sub>2</sub> nucleotide receptor, contains an RGD motif in the first extracellular loop that enables it to interact with  $\alpha V\beta 3$  and  $\alpha V\beta 5$  integrins (Erb et al., 2001). These interactions between the P2Y<sub>2</sub> nucleotide receptor and  $\alpha V$  integrins are necessary for the receptor to activate G<sub>o</sub> and to initiate G<sub>o</sub>-mediated signaling events leading to chemotaxis (Bagchi et al., 2005) and also are critical for astrocyte migration (Wang et al., 2005). Our studies provide the first evidence of the interaction of the BK B<sub>2</sub> receptor with  $\alpha 5\beta 1$  integrin and demonstrate that this interaction leads to both transactivation of EGFR and ERK phosphorylation in cultured mIMCD-3 cells. In conclusion, these studies demonstrate a novel mechanism of EGFR transactivation by the G<sub>q</sub>-coupled BK B<sub>2</sub> receptor that involves formation of a functional complex between  $\alpha 5\beta 1$  integrin, EGFR, and BK B<sub>2</sub> receptor.

## References

- Bagchi S, Liao Z, Gonzalez FA, Chorna NE, Seye CI, Weisman GA, and Erb L (2005) The P2Y<sub>2</sub> nucleotide receptor interacts with  $\alpha V$  integrins to activate G<sub>o</sub> and induce cell migration. *J Biol Chem* **280**:39050–39057.
- Berg KA, Zardeneta G, Hargreaves KM, Clarke WP, and Milam SB (2007) Integrins regulate opioid receptor signaling in trigeminal ganglion neurons. *Neuroscience* **144**:889–897.
- Brooks PC, Strömblad S, Sanders LC, von Schalscha TL, Aimes RT, Stetler-Stevenson WG, Quigley JP, and Cheresch DA (1996) Localization of matrix metalloproteinase MMP-2 to the surface of invasive cells by interaction with integrin  $\alpha v\beta 3$ . *Cell* **85**:683–693.
- Brooks PC, Silletti S, von Schalscha TL, Friedlander M, and Cheresch DA (1998) Disruption of angiogenesis by PEX, a noncatalytic metalloproteinase fragment with integrin binding activity. *Cell* **92**:391–400.
- Cabodi S, Moro L, Bergatto E, Boeri Erba E, Di Stefano P, Turco E, Tarone G, and Defilippi P (2004) Integrin regulation of epidermal growth factor (EGF) receptor and of EGF-dependent responses. *Biochem Soc Trans* **32**:438–442.
- Conant K, St Hillaire C, Nagase H, Visse R, Gary D, Haughey N, Anderson C, Turchan J, and Nath A (2004) Matrix metalloproteinase 1 interacts with neuronal integrins and stimulates dephosphorylation of Akt. *J Biol Chem* **279**:8056–8062.
- Davidson L, Pawson AJ, Millar RP, and Maudsley S (2004) Cytoskeletal reorganization dependence of signaling by the gonadotropin-releasing hormone receptor. *J Biol Chem* **279**:1980–1993.
- Della Rocca GJ, Maudsley S, Daaka Y, Lefkowitz RJ, and Luttrell LM (1999) Pleiotropic coupling of G protein-coupled receptors to the mitogen-activated protein kinase cascade. Role of focal adhesions and receptor tyrosine kinases. *J Biol Chem* **274**:13978–13984.
- Dumin JA, Dickeson SK, Stricker TP, Bhattacharyya-Pakrasi M, Roby JD, Santoro SA, and Parks WC (2001) Pro-collagenase-1 (matrix metalloproteinase-1) binds the  $\alpha 2\beta 1$  integrin upon release from keratinocytes migrating on type I collagen. *J Biol Chem* **276**:29368–29374.
- Eisinger DA and Ammer H (2008)  $\delta$ -Opioid receptors activate ERK/MAP kinase via integrin-stimulated receptor tyrosine kinases. *Cell Signal* **20**:2324–2331.
- Erb L, Liu J, Ockerhausen J, Kong Q, Garrad RC, Griffin K, Neal C, Krugh B, Santiago-Pérez LI, González FA, et al. (2001) An RGD sequence in the P2Y(2) receptor interacts with  $\alpha(V)\beta(3)$  integrins and is required for G(o)-mediated signal transduction. *J Cell Biol* **153**:491–501.
- Golser R, Gorren AC, Leber A, Andrew P, Habisch HJ, Werner ER, Schmidt K, Venema RC, and Mayer B (2000) Interaction of endothelial and neuronal nitric-oxide synthases with the bradykinin B2 receptor. *J Biol Chem* **275**:5291–5296.
- Humphries JD, Byron A, and Humphries MJ (2006) Integrin ligands at a glance. *J Cell Sci* **119**:3901–3903.
- Hunger-Glaser I, Salazar EP, Sinnott-Smith J, and Rozenfurt E (2003) Bombesin, lysophosphatidic acid, and epidermal growth factor rapidly stimulate focal adhesion kinase phosphorylation at Ser-910: requirement for ERK activation. *J Biol Chem* **278**:22631–22643.
- Juliano RL (2002) Signal transduction by cell adhesion receptors and the cytoskeleton: functions of integrins, cadherins, selectins, and immunoglobulin-superfamily members. *Annu Rev Pharmacol Toxicol* **42**:283–323.
- Kramarenko II, Bunni MA, Morinelli TA, Raymond JR, and Garnovskaya MN. (2009) Identification of a functional bradykinin B<sub>2</sub> receptors endogenously expressed in HEK293 cells. *Biochem Pharmacol* **77**:269–276.
- Litvak V, Tian D, Shaull YD, and Lev S (2000) Targeting of PYK2 to focal adhesions as a cellular mechanism for convergence between integrins and G protein-coupled receptor signaling cascades. *J Biol Chem* **275**:32736–32746.
- Miranti CK and Brugge JS (2002) Sensing the environment: a historical perspective on integrin signal transduction. *Nat Cell Biol* **4**:E83–E90.
- Moro L, Venturino M, Bozzo C, Silengo L, Altruda F, Beguinot L, Tarone G, and Defilippi P (1998) Integrins induce activation of EGF receptor: role in MAP kinase induction and adhesion-dependent cell survival. *EMBO J* **17**:6622–6632.
- Mukai H, Fitzgibbon WR, Bozeman G, Margolius HS, and Ploth DW (1996) Bradykinin B<sub>2</sub> receptor antagonist increases chloride and water absorption in rat medullary collecting duct. *Am J Physiol* **271**:R352–R360.
- Mukhin YV, Garnovsky EA, Ullian ME, and Garnovskaya MN (2003) Bradykinin B<sub>2</sub> receptor activates extra-cellular signal regulated protein kinase in mIMCD-3 cells via epidermal growth factor receptor transactivation. *J Pharmacol Exp Ther* **304**:968–977.
- Mukhin YV, Gooz M, Raymond JR, and Garnovskaya MN (2006) Collagenases 2 and 3 mediate epidermal growth factor receptor transactivation by Bradykinin B<sub>2</sub> receptor in kidney cells. *J Pharmacol Exp Ther* **318**:1033–1043.
- Plow EF, Haas TA, Zhang L, Loftus J, and Smith JW (2000) Ligand binding to integrins. *J Biol Chem* **275**:21785–21788.
- Prenzel N, Zwick E, Daub H, Leserer M, Abraham R, Wallasch C, and Ullrich A (1999) EGF receptor transactivation by G-protein-coupled receptors requires metalloproteinase cleavage of proHB-EGF. *Nature* **402**:884–888.
- Rolli M, Fransvea E, Pilch J, Saven A, and Felding-Habermann B (2003) Activated integrin  $\alpha v\beta 3$  cooperates with metalloproteinase MMP-9 in regulating migration of metastatic breast cancer cells. *Proc Natl Acad Sci USA* **100**:9482–9487.
- Schwartz MA and Ginsberg MH (2002) Networks and crosstalk: integrin signalling spreads. *Nat Cell Biol* **4**:E65–E68.
- Short SM, Boyer JL, and Juliano RL (2000) Integrins regulate the linkage between upstream and downstream events in G protein-coupled receptor signaling to mitogen-activated protein kinase. *J Biol Chem* **275**:12970–12977.
- Slack BE (1998) Tyrosine phosphorylation of paxillin and focal adhesion kinase by activation of muscarinic m3 receptors is dependent on integrin engagement by the extracellular matrix. *Proc Natl Acad Sci USA* **95**:7281–7286.
- Stefanidakis M, Björklund M, Ihanus E, Gahmberg CG, and Koivunen E (2003) Identification of a negatively charged peptide motif within the catalytic domain of progelatinases that mediates binding to leukocyte  $\beta 2$  integrins. *J Biol Chem* **278**:34674–34684.
- Stefanidakis M and Koivunen E (2006) Cell-surface association between matrix metalloproteinases and integrins: role of the complexes in leukocyte migration and cancer progression. *Blood* **108**:1441–1450.
- Stricker TP, Dumin JA, Dickeson SK, Chung L, Nagase H, Parks WC, and Santoro SA (2001) Structural analysis of the  $\alpha 2$  integrin I domain/procollagenase-1 (matrix metalloproteinase-1) interaction. *J Biol Chem* **276**:29375–29381.
- Tomita K, Pisano JJ, and Knepper MA (1985) Control of sodium and potassium transport in the cortical collecting duct of the rat. Effects of bradykinin, vasopressin, and deoxycorticosterone. *J Clin Invest* **76**:132–136.
- Wang M, Kong Q, Gonzalez FA, Sun G, Erb L, Seye C, and Weisman GA (2005) P2Y2 nucleotide receptor interaction with  $\alpha V$  integrin mediates astrocyte migration. *J Neurochem* **95**:630–640.
- Xie P, Browning DD, Hay N, Mackman N, and Ye RD (2000) Activation of NF- $\kappa$ B by Bradykinin through a G $\alpha_q$ - and G $\beta\gamma$ -dependent pathway that involves Phosphoinositide 3-Kinase and Akt. *J Biol Chem* **275**:24907–24914.
- Zeidel ML, Jabs K, Kikeri D, and Silva P (1990) Kinins inhibit conductive Na<sup>+</sup> uptake by rabbit inner medullary collecting duct cells. *Am J Physiol* **258**:F1584–F1591.

---

**Address correspondence to:** Dr. Maria N. Garnovskaya, Medical University of South Carolina, Department of Medicine, Division of Nephrology, 96 Jonathan Lucas St., MSC 629, Charleston, SC 29425-6290. E-mail: garnovsk@musc.edu

---