

Published in final edited form as:

J Am Chem Soc. 2010 July 21; 132(28): 9567–9569. doi:10.1021/ja1035495.


Total Synthesis of (±)-Aplykurodinone-1: Traceless Stereochemical Guidance

Yandong Zhang^a and Samuel J. Danishefsky^{a,b,*}

^aDepartment of Chemistry, Columbia University, Havemeyer Hall, 3000 Broadway, New York, NY 10027, USA

^bLaboratory for Bioorganic Chemistry, Sloan-Kettering Institute for Cancer Research, 1275 York Avenue, New York, NY 10065, USA

Abstract



The total synthesis of the highly degraded steroidal natural product, aplykurodinone-1 (**1**), has been accomplished. Key features include a one-flask hydrolysis/retro-aldol/iodolactonization sequence to excise the C₈ hydroxymethylene functionality with retention of stereochemistry, and the stereoselective installation of the C₁₃ methyl group through hydrogenation with homogeneous catalyst.

Of all of the classes of natural products, steroids have perhaps had the most enduring role in prompting new ideas in total synthesis.¹ A not insignificant body of organic chemistry has been discovered in the context of pursuits directed to the total synthesis of steroids. Apparently, this interest continues to the present day.²

One of the vexing problems in steroid total synthesis is that of exercising control of the configuration at C₂₀.³ The challenge is that of correlating the configuration of the presumably “freely rotating” C₂₀, with the resident stereochemistry of the polycyclic domain. Though ingenious solutions to the “C₂₀ problem” have been advanced, it is still an interesting and thought provoking matter.

It was in this setting that we took particular note of the aplykurodines, isolated from marine mollusks of the genus *Aplysia*.⁴ This family can well be viewed as a rare class of highly degraded marine steroids, where C₁₃ in the aplykurodine setting corresponds to C₂₀ in wild-type steroids. Also of note is the *cis* fusion of the hydrindane sector in contrast to the *trans* C-D fusion of hydrindanes, which is so prevalent in fully mature steroids.⁵

We note that various aplykurodines exhibit cytotoxic activity against a range of human cancer cell lines.⁶ Notwithstanding the potential elements of chemistry and medicinal chemistry-based points of interest, surprisingly little attention at the level of synthesis has

s-danishefsky@ski.mskcc.org.

Supporting Information Available Experimental procedures, copies of spectral data, and characterization data are available free of charge via the Internet at <http://pubs.acs.org>.

been directed to the aplykurodines. Our laboratory took particular note of a recently isolated member of the family, aplykurodinone-1 (**1**).⁷ Following its isolation from the sea hare *Symphonota geographica* in 2005, the structure of aplykurodinone-1 (**1**) was elucidated through a combination of spectroscopic methods, X-ray crystallography, and chemical correlation. As shown in Scheme 1, **1** was found to possess a *cis*-fused C-D ring with epimeric C₈ (steroidal numbering) and an unsaturated side chain (as compared to cholesterol).

On the basis of considerations advanced above, a total synthesis effort directed to this “degraded” steroid was undertaken. Below is described the inaugural total synthesis of (±)-**1**, resulting from this initiative. As will be seen, the program potentially encompassed several chemistry-level initiatives of broader heuristic value.

Using ideas stemming from analysis by pattern recognition,⁸ we envisioned that, in principle, the hypothetical hydrindanone **4** contains an array of coordinated functionality, which could well enable access to the eventual target system, **1**. The plan envisioned halolactonization as the instrumentality for fusing the γ -lactone to the *cis*-fused hydrindanone, thereby creating a ladder-like topography. Accordingly, we needed to gain secure control over the configurational relationship between strategic carbons 3 and 7 in **1** (vide infra). It occurred to us that an otherwise unnecessary quaternary function at C₈ could orchestrate the required relationship of the angular methyl group at C₇ with a soon-to-be-installed acetic acid moiety at C₃ (see structures **5** and **6**). Iodolactonization leading to the γ -lactone could then follow in the wake of excision of the now traceless C₁ fragment.

Given the general line of thinking adumbrated above, and a long-term proclivity for Diels-Alder reactions, we hoped to advance along such lines. In the event, it was discouraging to find that attempted cycloaddition between **29** and **310** failed under various thermal and Lewis acid-mediated conditions. However, reaction of **2** with methyllithium generated an operational lithium enolate,¹¹ which did undergo effective cycloaddition with **3**, thereby providing a 73% yield of **4**. It is well recognized that realization of the required cycloaddition does not, in and of itself, inform as to the limiting, mechanistic, extremi (two sequential Michael additions or an anionically mediated Diels-Alder reaction).¹²

The enone was regioselectively masked as a thioketal.¹³ The C₉ ketone was then reduced, as shown, and the resultant alcohol protected as a MOM ether. Upon reduction of the ester and removal of the dithiane ketal,^{13a}¹⁴ compound **10** was in hand. The primary alcohol was next converted to its diazoacetyl derivative, **11**. As shown in Scheme 2, intramolecular cyclopropanation onto the electron-deficient enone afforded the activated cyclopropane, **5**, though only in moderate yield (40%).¹⁵

The next phase entailed the opening of the cyclopropane ring, with concomitant installation of unsaturation at C₄–C₅. Toward this end, the C₅ ketone was converted to an iodide through a two-step sequence, commencing with Luche reduction,¹⁶ which provided a 2:1 (β : α) epimeric mixture of alcohols.¹⁷ These compounds were readily converted to iodide **12**. In the event, upon exposure of **12** to SmI₂,¹⁸ it readily suffered the hoped-for reductive cyclopropane opening, to deliver lactone **13** in good yield (85%). The latter was converted to **6** in a straightforward fashion, as shown.

The otherwise extraneous C₈ oxymethylene functionality had indeed been exploited as a very useful C₃ stereochemical linchpin (vide supra). Having served its function in mediating the stereoselective installation of the C₃ center, the C₁ fragment was cleaved via a hydrolysis–retro-aldol sequence following exposure of **6** to the action of K₂CO₃ in water at 100 °C. Subsequent iodolactonization¹⁹ of the resultant carboxylic acid (**14**) was accomplished in the same flask, to provide **15** in 75% yield from **6**. The re-emergence of the

cis-junction was confirmed through NOESY analysis of the methyl ester derivative of **14** and by X-ray diffraction of the subsequent deiodinated intermediate, **16**.²⁰ Finally, Saegusa oxidation²¹ of **16** afforded **7**, presenting the tricyclic core of aplykurodinone-1, in the sense shown.

With compound **7** in hand, the synthesis entered its terminal phase, i.e. installation of the C₁₁ side chain. A stepwise strategy was set in place. As outlined in Scheme 4, BF₃-mediated conjugate addition of the cuprate derived from vinyl bromide **1722** was accomplished with a high degree of facial selectivity (10:1 dr) to provide intermediate **18**. Given the difficulties surrounding the *heterogeneous* hydrogenation of the C₂₀–C₂₂ olefin as a means of installing the C₂₀ stereocenter of *trans*-fused mature steroids,²³ it was hoped that *homogeneous* hydrogenation protocols may be more productive.²⁴ Thus, following exposure to Crabtree catalyst,²⁵ under an atmospheric pressure of H₂, the tri-substituted olefin was reduced in a diastereoselective fashion (>5:1) to furnish compound **19**, which bears all of the stereocenters of aplykurodinone-1 (**1**). A single crystal of this advanced intermediate was obtained, and X-ray crystallographic analysis was found to support the proposed structure. Elaboration of the side chain was accomplished in a straightforward fashion, as shown. Thus, cleavage of the TBS group²⁶ and subsequent oxidation²⁷ delivered aldehyde **20**, which readily underwent modified Julia olefination²⁸ to furnish the target compound, aplykurodinone-1 (**1**). Its spectroscopic properties were the same as those reported for the isolated natural product. In the course of these synthetic studies, we also investigated a different route for the emplacement of the C₁₁ side chain. The fully elaborated dienyI fragment could be appended to the core (**7**) in a single step. It was hoped that the intermediate thus obtained could then be converted, in one step, to aplykurodinone-1 (**1**), through regio- and stereo-selective hydrogenation. In the event, the conjugate addition of **2129** to **7** was accomplished, to provide **22** in moderate yield. Upon exposure to Wilkinson's catalyst³⁰ under atmospheric H₂, the 13-*epi*-aplykurodinone-1 isomer (**23**) was selectively formed (>6:1 dr). Perhaps in each of the two hydrogenation transition states (**18**→**19**; **22**→**23**), the double bond projects from the β-face of the molecular backbone. The bulky catalyst then approaches the olefin from the unhindered of the olefin, to provide the observed stereochemical outcome.³¹

In summary, the total synthesis of aplykurodinone-1 (**1**) has been accomplished. A key feature of the effort involved an anionically-mediated cycloaddition of a metallo-enolate derived from **2** with **3** (see formation of **4**). The seemingly extraneous oxymethyl C₁ function at C₈, used to govern the configurational relationship between C₃ and C₇. The C₈ function is ultimately excised. The stereochemistry at C₁₃ is managed with good stereoselectivity based on suitable order of introduction of H and Me to this carbon.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.


Acknowledgments

Support was provided by the NIH (HL25848 and CA103823 to S.J.D.). Aaron Sattler, Wesley Sattler from the Parkin group (Columbia University) are thanked for their help with X-ray diffraction experiments (CHE-0619638 from the NSF). We thank Prof. M. Carbone (Istituto di Chimica Biomolecolare, CNR, Italy) for providing NMR spectra of the aplykurodinones. Valuable discussion with Ms. Rebecca Wilson throughout the course of this project is gratefully acknowledged.


References

1. (a) Kametani T, Nemoto H. *Tetrahedron*. 1981; 37:3–16. (b) Skoda-Foldes R, Kollar L. *Chem. Rev.* 2003; 103:4095–4129. [PubMed: 14531719] (c) Chapelon A-S, Moraleda D, Rodriguez R, Ollivier C, Santelli M. *Tetrahedron*. 2007; 63:11511–11616. (d) Ibrahim-Ouali M. *Steroids*. 2009; 74:133–162. [PubMed: 19022272]
2. (a) Yueng Y-Y, Chein R-J, Corey EJ. *J. Am. Chem. Soc.* 2007; 129:10346–10347. [PubMed: 17672468] (b) Giroux S, Corey EJ. *J. Am. Chem. Soc.* 2007; 129:9866–9867. [PubMed: 17658816]
3. (a) Piatak DM, Wicha J. *Chem. Rev.* 1978; 78:199–241. (b) Redpath J, Zeelen FJ. *Chem. Soc. Rev.* 1983; 12:75–98.
4. (a) Miyamoto T, Higuchi R, Komori T. *Tetrahedron Lett.* 1986; 27:1153–1156. (b) Spinella A, Gavagnin M, Crispino A, Cimino G, Martinez E, Ortea J, Sodano G. *J. Nat. Prod.* 1992; 55:989–993. [PubMed: 1402961] (c) Ortega MJ, Zubia E, Salva J. *J. Nat. Prod.* 1997; 60:488–489. [PubMed: 9170292]
5. Hill, RA.; Kirk, DN.; Makin, HLJ.; Murphy, GM. *Dictionary of Steroids, Chemical Data, Structures and Biographies*. London: Chapman and Hall; 1991. p. xiv-xxix.
6. (a) De Riccardis F, Spinella A, Izzo I, Giordano A, Sodano G. *Tetrahedron Lett.* 1995; 36:4303–4306. (b) Izzo I, Meduri G, Avallone E, De Riccardis F, Sodano G. *Eur. J. Org. Chem.* 2000:439–448.
7. Gavagnin M, Carbone M, Nappo M, Mollo E, Roussis V, Cimino G. *Tetrahedron*. 2005; 61:617–621.
8. For a discussion of the value of pattern analysis as a supplement to strategic bond disconnection in retrosynthetic analysis, see: Wilson RM, Danishefsky SJ. *J. Org. Chem.* 2007; 72:4293–4305. [PubMed: 17539594]
9. Danishefsky SJ. *Acc. Chem. Res.* 1981; 14:400–406.
10. Hua DH, Venkataraman S, King RCY, Paukstelis JV. *J. Am. Chem. Soc.* 1988; 110:4741–4748.
11. (a) Stork G, Hudrik PF. *J. Am. Chem. Soc.* 1968; 90:4464–4465. (b) Baker R, Selwood DL, Swain CJ, Webster NMH. *J. Chem. Soc., Perkin Trans 1*. 1988:471–480.
12. (a) Ihara M, Fukumoto K. *Angew. Chem. Int. Ed. Engl.* 1993; 32:1010–1022. (b) Kanemasa S, Kumegawa M, Wada E, Nomura M. *Bull. Chem. Soc. Jpn.* 1991; 64:2990–3004.
13. (a) Bosch MP, Camps F, Coll J, Guerrero A, Tatsuoka T, Meinwald J. *J. Org. Chem.* 1986; 51:773–784. (b) Golinski M, Brock CP, Watt DS. *J. Org. Chem.* 1993; 58:159–164.
14. Jones PS, Ley SV, Simpkins NS, Whittle AJ. *Tetrahedron*. 1986; 42:6519–6534.
15. Corey EJ, Myers AG. *Tetrahedron Lett.* 1984; 25:3559–3562. (b) An alternative approach to compound **12**, involving a high-yielding cyclopropanation of an electron rich olefin was also investigated. See Supporting Information for details.
16. Luche JL. *J. Am. Chem. Soc.* 1978; 100:2226–2227.
17. The stereochemistry of the 5- β -epimer was confirmed by X-ray analysis of its thiocarbonyl imidazole derivative. See Supporting Information. Barton DH, McCombie SW. *J. Chem. Soc., Perkin Trans. 1*. 1975:1574–1585.
18. Abad A, Agullo C, Cunat AC, Navarro I. *Tetrahedron Lett.* 2001; 42:8965–8968.
19. (a) Kitagawa O, Sato T, Taguchi T. *Chem. Lett.* 1991:177–180. (b) Devaux JF, O'Neil SV, Guillo N, Paquette LA. *Collect. Czech. Chem. Commun.* 2000; 65:490–510.
20. (a) Kamano Y, Pettit GR. *J. Org. Chem.* 1973; 38:2202–2204. (b) Piccinini P, Vidari G, Zanoni G. *J. Am. Chem. Soc.* 2004; 126:5088–5089. [PubMed: 15099090]
21. Ito Y, Hirao T, Saegusa T. *J. Org. Chem.* 1978; 43:1011–1013.
22. (a) Le Monez P, Fageas V, Poisson J, Ardisson J. *Tetrahedron Lett.* 1994; 35:7767–7770. (b) Tiefenbacher K, Arion VB, Mulzer J. *Angew. Chem. Int. Ed.* 2007; 46:2690–2693.
23. (a) Narwid TA, Cooney KE, Uskokovic MR. *Helv. Chim. Acta.* 1974; 57:771–781. [PubMed: 4364577] (b) Schow SR, McMorris TC. *J. Org. Chem.* 1979; 44:3760–3765. (c) Kametani T, Suzuki K, Nemoto H. *J. Am. Chem. Soc.* 1981; 103:2891–2892.
24. Harmon RE, Gupta SK, Brown DJ. *Chem. Rev.* 1973; 73:21–52.
25. Crabtree R. *Acc. Chem. Res.* 1979; 12:331–337.


26. When Bu_4NF was used, extensive decomposition of the starting material was observed.
27. Dess DB, Martin JC. *J. Org. Chem.* 1983; 48:4155–4156.
28. (a) Blakemore PR, Cole WJ, Kocienski PJ, Morley A. *Synlett.* 1998:26–28. (b) Marti C, Carreira EM. *J. Am. Chem. Soc.* 2005; 127:11505–11515. [PubMed: 16089481]
29. Mushti CS, Kim JH, Corey EJ. *J. Am. Chem. Soc.* 2006; 128:14050–14052. [PubMed: 17061886]
30. (a) Young JF, Osborn JA, Jardine FH, Wilkinson G. *Chem. Commun.* 1965:131–132. (b) O'Connor C, Wilkinson G. *J. Chem. Soc. (A).* 1968:2665–2671.
31. Proposed transition states of the homogeneous hydrogenation with Crabtree and Wilkinson catalyst are shown below. In each case, the bulky catalyst is believed to approach the double bond from its less hindered face. However, as described above, different stereochemical outcomes were observed in these two systems.



For mechanistic study of these two types of hydrogenation reaction, See: (a) Brown JM, Derome AE, Hughes GD, Monaghan PK. *Aust. J. Chem.* 1992; 45:143–153. (b) Nelson DJ, Li R, Brammer C. *J. Org. Chem.* 2005; 70:761–767. [PubMed: 15675830]




Scheme 1.
Synthetic Strategy Toward Aplykurodinone-1 (1).


**Scheme 2.**

Synthesis of **6** En Route to Aplykurodinone-1 (**1**).



^aKey: (a) MeLi, DME/THF, $-50\text{ }^{\circ}\text{C}$; TFA, CHCl_3 , reflux, 73%; (b) $\text{HSCH}_2\text{CH}_2\text{SH}$, AcOH, p-TSA, 74%; (c) NaBH_4 , $\text{CH}_2\text{Cl}_2/\text{MeOH}$, $-78\text{ }^{\circ}\text{C} \rightarrow -40\text{ }^{\circ}\text{C}$; (d) MOMCl, $i\text{Pr}_2\text{NEt}$, CH_2Cl_2 , 84% over 2 steps; (e) DIBAL-H, CH_2Cl_2 , $-78\text{ }^{\circ}\text{C}$; (f) $\text{Ti}(\text{NO}_3)_3$, $\text{MeOH}/\text{THF}/\text{H}_2\text{O}$, 95% over 2 steps; (g) $\text{TsNHN}=\text{CHCOCl}$, PhNMe_2 , Et_3N , CH_2Cl_2 , $0\text{ }^{\circ}\text{C}$, 88%; (h) bis-(*N*-*t*-butylsalicylaldiminato) copper (II), PhMe, reflux, 40%; (i) NaBH_4 , $\text{CeCl}_3 \cdot 7\text{H}_2\text{O}$; (j) I_2 , PPh_3 , imidazole, PhMe, reflux, 78% over 2 steps; (k) SmI_2 , THF/MeOH , 85%; (l) HCl, $\text{THF}/\text{H}_2\text{O}$; (m) DMP, CH_2Cl_2 , 80% over 2 steps.

**Scheme 3.**

^aKey: (a) K_2CO_3 , H_2O , $100\text{ }^\circ\text{C}$; (b) NIS, CH_2Cl_2 , 75%; (c) Ra-Ni, $CH_2Cl_2/EtOH$, 90%; (d) TMSOTf, Et_3N , CH_2Cl_2 , $0\text{ }^\circ\text{C}$; Pd(OAc)₂, CH_3CN , 76% (90% brsm).

**Scheme 4.**

^aKey: (a) **17**, *t*-BuLi, CuCN, BF₃•OEt₂, -78 °C → -98 °C, 73%, 10:1 dr; (b) Crabtree catalyst, H₂, CH₂Cl₂, 75% conversion, 50% yield (pure); (c) HF, CH₃CN/THF; (d) DMP, CH₂Cl₂, 87% over two steps; (e) THF, LiHMDS, -78 °C, 68%.

**Scheme 5.**

^aKey: (a) **17**, t-BuLi, CuCN, Et₂O, -78 °C, 51%; (b) Wilkinson catalyst, H₂, benzene, 67%.