



RESEARCH

Open Access

Two pore domain potassium channels in cerebral ischemia: a focus on $K_{2p}9.1$ (TASK3, KCNK9)

Petra Ehling¹, Stefan Bittner², Nicole Bobak³, Tobias Schwarz², Heinz Wiendl³, Thomas Budde¹, Christoph Kleinschnitz^{*†2} and Sven G Meuth^{*†2,3}

Abstract

Background: Recently, members of the two-pore domain potassium channel family (K_{2p} channels) could be shown to be involved in mechanisms contributing to neuronal damage after cerebral ischemia. $K_{2p}3.1^{-/-}$ animals showed larger infarct volumes and a worse functional outcome following experimentally induced ischemic stroke. Here, we question the role of the closely related K_{2p} channel $K_{2p}9.1$.

Methods: We combine electrophysiological recordings in brain-slice preparations of wildtype and $K_{2p}9.1^{-/-}$ mice with an in vivo model of cerebral ischemia (transient middle cerebral artery occlusion (tMCAO)) to depict a functional impact of $K_{2p}9.1$ in stroke formation.

Results: Patch-clamp recordings reveal that currents mediated through $K_{2p}9.1$ can be obtained in slice preparations of the dorsal lateral geniculate nucleus (dLGN) as a model of central nervous relay neurons. Current characteristics are indicative of $K_{2p}9.1$ as they display an increase upon removal of extracellular divalent cations, an outward rectification and a reversal potential close to the potassium equilibrium potential. Lowering extracellular pH values from 7.35 to 6.0 showed comparable current reductions in neurons from wildtype and $K_{2p}9.1^{-/-}$ mice ($68.31 \pm 9.80\%$ and $69.92 \pm 11.65\%$, respectively). These results could be translated in an in vivo model of cerebral ischemia where infarct volumes and functional outcomes showed a none significant tendency towards smaller infarct volumes in $K_{2p}9.1^{-/-}$ animals compared to wildtype mice 24 hours after 60 min of tMCAO induction ($60.50 \pm 17.31 \text{ mm}^3$ and $47.10 \pm 19.26 \text{ mm}^3$, respectively).

Conclusions: Together with findings from earlier studies on $K_{2p}2.1^{-/-}$ and $K_{2p}3.1^{-/-}$ mice, the results of the present study on $K_{2p}9.1^{-/-}$ mice indicate a differential contribution of K_{2p} channel subtypes to the diverse and complex in vivo effects in rodent models of cerebral ischemia.

Background

Although ischemic stroke represents a major health care problem with a high rate of permanent disability or even death, the underlying molecular mechanisms leading to neuronal death are still poorly understood [1]. However, ion channels which can influence basal cellular parameters are thought to play a major role within this context.

Activation of potassium channels results in membrane hyperpolarization thereby decreasing neuronal activity and cell death under pathophysiological conditions. Additionally, K^+ channels (e.g. large conductance Ca^{2+} -activated K^+ channels and ATP-sensitive K^+ channels [2,3]) might be neuroprotective as they counterbalance a prolonged harmful influx of Ca^{2+} ions via different pathways including a reversal of the Na^+/Ca^{2+} antiporter and voltage-dependent Ca^{2+} channels. Furthermore, an enhancement of the Mg^{2+} block of NMDA receptors (N-methyl D-aspartate) in postsynaptic neurons [4] is thought to protect against glutamate excitotoxicity [5,6].

* Correspondence: christoph.kleinschnitz@mail.uni-wuerzburg.de, sven.meuth@gmx.de

² University of Wuerzburg, Department of Neurology, Josef-Schneider Str. 11, 97080 Wuerzburg, Germany

³ University of Muenster, Neurological Clinic - Inflammatory Disorders of the Central Nervous System and Neurooncology, Mendelstr. 7, 48149 Muenster, Germany

† Contributed equally

Full list of author information is available at the end of the article



Concerning the recently identified family of two-pore domain potassium channels (K_{2p} channels), several members have been shown to play a major role in critical conditions leading to cerebral ischemia. $K_{2p2.1}^{-/-}$ mice displayed significantly less neuronal survival rates in a model of cerebral ischemia [7]. These data were confirmed by the neuroprotective effect of several $K_{2p2.1}$ channel activators (e.g. alpha linolenic acid or riluzole [8-10]). On the other hand, genetic depletion of another family member, namely $K_{2p3.1}$, resulted in increased infarct volumes following transient or permanent middle cerebral artery occlusion (MCAO) [11,12]. Based on sequence homologies and similar biophysical properties, it was suggested that related channel family members might also be of importance under these circumstances. We challenged the role of $K_{2p9.1}$ (TASK3; KCNK9) in a tMCAO model using previously described $K_{2p9.1}^{-/-}$ mice [13].

Methods

Slice preparation

Thalamic tissue slices including the dorsal lateral geniculate nucleus (dLGN) were prepared from 14 - 22 days old male C57BL/6 or $K_{2p9.1}^{-/-}$ mice [13] as described earlier [14]. Coronal sections were cut on a vibratome (Vibratome[®], Series 1000 Classic, St. Louis, USA) in an ice-chilled solution containing (mM): Sucrose, 200; PIPES, 20; KCl, 2.5; NaH_2PO_4 , 1.25; MgSO_4 , 10; CaCl_2 , 0.5; dextrose, 10; pH 7.35 adjusted with NaOH. Prior to the recording procedure, slices were kept submerged in artificial cerebrospinal fluid (ACSF, mM): NaCl, 125; KCl, 2.5; NaH_2PO_4 , 1.25; NaHCO_3 , 24; MgSO_4 , 2; CaCl_2 , 2; dextrose, 10; pH adjusted to 7.35 by bubbling with a mixture of 95% O_2 and 5% CO_2 .

Electrophysiology

Slices were transferred in a recording chamber and thalamic neurons of the dLGN were visualized with a microscope equipped with infrared-differential interference contrast optics [15]. Whole-cell recording pipettes were fabricated from borosilicate glass (GT150T-10, Clark Electromedical Instruments, Pangbourne, UK; typical resistance 2-3 M Ω) and filled with an intracellular solution containing (in mM): K-gluconate, 88; K_3 -citrate, 20; NaCl, 10; HEPES, 10; MgCl_2 , 1; CaCl_2 , 0.5; BAPTA, 3; phosphocreatin, 15; Mg-ATP, 3; Na-GTP, 0.5. The internal solution was set to a pH value of 7.25 using KOH and an osmolarity of 295 mOsm/kg. Slices were continuously superfused with a solution containing NaCl 125 mM, KCl 2.5 mM, NaH_2PO_4 1.25 mM, HEPES 30 mM, MgSO_4 2 mM, CaCl_2 2 mM and dextrose 10 mM. Whole-cell patch-clamp recordings were measured from relay neu-

rons of the dLGN with an EPC-10 amplifier (HEKA Elektronik, Lamprecht, Germany) and digitally analyzed using Pulse software (HEKA Elektronik; [16]). pH was adjusted to 7.35 or 6.0 with HCl. For divalent-cation-free conditions we switched from control solution to a solution containing 0 mM Mg^{2+} and 0 mM Ca^{2+} ; the osmolality was kept constant at 305 mosmol kg^{-1} by adding 4 mM NaCl [17]. All cells had a resting membrane potential negative to -65 mV, the access resistance was in the range of 5-15 M Ω and series resistance compensation of more than 40% was routinely used.

Induction of cerebral ischemia

Animal experiments were approved by governmental agencies for animal research and conducted according to the recommendations for research in mechanism-driven basic stroke studies [18]. Focal cerebral ischemia was induced in 6-8 weeks old male C57BL/6 and $K_{2p9.1}^{-/-}$ mice [13] weighing 20-25 g by transient middle cerebral artery occlusion (tMCAO) as described previously [19,20]. Briefly, mice were anesthetized with 2.5% isoflurane (Abbott, Wiesbaden, Germany) in a 70% $\text{N}_2\text{O}/30\%$ O_2 mixture. Core body temperature was maintained at 37°C throughout surgery using a feedback-controlled heating device. Following a midline skin incision in the neck, the proximal common carotid artery and the external carotid artery were ligated and a standardized silicon rubber-coated 6.0 nylon monofilament (6021; Doccol Corp., CA, USA) was inserted and advanced via the right internal carotid artery to occlude the origin of the right MCA. The intraluminal suture was left *in situ* for 1 hour, respectively. Then animals were re-anesthetized and the occluding monofilament was withdrawn to allow reperfusion. After 24 hours neurological deficits were scored by two blinded investigators and quantified according to Bederson [21]: 0, no deficit; 1, forelimb flexion; 2, as for 1, plus decreased resistance to lateral push; 3, unidirectional circling; 4, longitudinal spinning; 5, no movement. For the gript test, the mouse was placed midway on a string between two supports and rated as follows: 0, falls off; 1, hangs onto string by one or both forepaws; 2, as for 1, and attempts to climb onto string; 3, hangs onto string by one or both forepaws plus one or both hindpaws; 4, hangs onto string by fore- and hindpaws plus tail wrapped around string; 5, escape (to the supports).

Laser doppler flowmetry (Moor Instruments, Axminster, United Kingdom) was used to monitor cerebral blood flow [22] in wildtype, $K_{2p9.1}^{-/-}$ and sham-treated animals ($n = 4/\text{group}$) before surgery (baseline), immediately after MCA occlusion, and 5 minutes after removal of the occluding monofilament (reperfusion). Cerebral perfusion did not differ between the groups at any time point (Additional File 1, Figure S1).

Determination of infarct size

Mice were sacrificed 24 hours after tMCAO, respectively. Brains were quickly removed and cut in 2 mm thick coronal sections using a mouse brain slice matrix. The slices were stained with 2% 2,3,5-triphenyltetrazolium chloride (TTC; Sigma-Aldrich, St. Louis, MO) in PBS to visualize the infarctions. Planimetric measurements (ImageJ software, National Institutes of Health, Bethesda, MD) blinded to the treatment groups were used to calculate lesion volumes, which were corrected for brain edema as described [23,24].


Statistical analysis

Electrophysiological data and results from the animal experiments were analyzed by a modified student's t test for small samples [25] or by a Bonferroni-corrected One-way ANOVA in case of multiple comparisons using PrismGraph 4.0 software (GraphPad Software, San Diego, CA) or Origin® (Microcal). P-values < 0.05 were considered statistically significant.

Results

Thalamic relay neurons as a model system of central nervous system neurons display electrophysiological properties indicative of currents through $K_{2p9.1}$ channels

$K_{2p9.1}$ -like currents have been demonstrated in a number of different central nervous system neurons [14,26-28]. As highly specific inhibitors for K_{2p} channel subtypes are not available, different semi-selective blockers and experimental strategies to distinguish between these channels were established. Among them, extracellular reduction of divalent cations was introduced to increase potassium outward currents through $K_{2p9.1}$ channels [17]. Current-voltage relationships (I/V) of the standing outward current of wildtype and $K_{2p9.1}^{-/-}$ mice were investigated by ramping the membrane potential from -35 mV to -125 mV over 800 ms (Fig. 1A, inset; [29,30]). Under control conditions a standing outward current (I_{SO}) of 322.33 ± 30.20 pA was measured at -35 mV (Fig. 1A). Application of hyperpolarizing voltage ramps induced a complex current response. The wave form of this response was indicative for the contribution of current through outwardly rectifying TASK channels as well as inwardly rectifying K^+ channels (Fig. 1A, black trace). Removal of extracellular divalent cations resulted in a significant increase of I_{SO} by $35.47 \pm 9.59\%$ compared to control conditions ($n = 6$, $p = 0.007$; Fig. 1A). Ramp responses revealed a clear increase in the outwardly rectifying component (Fig. 1A, gray trace). The current sensitive to administering cation-free conditions was calculated by numerical subtraction of control currents from currents recorded under cation-free conditions [14]. The I/V relationship of the cation-sensitive current was typical of TASK channels with a strong outward rectification and a reversal potential close



to the expected potassium equilibrium potential (Fig. 1B; $E_K = -104$ mV). These findings indicate a strong contribution of $K_{2p9.1}$ channels to the I_{SO} of thalamocortical (TC) neurons in wildtype mice.

Neurons from $K_{2p9.1}^{-/-}$ and wildtype animals show no significant differences upon extracellular acidification

Sensitivity to extracellular acidification is a hallmark of TASK channels and a reduction of the extracellular pH

value can be typically observed under ischemic conditions. In a next experimental step we therefore mimicked cerebral ischemia by lowering the extracellular pH from control conditions (7.35) to 6.0. This maneuver resulted in a significant ($p < 0.05$) reduction of I_{SO} amplitudes in both genotypes (Fig. 2A). The degree of I_{SO} reduction was not different in wildtype ($68.31 \pm 9.80\%$) and $K_{2p9.1}^{-/-}$ neurons ($69.92 \pm 11.65\%$; $n = 5$; $p = 0.91$; Fig. 2B).

Genetic ablation of $K_{2p9.1}$ channels trends to result in a not significant reduction of stroke development after tMCAO

Stroke volumes of wildtype and $K_{2p9.1}^{-/-}$ mice were determined 24 hours after animals subjected to 60 min of tMCAO. Wildtype animals showed stroke volumes of $60.50 \pm 17.31 \text{ mm}^3$ while $K_{2p9.1}^{-/-}$ mice displayed infarct areas of $47.10 \pm 19.26 \text{ mm}^3$ ($n = 10$ and 8 ; $p = 0.23$; Fig. 3A and 3B). In accordance with this tendency towards none

significantly smaller infarct sizes in $K_{2p9.1}^{-/-}$, no functionally relevant differences could be found for the Bederson score (WT: 1.83 ± 0.98 ; $K_{2p9.1}^{-/-}$: 2.14 ± 0.80 ; $n = 6$; $p = 0.55$; Fig. 3C) and the grip test (WT: 3.17 ± 1.13 ; $K_{2p9.1}^{-/-}$: 4.29 ± 0.64 ; $n = 6$; $p = 0.09$; Fig. 3D).

Discussion

The results of the present study can be summarized as follows: (1) A pH- and divalent cation-sensitive I_{SO} is present in TC neurons of the dLGN. (2) The divalent cation-sensitive component is characterized by outward rectification and a reversal potential close to the potassium equilibrium potential. (3) The I_{SO} of neurons recorded from brain slices of $K_{2p9.1}^{-/-}$ mice and wildtype mice showed comparable pH-sensitivity during extracellular pH changes from 7.35 to 6.0. (4) In a model of cerebral ischemia, $K_{2p9.1}^{-/-}$ animals showed a tendency to reduced





Figure 2 pH dependence of the standing outward currents in $K_{2p9.1}^{-/-}$ neurons shows no difference compared to wildtype mice (pH 7.35 T pH 6.0). (A) Ramp currents under control conditions (pH 7.35) and after extracellular acidification (pH 6.0) in wildtype (WT; left panel) and $K_{2p9.1}^{-/-}$ mice (right panel). (B) Bar graph representation of current reduction after extracellular pH lowering in wildtype (WT) and $K_{2p9.1}^{-/-}$ mice.



infarct volumes 24 hours after undergoing 60 min of tMCAO compared to wildtype controls although these results were not statistically significant. (5) It is concluded that K_{2p}9.1-containing homodimeric and heterodimeric channels significantly contribute to I_{SO} in TC neurons from wildtype mice and that K_{2p}9.1 channels have only a minor impact on infarct volume and motor function following tMCAO compared to other members of the K_{2p} channel family.

Contribution of TASK channel subtypes to I_{SO} in TC neurons

During development, the mouse thalamus is characterized by high K_{2p}3.1 gene expression at P0 and displays moderate expression levels throughout postnatal stages [31]. K_{2p}9.1 expression in many thalamic nuclei is rather moderate for all developmental stages but is strong in dLGN from P14 to adult stages. Functional TASK chan-

nels can be K_{2p}3.1 homodimers, K_{2p}9.1 homodimers, and K_{2p}3.1/K_{2p}9.1 heterodimers [32-35]. Although K_{2p}3.1 and K_{2p}9.1 show high sequence homology, they differ in their sensitivity to extracellular divalent cations (Mg²⁺, Ca²⁺) based on the presence of a glutamate residue at position 70 in K_{2p}9.1 channels [17]. While the conductance of K_{2p}3.1 homodimeric channels is unaffected, the conductance of K_{2p}9.1 homodimeric and K_{2p}3.1/K_{2p}9.1 heterodimeric channels is strongly reduced in the presence of divalent cations [17,33]. Therefore the increase in I_{SO} following removal of extracellular divalent cations which was found in cells from different rodent strain (Long Evans rats, wildtype mice, K_{2p}3.1^{-/-} mice) point to the functional expression of K_{2p}9.1 homodimeric and K_{2p}3.1/K_{2p}9.1 heterodimeric channels in TC neurons.

Homodimeric and heterodimeric TASK channels also differ in their pH-sensitivity. While $K_{2p3.1}/K_{2p9.1}$ channel constructs have a pH-sensitivity (pK approximately 7.3) in the physiological range which is closer to that of $K_{2p3.1}$ channels (pK approximately 7.5) than $K_{2p9.1}$ channels (pK approximately 6.8) [34]. In the present study no significant difference was found for the decrease in I_{SO} amplitude when the pH was shifted to a value of 6.0. Therefore the pH- and divalent cation- sensitivities of native TASK-like currents in TC neurons is best represented by $K_{2p3.1}/K_{2p9.1}$ heterodimeric channels. However, additional modulators (isoflurane, Zn^{2+} , ruthenium red) have to be tested to get more indications for the ratio of homodimeric to heterodimeric TASK channels in these neurons.

The role of TASK channel subtypes in ischemic insults

It has been shown before that $K_{2p3.1}^{-/-}$ mice reveal larger tMCAO lesions in comparison to wildtype mice probably through a combination of direct neuronal effects and due to blood pressure/aldosterone effects [11,12]. Based on its physiological properties and expression pattern, it seemed reasonable to expect an at least similar phenotype of $K_{2p9.1}^{-/-}$ mice compared to $K_{2p3.1}^{-/-}$ mice. The reason for the unexpected results presented here remains unclear but may involve one or more of the following considerations: (1) The cell type-specific expression and the exact conditions of the cellular environment of TASK channels have to be taken into account [36]. (2) Compensatory mechanisms, e.g. upregulation of other K_{2p} channel family members, differences in oxygen sensitivity or yet unknown K_{2p} channel properties may play a role. (2) In GABAergic interneurons of the entorhinal cortex membrane depolarization mediated by inhibition of $K_{2p9.1}$ channels induce an increase in action potential firing [37]. In consequence, an increase in the release of GABA by interneurons results in a decrease in pyramidal cell activity thereby limiting the injurious effects of ischemia. Assuming that this type of network interaction is found in brain regions affected by tMCAO, the neuroprotection by $K_{2p9.1}^{-/-}$ channels is missing in knock out mice. (4) Gender differences should be taken into account [12]. (5) It should also be kept in mind that ischemic conditions may also affect a variety of other target structures including several ion channel, e.g. TRPV1 or ASICs as well as NMDA receptors [36].

To unravel the complex scenario of cerebral ischemia and to define the exact functional contribution of a particular K_{2p} channel family member, further pharmacological and genetic tools are warranted, e.g. cell-type specific or conditional $K_{2p3.1}^{-/-}$, $K_{2p9.1}^{-/-}$ or $K_{2p10.1}^{-/-}$ mice. Especially the development of highly specific channel inhibi-

tors or activators might open up the opportunity to proceed these research efforts.

Taken together, results from $K_{2p2.1}^{-/-}$ (enhancement of ischemic damage [7]), $K_{2p3.1}^{-/-}$ (increase in infarct volumes [11,12]) and $K_{2p9.1}^{-/-}$ (no significant change ([12] or a tendency towards none significant reduced infarct volumes: this work)) mice underline the fact that there are differential effects of different K_{2p} channel subtypes on cerebral ischemia, not allowing to reason a uniform influence of this intriguing channel family on stroke formation.

Additional material

Additional file 1 Figure S1 - rCBF does not differ between wildtype mice and $K_{2p9.1}^{-/-}$ mice. Determination of regional cerebral blood flow (rCBF) using Laser Doppler flowmetry before the occlusion of the middle cerebral artery (baseline), 10 min after the occlusion (ischemia) and again 10 min after the removal of the filament (reperfusion) in wildtype mice and $K_{2p9.1}^{-/-}$ mice (n = 3/group). No significant differences in rCBF were observed between the two groups. One-way ANOVA, Bonferroni post hoc test.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

All authors have read and approved the final manuscript. PE, SB, NB and TB performed and analyzed the electrophysiological recordings. CK and TS operated the animals, assessed the functional scores and interpreted the data. HW, CK, TB, SB and SGM conceived the experiments, analyzed data, funded the project and wrote the manuscript.

Acknowledgements

We are grateful to Ms. Melanie Glaser for excellent technical assistance. We are grateful to Douglas A. Bayliss for providing $K_{2p9.1}^{-/-}$ animals. This work was supported by Interdisziplinäres Zentrum für klinische Forschung (IZKF A-54-1) to SGM and HW, the DFG (SFB 581, TP: A10) to SGM and DFG K2P-Forscherguppe 1086 (BU1019/9-1 to TB and ME3283/1-1 to SGM). PE was a member of the Otto-Creutzfeldt-Center for Cognitive and Behavioral Neuroscience Muenster (OCC).

Author Details

¹Westfaelische Wilhelms-University Muenster, Institute of Physiology I, Robert-Koch Str. 27a, 48149 Muenster, Germany, ²University of Wuerzburg, Department of Neurology, Josef-Schneider Str. 11, 97080 Wuerzburg, Germany and ³University of Muenster, Neurological Clinic - Inflammatory Disorders of the Central Nervous System and Neurooncology, Mendelstr. 7, 48149 Muenster, Germany

Received: 29 April 2010 Accepted: 20 July 2010

Published: 20 July 2010

References

1. Dirnagl U, Iadecola C, Moskowitz MA: **Pathobiology of ischaemic stroke: an integrated view.** *Trends Neurosci* 1999, **22**:391-397.
2. Heurteaux C, Bertaina V, Widmann C, Lazdunski M: **K⁺ channel openers prevent global ischemia-induced expression of c-fos, c-jun, heat shock protein, and amyloid beta-protein precursor genes and neuronal death in rat hippocampus.** *Proc Natl Acad Sci USA* 1993, **90**:9431-9435.
3. Lauritzen I, De Weille JR, Lazdunski M: **The potassium channel opener (-)-cromakalim prevents glutamate-induced cell death in hippocampal neurons.** *J Neurochem* 1997, **69**:1570-1579.

4. Mayer ML, Westbrook GL, Guthrie PB: **Voltage-dependent block by Mg²⁺ of NMDA responses in spinal cord neurons.** *Nature* 1984, **309**:261-263.
5. Honore E: **The neuronal background K_{2P} channels: focus on TREK1.** *Nat Rev Neurosci* 2007, **8**:251-261.
6. Franks NP, Honore E: **The TREK K_{2P} channels and their role in general anaesthesia and neuroprotection.** *Trends Pharmacol Sci* 2004, **25**:601-608.
7. Heurteaux C, Guy N, Laigle C, Blondeau N, Duprat F, Mazzuca M, Lang-Lazdunski L, Widmann C, Zanzouri M, Romey G, Lazdunski M: **TREK-1, a K⁺ channel involved in neuroprotection and general anesthesia.** *EMBO J* 2004, **23**:2684-2695.
8. Lauritzen I, Blondeau N, Heurteaux C, Widmann C, Romey G, Lazdunski M: **Polyunsaturated fatty acids are potent neuroprotectors.** *EMBO J* 2000, **19**:1784-1793.
9. Heurteaux C, Laigle C, Blondeau N, Jarretou G, Lazdunski M: **Alpha-linolenic acid and riluzole treatment confer cerebral protection and improve survival after focal brain ischemia.** *Neuroscience* 2006, **137**:241-251.
10. Duprat F, Lesage F, Patel AJ, Fink M, Romey G, Lazdunski M: **The neuroprotective agent riluzole activates the two P domain K(+) channels TREK-1 and TRAAK.** *Mol Pharmacol* 2000, **57**:906-912.
11. Meuth SG, Kleinschnitz C, Broicher T, Austinat M, Braeuningner S, Bittner S, Fischer S, Bayliss DA, Budde T, Stoll G, Wiendl H: **The neuroprotective impact of the leak potassium channel TASK1 on stroke development in mice.** *Neurobiol Dis* 2009, **33**:1-11.
12. Muhammad S, Aller MI, Maser-Gluth C, Schwaninger M, Wisden W: **Expression of the kcnk3 potassium channel gene lessens the injury from cerebral ischemia, most likely by a general influence on blood pressure.** *Neuroscience* 2007, **147**:758-764.
13. Mulkey DK, Talley EM, Stornetta RL, Siegel AR, West GH, Chen X, Sen N, Mistry AM, Guyenet PG, Bayliss DA: **TASK channels determine pH sensitivity in select respiratory neurons but do not contribute to central respiratory chemosensitivity.** *J Neurosci* 2007, **27**:14049-14058.
14. Meuth SG, Budde T, Kanyshkova T, Broicher T, Munsch T, Pape HC: **Contribution of TWIK-related acid-sensitive K⁺ channel 1 (TASK1) and TASK3 channels to the control of activity modes in thalamocortical neurons.** *J Neurosci* 2003, **23**:6460-6469.
15. Dodt HU, Ziegglansberger W: **Visualizing unstained neurons in living brain slices by infrared DIC-videomicroscopy.** *Brain Res* 1990, **537**:333-336.
16. Meuth SG, Kanyshkova T, Landgraf P, Pape HC, Budde T: **Influence of Ca²⁺-binding proteins and the cytoskeleton on Ca²⁺-dependent inactivation of high-voltage activated Ca²⁺ currents in thalamocortical relay neurons.** *Pflügers Arch* 2005, **450**:111-122.
17. Musset B, Meuth SG, Liu GX, Derst C, Wegner S, Pape HC, Budde T, Preisig-Muller R, Daut J: **Effects of divalent cations and spermine on the K⁺ channel TASK-3 and on the outward current in thalamic neurons.** *J Physiol* 2006, **572**:639-657.
18. Dirnagl U: **Bench to bedside: the quest for quality in experimental stroke research.** *J Cereb Blood Flow Metab* 2006, **26**:1465-1478.
19. Kleinschnitz C, Pozgajova M, Pham M, Bendszus M, Nieswandt B, Stoll G: **Targeting platelets in acute experimental stroke: impact of glycoprotein Ib, VI, and IIb/IIIa blockade on infarct size, functional outcome, and intracranial bleeding.** *Circulation* 2007, **115**:2323-2330.
20. Kleinschnitz C, Schwab N, Kraft P, Hagedorn I, Dreykluft A, Schwarz T, Austinat M, Nieswandt B, Wiendl H, Stoll G: **Early detrimental T cell effects in experimental cerebral ischemia are neither related to adaptive immunity nor thrombus formation.** *Blood* 2010, **6**:3835-42.
21. Bederson JB, Pitts LH, Tsuji M, Nishimura MC, Davis RL, Bartkowski H: **Rat middle cerebral artery occlusion: evaluation of the model and development of a neurologic examination.** *Stroke* 1986, **17**:472-476.
22. Connolly ES, Winfree CJ, Springer TA, Naka Y, Liao H, Yan SD, Stern DM, Solomon RA, Gutierrez-Ramos JC, Pinsky DJ: **Cerebral protection in homozygous null ICAM-1 mice after middle cerebral artery occlusion. Role of neutrophil adhesion in the pathogenesis of stroke.** *J Clin Invest* 1996, **97**:209-216.
23. Swanson RA, Morton MT, Tsao-Wu G, Savalos RA, Davidson C, Sharp FR: **A semiautomated method for measuring brain infarct volume.** *J Cereb Blood Flow Metab* 1990, **10**:290-293.
24. Ginsberg MD, Becker DA, Busto R, Belayev A, Zhang Y, Khoutorova L, Ley JJ, Zhao W, Belayev L: **Stilbazulenylnitron, a novel antioxidant, is highly neuroprotective in focal ischemia.** *Ann Neurol* 2003, **54**:330-342.
25. Dixon W, Massey F: **Introduction to statistical analysis.** McGraw-Hill, New York; 1969.
26. Bean BP: **The action potential in mammalian central neurons.** *Nat Rev Neurosci* 2007, **8**:451-465.
27. Gonzalez JA, Jensen LT, Doyle SE, Miranda-Anaya M, Menaker M, Fugger L, Bayliss DA, Burdakov D: **Deletion of TASK1 and TASK3 channels disrupts intrinsic excitability but does not abolish glucose or pH responses of orexin/hypocretin neurons.** *Eur J Neurosci* 2009, **30**:57-64.
28. Brickley SG, Aller MI, Sandu C, Veale EL, Alder FG, Sambhi H, Mathie A, Wisden W: **TASK-3 two-pore domain potassium channels enable sustained high-frequency firing in cerebellar granule neurons.** *J Neurosci* 2007, **27**:9329-9340.
29. Watkins CS, Mathie A: **A non-inactivating K⁺ current sensitive to muscarinic receptor activation in rat cultured cerebellar granule neurons.** *J Physiol* 1996, **491**(Pt 2):401-412.
30. Millar JA, Barratt L, Southan AP, Page KM, Fyffe RE, Robertson B, Mathie A: **A functional role for the two-pore domain potassium channel TASK-1 in cerebellar granule neurons.** *Proc Natl Acad Sci USA* 2000, **97**:3614-3618.
31. Aller MI, Wisden W: **Changes in expression of some two-pore domain potassium channel genes (KCNK) in selected brain regions of developing mice.** *Neuroscience* 2008, **151**:1154-1172.
32. Czirjak G, Enyedi P: **Formation of functional heterodimers between the TASK-1 and TASK-3 two-pore domain potassium channel subunits.** *J Biol Chem* 2002, **277**:5426-5432.
33. Kim D, Cavanaugh EJ, Kim I, Carroll JL: **Heteromeric TASK-1/TASK-3 is the major oxygen-sensitive background K⁺ channel in rat carotid body glomus cells.** *J Physiol* 2009, **587**:2963-2975.
34. Berg AP, Talley EM, Manger JP, Bayliss DA: **Motoneurons express heteromeric TWIK-related acid-sensitive K⁺ (TASK) channels containing TASK-1 (KCNK3) and TASK-3 (KCNK9) subunits.** *J Neurosci* 2004, **24**:6693-6702.
35. Aller MI, Veale EL, Linden AM, Sandu C, Schwaninger M, Evans LJ, Korpi ER, Mathie A, Wisden W, Brickley SG: **Modifying the subunit composition of TASK channels alters the modulation of a leak conductance in cerebellar granule neurons.** *J Neurosci* 2005, **25**:11455-11467.
36. Bittner S, Budde T, Wiendl H, Meuth SG: **From the background to the spotlight: TASK channels in pathological conditions.** *Brain Pathology* in press.
37. Deng PY, Lei S: **Serotonin increases GABA release in rat entorhinal cortex by inhibiting interneuron TASK-3 K⁺ channels.** *Mol Cell Neurosci* 2008, **39**:273-284.

doi: 10.1186/2040-7378-2-14

Cite this article as: Ehling et al., Two pore domain potassium channels in cerebral ischemia: a focus on K_{2P}.1 (TASK3, KCNK9) *Experimental & Translational Stroke Medicine* 2010, **2**:14

Submit your next manuscript to BioMed Central and take full advantage of:

- [Convenient online submission](#)
- [Thorough peer review](#)
- [No space constraints or color figure charges](#)
- [Immediate publication on acceptance](#)
- [Inclusion in PubMed, CAS, Scopus and Google Scholar](#)
- [Research which is freely available for redistribution](#)

Submit your manuscript at
www.biomedcentral.com/submit

