

Published in final edited form as:

Nature. 1993 January 28; 361(6410): 362–365. doi:10.1038/361362a0.

Pancreatic beta-cells are rendered glucose-competent by the insulinotropic hormone glucagon-like peptide-1(7-37)

George G. Holz IV^{*}, Willem M. Kühtreiber[†], and Joel F. Habener^{*}

^{*} Laboratory of Molecular Endocrinology, Massachusetts General Hospital, Howard Hughes Medical Institute, Harvard Medical School, Boston, Massachusetts 02114, USA

Abstract

Non-insulin-dependent diabetes mellitus (NIDDM, type 2 diabetes) is a disorder of glucose homeostasis characterized by hyperglycaemia, peripheral insulin resistance, impaired hepatic glucose metabolism, and diminished glucose-dependent secretion of insulin from pancreatic β -cells¹. Glucagon-like-peptide-1(7-37) (GLP-1)² is an intestinally derived hormone that may be useful for the treatment of NIDDM because it acts *in vivo* to increase the level of circulating insulin, and thus lower the concentration of blood glucose^{3,4}. This therapeutic effect may result from the ability of GLP-1 to compensate for a defect in the glucose signalling pathway that regulates insulin secretion from β -cells. In support of this concept we report here that GLP-1 confers glucose sensitivity to glucose-resistant β -cells, a phenomenon we term glucose competence. Induction of glucose competence by GLP-1 results from its synergistic interaction with glucose to inhibit metabolically regulated potassium channels that are also targeted for inhibition by sulphonylurea drugs commonly used in the treatment of NIDDM⁵. Glucose competence allows membrane depolarization, the generation of action potentials, and Ca²⁺ influx, events that are known to trigger insulin secretion^{6,7}.

GLP-1 exerts diverse insulinotropic actions on β -cells that include stimulation of cyclic AMP formation⁸, insulin secretion⁹, insulin biosynthesis and proinsulin gene expression¹⁰. Because all insulinotropic actions of GLP-1 are dependent on simultaneous exposure of β -cells to glucose^{10,11}, GLP-1 may act as a hormonal regulator, or modulator, of the β -cell glucose signalling system. To investigate this possibility, perforated-patch¹²⁻¹⁴ and cell-attached-patch¹⁵ recordings were obtained from solitary β -cells isolated from dispersed rat islets of Langerhans and maintained in short-term primary cell culture¹⁶. For unknown reasons, these single, isolated β -cells exhibit reduced sensitivity to glucose as measured by glucose-induced insulin secretion¹⁷, electrical activity¹⁸ or calcium signalling¹⁹. When perforated-patch recordings were obtained from such cells under conditions in which the bathing solution contained no glucose, the depolarizing responses to 10 mM glucose were often ≤ 10 –15 mV in amplitude, and glucose did not always initiate repetitive spiking activity as is frequently observed in intact islets⁶ (Fig. 1a trace 1). In addition, in the absence of glucose very little change in membrane potential was recorded in response to 10 nM GLP-1 (Fig. 1a left of trace 2). In marked contrast, a 30–35 mV depolarization accompanied by repetitive spiking activity was often observed when glucose and GLP-1 were applied simultaneously (Fig. 1a right of trace 2). This synergistic interaction indicates that GLP-1 acts in a manner analogous to that of a modulatory transmitter. It enhances the responsiveness of β -cells to glucose, yet is without effect in the absence of glucose.

The interaction of GLP-1 and glucose is remarkable in that the timing of the application of the two substances need not be simultaneous. Glucose-insensitive β -cells were rendered glucose-competent (capable of responding to glucose) by pretreatment with GLP-1 (Fig. 1b trace 1). Conversely, GLP-1-insensitive cells were rendered GLP-1-sensitive by prior application of glucose (Fig. 1b trace 2). These 'priming' effects associated with GLP-1 or glucose pretreatment may reflect either the prolonged action of cytosolic second messengers and/or slow dissociation of GLP-1 from its receptor.

Synergism between glucose and GLP-1 was dose-dependent (effector concentration for half maximum response, EC_{50} , 1 nM), and exhibited pharmacological specificity because deletion of a single amino acid at the amino terminus to yield GLP-1(8-37)²¹ abrogated activity (Fig. 1c, left). For these reasons, the action of GLP-1 is mediated by a receptor with pharmacological properties similar to that which mediates GLP-1-induced stimulation of insulin secretion from insulinoma cell lines^{10,22,23}. That the action of GLP-1 is in fact a receptor-mediated process is supported by a recent report of the complementary DNA cloning of a β -cell GLP-1 receptor that binds GLP-1 (but not glucagon) with high affinity and which stimulates cAMP production²⁴.

The induction of glucose competence by GLP-1 was specific for a distinct subpopulation of β -cells (Fig. 2a, b). When initially challenged with 10 mM glucose for 30 s, most cells exhibited either no response or a small (≤ 15 mV) depolarizing response (Fig. 2a, cross-hatched bars). The glucose-sensitivity of individual β -cells therefore is diminished in comparison to β -cells of whole, intact islets. In contrast, when glucose-insensitive β -cells were subsequently challenged with a combined application of glucose and GLP-1, most cells were depolarized by ≥ 30 mV (Fig. 2a, cross-hatched bars). Scatter plot analysis revealed that the interaction between GLP-1 and glucose segregated among three distinct sets of observations (Fig. 2b, sets 1–3). Cells included in set 2 comprised 28% of all cells tested and were fully responsive to glucose (that is, constitutively glucose-competent). In contrast, cells included in sets 1 and 3 exhibited very little response when initially challenged with glucose, and these comprised 64% of all cells tested. When the cells that showed a poor initial response to glucose were subsequently challenged with glucose and GLP-1, introduction of glucose competence was observed in 40% of all cells tested, and it was this subpopulation of β -cells that comprised set 3. These findings are consistent with previous reports of micro-heterogeneity among β -cells^{16,17,19}.

To begin to define the signalling system by which GLP-1 acts, the actions of membrane-permeant cAMP analogues were tested for their effects on the induction of glucose competence. As summarized in Table 1, in β -cells that initially exhibited a weak depolarizing response (< 15 mV) to 10 mM glucose, the application of 10 μ M Rp-cAMPS, an antagonist of endogenous cAMP signalling pathways, inhibited the synergistic depolarizing interaction between glucose and GLP-1. In contrast, 10 μ M of the cAMP agonist Sp-cAMPS potentiated the depolarizing response to glucose even in the absence of GLP-1. These findings suggest that cAMP-mediated signalling systems are necessary for the induction of glucose competence by GLP-1. Previous studies have implicated cAMP-mediated signalling systems in the regulation of the β -cell membrane potential^{25,26}, possibly by controlling the activity of ATP-sensitive potassium channels (I_{KATP}) and/or voltage-sensitive calcium channels^{14,27-29}.

To begin to define the biophysical basis for induction of glucose competence by GLP-1, attention was focused on β -cells that exhibited a ≤ 10 mV depolarizing response when initially challenged with glucose. Voltage clamp analysis was done in the perforated patch configuration and the effects of glucose and GLP-1 on the resting membrane conductance were monitored under conditions in which the bath solution contained no glucose. Brief

application of 10 mM glucose or 10 nM GLP-1 was without significant effect on membrane conductance, whereas the conductance was reversibly inhibited by combined application of both test substances (Fig. 3a). A very similar inhibitory action was also recorded during application of 10 nM glyburide (a sulphonylurea that blocks I_{KATP} with high specificity^{6,20}) to these same cells (data not shown). These findings suggest that glucose and GLP-1 synergize to depolarize β -cells by closing sulphonylurea-sensitive potassium channels (I_{KATP} channels) that are open at or near the resting potential.

To characterize the channels inhibited by glucose and GLP-1, cell-attached patch recordings were obtained under conditions that allow analysis of inwardly-directed I_{KATP} currents. Baseline activity of single I_{KATP} channels exhibited a mean unitary current amplitude of 6.0 pA (pipette potential +50 mV) when the bath solution contained no added glucose (Fig. 3b trace 1), and the single channel conductance inferred from the unitary current amplitude as a function of voltage (I - V) relationship was 60 pS (Fig. 3c), as expected for I_{KATP} ^{6,7}. In 6 of 10 cells tested, application of 10–20 mM glucose had very little effect on channel activity (Fig. 3b trace 2), whereas a decrease in channel activity similar to that previously reported^{6,30} was recorded from 4 cells (data not shown). When the initially glucose-insensitive cells were challenged with a combined application of 20 mM glucose and 10 nM GLP-1, the channel activity was decreased in two cells and was nearly eliminated in four cells (Fig. 3b trace 3). In addition, the unitary current amplitude decreased to 4.2 pA, as expected if glucose/GLP-1-induced whole-cell depolarization resulted in a decreased driving force for K^+ . Channel activity recovered (Fig. 3b trace 4) and was then blocked by glyburide (Fig. 3b trace 5), indicating that these currents are in fact attributable to I_{KATP} . This experiment is summarized in Fig. 3d which illustrates the maximal inhibitory effects of glucose, GLP-1, and glyburide on I_{KATP} activity, expressed as the percentage of 500 ms oscilloscope traces ($n = 50$ traces per treatment) that exhibited no channel activity (blanks). Because I_{KATP} channels within the patch were inhibited by application of glucose and GLP-1 to the membrane outside the patch, a cytosolic signalling mechanism is implicated. Whether cAMP-dependent protein phosphorylation is responsible for the inhibition of I_{KATP} will require further analysis given that conflicting reports exist concerning how phosphorylation influences the activity of these channels³⁰⁻³².

The inhibitory action of glucose on I_{KATP} , as observed in the intact islet, is proposed to result from a glucose signalling system whereby uptake of glucose is mediated by a facilitative glucose transporter, intracellular glucose is converted to glucose-6-phosphate by glucokinase, and aerobic glycolysis generates ATP that is required to inhibit channel function^{6,7}. In contrast, the best evidence available to date indicates that GLP-1 exerts its effects on β -cells by stimulating the production of cAMP^{8,21,24}. The action of GLP-1 to induce glucose competence is therefore suggestive of crosstalk between the cAMP and glucose signalling systems. The consequences of crosstalk are likely to be manifest not only at the level of I_{KATP} modulation, but possibly also at more proximal components (glucose transporter, glucokinase) or distal components (voltage-sensitive calcium channels, vesicular insulin secretion) in the glucose signalling system.

These findings indicate that GLP-1, a circulating hormone, synergizes with glucose to inhibit I_{KATP} , thereby allowing β -cells to depolarize and generate action potentials, events that are necessary prerequisites to glucose-stimulated insulin secretion. Induction of glucose competence by GLP-1 is a receptor-mediated process that is distinct from the more direct inhibitory action of sulphonylureas on I_{KATP} , and provides a straightforward explanation for the ability of GLP-1 to stimulate insulin secretion from the pancreas. Furthermore, as is the case for its insulinotropic action *in vivo*, the inhibition of I_{KATP} by GLP-1 exhibits an absolute requirement for glucose. It is for these reasons that GLP-1 may offer distinct therapeutic advantages over sulphonylureas when it is used for treatment of NIDDM.

Administration of GLP-1 by intravenous infusion produces a rise in circulating insulin accompanied by a lowering of blood glucose, a process that is self-terminating³, as expected because the insulinotropic actions of GLP-1 are glucose-dependent. The likelihood for development of hypoglycaemia during treatment with GLP-1 is therefore reduced, and an extra margin of safety is provided that sulphonylureas or insulin do not offer.


Acknowledgments

We thank R. Kramer for critical reading of the manuscript, and C. P. Miller for advice regarding preparation of islet cell cultures. This work was supported by a USPHS grant J.F.H. is an Investigator with the Howard Hughes Medical Institution

References


1. Unger, RH.; Foster, DW. Williams Textbook of Endocrinology. Wilson, JD.; Foster, DW., editors. Saunders; Philadelphia: 1992. p. 1255-1333.
2. Mojsov S, et al. J. biol. Chem 1986;261:11880–11889. [PubMed: 3528148]
3. Nathan DM, Schreiber E, Fogel H, Mojsov S, Habener JF. Diabetes Care 1992;15:270–276. [PubMed: 1547685]
4. Gutniak M, Orskov C, Holst JJ, Ahren B, Efendic S. New Eng. J. Med 1992;326:1316–1322. [PubMed: 1348845]
5. Gerich JE. New Engl. J. Med 1989;321:1231–1245. [PubMed: 2677730]
6. Ashcroft FM, Rorsman P. Prog. Biophys. molec. Biol 1991;54:87–143. [PubMed: 2484976]
7. Rajan AS, et al. Diabetes Care 1990;13:340–363. [PubMed: 1689632]
8. Drucker DJ, Philippe J, Mojsov S, Chick WL, Habener JF. Proc. natn. Acad. Sci. U.S.A 1987;84:3434–3438.
9. Mojsov S, Weir GC, Habener JF. J. clin. Invest 1987;79:616–619. [PubMed: 3543057]
10. Fehmann HC, Habener JF. Endocrinology 1992;130:159–166. [PubMed: 1309325]
11. Weir GC, Mojsov S, Hendrick GK, Habener JF. Diabetes 1989;38:338–342. [PubMed: 2645190]
12. Horn R, Marty A. J. gen. Physiol 1988;92:145–159. [PubMed: 2459299]
13. Falke LC, Gillis KD, Pressel DM, Mislser S. FEBS Lett 1989;251:167–172. [PubMed: 2473925]
14. Smith PA, Ashcroft FM, Rorsman P. FEBS Lett 1990;261:187–190. [PubMed: 2407553]
15. Hamill OP, Marty A, Neher E, Sakman B, Sigworth F. Pflügers Arch 1981;391:85–100.
16. Hiriart M, Matteson DR. J. gen. Physiol 1988;91:617–639. [PubMed: 2458427]
17. Pipeleers D, In't Veld P, Maes E, Van De Winkel M. Proc. natn. Acad. Sci. U.S.A 1982;79:7322–7325.
18. Sherman A, Rinzel J, Keizer J. Biophys. J 1988;54:411–425. [PubMed: 2850029]
19. Wang J, Baimbridge KG, Brown JC. Endocrinology 1992;131:146–152. [PubMed: 1611994]
20. Ashcroft FM. A. Rev. Neurosci 1988;11:97–118.
21. Blackmore PF, Mojsov S, Exton JH, Habener JF. FEBS Lett 1991;283:7–10. [PubMed: 1645298]
22. Goke R, Conlon JM. J. Endocrinol 1988;116:357–362. [PubMed: 2832504]
23. Fehmann HC, Habener JF. Endocrinology 1991;128:2880–2888. [PubMed: 1645253]
24. Thorens B. Proc. natn. Acad. Sci. U.S.A 1992;89:8641–8645.
25. Henquin JC, Meissner HP. Endocrinology 1984;115:1125–1134. [PubMed: 6086286]
26. Eddlestone GT, Oldham SB, Lipson LG, Premdas FH, Beigelman PM. Am. J. Physiol 1985;248:C145–C153. [PubMed: 2578253]
27. Smith PA, Rorsman P, Ashcroft FM. Nature 1989;342:550–553. [PubMed: 2479839]
28. Prentki M, Glennon MC, Geschwind JF, Matschinsky FM, Corkey BE. FEBS Lett 1987;220:103–107. [PubMed: 2440720]
29. Rajan AS, Hill RS, Boyd AE. Diabetes 1989;38:874–880. [PubMed: 2472299]
30. Ribalet B, Ciani S, Edelstone GT. J. gen. Physiol 1989;94:693–717. [PubMed: 2693587]

31. Nichols CG, Lederer WJ. *J. gen. Physiol* 1991;97:1095–1098. [PubMed: 1865175]
32. Wang W, Giebisch G. *Proc. natn. Acad. Sci. U.S.A* 1991;88:9722–9725.

**FIG. 1.**

Synergism of glucose and GLP-1 to depolarize β -cells. *a*, Trace 1: membrane potential recordings from a β -cell exhibiting weak sensitivity to 10 mM glucose repeatedly applied for 30 s as indicated. Trace 2: recordings from the same cell during application of 10 nM GLP-1 (left) or combined application of 10 mM glucose and 10 nM GLP-1 (right). Initial resting potential (V_m) = -62 mV. *b*, trace 1: a cell that initially failed to respond to 10 mM glucose, but which responded to 10 mM glucose after pretreatment with 10 nM GLP-1 (V_m = -66 mV). Trace 2: a cell that initially failed to respond to 10 nM GLP-1, but which responded to 10 nM GLP-1 after pretreatment with 10 mM glucose (V_m = -60 mV). The initial insensitivity of β -cells to glucose or GLP-1 did not result from their failure to express functional I_{KATP} channels because application of 10 nM glyburide resulted in rapid membrane depolarization (42 ± 11 mV, \pm s.e.m., $n = 10$ cells), as illustrated (*b*, traces 1,2). *b*, -5 pA of hyperpolarizing pipette current was applied to repolarize the membrane after application of 10 nM glyburide. Repolarization confirms the integrity of the membrane/pipette seal. *c*, Cumulative dose-response analysis summarizing the interaction of glucose and GLP-1 to depolarize β -cells. The action of GLP-1(7-37) but not GLP-1(8-37) exhibited dose-dependence under conditions in which the test solution also contained 10 mM glucose (left). In contrast, the action of GLP-1(7-37) but not 10 nM glyburide was abrogated when the test solution contained no added glucose (right). Statistical significance was determined by Student's *t*-test. Probability values (* $P \leq 0.05$, ** $P \leq 0.005$) are expressed relative to control (10 mM glucose and no GLP-1). *c*, The number of cells tested was 50 for GLP-1(7-37) and 10 mM glucose, 25 for GLP-1(8-37) and 10 mM glucose, 20 for GLP-1(7-37) and 0 mM glucose, and 20 for glyburide and 0 mM glucose. The EC_{50} value for GLP-1(7-37) was calculated as 1 nM by establishing a cumulative dose-response relationship in which the action of the peptide was assessed over a concentration range of 0.01–100 nM. The EC_{50} was defined as the concentration of GLP-1(7-37) that depolarized 50% of the cells by ≥ 20 mV when the cells were simultaneously challenged with 10 mM glucose. **METHODS.** Islets were isolated from pancreata of male rats (200–250 g) by collagenase digestion, suspended in culture medium (RPMI 1640 containing 11.1 mM glucose, 10% fetal bovine serum, 100 μ g ml⁻¹ streptomycin, 100 U ml⁻¹ penicillin G), and maintained in culture 1–4 days. Single-cell suspensions were prepared by trypsin-EDTA digestion and mechanical dispersion of cultured islets, and cells were plated on Falcon tissue culture dishes coated with

concanavalin-A for short-term (≥ 16 h) culture. Under these conditions, $\geq 90\%$ of the 10–15 μm diameter cells secrete insulin, as determined by the reverse haemolytic plaque assay¹⁶, and are therefore considered to be β -cells. The fact that $\sim 80\%$ of our cells responded to glyburide indicates that the primary rat cell cultures are indeed enriched with β -cells. For perforated-patch recordings the pipette solution contained 240 $\mu\text{g ml}^{-1}$ nystatin and (in mM): 10KCl, 10NaCl, 70K₂SO₄, 2MgCl₂, 10HEPES (pH 7.35, 295 mOsm). In Figs 1-3 the bath solution contained 138 NaCl, 5.6 KCl, 1.2 MgCl₂, 2.6 CaCl₂, 10 HEPES (pH 7.35, 300 mOsm), and was maintained at 32 °C by a peltier device mounted on the stage of an inverted microscope. All recordings were obtained from solitary cells not organized in clusters, and recordings were initiated within 30 min after transfer of the cells to the glucose-free solution. Pipettes were prepared from borosilicate glass capillary tubes and fire polished to tip resistances of 2–8 M Ω . Measurements were made using an EPC-9 amplifier (bandwidth 10 kHz) interfaced with the Instrutech Data Acquisition and Analysis system (Instrutech Corp., Mineola, NY). The signal was stored on videotape, low-pass filtered (0.5–3.0 kHz, 4-pole Bessel filter, –3 dB attenuation), digitized (100 Hz for recordings of the membrane potential, 5–10 kHz for unitary current recordings), and selected recordings were sampled. Following seal formation, perforation was achieved within 5–10 min, at which time values for the series resistance (R_s), slow capacitance compensation (C_s), and the resting membrane potential were 10–20 M Ω , 6–8 pF and -61 ± 5 mV ($n = 50$), respectively. No correction was made for liquid junction potentials. Test substances were applied by pressure ejection from ‘puffer’ pipettes. The kinetics of this delivery system were established by measuring the rates of onset (time constant, τ , 1.5 s) and offset (τ , 10 s) of 60 mM KCl-induced depolarizations. The change in membrane potential in response to glucose or GLP-1 was measured as the difference between the resting potential and the plateau potential. GLP-1 was obtained from Scios Nova.

**FIG. 3.**

Glucose and GLP-1 synergize to decrease the resting membrane conductance and inhibit the activity of single I_{KATP} channels, *a*, The membrane conductance was monitored by perforated-patch recording in the voltage clamp mode under conditions in which the bath solution contained no added glucose. Application for 30 s of either 10 mM glucose or 10 nM GLP-1 was without significant effect on the magnitude of evoked current responses to ± 10 mV voltage steps from a holding potential of -80 mV, whereas the evoked currents were reversibly inhibited by simultaneous application of 10 mM glucose and 10 nM GLP-1. The conductance decreased 80% from 2.0 nS to 0.4 nS and no shift in the holding current was observed. Outward currents are indicated by upward deflections, *b*, Cell-attached patch recordings of unitary inward currents measured when the bath solution contained no added glucose (traces 1 and 4), 20 mM glucose (trace 2), 20 mM glucose and 10 nM GLP-1 (trace 3), or 10 nM glyburide (trace 5), each applied for 30 s. Inward currents are indicated by downward deflections from a closed level (C) to three superimposed levels of openings (0_1 – 0_3). Filter 1 kHz, sample rate 5kHz. *c*, The I – V relationship for unitary currents recorded in the cell-attached patch configuration. The slope of the I – V relationship decreased at very negative pipette potentials (inward rectification), reversed direction when the pipette potential (V_p) was more negative than -70 mV, and the single channel conductance inferred from the slope of the linear portion of the I – V relationship indicated a value of 60 pS, as expected for I_{KATP} ^{6,20} *d*, Histogram analysis summarizing the actions of glucose, GLP-1, and glyburide, as illustrated in *b*, traces 1–5, to inhibit I_{KATP} . The effects of these test substances were assessed by determining the frequency of occurrence of 500-ms

oscilloscope traces that exhibited no channel activity (blanks). Fifty traces were recorded before, during the peak effect, and after recovery from each test substance.

METHODS, *a*, For measurements of the resting membrane conductance the command potential was shifted by ± 10 mV for 1.5 s at a frequency of 0.1 Hz. The conductance was monitored while simultaneously compensating for the series resistance by 80%. Glucose and GLP-1 were applied at 4-min intervals to avoid priming effects that are observed at shorter intervals, *b*, For cell-attached-patch recordings the pipette solution contained (in mM): 140 KCl, 5CaCl₂, 5MgCl₂, 10 HEPES (pH adjusted to 7.35 with KOH, 305 mOsm) and V_p was +50 mV. Test substances were applied at 10-min intervals, and the current traces illustrated are representative of the peak effects observed. In *b*, trace 2, the relatively small effect of 20 mM glucose on channel activity was accompanied by a decreased unitary current amplitude, suggestive of a decreased driving force for K⁺, possibly due to a depolarizing action of glucose at the whole-cell level. The decreased unitary current amplitude observed in trace 2 may therefore suggest an action of glucose to depolarize β -cells, not simply by inhibiting I_{KATP} channels, but also by inducing an uncharacterized conductance change occurring in the membrane outside the patch.

TABLE 1

Effects of membrane-permeant analogues of cAMP on the interaction of glucose and GLP-1(7-37) to depolarize rat pancreatic β -cells

Treatment	Number of cells tested	mV Change in membrane potential (\pm s.e.m.)
10 mM glucose	10	12 \pm 5
10 mM glucose	5	35 \pm 10*
10 nM GLP-1(7-37)		
10 mM glucose	5	8 \pm 6
10 μ M Rp-cAMPS		
10 mM glucose	5	10 \pm 7
10 nM GLP-1(7-37)		
10 μ M Rp-cAMPS		
10 mM glucose	5	40 \pm 12*
10 μ M Sp-cAMPS		

Sp- and Rp-cAMPS were obtained from BioLog Life Science Institute, La Jolla, CA and prepared as 1 mM stock solutions in distilled water. The analogues were then diluted to a final concentration of 10 μ M in the standard extracellular recording solution described in Fig. 1. Relatively glucose-insensitive β -cells were initially identified by testing for their inability to generate a substantial depolarizing response to 10 mM glucose applied for 30 s. After allowing 5 min recovery, such cells were subsequently challenged with a 30 s application of test solutions containing the indicated concentrations of glucose, GLP-1(7-37), and Sp-cAMPS. For those experiments examining the action of Rp-cAMPS the cells were pretreated in 10 μ M Rp-cAMPS for 30 min at 37 °C. The cells were then challenged with the indicated test solutions containing glucose, GLP-1(7-37), and Rp-cAMPS. Statistical significance was evaluated by Student's *t*-test.

* A value that is significantly different ($P \leq 0.05$) from control (10 mM glucose alone).