

## Symptom Presentation in Patients Hospitalized with Acute Heart Failure

Robert J. Goldberg, PhD; Frederick A. Spencer, MD; Mariana Szklo-Coxe, PhD, MHS; Mayra Tisminetzky, MD; Jorge Yarzebski, MD, MPH; Darleen Lessard, MS; Joel M. Gore, MD; William Gaasch, MD

Department of Medicine, Division of Cardiovascular Medicine, University of Massachusetts Medical School (Goldberg, Tisminetzky, Yarzebski, Lessard, Gore), Worcester, Massachusetts; Department of Medicine, McMaster University (Spencer), Hamilton, Ontario, Canada; School of Community and Environmental Health, Old Dominion University (Szklo-Coxe), Norfolk, Virginia; Department of Cardiology, Lahey Medical Center (Gaasch), Burlington, Massachusetts

### ABSTRACT

**Objectives:** The objectives of this study were to examine the type and frequency of symptoms in patients hospitalized with acute heart failure (HF) as well as the relationship between symptom patterns and patient characteristics, treatment practices, and hospital outcomes in patients hospitalized with decompensated HF.

**Methods:** The study sample consisted of 4537 residents of the Worcester, MA metropolitan area hospitalized for decompensated HF at 11 greater Worcester medical centers in 1995 and 2000.

**Results:** The average age of the study sample was 76 years; the majority (57%) were women, and three-quarters of our patient population had been previously diagnosed with HF. Dyspnea (93%) was the most frequent complaint reported by patients followed by the presence of peripheral edema (70%), cough (51%), orthopnea (37%), and chest pain/discomfort (30%). Patients reporting few cardiac symptoms were less likely to be treated with effective cardiac therapies during hospitalization than patients with multiple cardiac signs and symptoms and experienced higher hospital (9.7% vs. 7.7%) as well as 30-day (17.1% vs. 10.2%) death rates ( $P < 0.05$ ).

**Conclusions:** The results of this study in residents of a large New England community suggest that patients with fewer reported symptoms of decompensated HF were less likely to receive effective cardiac treatments and had worse short-term outcomes. Reasons for these differences in treatment practices and short-term outcomes need to be elucidated and attention directed to these high-risk patients.

### Introduction

Despite the increasing prevalence of heart failure (HF) in the United States and the considerable morbidity, mortality, and functional disability associated with this clinical syndrome, there is limited literature describing the symptoms associated with acute HF. The association between patients' symptoms of acute HF, their demographic and clinical characteristics, treatment practices, and short-term outcomes has also been infrequently examined, particularly from the more generalizable perspective of a population-based investigation. Knowledge and awareness of the signs and symptoms of HF by patients, their relatives and close acquaintances, as well as their healthcare providers is important for enhancing patients' care seeking behavior and the institution of early treatment approaches to reduce the exacerbations of, and complications associated with, HF.

The purpose of this study was to describe the type and frequency of various signs and symptoms related to HF in residents of a large New England metropolitan area hospitalized with decompensated HF at all area medical centers.<sup>1-3</sup> Our secondary study objectives were to examine the association between signs and symptoms of acute HF with patients' demographic and clinical characteristics as well as with hospital treatment practices and short-term death rates.

### Methods

Residents of the Worcester, MA metropolitan area (2000 census estimate = 478 000) hospitalized for possible HF at all 11 greater Worcester medical centers during 1995 and 2000 comprised the study sample.<sup>1-3</sup> In brief, the medical records of patients with primary and/or secondary International Classification of Disease (ICD)-9 discharge diagnoses consistent with the possible presence of decompensated HF were reviewed by trained nurse and physician abstractors.<sup>1-3</sup> The primary ICD-9 code reviewed for the identification of cases of acute HF was

Grant support for this project was provided by the National Heart, Lung, and Blood Institute (R37 HL69874).

ICD code 428. Confirmation of the diagnosis of HF, based on use of the Framingham criteria, required the presence of 2 major criteria or 1 major and 2 minor criteria (Appendix).<sup>4,5</sup> Examples of the major criteria included paroxysmal nocturnal dyspnea, cardiomegaly, and neck vein distention, while examples of the minor criteria included bilateral ankle edema and a night cough.

Information was collected about patients' demographic and clinical characteristics through the review of hospital medical records. Information about prior comorbidities, including a history of HF, was obtained through the review of hospital charts as noted by the attending physicians. Emergency department notes by physicians and nurses were reviewed to identify patients' presenting signs and symptoms. Information on a total of 14 acute clinical signs and symptoms was collected. These included abdominal pain, angina/chest pain, ascites, cough, dyspnea, edema, fatigue, mental obtundation, nausea/vomiting, nocturnal paroxysms, orthopnea, palpitations, weakness, and weight gain. Classification of these signs and symptoms was based on any mention of its presence as opposed to the documentation of the absence of said finding. Information about hospital treatment practices, lifestyle recommendations by health care providers, and hospital case-fatality rates was also obtained through the retrospective review of hospital records.<sup>2</sup> Patients who developed HF secondary to admission for another illness (eg, acute myocardial infarction) or after an interventional procedure (eg, percutaneous coronary intervention) were not included.

Differences in the characteristics, as well as hospital treatment practices and short-term (hospital and 30 d after admission) death rates, of patients presenting with few ( $\leq 2$  of the 5 most frequent signs and symptoms reported by patients and  $\leq 3$  of the total number of 14 signs and symptoms examined) vs. multiple acute HF-related symptoms were examined through the use of  $\chi^2$  tests and *t* tests for discrete and continuous variables, respectively.

Logistic regression analyses were utilized to examine the association between number of signs and symptoms of acute HF at the time of hospital presentation with a variety of patient associated characteristics as well as with hospital treatment regimens and short-term death rates. Factors included for consideration in this analysis were included in our regression models based on their associations with prognosis in the setting of HF from prior publications as well as on the basis of their univariate association ( $P < .20$ ) with hospital and 30-day death rates and differences between patients with few and multiple signs and symptoms of HF (Table 1). For each of the physiologic variables examined, we created quartiles based on commonly utilized cut points to make our findings more interpretable. We categorized the individual comorbid conditions into a frequency count of these aggregated comorbidities (eg, HF, diabetes, coronary heart disease) to make our analytical models more robust

and our findings more easily interpretable. Despite the nonrandomized nature of the present study and potential for confounding by medication indication, we also controlled for the receipt of hospital treatment practices in our regression analyses. Collection of information about 30-day post admission survival status was obtained through the review of hospital medical records and review of state and local death certificates on all study patients.

## Results

A total of 4537 residents of the Worcester metropolitan area were hospitalized with confirmed acute HF at all greater Worcester medical centers during the two study years of 1995 and 2000. The average age of study patients was 76 years, more than one-half were women, and the vast majority were Caucasian (Table 2). Approximately three-quarters of the study sample had been previously diagnosed with HF.

### Frequency of Signs and Symptoms of Decompensated HF

Dyspnea was reported by the vast majority (93%) of study patients as documented in hospital charts (Figure). The presence of edema (70%), cough (51%), orthopnea (37%), and chest pain/discomfort (30%) were the next most commonly reported symptoms. The frequency of these symptoms was not appreciably changed when we examined the occurrence of these symptoms in patients with a first episode of HF (Figure).

We examined the frequency of multiple signs and symptoms associated with acute HF based on information contained in hospital medical records. Approximately 1 in every 20 patients reported having only 1 acute sign or symptom present, 1 in 4 experienced any 3 acute symptoms, and approximately 1 in 3 reporting experiencing any 5 or more of the 14 signs or symptoms examined; only 15% of hospitalized patients reported experiencing 6 or more of these symptoms. A similar profile of signs and symptoms was observed in patients with a first hospitalized episode of HF.

When we restricted our analysis to the 5 most commonly reported symptoms of decompensated HF, namely shortness of breath, edema, cough, orthopnea, and chest discomfort, 29% of all patients reported the presence of any 2 of these symptoms, 35% reported any 3 of these symptoms, while 24% reported 4 or more of these signs and symptoms.

### Characteristics of Patients According to Frequency of Acute Signs and Symptoms

In examining the characteristics of patients experiencing relatively few acute symptoms ( $\leq 2$  of the 5 most frequent signs and symptoms) as compared to those experiencing multiple ( $\geq 3$ ) of the 5 most frequent symptoms reported, patients reporting few acute symptoms were relatively older, more likely to be female, but less likely to be overweight

**Table 1.** Characteristics of Patients Hospitalized With Acute Heart Failure According to Frequency of Acute Symptoms

Characteristic	Five Most Commonly Reported Symptoms			All Symptoms Examined		
	Few Acute Symptoms <sup>a</sup> (n = 1836)	Multiple Acute Symptoms (n = 2701)	P Value	Few Acute Symptoms <sup>b</sup> (n = 2028)	Multiple Acute Symptoms (n = 2509)	P Value
Age (mean, yrs)	77.4	75.1	<0.001	76.7	75.6	<0.005
Male (%)	39.7	45.5	<0.001	41.9	44.2	0.13
Caucasian (%)	95.0	93.6	0.43	94.6	93.8	0.53
Body mass index (%)						
25–29.9	27.0	27.1	<0.001	26.7	27.3	<0.001
≥30	20.1	32.7		24.5	30.3	
Comorbid condition count (%)						
0–2	26.9	22.0		25.6	22.6	
3–5	57.4	58.9	<0.001	57.4	59.0	<0.05
≥6	15.7	19.1		17.0	18.4	
Hospital admission findings (mean)						
Systolic blood pressure (mm Hg)	142.8	145.9	<0.05	144.8	144.5	0.70
Diastolic blood pressure (mm Hg)	75.7	77.5	<0.01	76.9	76.7	0.68
Heart rate (bpm)	91.0	90.5	0.48	91.1	90.4	0.34
Laboratory measures (mean, mg/dL)						
Blood urea nitrogen	36.0	33.3	<0.005	34.2	34.6	0.62
Creatinine	1.7	1.6	<0.005	1.6	1.6	0.59
Sodium	137.4	137.2	0.18	137.4	137.1	0.13

<sup>a</sup> ≤2 vs ≥3 of the 5 most commonly reported acute symptoms.  
<sup>b</sup> ≤3 vs ≥4 of the 14 total acute symptoms examined.

or to have had selected comorbidities previously diagnosed (Table 1). Patients who reported the presence of 2 or fewer of these symptoms were significantly more likely to have presented with lower blood pressure and have higher serum levels of blood urea nitrogen and creatinine than patients with multiple acute signs and symptoms. Relatively similar trends were observed when we examined the distribution of these factors in patients who experienced 3 or fewer of all acute signs and symptoms examined in comparison to those who reported 4 or more of these 14 signs and symptoms (Table 1).

The results of a logistic regression analysis, which included factors associated with the symptoms under study based on their univariate associations and prior findings from the published literature, suggested that relatively older patients, women, those of normal or low body weight, and

those with fewer comorbidities present were significantly more likely to report fewer acute symptoms (of the 5 most common symptoms reported) than respective comparison groups (Table 3). Patients with higher levels of systolic blood pressure and those with higher estimated glomerular filtration rate (eGFR) findings were less likely to report fewer acute symptoms. Relatively similar factors were associated with the reporting of 3 or fewer, compared to 4 or more, of the 14 total symptoms examined, though the statistical significance of several of these factors was attenuated.

#### Hospital Treatment Practices and Frequency of HF Signs and Symptoms

Residents of the Worcester metropolitan area who presented to central Massachusetts medical centers with 2 or fewer

Table 2. Demographic and Clinical Characteristics of the Study Sample

Characteristic	n	%
Age (mean, yrs)	4534	76.0
Age (yrs)		
<65	675	14.9
65–74	1040	22.9
75–84	1731	38.2
≥85	1088	24.0
Male	4537	43.2
Caucasian	4537	94.2
Body mass index (mean)	3722	27.7
Comorbidities		
Anemia	4537	23.5
Chronic lung disease	4537	34.8
Coronary heart disease	4537	56.8
Diabetes	4537	39.9
Heart failure	4537	74.6
Hyperlipidemia	4537	20.1
Hypertension	4537	65.2
Myocardial infarction	4537	47.9
Renal failure/disease	4537	23.3
Stroke	4537	14.6
Physiologic findings (mean)		
Systolic blood pressure (mm Hg)	4524	144.6
Diastolic blood pressure (mm Hg)	4470	76.8
Heart rate (bpm)	4532	90.7
eGFR	4472	63.1
Blood urea nitrogen (mg/dL)	4482	34.4
Serum sodium (mg/dL)	4495	137.2
Died during hospitalization	4537	6.8
Abbreviations: eGFR, estimated glomerular filtration rate.		

of the 5 most frequent symptoms examined were less likely to be treated with effective cardiac medications than patients who reported a greater number of acute symptoms (Table 4). These patients were also less likely to have been told by their health care providers to carry out various lifestyle changes than patients who presented with




Figure 1. Distribution of selected signs and symptoms in patients with acute heart failure.

multiple HF related symptoms. Relatively similar treatment and lifestyle practice recommendations were observed in patients who reported 3 or fewer of the total number of acute symptoms examined (Table 4).

#### Hospital and 30-Day Death Rates

Hospital death rates were significantly higher in those who reported few ( $\leq 2$  of the 5 most common signs and symptoms reported or  $\leq 3$  of all signs and symptoms examined) acute symptoms (9.7% and 7.7%, respectively) as compared to those who reported the presence of more of these symptoms (4.8% and 6.1%, respectively). The 30-day death rates after hospital admission were also higher in patients who reported fewer of the most common symptoms reported as compared to those who reported many of the most common symptoms examined (17.1% vs. 10.2%;  $P < .05$ ).

In addition to the univariate association between the various signs and symptoms of acute HF with hospital and 30-day death rates, we carried out a series of multivariable adjusted regression analyses for purposes of controlling for selected covariates and more systematically examine this association. These variables were controlled for either because they differed between our primary comparison groups (e.g, few vs. more frequent symptoms) or because they had been shown to be of prognostic importance based on findings from the published literature. The results of our regression analyses showed that hospital death rates remained significantly higher in those who reported few, as compared to those who reported the presence of multiple, acute symptoms, with the strongest associations observed when we examined the 5 most commonly reported acute symptoms (Table 5). Similar associations were observed when we examined 30-day death rates, particularly among those who reported  $\leq 2$ , as compared to 3 or more, of the 5 most commonly reported symptoms. There were no significant differences in the 30-day death rates of those who reported few, as compared to those who reported more, of the 14 total acute symptoms examined (Table 5). Adjustment

Table 3. Factors Associated with Multiple Heart Failure Related Symptoms

Characteristic <sup>c</sup>	Five Most Commonly Reported Symptoms <sup>a</sup>		All Symptoms Examined <sup>b</sup>	
	Adjusted OR	95% CI	Adjusted OR	95% CI
Age (yrs)				
65–74	1.25	1.01, 1.54	1.23	1.01, 1.51
75–84	1.33	1.09, 1.62	1.16	0.96, 1.39
≥85	1.38	1.11, 1.71	1.20	0.98, 1.47
Male	0.76	0.67, 0.86	...	...
Length of hospital stay (≤4 or 5 d)	0.83	0.73, 0.94	...	...
Body mass index				
25–29.9	0.79	0.68, 0.92	0.78	0.67, 0.90
≥30	0.50	0.42, 0.58	0.66	0.56, 0.77
Comorbid condition count (%)				
3–5	0.77	0.66, 0.89	0.86	0.74, 0.99
≥6	0.64	0.51, 0.77	0.83	0.69, 1.00
Hospital admission findings				
Systolic blood pressure (mm Hg)				
120–139	0.96	0.80, 1.16		
140–159	0.90	0.74, 1.09		
≥160	0.81	0.66, 0.99		
Diastolic blood pressure (mm Hg)				
60–79	0.80	0.67, 0.96		
80–99	0.86	0.71, 1.05		
≥100	0.94	0.72, 1.22		
BUN (mg/dL)				
30–39	0.95	0.78, 1.15		
40–49	0.86	0.68, 1.10		
≥50(mL/min/1.73m <sup>2</sup> )	1.04	0.84, 1.30		
eGFR				
30–59	0.80	0.65, 0.98		
≥60	0.68	0.54, 0.85		

Abbreviations: BUN, blood urea nitrogen; CI, confidence interval; eGFR, estimated glomerular filtration rate; OR, odds ratio.

<sup>a</sup> 2 or fewer of the 5 most commonly reported acute symptoms.

<sup>b</sup> 3 or fewer of the 14 total acute signs and symptoms examined.

<sup>c</sup> Reference groups: age <65 years, female, length of hospital stay >5 days, body mass index < 25, 0–2 comorbid conditions, systolic blood pressure < 120 mm Hg, diastolic blood pressure < 60 mm Hg, BUN < 30 mg/dL, eGFR < 30 (mL/min/1.73m<sup>2</sup>).

Table 4. Hospital Treatment Practices According to Frequency of Symptoms of Acute Heart Failure

	Five Most Commonly Reported Symptoms		All Symptoms Examined	
	Few Acute Symptoms <sup>a</sup> (n = 1836)	Multiple Acute Symptoms (n = 2701)	Few Acute Symptoms <sup>b</sup> (n = 2028)	Multiple Acute Symptoms (n = 2509)
<b>Medications (%)</b>				
ACE inhibitors/ARBs	50.5 <sup>d</sup>	55.6	51.5 <sup>c</sup>	55.2
β-Blockers	39.1 <sup>c</sup>	42.0	40.0	41.5
Both therapies	21.5 <sup>c</sup>	24.4	22.7	23.6
Digoxin	49.3	51.3	48.9	51.8
Diuretics	96.9 <sup>d</sup>	98.3	97.1 <sup>c</sup>	98.3
<b>Lifestyle Recommendations (%)</b>				
Limit salt intake	72.2 <sup>e</sup>	85.7	74.9 <sup>e</sup>	84.6
Fluid restriction	17.6 <sup>e</sup>	23.8	17.8 <sup>e</sup>	24.2
Smoking cessation	0.7 <sup>d</sup>	1.5	0.6 <sup>d</sup>	1.6
Increase physical activity	38.1 <sup>e</sup>	44.0	39.7 <sup>c</sup>	43.1
Limit alcohol consumption	0.3 <sup>d</sup>	1.0	0.4 <sup>c</sup>	1.0

Abbreviations: ACE, angiotensin-converting enzyme; ARB, angiotensin receptor blockers.  
<sup>a</sup> ≤2 vs ≥3 of the 5 most commonly reported acute symptoms.  
<sup>b</sup> ≤3 vs ≥4 of the 14 total acute symptoms examined.  
<sup>c</sup> P < .05.  
<sup>d</sup> P < .01.  
<sup>e</sup> P < .001.

for hospital treatment practices in each of our regression models resulted in considerable attenuation of our observed findings.

**Discussion**

The results of our study in residents of a large central New England metropolitan area provide insights to the clinical symptoms associated with decompensated HF and the relation of these signs and symptoms to patients’ demographic and clinical characteristics, treatment practices, and short-term outcomes. Our results suggest that many patients with HF present with multiple acute signs and symptoms and that the vast majority present with breathlessness and signs of congestion, rather than signs and symptoms indicative of low cardiac output. That said, approximately 40% of patients reported 2 or fewer common symptoms of HF. Patients reporting fewer HF-related symptoms were older and more likely to be female, less likely to have serious accompanying comorbidities, and less likely to be treated with effective cardiac therapies. Symptom presentation was similar for patients with an initial or previously documented episode of HF and whether

we examined the 5 most common symptoms reported by patients or all 14 of the acute signs and symptoms examined.

The profile of different symptoms, extent of delay in seeking medical care, and relationship of acute symptoms to medical treatment and hospital outcomes in hospitalized patients with decompensated HF has not been examined in detail in the published literature. In prior studies based on the review of information contained in hospital medical records, an approach similar to the approach utilized in the present investigation, the frequency and type of symptoms reported by patients with acute HF were similar to those observed in our population-based investigation.<sup>6–10</sup> For example, in a study based on the review of the medical records of 753 patients with HF who were hospitalized at a Veteran’s Administration hospital, the most common symptoms reported by patients at the time of hospital admission were dyspnea, edema, and fatigue.<sup>7</sup>

Heart failure is typically associated with multiple clinical signs and symptoms that can interact, thereby affecting a patient’s perception of some or all of these acute symptoms. The intensity, timing, level of distress, and quality of symptoms, as well as mismatch between the symptoms



Table 5. Frequency of Symptoms of Acute Heart Failure and Short-Term Mortality

	Hospital Death Rates		30-day Death Rates	
	Few Acute Symptoms <sup>a</sup>	Few Acute Symptoms <sup>b</sup>	Few Acute Symptoms <sup>a</sup>	Few Acute Symptoms <sup>b</sup>
Controlling variables				
Age and sex	2.06 (1.62, 2.61)	1.27 (1.00, 1.60)	1.73 (1.45, 2.06)	1.18 (0.99, 1.40)
Age, sex, and prior comorbidities <sup>c</sup>	1.82 (1.43, 2.33)	1.16 (0.91, 1.47)	1.55 (1.29, 1.86)	1.08 (0.90, 1.29)
Age, sex, prior comorbidities, and physiologic findings <sup>d</sup>	1.74 (1.35, 2.26)	1.12 (0.87, 1.45)	1.45 (1.19, 1.76)	1.04 (0.86, 1.26)
Age, sex, prior comorbidities, physiologic findings, and hospital treatment practices <sup>e</sup>	1.44 (1.80, 1.88)	0.98 (0.75, 1.27)	1.28 (1.05, 1.56)	0.94 (0.77, 1.15)
All variables univariately related to death	1.78 (1.40, 2.27)	1.15 (0.90, 1.45)	1.52 (1.27, 1.83)	1.07 (0.89, 1.28)

Abbreviations: ACE, angiotensin-converting enzyme; ARB, angiotensin receptor blockers; BMI, body mass index; BUN, blood urea nitrogen; CHD, coronary heart disease; COPD; chronic obstructive pulmonary disease; DNR, do not resuscitate; eGFR, estimated glomerular filtration rate; HF, heart failure; MI; myocardial infarction.  
Hospital death rates controlling variables = age, race, BMI, prior angina, CHD, diabetes, hyperlipidemia, hypertension, MI, and DNR orders.  
30-day death rates controlling variables = age, race, BMI, prior anemia, angina, CHD, diabetes, CHF, hyperlipidemia, hypertension, MI, stroke, and DNR orders.  
<sup>a</sup>  $\leq 2$  vs  $\geq 3$  of the 5 most commonly reported acute symptoms.  
<sup>b</sup>  $\leq 3$  vs  $\geq 4$  of the 14 total acute symptoms examined.  
<sup>c</sup> Prior comorbidities = BMI, prior anemia, angina, COPD, CHD, diabetes, HF, hyperlipidemia, MI, renal disease, stroke, and DNR orders.  
<sup>d</sup> Physiologic findings = Factors listed in regression <sup>c</sup> plus admission systolic and diastolic blood pressure, heart rate, eGFR, BUN, serum sodium, and white blood cell count.  
<sup>e</sup> Hospital treatment practices = Factors listed in regression <sup>d</sup> plus use of ACE inhibitors, ARB blockers,  $\beta$ -blockers, digoxin, diuretics, recommendation to reduce salt intake, fluid restriction, smoking cessation, increased physical activity, and alcohol reduction.

expected and those actually experienced, may each affect a patient's perceptions of, and responses to, his or her acute symptomatology.<sup>11</sup> Psychologic as well as situational factors may also influence the acute symptom experience and its interpretation. In our study, we found that patients' reporting of symptoms may also impact how they are treated once they seek medical attention.

Patients who reported fewer symptoms of acute HF were less likely to be treated with effective treatment regimens and use of different nonpharmacologic regimens. In addition, in both crude and multivariable adjusted analyses, our findings suggest that patients with fewer signs and symptoms of decompensated HF, particularly those related to the 5 most commonly reported symptoms, are at greater risk for dying during their acute hospitalization as

well as during the first month of their hospital admission. On the interpretation of our findings, that after adjustment. The nonrandomized nature of the present study places appropriate caveats on the interpretation of our findings. However, it needs to be noted that the association between hospital and 30 day death rates became increasingly attenuated (especially for the 5 most commonly reported acute symptoms) after adjustment for the receipt of various hospital treatment practices. These findings suggest that the failure to treat these patients with effective cardiac medications may lead to increased short-term death rates in this relatively under-recognized (eg, fewer acute symptoms) patient population.

Numerous hypotheses can be offered for our observed findings. It is possible that patients reporting fewer

acute symptoms may have experienced longer delays to hospital presentation, more atypical presentation, and more diagnostic uncertainty leading to possible delays in appropriate treatment and higher death rates. The observed differences in hospital treatment practices based on the number of symptoms reported by patients suggest that health care providers respond differently based on symptom profile of patients. It is also possible that the reporting of fewer symptoms is a marker for patients with more severe comorbidities or even impaired cognition, that may impact subsequent survival. Further studies need to be carried out to understand the reasons for these observed associations including a more detailed assessment of the frequency, intensity, and duration of patient's symptoms and actions taken to seek acute medical care.

### Study Strengths and Limitations

The strengths of the present study are the inclusion of all hospitalized patients with decompensated HF from a large New England metropolitan area, use of standardized criteria for confirmation of the diagnosis of HF, and inclusion of a wide variety of possible symptoms of HF. The limitations of this study include the reliance of information contained in hospital medical records and inability to separate the chief symptom complaint from more ancillary symptoms. Our study also does not necessarily confirm the absence of selected symptoms of acute HF, but merely the absence of documentation of the various symptoms examined. Due to the large amount of missing data on ejection fraction, we were unable to further classify patients into those with systolic from those with diastolic HF. We were unable to examine factors such as socioeconomic status, noncardiac disabilities, usual activity levels, prior treatment history, cognitive function, and a history of depression in the present study. Our study sample was primarily white, and the present findings may not apply to patients of other races or ethnicities.

### Conclusions

The results of the present population-based study suggest that patients with acute HF present with many signs and symptoms of decompensation. Patients who report fewer symptoms tend to be older, female, have fewer comorbidities present, have lower prescription rates of HF therapy, and have worse short-term outcomes in comparison to patients who report the presence of multiple acute symptoms. These findings suggest that health care providers should pay particular attention to patients with acute HF who present with fewer symptoms of decompensated HF and increase their efforts of surveillance and targeted therapeutic approaches in these high risk patients.

## Appendix

### Framingham Heart Study Criteria

Major	Minor
Paroxysmal nocturnal dyspnea	Bilateral ankle edema
Neck vein distention	Nocturnal cough
Rales	Dyspnea on exertion
Pleural effusion by chest x-ray consistent with cardiomegaly	Hepatomegaly
Acute pulmonary edema	Tachycardias 120 beats per min
S3 gallop	Pulmonary vascular engorgement by chest x-ray
Increased venous pressure (>16 cm H <sup>2</sup> o)	
Positive hepatojugular reflex	
Major or Minor	
Weight loss >4.5 kg (10 lbs) over 5 d of treatment	

## References

- Goldberg RJ, Spencer FA, Farmer C, et al. Incidence and hospital death rates associated with heart failure: a community-wide perspective. *Am J Med.* 2005;118:728–734.
- Goldberg RJ, Spencer FA, Farmer C, et al. Use of disease-modifying therapies in patients hospitalized with heart failure: a population-based perspective. *Am J Med.* 2007;120:98.e1–8.
- Goldberg RJ, Ciampa J, Lessard D, et al. Long-term survival after heart failure: a contemporary population-based perspective. *Arch Intern Med.* 2007;167:490–496.
- Ho KKL, Pinsky JL, Kannel WB, et al. The epidemiology of heart failure: the Framingham Study. *J Am Coll Cardiol.* 1993; 22(Suppl A):6A–13A.
- Vasan RS, Larson MG, Benjamin EJ, et al. Congestive heart failure in subjects with normal versus reduced left ventricular ejection fraction: prevalence and mortality in a population-based cohort. *J Am Coll Cardiol.* 1999;33: 1948–1955.
- Friedman M. Older adults' symptoms and their duration before hospitalization for heart failure. *Heart Lung.* 1997;26:169–176.
- Evangelista L, Dracup K, Doering L. Treatment-seeking delays in heart failure patients. *J Heart Lung Transplant.* 2000;19:932–938.
- Schiff GD, Fung S, Speroff T, et al. Decompensated heart failure: symptoms, patterns of onset, and contributing factors. *Am J Med.* 2003;114:625–630.
- Parshall MB, Welsh JD, Brockopp DY, et al. Dyspnea duration, distress and intensity in emergency department visits for heart failure. *Heart Lung.* 2001;30:47–56.
- Bekelman DB, Havranek EP, Becker DM, et al. Symptoms, depression, and quality of life in patients with heart failure. *J Card Fail.* 2007;13:643–648.
- Horne R, James D, Petrie K, et al. Patients' interpretation of symptoms as a cause of delay in reaching hospital during acute myocardial infarction. *Heart.* 2000;83:388–393.