

Simulation in surgical education

Vanessa N. Palter MD, Teodor P. Grantcharov MD PhD

Previously published at www.cmaj.ca

In recent years, simulation has come to the foreground as a method of teaching technical skills to surgical trainees. Rapid changes in health care systems underlie the shift to simulation-based education in many countries. Pressures include a decrease in work hours for residents, a shortening of training programs, a decrease in available operating-room time, and ethical imperatives to protect patients from harm.¹⁻³ The expectation that trainees will acquire all necessary skills in a purely clinical environment is no longer realistic. Simulation offers the benefits of a safe environment for practice and error, opportunities for feedback and assessment, and standardized experience for trainees.^{1,3-5} It also realizes the concept of the “pretrained” novice, whereby a resident learns basic technical skills in a simulated environment so that when faced with an analogous clinical situation, he or she has already completed the early phase of the learning curve. The result is improved patient safety and learning efficiency.^{6,7}

Although acquisition of technical skills is undoubtedly essential for surgical trainees, non-technical factors such as communication, leadership and teamwork have also been shown to play a substantial role in operative success. A breakdown in these non-technical factors has been shown to increase the rate of errors in the operating room.⁸⁻¹¹ In addition to their role in acquisition of technical skills, simulators also have potential for use in teaching and assessment of non-technical skills.

We review the simulators currently available to surgical educators, discuss the potential uses for simulation technology in surgery and review the current evidence for the role of simulation in both the teaching and assessment of technical and non-technical skills.

Methods

We searched MEDLINE and the Cochrane Database of Systematic Reviews to identify studies of potential relevance. The Cochrane search yielded one relevant article. In the MEDLINE search, we used the Medical Subject Headings (MeSH) “computer simulation” and “education, medical, graduate” (retrieved articles 143, relevant articles 22) and the headings “laparoscopy” and “computer simulation” (retrieved 229, relevant 23). The following keywords were used in the MEDLINE search: “technical skills” and “assessment” (retrieved 283, relevant 64) and “non-technical skills” (retrieved 41, relevant 15). We focused the search on articles published from 1996 onwards, and we limited it to articles published in English. We did not include case reports and data from abstracts in our data synthesis. All of the abstracts of identified articles

Key points

- Both technical and non-technical skills are essential for surgeons in training.
- Technical skills learned on low-fidelity benchtop models, video-based trainers and virtual-reality models are transferable to the operating room.
- Current evidence does not show a clear advantage of either virtual reality or synthetic models in teaching technical skills.
- Evidence is lacking on how best to integrate each type of simulator into a holistic, proficiency-based curriculum.

were examined for relevance. Retrieved studies were screened for duplication, and additional studies were identified using a manual search of the reference lists of the relevant included articles. Given that our review is focused on the role of simulation in surgery, our search strategy was limited to identifying articles focusing on surgical education.

Available simulation models

A wide variety of models is currently available for teaching technical skills. These models range from high-fidelity animal or cadaveric models to virtual-reality simulators.

Animals and human cadavers

Some of the original models used to teach procedural skills are human cadavers and animals (e.g., the porcine model for bronchoscopy, endoscopic retrograde cholangiopancreatography, laparoscopic cholecystectomy, or the canine model for coronary bypass).¹² The advantages of animal models include realism and opportunities to mimic complications. However, they have been criticized for their expense, the fact that their anatomy can differ from that of humans, and ethical reasons.¹²⁻¹⁵ Cadavers are infrequently used in surgical education because of cost, limited availability and inability to simulate complications such as bleeding.^{12,15}

Synthetic benchtop models and tower trainers

Synthetic models include those designed to teach open surgical procedures (commonly referred to as benchtop models) and

From the Department of Surgery, University of Toronto, Division of General Surgery, St. Michael's Hospital, Toronto, Ont.

CMAJ 2010. DOI:10.1503/cmaj.091743

those used for teaching minimally invasive procedures (referred to as tower trainers or video-box trainers). Numerous benchtop models are available for the simulation of a variety of procedures, including knot-tying, fascial closure, and suturing¹⁴ (Figure 1). The most obvious limitation of such models is that they teach only one surgical technique rather than an entire operation.⁵ Their use requires the presence of an expert to demonstrate the procedure and provide feedback on performance.¹⁵

Tower trainers include a box with slits on the superior surface for trocar insertion. Real laparoscopic instruments are inserted through the trocars into the box, where the procedure is simulated. A camera inside the box provides video output to a monitor on which trainees can watch their own movements. These models can simulate a variety of techniques, including laparoscopic suturing, knot-tying, clip-applying and coordination drills.¹⁴ One of the most well-described tower trainers is the physical laparoscopy trainer known as the McGill Inanimate System for Training and Evaluation of Laparoscopic Skills. This system consists of five tasks performed on an endotrainer box.¹⁶ The tasks include peg transfer, cutting, placing a ligating loop and intracorporeal suturing.

Virtual-reality simulation

Training by virtual-reality simulation encompasses systems designed to teach laparoscopic, endoscopic and percutaneous interventions.¹³ The fidelity of virtual-reality simulators ranges widely, from the Minimally Invasive Surgical Trainer – Virtual Reality (MIST-VR) system, which is a low-fidelity system designed to teach general laparoscopic proficiency, to systems that teach component procedural skills and other systems that replicate entire operations¹⁷ (Figure 2 and Videos 1 and 2, available at www.cmaj.ca/cgi/content/full/cmaj.091743/DC1). Unlike bench-model learning, the presence of an expert is not necessary when virtual-reality simulation is used. Such models allow for practice at varying levels of difficulty, can simulate complications and automatically provide objective measures of assessment, allowing for both formative and summative trainee assessment.¹⁸ Virtual-reality mod-



Figure 1: A synthetic benchtop model for teaching abdominal wall closure.

els have been criticized, however, for high initial cost of system acquisition, limited force-feedback with use of surgical instruments and lack of realism of graphics.^{14,19}

The virtual operating room

Benchtop models, virtual-reality models, and animal and cadaveric simulators are all focused on teaching the technical skill set necessary to perform a wide range of surgical procedures. To both teach and assess non-technical and technical aspects of performance, virtual operating rooms have been developed.^{4,20-22} These rooms consist of an operating table, lights, suction equipment and an anesthesia machine. They contain a simulator (either synthetic or virtual reality) and a control room, which is separated from the operating room by a one-way mirror. Virtual operating rooms are designed to take into account the technical skills required to perform an operation, along with the intangible factors, such as interpersonal dynamics or teamwork, that are also responsible for operative success.

The role of simulators in surgical education

A critical role of simulators is to teach trainees a specific skill set. When assessing simulators for utility in a particular curriculum, an examination of the learning curve on a particular simulator and evidence for transfer validity of the taught skills are important. Learning curves refer to improvement of



Figure 2: A virtual reality simulator. (Image reproduced with permission from Surgical Science, Inc.)

performance as measured on a particular simulator. Transfer validity, or the degree to which skills learned on the simulator result in an improvement in skills in an analogous clinical situation, is essential.⁷

Parallel to the development of simulators for technical skills training has been that of assessment tools for measuring these acquired skills. Assessment allows for monitoring of skills acquisition of surgical trainees and provides a base for constructive feedback along the learning curve for new procedures.²³ Objective measures of performance are crucial for determining advancement to the next stage of training by surgical trainees or re-entry after a career break by staff surgeons.²³ Simulators must demonstrate acceptable reliability and validity before they are integrated into high-stakes assessment.

Assessment tools can be classified into three categories: observational tools, virtual reality and motion analysis (Table 1). Observational tools rely on an expert observer who assesses technical ability according to defined criteria on either a global rating scale or a procedure-specific checklist. Virtual-reality simulators provide immediate and automatic assessment by recording metrics such as time taken and error score. Analyses of dexterity equate technical ability with the number and speed of a surgeon's hand movements.

Several assessment tools for non-technical scales have also been designed for surgeons. Yule and colleagues developed a taxonomy and behaviour-rating system for surgeons termed Non-Technical Skills for Surgeons. This rating scale consists of five categories: situation awareness, decision-making, task management, communication and teamwork, and leadership.^{41,42} The inter-rater reliability and intra-class correlation coefficients of

this tool have been demonstrated in initial validation studies.⁴¹ A similar tool, known as the Non-Technical Skills Scale, has been adapted from the aviation industry and shown acceptable reliability and validity for use in surgery.^{43,44} Using these assessment tools effectively, however, requires some degree of training.

Effectiveness in teaching technical skills

Ultimately, the goal of learning technical skills on simulators is to improve technical performance in the operating room. Technical skills learned both on low-fidelity benchtop simulators and on high-fidelity virtual-reality systems have been shown to transfer to the operating room. Assessing which simulator is more useful in which type of training situation, however, is more difficult to determine.

Animal and cadaveric models

Few trials have examined the role that animal or cadaveric models play in a surgical skills curriculum. Several studies have compared these high-fidelity models to their low-fidelity counterparts in teaching technical skill. In a randomized single-blinded trial, Grober and colleagues demonstrated that microsurgical training on the vas deferens of a live rat compared with use of a low-fidelity silicone tubing model produced an equivalent improvement in technical skills on both models using a previously validated checklist and global rating scale.⁴⁵

These findings are similar to those of an earlier randomized single-blinded study by Anastakis and colleagues, in which training on a cadaver was found to be equivalent to low-fidelity benchtop training.¹⁵ Among trainees with increased

Table 1: Tools for assessing technical skill, by level of validity

Assessment tool	Examples of specific tools	Evidence for validity	Advantages	Disadvantages
High validity				
Global rating scales	<ul style="list-style-type: none"> Objective structured assessment of technical skills (OSATS) Global operative assessment of laparoscopic skills (GOALS) 	Observational studies ²³⁻²⁸	Applicable in a wide range of situations (open surgery, bench-station exams laparoscopic surgery, video-based assessment)	Requires time of an expert assessor
Virtual-reality simulators	<ul style="list-style-type: none"> Laparoscopy Bronchoscopy Endovascular 	Observational studies ³⁶⁻³⁸	Automatic, non-biased assessment	Feedback not useful for the trainee
Moderate validity				
Checklist scales	<ul style="list-style-type: none"> Laparoscopic cholecystectomy Carotid endarterectomy 	Observational studies ³²⁻³⁵	Novices can see specifically where they need to improve their skills	Requires time of an expert assessor. Binary assessment.
Analysis of dexterity	<ul style="list-style-type: none"> Imperial College surgical assessment device (ICSAD) Advanced Dundee endoscopic psychomotor tester (ADEPT) 	Observational studies ³⁹⁻⁴⁰	Automatic, nonbiased assessment	Feedback not useful for the trainee. Feasibility-related concerns.
Procedure-specific checklists	<ul style="list-style-type: none"> Laparoscopic colorectal surgery Laparoscopic cholecystectomy Gastric bypass 	Observational studies ²⁹⁻³¹	Novices can see specifically where they need to improve their skills	Requires time of an expert assessor

technical experience, however, learning on higher-fidelity models (i.e., animal or cadaveric) appear to result in greater improvement in technical proficiency compared with training on low-fidelity models.⁴⁶ These studies illustrate that inexpensive, reliable and valid models exist that are equivalent to animal or cadaveric models for training novice surgeons.

Synthetic models

The evidence for effective learning using synthetic training tools is well established. In a blinded, randomized controlled trial, Traxer and colleagues found that practice on a video-trainer resulted in a 51% reduction in time (as measured on the simulator, $p = 0.003$) and an improvement in technical ability (measured by a validated global assessment tool in a porcine model $p = 0.0008$) as compared with a no-training control group.⁴⁷ Similarly, Fried and colleagues established the educational value of the McGill Inanimate System for Training and Evaluation of Laparoscopic Skills by showing evidence of its transfer validity to an animal model.^{16,48,49} Transfer validity to the operating room has also been shown in two randomized controlled trials.^{50,51} Finally, a 2006 systematic review of 14 studies involving a total of 261 participants found that video-based tower training is superior to either no training or conventional training.¹⁹

Not all training on synthetic models is equivalent, and surgical educators must therefore determine the training schedule for a particular curriculum thoughtfully. The quality and amount of expert feedback provided during the learning process plays an essential role in the acquisition of technical skills. Summary feedback (i.e., given at the completion of a task) is reported to be most useful for learning as compared to concurrent feedback (given as the task is occurring).⁵²⁻⁵⁵ Moreover, distributed (as opposed to massed) practice significantly improves learning.⁵⁶ Finally, as with the equivalence shown between high-fidelity animal models and low-fidelity benchtop models, high-fidelity benchtop models are equivalent to low-fidelity benchtop models in acquisition of technical skills by novices.⁵⁷

Virtual reality

Novices attain expert levels of proficiency after at least six trials on a simulator, whereas experts and intermediates tend to plateau much faster, according to observational studies.^{36,58-65} Individual variations exist in the rate of acquisition of technical skills, and therefore training to expert proficiency (as opposed to training for a certain duration) may be the most efficient means by which to enhance learning of technical skills on virtual-reality simulators.^{66,67}

Two randomized unblinded trials showed that, compared with participants with no training, those trained using virtual-reality tools showed significantly improved learning, as measured by both previously validated global assessment measures and automatic simulator-generated parameters.^{68,69} The technical skills acquired during virtual-reality training have been shown to be transferable both to an animal model and to patients in the operating room.⁷⁰⁻⁷³ In a randomized double-blinded study, Seymour and colleagues showed that participants who received low-fidelity virtual-reality training showed improved time to complete laparoscopic cholecystectomy (29% faster than residents not trained using virtual reality),

were five times more likely to make progress and six times less likely to make errors compared with residents not trained using virtual reality.⁷⁴ These results were replicated in a later randomized blinded controlled trial by Grantcharov and colleagues.²⁴ Moreover, the effect of virtual-reality training was shown to persist for the first 10 laparoscopic cholecystectomies performed by a novice resident.⁷⁵ Finally, Larsen and colleagues reported improved performance after virtual-reality training compared with a control group in operative salpingectomy with respect to time ($p < 0.0001$) and on a global rating scale ($p < 0.0001$), in a randomized blinded study using previously validated assessment measures.⁷⁶

Both video-trainers and virtual-reality models have been shown to be effective for learning technical skills, and these skills have been shown to transfer to real clinical situations. In several randomized controlled trials, as well as a recent systematic review looking only at novice surgical trainees, virtual-reality training was shown to be at least as effective as benchtop training in the acquisition of technical skills.^{65,77-80} Notably, many of the identified studies had methodological limitations, including small sample sizes, inadequate blinding, disparate interventions and use of surrogate outcomes.^{19,81,82}

A small number of studies, however, suggest that virtual reality training may be superior to benchtop-model training. In a randomized trial, Youngblood and colleagues reported that surgically naïve medical students who trained on a virtual-reality simulator, as opposed to a tower trainer, showed improved time, accuracy and global rating scores, as measured in a porcine model.⁸³ The assessment measures used in this study, however, were not validated. Hamilton and colleagues found that surgical residents who received virtual-reality (as opposed to tower) training showed improved operative performance during laparoscopic cholecystectomy, as measured by a global rating scale completed by blinded faculty observers.⁸⁴ Although these findings suggest that virtual-reality training may provide more benefit than benchtop training, more well-designed randomized trials need to be performed before firm conclusions can be drawn.

Each virtual-reality system has various levels of realism and capabilities for haptics. Haptics allow for trainees practicing on virtual-reality simulators to feel force on their instruments, thus providing a higher degree of realism to the device. These features add substantial cost to the device. In a crossover study, Panait and colleagues investigated the role of haptics in technical skills acquisition. Students learned tasks on a virtual-reality model with haptics either present or absent.⁸⁵ The results showed a trend toward improved skill acquisition for complex tasks only when haptics were enabled, suggesting that the cost of haptics-enhanced simulation may be more worthwhile for complex tasks, such as learning operations in their entirety.

The evidence suggests that trainees should learn technical skills in a graded manner on simulators. Early learning could occur on low-fidelity synthetic or virtual-reality models. Trainees could then progress to training on high-fidelity synthetic or animal models, or to virtual-reality systems that simulate complex procedures with the added effect of force-feedback.

Effectiveness in teaching non-technical skills

The development and validation of a simulated operating room where nurses, anesthesiologists and surgeons can work through a scenario designed to assess and teach both technical and non-technical skills has been described by Moorthy and colleagues and by Undre and colleagues.^{4,22} Both of these studies showed the feasibility of interdisciplinary training and also confirmed participant satisfaction in the virtual operating-room experience. Construct validity (i.e., ability to distinguish between experts and novices) was similarly demonstrated for a laparoscopic-crisis scenario in a simulated operating room.²¹

The results are ambiguous, however, with respect to using these simulated environments to specifically teach complex non-technical skills. While simulated operating rooms are generally thought to be useful for training members of the operating-room team, they are complex, and expensive to maintain. By contrast, the Integrated Procedural Performance Instrument developed by Kneebone and colleagues at the Imperial College in London^{86,87} is a simulation model that combines use of physical benchtop simulators to test technical skill with use of standardized patients to assess non-technical skills. Using this model, LeBlanc and colleagues found in an observational study that communication skills are not correlated with technical skill, thus emphasizing the need for specific training in non-technical skills in surgical training programs.⁸⁸

Gaps in knowledge

The goal of implementing simulation-based training in surgery is to provide a complementary experience that accelerates the learning curve and enhances patient encounters.¹⁴ Although the literature supports both synthetic and virtual-reality training in technical skills acquisition, there are some weaknesses in this body of literature. First, although a large variety of virtual-reality simulators is available on the market, many of the existing studies investigated the earlier virtual-reality models. Second, many of the new procedural models lack evidence for validity, which raises concern that the field will be dominated by technology rather than educational principles. Finally, the studies describe disparate interventions, sometimes on the same simulator, making comparisons difficult. Further research in the field should therefore be directed toward the newer procedural simulators, because they are substantially more costly than the lower-fidelity models, and should target surgical trainees rather than medical students.

Simulators are only part of the solution to training staff and residents outside of the operating room. As Gallagher stated, "their power can only be truly realized if they are integrated into a validated comprehensive curriculum."⁷⁷ Although there are studies looking at specific tasks within a proficiency-based curriculum,^{24,89-91} a more holistic curriculum that looks at teaching and assessing both technical and non-technical skills needs development and validation. To overcome the focus in the literature on validating specific simulators for specific surgical tasks, Schout and colleagues recommend designing and evaluating a comprehensive training program rather than validating only one aspect of the curriculum that can be performed on a simulator.¹⁸

This article has been peer reviewed.

Competing interests: None declared.

Contributors: Both of the authors were involved in the conceptualization of the article. Vanessa Palter drafted the manuscript and Teodor Grantcharov critically revised it. Both of the authors approved the final version submitted for publication.

REFERENCES

1. Kneebone R, Nestel D, Wetzel C, et al. The human face of simulation: patient-focused simulation training. *Acad Med* 2006;81:919-24.
2. Haluck RS, Krummel TM. Computers and virtual reality for surgical education in the 21st century. *Arch Surg* 2000;135:786-92.
3. Tavakol M, Mohagheghi MA, Dennick R. Assessing the skills of surgical residents using simulation. *J Surg Educ* 2008;65:77-83.
4. Moorthy K, Munz Y, Forrest D, et al. Surgical crisis management skills training and assessment: a simulation[corrected]-based approach to enhancing operating room performance. *Ann Surg* 2006;244:139-47.
5. Kneebone R. Simulation in surgical training: educational issues and practical implications. *Med Educ* 2003;37:267-77.
6. Van Sickle KR, Ritter EM, Smith CD. The pretrained novice: using simulation-based training to improve learning in the operating room. *Surg Innov* 2006;13:198-204.
7. Gallagher AG, Ritter EM, Champion H, et al. Virtual reality simulation for the operating room: proficiency-based training as a paradigm shift in surgical skills training. *Ann Surg* 2005;241:364-72.
8. Christian CK, Gustafson ML, Roth EM, et al. A prospective study of patient safety in the operating room. *Surgery* 2006;139:159-73.
9. Mishra A, Catchpole K, Dale T, et al. The influence of non-technical performance on technical outcome in laparoscopic cholecystectomy. *Surg Endosc* 2008;22:68-73.
10. Stevenson KS, Gibson SC, MacDonald D, et al. Measurement of process as quality control in the management of acute surgical emergencies. *Br J Surg* 2007;94:376-81.
11. Yule S, Flin R, Paterson-Brown S, et al. Non-technical skills for surgeons in the operating room: a review of the literature. *Surgery* 2006;139:140-9.
12. Sarker SK, Patel B. Simulation and surgical training. *Int J Clin Pract* 2007;61:2120-5.
13. Aggarwal R, Grantcharov TP, Darzi A. Framework for systematic training and assessment of technical skills. *J Am Coll Surg* 2007;204:697-705.
14. Hammoud MM, Nuthalapaty FS, Goepfert AR, et al. To the point: medical education review of the role of simulators in surgical training. *Am J Obstet Gynecol* 2008;199:338-43.
15. Anastakis DJ, Regehr G, Reznick RK, et al. Assessment of technical skills transfer from the bench training model to the human model. *Am J Surg* 1999;177:167-70.
16. Fried GM, Feldman LS, Vassiliou MC, et al. Proving the value of simulation in laparoscopic surgery. *Ann Surg* 2004;240:518-25; discussion 525-8.
17. Dunkin B, Adrales GL, Apelgren K, et al. Surgical simulation: a current review. *Surg Endosc* 2007;21:357-66.
18. Schout BM, Hendriks AJ, Scheele F, et al. Validation and implementation of surgical simulators: a critical review of present, past, and future. *Surg Endosc* 2010;24:536-46.
19. Sutherland LM, Middleton PF, Anthony A, et al. Surgical simulation: a systematic review. *Ann Surg* 2006;243:291-300.
20. Moorthy K, Munz Y, Adams S, et al. A human factors analysis of technical and team skills among surgical trainees during procedural simulations in a simulated operating theatre. *Ann Surg* 2005;242:631-9.
21. Powers KA, Rehrig ST, Irias N, et al. Simulated laparoscopic operating room crisis: An approach to enhance the surgical team performance. *Surg Endosc* 2008;22:885-900.
22. Undre S, Koutantji M, Sevdalis N, et al. Multidisciplinary crisis simulations: the way forward for training surgical teams. *World J Surg* 2007;31:1843-53.
23. Aggarwal R, Grantcharov T, Moorthy K, et al. Toward feasible, valid, and reliable video-based assessments of technical surgical skills in the operating room. *Ann Surg* 2008;247:372-9.
24. Grantcharov TP, Kristiansen VB, Bendix J, et al. Randomized clinical trial of virtual reality simulation for laparoscopic skills training. *Br J Surg* 2004;91:146-50.
25. Datta V, Bann S, Aggarwal R, et al. Technical skills examination for general surgical trainees. *Br J Surg* 2006;93:1139-46.
26. Reznick R, Regehr G, MacRae H, et al. Testing technical skill via an innovative "bench station" examination. *Am J Surg* 1997;173:226-30.
27. Faulkner H, Regehr G, Martin J, et al. Validation of an objective structured assessment of technical skill for surgical residents. *Acad Med* 1996;71:1363-5.
28. Ault G, Reznick R, MacRae H, et al. Exporting a technical skills evaluation technology to other sites. *Am J Surg* 2001;182:254-6.
29. Aggarwal R, Boza C, Hance J, et al. Skills acquisition for laparoscopic gastric bypass in the training laboratory: an innovative approach. *Obes Surg* 2007;17:19-27.
30. Palter V.N., MacRae H.M. T.P. G. Development of an objective evaluation tool to assess technical skill in laparoscopic colorectal surgery: a Delphi methodology. *Am J Surg*. In press.
31. Sarker SK, Chang A, Vincent C, et al. Development of assessing generic and specific technical skills in laparoscopic surgery. *Am J Surg* 2006;191:238-44.
32. Insel A, Carofino B, Leger R, et al. The development of an objective model to assess arthroscopic performance. *J Bone Joint Surg Am* 2009;91:2287-95.
33. Eubanks TR, Clements RH, Pohl D, et al. An objective scoring system for laparoscopic cholecystectomy. *J Am Coll Surg* 1999;189:566-74.

34. Sarker SK, Chang A, Vincent C. Technical and technological skills assessment in laparoscopic surgery. *JLSLS* 2006;10:284-92.
35. Beard JD, Choksy S, Khan S. Assessment of operative competence during carotid endarterectomy. *Br J Surg* 2007;94:726-30.
36. Grantcharov TP, Bardram L, Funch-Jensen P, et al. Learning curves and impact of previous operative experience on performance on a virtual reality simulator to test laparoscopic surgical skills. *Am J Surg* 2003;185:146-9.
37. Larsen CR, Grantcharov T, Aggarwal R, et al. Objective assessment of gynecologic laparoscopic skills using the LapSimGyn virtual reality simulator. *Surg Endosc* 2006;20:1460-6.
38. Gallagher AG, Satava RM. Virtual reality as a metric for the assessment of laparoscopic psychomotor skills. Learning curves and reliability measures. *Surg Endosc* 2002;16:1746-52.
39. Moorthy Y, Munz A, Dosis F, et al. Motion analysis in the training and assessment of minimally invasive surgery. *Minim Invasive Ther Allied Technol* 2003;12:137-42.
40. Francis NK, Hanna GB, Cuschieri A. The performance of master surgeons on the Advanced Dundee Endoscopic Psychomotor Tester: contrast validity study. *Arch Surg* 2002;137:841-4.
41. Yule S, Flin R, Maran N, et al. Surgeons' non-technical skills in the operating room: reliability testing of the NOTSS behavior rating system. *World J Surg* 2008;32:548-56.
42. Yule S, Flin R, Paterson-Brown S, et al. Development of a rating system for surgeons' non-technical skills. *Med Educ* 2006;40:1098-104.
43. Sevdalis N, Davis R, Koutantji M, et al. Reliability of a revised NOTECHS scale for use in surgical teams. *Am J Surg* 2008;196:184-90.
44. Mishra A, Catchpole K, McCulloch P. The Oxford NOTECHS System: reliability and validity of a tool for measuring teamwork behaviour in the operating theatre. *Qual Saf Health Care* 2009;18:104-8.
45. Grober ED, Hamstra SJ, Wanzel KR, et al. The educational impact of bench model fidelity on the acquisition of technical skill: the use of clinically relevant outcome measures. *Ann Surg* 2004;240:374-81.
46. Sidhu RS, Park J, Brydges R, et al. Laboratory-based vascular anastomosis training: a randomized controlled trial evaluating the effects of bench model fidelity and level of training on skill acquisition. *J Vasc Surg* 2007;45:343-9.
47. Traxer O, Gettman MT, Napper CA, et al. The impact of intense laparoscopic skills training on the operative performance of urology residents. *J Urol* 2001;166:1658-61.
48. Fried GM, Derossis AM, Bothwell J, et al. Comparison of laparoscopic performance in vivo with performance measured in a laparoscopic simulator. *Surg Endosc* 1999;13:1077-81, discussion 1082.
49. Stelzer MK, Abdel MP, Sloan MP, et al. Dry lab practice leads to improved laparoscopic performance in the operating room. *J Surg Res* 2009;154:163-6.
50. Scott DJ, Bergen PC, Rege RV, et al. Laparoscopic training on bench models: better and more cost effective than operating room experience? *J Am Coll Surg* 2000;191:272-83.
51. Hamilton EC, Scott DJ, Kapoor A, et al. Improving operative performance using a laparoscopic hernia simulator. *Am J Surg* 2001;182:725-8.
52. Risucci D, Cohen JA, Garbus JE, et al. The effects of practice and instruction on speed and accuracy during resident acquisition of simulated laparoscopic skills. *Curr Surg* 2001;58:230-5.
53. Stefanidis D, Korndorffer Jr JR, Heniford BT, et al. Limited feedback and video tutorials optimize learning and resource utilization during laparoscopic simulator training. *Surgery* 2007;142:202-6.
54. Porte MC, Xeroulis G, Reznick RK, et al. Verbal feedback from an expert is more effective than self-accessed feedback about motion efficiency in learning new surgical skills. *Am J Surg* 2007;193:105-10.
55. Xeroulis GJ, Park J, Moulton CA, et al. Teaching suturing and knot-tying skills to medical students: a randomized controlled study comparing computer-based video instruction and (concurrent and summary) expert feedback. *Surgery* 2007;141:442-9.
56. Moulton CA, Dubrowski A, Macrae H, et al. Teaching surgical skills: what kind of practice makes perfect?: a randomized, controlled trial. *Ann Surg* 2006;244:400-9.
57. Matsumoto ED, Hamstra SJ, Radomski SB, et al. The effect of bench model fidelity on endourological skills: a randomized controlled study. *J Urol* 2002;167:1243-7.
58. Sherman V, Feldman LS, Stanbridge D, et al. Assessing the learning curve for the acquisition of laparoscopic skills on a virtual reality simulator. *Surg Endosc* 2005;19:678-82.
59. Watterson JD, Beiko DT, Kuan JK, et al. Randomized prospective blinded study validating acquisition of ureteroscopy skills using computer based virtual reality endourological simulator. *J Urol* 2002;168:1928-32.
60. Jacomides L, Ogan K, Cadeddu JA, et al. Use of a virtual reality simulator for ureteroscopy training. *J Urol* 2004;171:320-3, discussion 323.
61. Knoll T, Trojan L, Haecker A, et al. Validation of computer-based training in ureterorenoscopy. *BJU Int* 2005;95:1276-9.
62. Schreuder HW, van Dongen KW, Roeleveld SJ, et al. Face and construct validity of virtual reality simulation of laparoscopic gynecologic surgery. *Am J Obstet Gynecol* 2009 May;200(5):540.e1-8.
63. Aggarwal R, Tully A, Grantcharov T, et al. Virtual reality simulation training can improve technical skills during laparoscopic salpingectomy for ectopic pregnancy. *BJOG* 2006;113:1382-7.
64. Gor M, McCloy R, Stone R, et al. Virtual reality laparoscopic simulator for assessment in gynaecology. *BJOG* 2003;110:181-7.
65. Pearson AM, Gallagher AG, Rosser JC, et al. Evaluation of structured and quantitative training methods for teaching intracorporeal knot tying. *Surg Endosc* 2002;16:130-7.
66. Grantcharov TP, Funch-Jensen P. Can everyone achieve proficiency with the laparoscopic technique? Learning curve patterns in technical skills acquisition. *Am J Surg* 2009;197:447-9.
67. Gallagher AG, Lederman AB, McGlade K, et al. Discriminative validity of the Minimally Invasive Surgical Trainer in Virtual Reality (MIST-VR) using criteria levels based on expert performance. *Surg Endosc* 2004;18:660-5.
68. Wilhelm DM, Ogan K, Roehrborn CG, et al. Assessment of basic endoscopic performance using a virtual reality simulator. *J Am Coll Surg* 2002;195:675-81.
69. Knudsen BE, Matsumoto ED, Chew BH, et al. A randomized, controlled, prospective study validating the acquisition of percutaneous renal collecting system access skills using a computer based hybrid virtual reality surgical simulator: phase I. *J Urol* 2006;176:2173-8.
70. Hart R, Doherty DA, Karthigasu K, et al. The value of virtual reality-simulator training in the development of laparoscopic surgical skills. *J Minim Invasive Gynecol* 2006;13:126-33.
71. Lucas SM, Zeltser IS, Bensalah K, et al. Training on a virtual reality laparoscopic simulator improves performance of an unfamiliar live laparoscopic procedure. *J Urol* 2008;180:2588-91, discussion 2591.
72. Hyltander A, Liljegren E, Rhodin PH, et al. The transfer of basic skills learned in a laparoscopic simulator to the operating room. *Surg Endosc* 2002;16:1324-8.
73. Ahlberg G, Heikkinen T, Iselius L, et al. Does training in a virtual reality simulator improve surgical performance? *Surg Endosc* 2002;16:126-9.
74. Seymour NE, Gallagher AG, Roman SA, et al. Virtual reality training improves operating room performance: results of a randomized, double-blinded study. *Ann Surg* 2002;236:458-63; discussion 463-4.
75. Ahlberg G, Enochsson L, Gallagher AG, et al. Proficiency-based virtual reality training significantly reduces the error rate for residents during their first 10 laparoscopic cholecystectomies. *Am J Surg* 2007;193:797-804.
76. Larsen CR, Soerensen JL, Grantcharov TP, et al. Effect of virtual reality training on laparoscopic surgery: randomised controlled trial. *BMJ (Clinical research ed.)*. May 14 2009;338:b1802.
77. Kothari SN, Kaplan BJ, DeMaria EJ, et al. Training in laparoscopic suturing skills using a new computer-based virtual reality simulator (MIST-VR) provides results comparable to those with an established pelvic trainer system. *J Laparoendosc Adv Surg Tech A* 2002;12:167-73.
78. Munz Y, Kumar BD, Moorthy K, et al. Laparoscopic virtual reality and box trainers: is one superior to the other? *Surg Endosc* 2004;18:485-94.
79. Chou DS, Abdelshehid C, Clayman RV, et al. Comparison of results of virtual-reality simulator and training model for basic ureteroscopy training. *J Endourol* 2006;20:266-71.
80. McDougall EM, Kolla SB, Santos RT, et al. Preliminary study of virtual reality and model simulation for learning laparoscopic suturing skills. *J Urol* 2009;182:1018-25.
81. Gurusamy K, Aggarwal R, Palanivelu L, et al. Systematic review of randomized controlled trials on the effectiveness of virtual reality training for laparoscopic surgery. *Br J Surg* 2008;95:1088-97.
82. Gurusamy KS, Aggarwal R, Palanivelu L, et al. Virtual reality training for surgical trainees in laparoscopic surgery. *Cochrane Database Syst Rev* 2009; (1):CD006575.
83. Youngblood PL, Srivastava S, Curet M, et al. Comparison of training on two laparoscopic simulators and assessment of skills transfer to surgical performance. *J Am Coll Surg* 2005;200:546-51.
84. Hamilton EC, Scott DJ, Fleming JB, et al. Comparison of video trainer and virtual reality training systems on acquisition of laparoscopic skills. *Surg Endosc* 2002;16:406-11.
85. Panait L, Akkary E, Bell RL, et al. The role of haptic feedback in laparoscopic simulation training. *J Surg Res* 2009;156:312-6.
86. Kneebone RL, Kidd J, Nestel D, et al. Blurring the boundaries: scenario-based simulation in a clinical setting. *Med Educ* 2005;39:580-7.
87. Kneebone R, Nestel D, Yadollahi F, et al. Assessing procedural skills in context: Exploring the feasibility of an Integrated Procedural Performance Instrument (IPPI). *Med Educ* 2006;40:1105-14.
88. LeBlanc VR, Tabak D, Kneebone R, et al. Psychometric properties of an integrated assessment of technical and communication skills. *Am J Surg* 2009;197:96-101.
89. Panait L, Bell RL, Roberts KE, et al. Designing and validating a customized virtual reality-based laparoscopic skills curriculum. *J Surg Educ* 2008;65:413-7.
90. Munz Y, Almoudaris AM, Moorthy K, et al. Curriculum-based solo virtual reality training for laparoscopic intracorporeal knot tying: objective assessment of the transfer of skill from virtual reality to reality. *Am J Surg* 2007;193:774-83.
91. Aggarwal R, Grantcharov TP, Eriksen JR, et al. An evidence-based virtual reality training program for novice laparoscopic surgeons. *Ann Surg* 2006;244:310-4.

Correspondence to: Dr. Vanessa Palter, Division of General Surgery, Department of Surgery, St. Michael's Hospital, University of Toronto, 30 Bond St., Toronto ON M5B 1W8; vanessa.palter@utoronto.ca



The following videos are available online:
Virtual reality simulator for laparoscopic cholecystectomy:
www.cmaj.ca/video/chole1.wmv
Virtual reality simulator for tubal occlusion:
www.cmaj.ca/video/tubalocclusion.wmv