

Effects of Vasopressin and Prostaglandin E₁ on the Adenyl Cyclase–Cyclic 3',5'-Adenosine Monophosphate System of the Renal Medulla of the Rat

NAMA P. BECK, TOSHIO KANEKO, URIEL ZOR, JAMES B. FIELD, and
BERNARD B. DAVIS

From the Department of Medicine, The Oakland Veterans Administration Hospital, University of Pittsburgh School of Medicine, Pittsburgh, Pennsylvania 15213

ABSTRACT Vasopressin increased adenyl cyclase activity in homogenates of both inner and outer renal medulla of the rat. It also increased the concentration of cyclic 3',5'-adenosine monophosphate (AMP) in slices of both inner and outer medulla but not in renal cortex. In the inner medulla, a concentration of prostaglandin E₁ (PGE₁), which was ineffective by itself significantly reduced the stimulation of adenyl cyclase activity and cyclic AMP concentration induced by vasopressin. These results are consistent with the hypothesis that PGE₁ can compete with vasopressin for adenyl cyclase-binding sites. However, the findings in the outer medulla suggest the situation is more complex. Although 10⁻⁸ M PGE₁ had no effect by itself and inhibited the vasopressin-induced elevation of cyclic AMP, larger amounts of PGE₁ increased both adenyl cyclase activity and cyclic AMP levels. The maximum effect on the latter parameter was at least 6 times as great as that of maximum amounts of vasopressin.

INTRODUCTION

It has been proposed that cyclic 3',5'-adenosine monophosphate (cyclic AMP)¹ is the intracellular mediator of the physiological effects of vasopressin. This suggestion, first made by Orloff and Handler (1), has received considerable experimental support. Vasopressin raises the concentration of cyclic AMP in the toad bladder (2) and increases the activity of adenyl cyclase in homogenates of the medulla of the rat kidney (3).

Received for publication 22 January 1971 and in revised form 28 April 1971.

¹Abbreviations used in this paper: cyclic AMP, cyclic 3',5'-adenosine monophosphate; PGE₁, prostaglandin E₁.

Furthermore cyclic AMP, like vasopressin, increases water permeability and active sodium transport in the urinary bladder of the toad (1) and increases water permeability of the isolated perfused collecting duct of the rabbit (4).

Grantham and Orloff (5) observed that prostaglandin E₁ (PGE₁), by itself, increases the permeability of perfused collecting ducts to water but inhibits the effects of submaximal amounts of vasopressin. Although the effects of theophylline and PGE₁ are additive, PGE₁ does not influence the cyclic AMP-induced increases in water permeability of the collecting duct. To explain these observations, Grantham and Orloff proposed that PGE₁ stimulated adenyl cyclase activity and increased production of cyclic AMP. Furthermore, by competing with vasopressin for receptors which influence adenyl cyclase, but being a less potent activator than vasopressin, PGE₁ decreased vasopressin-mediated increases in cyclic AMP production. This finding and this hypothesis were in general agreement with Bergstrom's proposal that prostaglandins might act as modulators of the adenyl cyclase reaction (6). The present investigation was undertaken to evaluate further this concept by measuring the effects of vasopressin and PGE₁, singly and in combination, on the concentration of cyclic AMP and on adenyl cyclase activity in the renal medulla.

METHODS

Male Sprague-Dawley rats weighing 140–250 g were sacrificed by decapitation. Tissue slices weighing 10–75 mg were prepared from outer red medulla, inner white medulla, or cortex using a Stadie-Riggs microtome. Slices were incubated for 15 min at 37°C in 2 ml Krebs-Ringer bicarbonate buffer containing 0.5 mg/ml glucose, 2 mg/ml albu-




FIGURE 1 Represents a dose-response relationship between concentration of vasopressin on the ordinate and adenylyl cyclase activity on the abscissa. The outer medulla is the solid line, and the broken line is the inner medulla.

min, and 10 mM theophylline. The gas phase was 95% O₂ and 5% CO₂. PGE₁, vasopressin, or both were added to the appropriate flasks. 1 mg of PGE₁ was dissolved in 0.1 ml of absolute ethanol. To this was added 0.9 ml Na₂CO₃ (0.2 mg/ml, pH 7). This stock solution was diluted with Krebs-Ringer bicarbonate buffer to give the appropriate concentration in the flask. The final concentration of ethanol never exceeded 0.1% (v/v). This diluent had no effect on cyclic AMP concentration in medulla, control 5.5 ± 0.5 (SE) nmoles/g wet tissue and diluent 5.8 ± 1.3. The final concentration of ethanol in the incubation flask varied between 0.1 and 0.001% (v/v). Bovine arginine-vasopressin in aqueous solution was used. The potency units referred to in the text are international pressor units (USP). A molar solution of vasopressin would be the equivalent of 433.6 U/ml. The concentration of the test substance is indicated in the Results section. The incubation was terminated by homogenizing the slices in cold 5% trichloroacetic acid. Cyclic AMP was extracted by the method of Krishna, Weiss, and Brodie (7) and assayed by the method of

Kaneko and Field (8). When adenylyl cyclase activity was assayed, the inner white or outer red medulla was separated and homogenized separately in 2 M sucrose-0.05 M Tris buffer, pH 7.4. After centrifugation at 4°C at 2000 g for 15 min the precipitate was resuspended in an equal volume of 0.05 M Tris and centrifuged again at the same speed and for the same period of time. Adenylyl cyclase activity of the final precipitate was measured by the method of Chase and Aurbach (3). The protein concentration of the preparation was determined by the Lowry, Rosebrough, Farr, and Randall technique (9).

RESULTS

Adenylyl cyclase activity. Vasopressin increased adenylyl cyclase activity in both inner and outer medulla. Fig. 1 demonstrates the dose-response relationships between vasopressin concentration and adenylyl cyclase activity in both the inner and outer medulla. In both significant increases were produced by 10⁻³ U/ml and 10⁻¹ U/ml gave a maximal response.

PGE₁ in concentrations as high as 10⁻⁴ M had no effect on adenylyl cyclase activity in inner medulla but caused a significant increase in activity in the outer medulla (Table I). In the inner medulla, the addition of 10⁻⁴ M PGE₁, which by itself had no effect on adenylyl cyclase activity, together with a submaximal concentration of 10⁻³ U/ml vasopressin, resulted in less of an increase in adenylyl cyclase activity than was observed with that amount of vasopressin alone (Table I). The mean difference in 22 paired incubations between vasopressin alone and the combination was 14.0 ± 5.0 (SE) pmole cyclic AMP/mg protein per 15 min (*P* < 0.01). Also noted in Table I is the fact that 10⁻⁴ M PGE₁ had no effect on concentrations of vasopressin as high as 5 × 10² U/ml. 10⁻⁴ M PGE₁ had no apparent effect on vasopressin, 10⁻³ U/ml, in the outer medulla. In a separate group of experiments in the outer medulla, control adenylyl cyclase activity was 24.8 ± 3.9 SE. Vasopressin 10⁻³ U/ml increased this value to 46.6 ± 4.4 (*P* < 0.01), but 10⁻⁸ M PGE₁ had no measurable effect 25.4 ± 4.1. The adenylyl cyclase activity with the com-

TABLE I
Effects of Vasopressin and PGE₁ on Adenylyl Cyclase Activity (picomole Cyclic AMP/milligram Protein per 15 min)

Media	Inner medulla	Outer medulla
Control	83 ± 7 (21)	71 ± 2 (10)
PGE ₁ , 10 ⁻⁴ M	87 ± 6 (21)	104 ± 7 (10)
Vasopressin, 10 ⁻³ U/ml	134 ± 7 (22)*	229 ± 7 (10)
Vasopressin, 10 ⁻³ U/ml, and PGE ₁ 10 ⁻⁴ M	120 ± 7 (22)*	254 ± 18 (10)
Vasopressin, 5 × 10 ⁻³ U/ml	323 ± 9 (13)	
Vasopressin, 5 × 10 ⁻² U/ml, and PGE ₁ , 10 ⁻⁴ M	327 ± 10 (13)	

Values are the mean ± SE. The number of experiments are represented in the parentheses.

* The mean difference of 14 ± 5 (*P* < 0.01).

bination of vasopressin 10^{-8} U/ml and 10^{-8} M PGE_1 , 53.9 ± 3.6 , was not measurably different than vasopressin alone.

Cyclic AMP concentration. Vasopressin increased cyclic AMP in both inner and outer medulla, and there was a dose-response relationship between the amount of vasopressin added and cyclic AMP concentration in both (Figs. 2 and 3). As with adenylyl cyclase activity, vasopressin 10^{-8} U/ml appeared to have submaximal effect in both inner and outer medulla. In the inner medulla PGE_1 in concentrations as high as 10^{-4} M had no measurable effect on cyclic AMP concentration (Table II). In the outer medulla PGE_1 increased cyclic AMP concentration, and as is apparent in Fig. 4 there was a dose-response relationship between the amount of PGE_1 added and the measured concentration of cyclic AMP. 10^{-7} M PGE_1 was the lowest concentration noted to effect tissue cyclic AMP concentration. The highest concentration of PGE_1 tested (10^{-8} M) raised tissue cyclic AMP concentration to 131.0 ± 10.8 , a value over 6-fold the maximum amount measured with vasopressin in the outer medulla.

In the inner medulla 10^{-6} M PGE_1 , which alone had no effect on cyclic AMP concentration, decreased the effect of submaximal vasopressin, 10^{-8} U/ml, to increase tissue cyclic AMP concentration, but had no effect on concentrations of vasopressin as high as 5×10^{-2} U/ml (Table III). The combination of 10^{-6} M PGE_1 , which by itself increased cyclic AMP concentration in the outer medulla, and vasopressin 10^{-8} U/ml increased cyclic AMP concentration more than vasopressin alone. However, 10^{-8} M PGE_1 , which by itself is ineffective, had an inhibitory effect on vasopressin, 10^{-8} U/ml. (Table III).

In contrast with the effects of vasopressin in the renal medulla, this hormone had no effect on the renal cortex where the mean control cyclic AMP concentra-




FIGURE 2 The dose-response curve of vasopressin concentration (ordinate) and cyclic AMP concentration (abscissa) in the outer medulla.

TABLE II
Effect of PGE_1 on Cyclic AMP Concentration (nanomoles/gram wet tissue) in Inner Medulla

Media	Inner medulla
Control	2.2 ± 0.3 (16)*
PGE_1 , 10^{-6} M	2.1 ± 0.2 (16)
Control	3.5 ± 0.7 (10)*
PGE_1 , 10^{-4} M	3.6 ± 0.3 (12)

Values are the mean \pm SE. The numbers of experiments are represented in the parentheses.

* Control values are different because experiments were done on separate days.

tion of 2.2 ± 0.3 was not significantly altered by vasopressin 10^{-2} U/ml (2.6 ± 0.4).

DISCUSSION

Handler, Butcher, Sutherland, and Orloff reported that vasopressin increased cyclic AMP concentration in the anuran bladder (2). Previous results in the kidney, however, have been variable. Brown, Clarke, Roux, and Sherman reported that vasopressin increased cyclic AMP concentration in the particulate fraction of a homogenate of the dog kidney (10). They noted, however, that the cortex was more sensitive to stimulation by vasopressin than the medulla. Senft, Hoffmann, Munske, and Schultz documented an increase in cortical and outer medullary cyclic AMP with dehydration, but the concentration in




FIGURE 3 The dose-response relationship between vasopressin concentration (ordinate) and cyclic AMP concentration (abscissa) in the inner medulla.




FIGURE 4 Represents the dose-response relationship between PGE₁ concentration on the ordinate and cyclic AMP concentration on the abscissa.

the inner medulla was below the sensitivity of their method (11). Chase and Aurbach studied hormonal effects on adenylyl cyclase activity in cortex and medulla in vitro (3). The medulla was much more responsive to vasopressin while the cortex was much more sensitive to parathyroid hormone.

In the present experiments vasopressin significantly increased adenylyl cyclase activity and tissue concentration of cyclic AMP in inner and outer medulla but had no effect on cyclic AMP concentration in cortex. These findings further support the concept of Orloff and Handler that the physiological actions of vasopressin on responsive epithelial structures are mediated by cyclic AMP (1).

The prostaglandins influence the adenylyl cyclase-cyclic AMP system in several tissues in various ways. PGE₁ increased tissue cyclic AMP concentration in pituitary (12) and thyroid (13) but inhibited the effects of epinephrine on cyclic AMP concentration and free fatty acid mobilization from adipose tissue (14). PGE₁ increases the permeability of the isolated perfused collecting duct to water (5). The PGE₁-induced increases in adenylyl cyclase activity and tissue cyclic AMP concentration in the outer medulla are consistent with this hydro-osmotic effect of PGE₁ being mediated by the adenylyl cyclase-cyclic AMP system. In the isolated collecting duct, PGE₁ decreases the effect of vasopressin to augment water permeability (5). The inhibition of vasopressin-induced increases in adenylyl cyclase activity and cyclic AMP concentration in the inner medulla suggests that PGE₁ inhibits the physiological effect of vasopressin by competing with vasopressin for adenylyl cyclase re-

TABLE III

Effects of Vasopressin and PGE₁ on Cyclic AMP Concentration (nanomoles/gram wet tissue)

Media	Inner medulla	Outer medulla
Control	3.5 ± 0.7 (10)	3.8 ± 0.5 (13)
PGE ₁ , 10 ⁻⁸ M		4.2 ± 0.8 (13)
PGE ₁ , 10 ⁻⁶ M	3.6 ± 0.3 (12)	13.0 ± 0.7 (5)
Vasopressin, 10 ⁻³ U/ml	10.6 ± 0.8 (14)	11.0 ± 1.3 (22)*
Vasopressin, 10 ⁻³ U/ml, and PGE ₁ , 10 ⁻⁸ M		7.0 ± 1.2 (22)*
Vasopressin, 10 ⁻³ U/ml, and PGE ₁ , 10 ⁻⁶ M	7.3 ± 0.9 (14)‡	15.8 ± 2.5 (6)§
Vasopressin, 5 × 10 ⁻² U/ml	36.8 ± 11.8 (12)	
Vasopressin, 5 × 10 ⁻² U/ml, and PGE ₁ , 10 ⁻⁶ M	39.1 ± 5.2 (12)	

Values are the mean ± SE. The number of experiments are represented in the parentheses.

* The increases from the control are $\Delta 7.2 \pm 0.8$ and $\Delta 4.8 \pm 1.4$, and the difference is significant $P < 0.01$.

‡ $P < 0.01$ when compared with vasopressin alone.

§ $P < 0.05$ when compared with vasopressin alone.

ceptors. In addition, the inhibition of vasopressin-induced increases in cyclic AMP concentration by an amount of 10^{-8} M PGE₁, which by itself had no effect on cyclic AMP concentration, is consistent with vasopressin-PGE₁ competition for adenylyl cyclase receptors. It should be emphasized that the PGE₁ inhibition of vasopressin-induced increases in adenylyl cyclase activity and cyclic AMP concentration was not complete. The amount of increase in cyclic AMP concentration caused by vasopressin even in the presence of PGE₁, may be sufficient to exert maximal physiological effect. Conversely, the PGE₁ inhibition of the hydro-osmotic effect of vasopressin is not complete (5). Providing these in vitro data have physiological significance all of these findings then support the Grantham and Orloff hypothesis with respect to both the mechanism of action of PGE₁ itself on the kidney and the means by which it interferes with the action of vasopressin. However, Grantham and Orloff also suggested PGE₁ was a less potent activator of adenylyl cyclase than vasopressin (5). The observation that in outer medulla PGE₁ caused as much as a 6-fold greater increase in cyclic AMP concentration than did maximal amounts of vasopressin does not indicate that PGE₁ is less effective in stimulating adenylyl cyclase than is vasopressin. In addition the combination of PGE₁ in amounts which had an effect on cyclic AMP concentration with vasopressin, actually caused a greater increase in cyclic AMP concentration than did either agent alone. These findings suggest that in outer medulla PGE₁ is either a more effective activator of adenylyl cyclase or there are at least two species of adenylyl cyclase receptors in the outer medulla. If the latter is true then at least one species of receptor might be responsive to PGE₁ but not to vasopressin, and the additive effects of vasopressin and PGE₁ in the outer medulla might not be inconsistent with the Grantham-Orloff hypothesis (5).

ACKNOWLEDGMENTS

This research was supported by U. S. Public Health Service grants AM206865 and AM-11911 and Health Research and Services Foundation grant M82.

REFERENCES

1. Orloff, J., and J. S. Handler. 1962. The similarity of effects of vasopressin, adenosine-3',5'-phosphate (cyclic AMP) and theophylline on the toad bladder. *J. Clin. Invest.* **41**: 702.
2. Handler, J. S., R. W. Butcher, E. W. Sutherland, and J. Orloff. 1965. The effect of vasopressin and of theophylline on the concentration of adenosine 3',5'-phosphate in the urinary bladder of the toad. *J. Biol. Chem.* **240**: 4524.
3. Chase, L. R., and G. D. Aurbach. 1968. Renal adenylyl cyclase: anatomically separate sites for parathyroid hormone and vasopressin. *Science (Washington)*. **159**: 545.
4. Grantham, J. J., and M. B. Burg. 1966. Effect of vasopressin and cyclic AMP on permeability of isolated collecting tubules. *Amer. J. Physiol.* **211**: 255.
5. Grantham, J. J., and J. Orloff. 1968. Effect of prostaglandin E₁ on the permeability response of the isolated collecting tubule to vasopressin, adenosine 3',5'-monophosphate and theophylline. *J. Clin. Invest.* **47**: 1154.
6. Bergström, S. 1967. Prostaglandins: members of a new hormonal system. *Science (Washington)*. **157**: 382.
7. Krishna, G., B. Weiss, and B. B. Brodie. 1968. A simple, sensitive method for the assay of adenylyl cyclase. *J. Pharmacol. Exp. Ther.* **163**: 379.
8. Kaneko, T., and J. B. Field. 1969. A method for determination of 3',5'-cyclic adenosine monophosphate based on adenosine triphosphate formation. *J. Lab. Clin. Med.* **74**: 682.
9. Lowry, O. H., N. J. Rosebrough, A. L. Farr, and R. J. Randall. 1951. Protein measurement with the Folin phenol reagent. *J. Biol. Chem.* **193**: 265.
10. Brown, E., D. L. Clarke, V. Roux, and G. H. Sherman. 1963. The stimulation of adenosine 3',5'-monophosphate production by antidiuretic factors. *J. Biol. Chem.* **238**: 852.
11. Senft, G., M. Hoffmann, K. Munske, and G. Schultz. 1968. Effects of hydration and dehydration on cyclic adenosine 3',5'-monophosphate concentration in the rat kidney. *Pfluegers Arch.* **297**: 348.
12. Zor, U., T. Kaneko, H. P. G. Schneider, S. M. McCann, and J. B. Field. 1970. Further studies of stimulation of anterior pituitary cyclic adenosine 3',5'-monophosphate formation by hypothalamic extract and prostaglandins. *J. Biol. Chem.* **245**: 2883.
13. Kaneko, T., U. Zor, and J. B. Field. 1969. Thyroid-stimulating hormone and prostaglandin E₁ stimulation of cyclic 3',5'-adenosine monophosphate in thyroid slices. *Science (Washington)*. **163**: 1062.
14. Butcher, R. W., and C. E. Baird. 1968. Effects of prostaglandins on adenosine 3',5'-monophosphate levels in fat and other tissues. *J. Biol. Chem.* **243**: 1713.