

Published in final edited form as:

Acc Chem Res. 2010 August 17; 43(8): 1125–1134. doi:10.1021/ar100030m.

Molecular Factors Controlling Photosynthetic Light-Harvesting by Carotenoids


Tomáš Polívka^{1,2} and Harry A. Frank³

¹Institute of Physical Biology, University of South Bohemia, Zámek 136, 373 33 Nové Hradý Czech Republic

²Biological Centre, Czech Academy of Sciences, České Budějovice Czech Republic

³Department of Chemistry, 55 North Eagleville Road, University of Connecticut, Storrs, CT 06269-3060 USA

Abstract



Carotenoids are naturally-occurring pigments that absorb light in the spectral region in which the sun irradiates maximally. These molecules transfer this energy to chlorophylls, initiating the primary photochemical events of photosynthesis. Carotenoids also regulate the flow of energy within the photosynthetic apparatus and protect it from photo-induced damage caused by excess light absorption. To carry out these functions in nature, carotenoids are bound in discrete pigment-protein complexes in close proximity to chlorophylls. A few 3D structures of these carotenoid complexes have been determined by X-ray crystallography. Thus, the stage is set for attempting to correlate the structural information with the spectroscopic properties of carotenoids to understand the molecular mechanism(s) of their function in photosynthetic systems.

In this Account, we summarize current spectroscopic data describing the excited state energies and ultrafast dynamics of purified carotenoids in solution and bound in light-harvesting complexes from purple bacteria, marine algae, and green plants. Many of these complexes can be modified using mutagenesis or pigment exchange which facilitates making the correlations between structure and function. We describe the structural and electronic factors controlling the function of carotenoids as energy donors. We also discuss unresolved issues related to the nature of spectroscopically dark excited states, which could play a role in light-harvesting.

To illustrate the interplay between structural determinations and spectroscopic investigations that exemplifies work in the field, we describe the spectroscopic properties of four light-harvesting complexes whose structures have been determined to atomic resolution. The first, the LH2 complex from the purple bacterium *Rhodospseudomonas acidophila*, contains the carotenoid, rhodopin glucoside. The second is the LHCII trimeric complex from higher plants which uses the carotenoids, lutein, neoxanthin and violaxanthin to transfer energy to chlorophyll. The third, the peridinin-chlorophyll-protein (PCP) from the dinoflagellate, *Amphidinium carterae*, is the only known complex where the bound carotenoid (peridinin) pigments outnumber the chlorophylls. The last is xanthorhodopsin from the eubacterium, *Salinibacter ruber*. This complex contains the carotenoid, salinixanthin, which transfers energy to a retinal chromophore. The carotenoids in

these pigment-protein complexes transfer energy with high efficiency by optimizing both the distance and orientation of the carotenoid donor and chlorophyll acceptor molecules.

Importantly, the versatility and robustness of carotenoids in these light-harvesting pigment-protein complexes have led to their incorporation in the design and synthesis of nanoscale antenna systems. In these bio-inspired systems, researchers are seeking to improve the light capture and use of energy from the solar emission spectrum.

Introduction

Carotenoids are a group of natural pigments whose central structural feature is a linear chain of alternating C-C and C=C bonds.¹ They differ in π -electron conjugation length (number of conjugated double bonds, N), and in the type and number of functional groups attached to the carbon backbone. Of the more than one thousand naturally-occurring carotenoids, only ~50 play a light-harvesting role in photosynthesis. The structures of four such carotenoids are shown in Figure 1.

The ability of carotenoids to act as light-harvesting agents is inextricably linked to their spectroscopic properties which are best described using a three-state model consisting of the ground state, S_0 , and two excited states denoted S_1 and S_2 (Figure 1). The symmetry of the fully-extended conjugated π -electron backbone places carotenoids in the C_{2h} point group, and a further consideration of the symmetry of the individual π orbitals assigns the S_0 and S_1 states to the A_g^- irreducible representation. The S_2 state has B_u^+ symmetry. Because one-photon transitions between electronic states of the same symmetry are forbidden by quantum mechanical selection rules, the $S_0 \rightarrow S_1$ transition is forbidden, and the lowest energy allowed transition is $S_0 \rightarrow S_2$.^{2,3} It should be noted that a few other states into which transitions from S_0 are forbidden may exist either between or in the vicinity of S_2 and S_1 , further complicating the carotenoid photophysics.⁴ After absorption of light via the $S_0 \rightarrow S_2$ transition, the S_2 state relaxes within a few hundred femtoseconds to the S_1 state whose lifetime depends on N and is between 1 and 200 ps for most natural carotenoids.³

At a first glance, carotenoids are not pigments that one would expect nature to choose as light-harvesting molecules because the lifetimes of the S_1 and S_2 states are significantly shorter than those of other naturally-occurring pigments such as (B) Chls; any possible energy transfer route using the S_1 or S_2 state of carotenoids as an energy donor must contend with these very short intrinsic relaxation times. Moreover, a transition between the ground state and S_1 state is forbidden, resulting in a negligible transition dipole moment, the consequence of which is that the S_1 state cannot participate in Förster-type dipole-dipole-mediated energy transfer. Thus, the sub-picosecond lifetime of the S_2 state and negligible dipole moment of the S_1 state would appear to hardly qualify carotenoids as effective light-harvesting molecules. However, essentially all photosynthetic organisms utilize carotenoids as light-harvesting pigments, and both the S_1 and S_2 states of carotenoids function as energy donors.

High efficiency of carotenoid-mediated energy transfer in light-harvesting proteins is achieved by optimization of the distance and orientation of the carotenoid donor and the (B)Chl acceptor. In most light-harvesting proteins, the distance between the conjugated systems of the donor and acceptor is between 3–10 Å (Figure 2). Combined with a proper orientation and a large dipole moment of the S_2 state, it was calculated that the S_2 -mediated carotenoid-to-(B) Chl energy transfer occurs in 100–300 fs via the Förster mechanism;⁵ i.e. clearly competitive with the intrinsic carotenoid S_2 lifetime. However, due to the fact that the donor-acceptor distance is smaller than dimensions of participating molecules, the donor-acceptor interaction term in the Förster formalism must be calculated using elaborate

quantum mechanical methods.⁶ Energy transfer from the carotenoid S_1 state can also be considered to proceed via a Förster mechanism, but due to the dipole-forbidden nature of this state it is necessary to compute the full Coulomb coupling between the donor and acceptor molecules to achieve reasonable agreement with experiment.⁶

Whether the S_1 or S_2 state acts as the energy donor depends in large part on N , because this is a primary factor in determining the energies of the states relative to those of the energy acceptor. Energy transfer according to the Förster mechanism will be optimized when the spectral overlap between the S_1 and/or S_2 donor emission and acceptor absorption is maximized. The N -dependence of the energies of the S_1 and S_2 states, together with energies of acceptors in various light-harvesting systems, is shown in Figure 3. It is clear from the figure that an energy transfer pathway between the carotenoid S_2 state and Q_x state of (B)Chl is favorable for a broad range of conjugation lengths. Indeed, the S_2 -mediated energy transfer operates in almost all carotenoid-containing light-harvesting systems regardless of the conjugation length of the carotenoid. In contrast, the S_1 state can act as an efficient energy donor only if the acceptor has a lower energy, as it does in light-harvesting complexes utilizing BChl-*a*. If the energy acceptor is Chl-*a*, only carotenoids having short π -electron conjugations will be able to use the S_1 pathway. Also, while the S_1 energy is nearly insensitive to the environment, the S_2 energy may be modulated by interaction with proteins. In addition, the effective conjugation length, and hence the excited state energies, of the carotenoid can be altered by protein binding site-induced structure changes to the molecule. This is particularly likely if the carotenoid has extended π -electron conjugation that may either be planarized with the linear chain or twisted in a way that decouples terminal π -bond(s) from the conjugation completely, resulting in changes in effective conjugation length. These effects produce diverse energy transfer pathways and efficiencies in light-harvesting proteins described in the following sections.

Purple bacterial antenna complexes

The exemplary system for energy transfer between carotenoids and BChl-*a* is the LH2 antenna of purple bacteria (Figure 2a).⁷ The structure of this complex revealed that the fundamental building block is an $\alpha\beta$ -polypeptide subunit pair that binds two strongly-coupled BChl-*a* molecules absorbing at ~ 850 nm (B850), one monomeric BChl-*a* molecule having an absorption band at 800 nm (B800), and one carotenoid molecule spanning the membrane in close contact with both the B800 and B850 molecules.

The carotenoid composition varies substantially among species of purple bacteria, but linear carotenoids with conjugation lengths $N=9-13$ are predominant. All LH2 complexes studied so far exhibit ultrafast energy transfer from the S_2 state as initially reported by Shreve et al.⁸ Many studies have demonstrated that the S_2 state has a sub-100 fs lifetime in LH2, corresponding to S_2 -mediated energy transfer route in LH2 operating with an efficiency of 40–60 % in LH2 complexes with carotenoids having $N=9-12$.^{5,9,10} Using LH2 complexes lacking the B800 BChl-*a*, a branching ratio of $\sim 2:3$ for energy transfer to B800 and B850 BChl-*a* molecules was determined in LH2 of *Rps. acidophila*.⁹ The experimentally-measured energy transfer rates were successfully reproduced by calculations invoking the BChl-*a* excited state associated with the Q_x absorption band as the energy acceptor.^{5,11}

An S_2 -mediated energy transfer channel was also reported in the LH1 complex from purple bacteria, which is the inner antenna system surrounding reaction center. In this complex S_2 -mediated energy transfer ranges from 60 to 70% for neurosporene ($N=9$), drops to about 50% for carotenoids with $N=10-11$, and further decreases to 40% and 30% for carotenoids with $N=12$ and $N=13$, respectively.¹²

The origin of this dependence lies in the fact that the intrinsic S_2 lifetime that becomes shorter with increasing N .¹³ Consequently, in longer carotenoids, S_2 -mediated energy transfer must compete with shorter intrinsic S_2 lifetimes, resulting in less efficient energy transfer, even though the energy transfer rates remain nearly independent of N .¹² Alternative hypotheses invoking so-called “dark” states in energy transfer have been also suggested (Figure 1),¹⁰ but the role of these states remains the subject of considerable debate. (For a recent review of this topic see Polívka and Sundström.⁴) An interesting issue is that only minor changes in spectral overlap between the carotenoid S_2 emission and Q_x absorption are expected at room temperature due to the broad S_2 emission profile. However, recent experiments at 10K demonstrated a significant decrease in S_2 energy transfer efficiency compared to the value at room temperature.¹⁴ This decrease was proposed to be caused by a narrowing of the emission and absorption bands at low temperature.

As suggested from Figure 3, energy transfer from the S_1 state is also possible. However, even though the S_1 energies of most carotenoids are higher than energies of the S_1 states of BChl-*a*, the S_1 -mediated energy transfer route drops off precipitously for carotenoids with $N > 10$.^{14–16} When neurosporene ($N=9$) is present in LH2, the efficiency is 92–95%. It drops below 90% for spheroidene ($N=10$) and is around 80% for spheroidenone which has the same number of C=C bonds as spheroidene, but with the conjugation extended to a carbonyl (C=O) group. The efficiency drops below 20% for $N=11$. A very similar N -dependence was found for LH1 reconstituted with different carotenoids.¹² Carotenoid-to-BChl energy transfer times involving the S_1 state were reported to vary between 1–2 ps,^{14–16} which are about order of magnitude slower than for the S_2 channel.

It is tempting to assign the absence of the S_1 channel in longer carotenoids as due to the B800 pigment being at too high an energy to act as an energy acceptor, but this explanation does not hold, because both B800 and B850 accept energy from the carotenoid S_1 state.¹⁷ In addition, the drop in energy transfer in going from $N=10$ to $N=11$ was also observed for LH1 complexes which have no B800.¹² Moreover, the S_1 channel exhibits essentially no temperature dependence,¹⁴ indicating that factors besides spectral overlap are involved. A possibility is that the sudden drop in energy transfer is caused by opening a new channel involving the carotenoid S^* state (Figure 1), which is able to transfer energy to BChl with only low efficiency, but it is also known to be a precursor of carotenoid triplet state formation.¹⁸ Since the triplet yield increases with increasing N ,¹⁹ the absence of significant S_1 -mediated energy transfer for $N > 10$ may be attributed to population being funneled to the S^* state.

Another interesting proposal was offered by Ritz et al.²⁰ who calculated carotenoid-BChl-*a* interaction energies and showed that strength of this interaction depends on whether the methyl groups of the carotenoids are positioned asymmetrically or symmetrically with respect to the center of the π -electron conjugation in the molecule. Symmetrical positioning of methyl groups, which occurs in carotenoids with $N=11$, is computed to diminish the interaction and lead to a drop in energy transfer efficiency. Although this finding has never been experimentally rationalized, it remains an intriguing idea regarding how energy transfer in purple bacterial antenna may be controlled.

Antenna systems of green plants

Green plants assemble a complicated network of Chl-*a*-containing antenna pigment-protein complexes. Energetic considerations (Figure 3) suggest that the S_1 pathway will be largely suppressed whereas energy transfer from the S_2 state to the Q_x state of Chl-*a* will be favored. Indeed, the lack of energy transfer via the S_1 state of carotenoids has been confirmed experimentally, but its absence is, in some Chl-*a*-based antenna, compensated by

a very efficient S_2 pathway. The most abundant antenna complex of this type is the LHCII protein (Figure 2) which resides on the outermost periphery of the Photosystem II reaction center. Besides Chl-*a*, LHCII contains Chl-*b*, two luteins (N=10), one neoxanthin (N=9) and one violaxanthin (N=9).²¹ In this complex, the S_2 pathway of carotenoid-to-Chl energy transfer is utilized almost exclusively. The S_2 lifetimes of lutein and neoxanthin in LHCII are in the sub-100 fs range, resulting in efficiencies of 60–70%.^{22–24} The key role of both lutein molecules in carotenoid-to-Chl energy transfer has been clearly established, and energy transfer from the S_2 state of neoxanthin has been also demonstrated, but reports of its efficiency span ranges from <10%²³ to >50%.²⁴ This uncertainty originates from the inability to selectively excite carotenoids independently from Chls in LHCII. Also, the question of whether Chl-*b* or Chl-*a* molecules are primary acceptors has been extensively debated.^{22–24} Carotenoid absorption overlapping the Soret band of Chl-*b* complicates the analysis. Nevertheless, it is now clear that some fraction of acceptors are Chl-*b*, but the precise ratio of Chl-*a* and Chl-*b* acceptors remains an open question. A very similar pattern of carotenoid-to-Chl energy transfer was described in the peripheral Lhca4 complex of PSI from plants, which is similar to LHCII but contains only lutein and violaxanthin.²⁵ Studies of PSI and PSII core antenna that contain only β -carotene have also revealed an active S_2 pathway.^{26–27} Tuning the interaction between β -carotene and Chl-*a* can change the efficiency of the S_2 pathway significantly, as shown by efficiencies ranging from nearly 60% in the PSI core to 30% in the inner antennae of PSII.^{26–27}

Energy transfer from the carotenoid S_1 state is less favorable due to the high energy of the Chl acceptor states. The S_1 pathway to Chl-*a* may be marginally active in complexes containing the shorter carotenoids, neoxanthin, violaxanthin and lutein. Efficiencies reported for energy transfer from the S_1 state of these carotenoids did not exceed 15% in LHCII and CP29 complexes.^{22–24} A slightly higher efficiency of ~20%, caused likely by the presence of low-energy Chl-*a* molecules in this complex, was found in Lhca4 peripheral antenna of plant PSI.²⁵ On the other hand, no evidence for the S_1 -mediated channel was found in LHCII complexes reconstituted with carotenoids having N=9–11.²⁸ Similarly, no energy transfer from the S_1 state was found in CP43 and CP47 complexes containing exclusively β -carotene.^{26–27} In the PSI core that also contains only β -carotene, but has low-energy Chl-*a* molecules, some activity of the S_1 -mediated channel was reported,^{27–29} but its efficiency did not exceed 20%. Interestingly, the inability to transfer energy from the S_1 state in some antenna complexes has been proposed to be compensated by an energy transfer channel using vibrationally hot S_1 as an energy donor. This pathway was suggested by two-photon excitation experiments which allow for direct excitation of the carotenoid into its S_1 state.³⁰ A sub-picosecond rise of Chl-*a* emission after two-photon excitation of the S_1 state in the LHCII complex was reported and assigned to energy transfer between a vibrationally hot S_1 state of lutein and Chl-*a*.³⁰ The same channel was also reported in the PSI core where it likely accounts for ~20% of the total energy transfer.²⁹

Although it seems that the role of the carotenoid S_1 state in light-harvesting is only minor in plants and green algae, it may have a key function in regulating energy flow within antenna complexes. It has been proposed that the S_1 state of some carotenoids may be low enough to quench excited Chl-*a* via energy transfer to the carotenoid S_1 state.³¹ Evidence of carotenoid S_1 state population after excitation of Chl-*a* has been reported in a transient absorption experiment,³² supporting the notion that a reverse energy transfer channel is possible. Whether this pathway constitutes the long-sought mechanism of non-photochemical quenching in higher plants remains an open question. Indeed, several alternatives have also been proposed.^{33–35}

Antenna of marine algae

Another group of proteins utilizing carotenoids as light-harvesting pigments derives from marine algae that employ predominantly peridinin and fucoxanthin which contain a conjugated carbonyl group. This moiety causes the spectroscopic features and excited state dynamics of the carotenoid to be dependent on the polarity of the environment.^{36,37} The dependence on polarity is caused by the electron-withdrawing nature of the conjugated carbonyl which leads to a stabilization of an intramolecular charge-transfer (ICT) state in polar environments.^{36,37} The major effect of the ICT state is that it modulates the lifetime of the S_1 state with which it is electronically coupled. Besides the effect on the S_1 lifetime which becomes shorter in a polar environment, the S_2 - S_1 energy gap decreases when a carbonyl group is introduced. The S_2 energy of carbonyl carotenoids is significantly lower than in their non-carbonyl counterparts, but the S_1 energy remains largely unaffected (Figure 3). Thus, the coupled S_1 /ICT state is high enough to keep favorable spectral overlap with the Q_y band of Chl-*a*, while the S_2 state is shifted to lower energies to allow efficient capturing of green light of vital importance for underwater organisms.³⁷

An excellent system utilizing this light-harvesting strategy is the water-soluble peridinin-Chl-*a* protein (PCP) from the dinoflagellate *Amphidinium carterae*.^{38,39} Its structure (Figure 2) has been refined recently to 1.5 Å.⁴⁰ Contrary to other light-harvesting systems, the pigment stoichiometry in PCP is dominated (4:1 carotenoid-to-Chl) by the carotenoid, peridinin. Peridinin absorbs light in the 450–550 nm region and transfer energy to Chl-*a* with an efficiency of ~90%.⁴¹ The S_2 state in PCP provides a channel that accounts for 20–30% of the total energy transfer efficiency.⁴² The dominant energy transfer channel is via the S_1 /ICT state which is more than 80% efficient.^{42,43} The lifetime of the S_1 /ICT state of peridinin in PCP is 2.5 ps⁴² which is shorter by about a factor of five than in polar solvents. Using these data the efficiency of the S_1 /ICT channel was determined from the total energy transfer efficiency measured by fluorescence excitation spectroscopy.⁴⁴ It was found that the S_1 /ICT-mediated energy transfer rate constant in native PCP is $\sim(3 \text{ ps})^{-1}$, which is consistent with an intrinsic S_1 /ICT lifetime of peridinin in the PCP complex of ~16 ps.⁴²

A major advantage of working with the PCP complex is that it is amenable to site-directed mutagenesis and reconstitution with different pigments.^{40,45} Reconstitution of PCP with various Chls alters the position of the Q_y band from 650 nm (Chl-*b*) to 790 nm (BChl-*a*) providing a systematic variation of spectral overlap.⁴⁶ Experiments on these reconstituted complexes confirmed the applicability of the Förster mechanism, and also showed that when the Q_y band of Chl is close to the maximum of the peridinin S_1 /ICT emission, the energy transfer efficiency is even better than in native PCP.^{44,46}

Site-directed mutagenesis carried out on specific amino acid residues in the PCP demonstrated that it is a robust system in that the peridinin-to-Chl energy transfer efficiency was found to be insensitive to changes in the local protein environment of the pigments.⁴⁰ Yet, while energy transfer efficiency remained largely unaffected, the change of a single amino acid (Asn-89 to Leu-89) dramatically altered the absorption spectrum of the complex: The absorption band of the longest wavelength-absorbing peridinin was shifted 24 nm to shorter wavelength despite no change in the structure of the PCP.⁴⁰ This clearly demonstrates how the pigment environment can be engineered by nature to achieve broad absorption that expands light-capture by the organism into the green spectral region.

The light-harvesting strategy of utilizing spectroscopic properties of carbonyl carotenoids also manifests itself in a membrane-bound antenna protein called LHC.⁴⁷ This protein is related to the Lhc family of plant protein complexes, but binds Chl-*a*, Chl-*c* and the carotenoids, peridinin and diadinoxanthin. In this complex, peridinin transfers energy to

Chl-*a* primarily via the S_1 /ICT state, and the energy transfer rate is essentially identical to that in PCP.⁴⁷ Very similar behavior was reported in a related protein from the diatom *Cyclotella meneghiniana* that binds fucoxanthin instead of peridinin.⁴⁸

Other systems

Besides the three examples of antenna proteins described above, carotenoids have also shown their light-harvesting capability in the large BChl-*c,d,e*-containing antenna of green sulfur bacteria known as chlorosomes. These contain predominantly the carotenoids, chlorobactene and isorenieratene possessing aryl terminal rings. Most likely due to tight packing of pigments within the chlorosomes, isorenieratene exhibits efficient energy transfer from the S_2 state. Efficiencies over 60% in *Chlorobium phaeobacteroides* have been reported.⁴⁹ No evidence for S_1 -mediated energy transfer has been found, perhaps due to the fact that spectral overlap between the S_1 emission of isorenieratene, which has nearly identical spectroscopic properties as its non-phenolic counterpart β -carotene,⁵⁰ and the Q_y absorption of BChl-*e* is small.

A completely different energy transfer scheme was reported for the B808-866 antenna complex of the green bacterium *Chloroflexus aurantiacus*. This antenna system, which contains BChl-*a* and γ -carotene, is similar to the LH2 and LH1 complexes of purple bacteria, but unlike these antennas, the efficiency of carotenoid-to-BChl energy transfer is only 15%. The energy transfer pathway occurs from the S_1 state of γ -carotene.⁵¹ Interestingly, no S_2 -mediated pathway was observed, making the B808-866 complex the only known antenna that employs exclusively the S_1 -pathway.

Light-harvesting by carotenoids was also described in a novel system that does not contain (B)Chl. This is the so-called xanthorhodopsin from *Salinibacter ruber* which belongs to the large group of retinal-based energy transducers.⁵² A few these proteins from archae showed association with carotenoids, but only in xanthorhodopsin which binds the carotenoid, salinixanthin, has it been demonstrated that the carotenoid is able to transfer energy to retinal.⁵² Due to the high energy of the acceptor state ($\sim 17,200\text{ cm}^{-1}$), only the S_2 channel is active, and it has an efficiency of 40%.⁵³

Synthetic antennas

The versatility and robustness of carotenoids in naturally-occurring light-harvesting systems, summarized in Figure 4, have not gone unnoticed in attempts to design and synthesize nanoscale systems seeking to mimic and perhaps even improve upon the efficiency of the natural systems in light-capture and solar energy conversion. A series of carotenoid-pyropheophorbide dyads having either zeaxanthin or fucoxanthin as the energy donor were reported to reproduce qualitatively the energy transfer efficiency observed in the natural systems.⁵⁴ For zeaxanthin only the S_2 pathway was active, and efficiencies up to 15% did not come close to matching those of natural systems. With fucoxanthin as an energy donor, which has a higher S_1 energy than zeaxanthin, pyropheophorbide became accessible via the S_1 channel. The resulting efficiencies were 15–45%, depending on the orientation of the donor and acceptor. Pyropheophorbide as an energy acceptor was used again in a subsequent study employing peridinin and fucoxanthin as energy donors.⁵⁵ In this work, solvents with different polarities were used and revealed the potential for tuning the energy transfer efficiency of these carotenoids in artificial systems by changing the solvent polarity. Peridinin was found to transfer energy from its S_1 /ICT state with 80% efficiency in benzene, which is nearly its efficiency in natural PCP. The efficiency decreased with increasing solvent polarity and became as low as 20% in the polar solvent, acetonitrile. The same trend was observed for fucoxanthin, but its efficiency was tunable over a narrower range between 13 and 23%.⁵⁵

A synthetic carotenoid having $N=11$ covalently bound to purpurin in a dyad system provided an S_2 channel with an efficiency greater than 70%, thus exceeding that in a natural system.⁵⁶ Another version of the dyad employed two shorter ($N=10$) carotenoids attached axially to the central Si atom of the Si-phthalocyanine. In this arrangement, the efficiency of the S_2 channel dropped below 70%, but the S_1 channel operated with nearly 90% efficiency, making this system competitive with the most efficient carotenoid-based antenna found in nature.⁵⁷ Moreover, the properties of this carotenoid-phthalocyanine dyad are tunable by solvent polarity as evidenced by the fact that in nonpolar solvents, energy transfer occurs, whereas in polar solvents electron transfer takes place.

Switching between energy and electron transfer was also achieved in a series of carotenoid-phthalocyanine dyads, in which a single carotenoid with $N=9$, 10 or 11 was attached to the tetrapyrrole macrocycle.⁵⁸ For the shortest carotenoid, both the S_2 and S_1 states were active in energy transfer, with efficiencies of 70 and 20%, respectively, whereas for carotenoids with $N=10$ and $N=11$, only the S_2 pathway remained active, and the efficiency dropped to 60 and 24%, respectively. To a certain extent this is reminiscent of the N -dependence of energy transfer observed in purple bacterial antenna, but in these cases the S_1 pathway is inhibited for even shorter π -electron conjugated molecules due to the high S_1 energy of the acceptor. Remarkably, these dyads were capable of mimicking the critical photoprotective function of higher plants evidenced by the two longer carotenoids quenching the excited S_1 state of phthalocyanine.⁵⁹ Thus, these systems hold great promise for the design and construction of novel synthetic solar energy conversion devices that not only carry out light-harvesting as described in this account, but also dissipate excess absorbed energy not required for the specific photochemical process at hand.

Acknowledgments

We thank Miriam Enriquez for assistance with the graphics. TP acknowledges financial support from the Czech Ministry of Education (grants No. MSM6007665808, AV0Z50510513 and ME09037). Work in the laboratory of HAF is funded by grants from the NSF (MCB-0913022, MCB-0842500), the NIH (GM-30353), NASA (NNX08AX20G), and the University of Connecticut Research Foundation.

References

1. Isler, O., editor. Carotenoids. Basel: 1971.
2. Hudson BS, Kohler BE. A low-lying weak transition in polyene α,ω -diphenyloctatetraene. Chem. Phys. Lett. 1972; 14:299–304.
3. Polívka T, Sundström V. Ultrafast dynamics of carotenoid excited states - From solution to natural and artificial systems. Chem. Rev. 2004; 104:2021–2071. [PubMed: 15080720]
4. Polívka T, Sundström V. Dark excited states of carotenoids: Consensus and controversy. Chem. Phys. Lett. 2009; 477:1–11.
5. Krueger BP, Scholes GD, Jimenez R, Fleming GR. Electronic excitation transfer from carotenoid to bacteriochlorophyll in the purple bacterium *Rhodospseudomonas acidophila*. J. Phys. Chem. B. 1998; 102:2284–2292.
6. Scholes GD. Long-range resonance energy transfer in molecular systems. Annu. Rev. Phys. Chem. 2003; 54:57–87. [PubMed: 12471171]
7. McDermott G, Prince SM, Freer AA, Hawthornwaite-Lawless AM, Papiz MZ, Cogdell RJ, Isaacs NW. Crystal structure of an integral membrane light-harvesting complex from photosynthetic bacteria. Nature. 1995; 374:517–521.
8. Shreve AP, Trautman JK, Frank HA, Owens TG, Albrecht AC. Femtosecond energy transfer processes in the B800–850 light-harvesting complex of *Rhodobacter sphaeroides* 2.4.1. Biochim. Biophys. Acta. 1991; 1058:280–288. [PubMed: 2049375]
9. Macpherson AN, Arellano JB, Fraser NJ, Cogdell RJ, Gillbro T. Efficient energy transfer from the carotenoid S_2 state in a photosynthetic light-harvesting complex.

10. Koyama Y, Rondonuwu FS, Fujii R, Watanabe Y. Light-harvesting function of carotenoids in photosynthesis: The roles of the newly found $1B_u^-$ state. *Biopolymers*. 2004; 74:2–18. [PubMed: 15137086]
11. Krueger BP, Scholes GD, Fleming GR. Calculation of couplings and energy transfer pathways between the pigments of LH2 by the ab initio transition density cube method. *J. Phys. Chem. B*. 1998; 102:5378–5386.
12. Akahane J, Rondonuwu FS, Fiedor L, Watanabe Y, Koyama Y. Dependence of singlet-energy transfer on the conjugation length of carotenoids reconstituted into the LH1 complex from *Rhodospirillum rubrum* G9. *Chem. Phys. Lett*. 2004; 393:184–191.
13. Kosumi D, Fujiwara M, Fujii R, Cogdell RJ, Hashimoto H, Yoshizawa M. The dependence of the ultrafast relaxation kinetics of the S_2 and S_1 states in (β -carotene homologs and lycopene on conjugation length studied by femtosecond time-resolved absorption and Kerr-gate fluorescence spectroscopies. *J. Chem. Phys.* 2009; 130:214–506.
14. Cong H, Niedzwiedzki DM, Gibson GN, LaFountain AM, Kelsh RM, Gardiner AT, Cogdell RJ, Frank HA. Ultrafast time-resolved carotenoid to-bacteriochlorophyll energy transfer in LH2 complexes from photosynthetic bacteria. *J. Phys. Chem. B*. 2008; 112:10689–10703. [PubMed: 18671366]
15. Zhang JP, Fujii R, Qian P, Inaba T, Mizoguchi T, Koyama Y, Onaka K, Watanabe Y, Nagae H. Mechanism of the carotenoid-to-bacteriochlorophyll energy transfer via the S_1 state in the LH2 complexes from purple bacteria. *J. Phys. Chem. B*. 2000; 704:3683–3691.
16. Polívka T, Pullerits T, Frank HA, Cogdell RJ, Sundström V. Ultrafast formation of a carotenoid radical in LH2 antenna complexes of purple bacteria. *J. Phys. Chem. B*. 2004; 108:15398–15407.
17. Polívka T, Niedzwiedzki D, Fuciman M, Sundström V, Frank HA. Role of B800 in carotenoid-bacteriochlorophyll energy and electron transfer in LH2 complexes from the purple bacterium *Rhodobacter sphaeroides*. *J. Phys. Chem. B*. 2007; 111:7422–7431. [PubMed: 17547450]
18. Gradinaru CC, Kennis JTM, Papagiannakis E, van Stokkum IHM, Cogdell RJ, Fleming GR, Niederman RA, van Grondelle R. An unusual pathway of excitation energy deactivation in carotenoids: Singlet-to-triplet conversion on an ultrafast timescale in a photosynthetic antenna. *Proc. Natl. Acad. Sci. U.S.A.* 2001; 98:2364–2369. [PubMed: 11226245]
19. Papagiannakis E, Kennis JTM, van Stokkum IHM, Cogdell RJ, van Grondelle R. An alternative carotenoid-to-bacteriochlorophyll energy transfer pathway in photosynthetic light harvesting. *Proc. Natl. Acad. Sci. U.S.A.* 2002; 99:6017–6022. [PubMed: 11972067]
20. Ritz T, Damjanovic A, Schulten K, Zhang JP, Koyama Y. Efficient light harvesting through carotenoids. *Photosynth. Res.* 2000; 66:125–144. [PubMed: 16228415]
21. Liu ZF, Yan HC, Wang KB, Kuang TY, Zhang JP, Gui LL, An XM, Chang WR. Crystal structure of spinach major light-harvesting complex at 2.72 Å resolution. *Nature*. 2004; 428:287–292. [PubMed: 15029188]
22. Gradinaru CC, van Stokkum IHM, Pascal AA, van Grondelle R, van Amerongen H. Identifying the pathways of energy transfer between carotenoids and chlorophylls in LHCII and CP29. A multicolor, femtosecond pump-probe study. *J. Phys. Chem. B*. 2000; 104:9330–9342.
23. Akimoto S, Yokono M, Ohmae M, Yamazaki I, Tanaka A, Higuchi M, Tsuchiya T, Miyashita H, Mimuro M. Ultrafast excitation relaxation dynamics of lutein in solution and in the light-harvesting complexes II isolated from *Arabidopsis thaliana*. *J. Phys. Chem. B*. 2005; 109:12612–12619. [PubMed: 16852560]
24. Croce R, Müller MG, Bassi R, Holzwarth AR. Carotenoid-to-chlorophyll energy transfer in recombinant major light-harvesting complex (LHCII) of higher plants. Femtosecond transient absorption measurements. *Biophys. J.* 2001; 80:901–915. [PubMed: 11159457]
25. Gibasiewicz K, Croce R, Morosinotto T, Ihalainen JA, van Stokkum IHM, Dekker JP, Bassi R, van Grondelle R. Excitation energy transfer pathways in Lhca4. *Biophys. J.* 2005; 88:1959–1969. [PubMed: 15653744]
26. Holt NE, Kennis JTM, Fleming GR. Femtosecond fluorescence upconversion studies of light harvesting by beta-carotene in oxygenic photosynthetic core proteins. *J. Phys. Chem. B*. 2004; 108:19029–19035.

27. de Weerd FL, Kennis JTM, Dekker JP, van Grondelle R. β -carotene to chlorophyll singlet energy transfer in the photosystem I core of *Synechococcus elongatus* proceeds via the β -carotene S_2 and S_1 states. *J. Phys. Chem. B.* 2003; 107:5995–6002.
28. Polívka T, Zigmantas D, Sundström V, Formaggio E, Cinque G, Bassi R. Carotenoid S_1 state in a recombinant light-harvesting complex of photosystem II. *Biochemistry.* 2002; 41:439–450. [PubMed: 11781082]
29. Hilbert M, Wehling A, Schlodder E, Walla PJ. Two-photon-sensitized fluorescence and excitation spectra of photosystem I of *Thermosynechococcus elongates*. *J. Phys. Chem. B.* 2004; 108:13022–13030.
30. Walla PJ, Linden PA, Ohta K, Fleming GR. Excited-state kinetics of the carotenoid S_1 state in LHC II and two-photon excitation spectra of lutein and β -carotene in solution: Efficient car S_1 -Chl electronic energy transfer via hot S_1 states? *J. Phys. Chem. A.* 2002; 106:1909–1916.
31. Frank HA, Cua A, Chynwat V, Young A, Gosztola D, Wasielewski MR. Photophysics of the carotenoids associated with the xanthophyll cycle in photosynthesis. *Photosynth. Res.* 1994; 41:389–395.
32. Ruban AV, Berera R, Iliaia C, van Stokkum IHM, Kennis JTM, Pascal AA, van Amerongen H, Robert B, Horton P, van Grondelle R. Identification of a mechanism of photoprotective energy dissipation in higher plants. *Nature.* 2007; 450:575–578. [PubMed: 18033302]
33. Holt NE, Zigmantas D, Valkunas L, Li X, Niyogi KK, Fleming GR. Carotenoid cation formation and the regulation of photosynthetic light harvesting. *Science.* 2005; 307:433–436. [PubMed: 15662017]
34. Bode S, Quentmeier CC, Liao PN, Hafi N, Barros T, Wilk L, Bittner F, Walla PJ. On the regulation of photosynthesis by excitonic interactions between carotenoids and chlorophylls. *Proc. Natl. Acad. Sci. U.S.A.* 2009; 106:12311–12316. [PubMed: 19617542]
35. Holzwarth AR, Miloslavina Y, Nilkens M, Jahns P. Identification of two quenching sites active in the regulation of photosynthetic light-harvesting studied by time-resolved fluorescence. *Chem. Phys. Lett.* 2009; 483:262–267.
36. Frank HA, Bautista JA, Josue J, Pendon Z, Hiller RG, Sharples FP, Gosztola D, Wasielewski MR. Effect of the solvent environment on the spectroscopic properties and dynamics of the lowest excited states of carotenoids. *J. Phys. Chem. B.* 2000; 104:4569–4577.
37. Zigmantas D, Hiller RG, Sharples FP, Frank HA, Sundström V, Polívka T. Effect of a conjugated carbonyl group on the photophysical properties of carotenoids. *Phys. Chem. Phys.* 2004; 6:3009–3016.
38. Polívka T, Hiller RG, Frank HA. Spectroscopy of the peridinin-chlorophyll-a protein: Insight into light-harvesting strategy of marine algae. *Arch. Biochem. Biophys.* 2007; 458:111–120. [PubMed: 17098207]
39. Hofmann E, Wrench PM, Sharples FP, Hiller RG, Welte W, Diederichs K. Structural basis of light harvesting by carotenoids: Peridinin-chlorophyll-protein from *Amphidinium carterae*. *Science.* 1996; 272:1788–1791. [PubMed: 8650577]
40. Schulte T, Niedzwiedzki DM, Birge RR, Hiller RG, Polívka T, Hofmann E, Frank HA. Identification of a single peridinin sensing Chl-a excitation in reconstituted peridinin-chlorophyll a-proteins (PCP) by crystallography and spectroscopy. *Proc. Natl. Acad. Sci. U.S.A.* 2009; 106:20764–20769. [PubMed: 19934052]
41. Bautista JA, Hiller RG, Sharples FP, Gosztola D, Wasielewski M, Frank HA. Singlet and triplet energy transfer in the peridinin-chlorophyll a protein from *Amphidinium carterae*. *J. Phys. Chem. A.* 1999; 103:2267–2273.
42. Zigmantas D, Hiller RG, Sundström V, Polívka T. Carotenoid to chlorophyll energy transfer in the peridinin-chlorophyll-a-protein complex involves an intramolecular charge transfer state. *Proc. Natl. Acad. Sci. U.S.A.* 2002; 99:16760–16765. [PubMed: 12486228]
43. Krueger BP, Lampoura SS, van Stokkum IHM, Papagiannakis E, Salverda JM, Gradinaru CC, Rutkauskas D, Hiller RG, van Grondelle R. Energy transfer in the peridinin chlorophyll-a protein of *Amphidinium carterae* studied by polarized transient absorption and target analysis. *Biophys. J.* 2001; 80:2843–2855. [PubMed: 11371458]

44. Ilagan RP, Chapp TW, Hiller RG, Sharples FP, Polívka T, Frank HA. Optical spectroscopic studies of light-harvesting by pigment-reconstituted peridinin-chlorophyll-proteins at cryogenic temperatures. *Photosynth. Res.* 2006; 90:5–15. [PubMed: 17361463]
45. Miller DJ, Catmull J, Puskeiler R, Tweedale H, Sharples FP, Hiller RG. Reconstitution of the peridinin-chlorophyll a protein (PCP): Evidence for functional flexibility in chlorophyll binding. *Photosynth. Res.* 2005; 86:229–240. [PubMed: 16172941]
46. Polívka T, Pascher T, Sundström V, Hiller RG. Tuning energy transfer in the peridinin-chlorophyll complex by reconstitution with different chlorophylls. *Photosynth. Res.* 2005; 86:217–227. [PubMed: 16172940]
47. Polívka T, van Stokkum HM, Zigmantas D, van Grondelle R, Sundström V, Hiller RG. Energy transfer in the major intrinsic light-harvesting complex from *Amphidinium carterae*. *Biochemistry.* 2006; 45:8516–8526. [PubMed: 16834325]
48. Papagiannakis E, van Stokkum IHM, Fey H, Büchel C, van Grondelle R. Spectroscopic characterization of the excitation energy transfer in the fucoxanthin-chlorophyll protein of diatoms. *Photosynth. Res.* 2005; 86:241–250. [PubMed: 16172942]
49. Pšenčík J, Ma YZ, Arellano JB, Garcia-Gil J, Holzwarth AR, Gillbro T. Excitation energy transfer in chlorosomes of *Chlorobium phaeobacteroides* strain CL1401: the role of carotenoids. *Photosynth. Res.* 2002; 71:5–18. [PubMed: 16228497]
50. Fuciman M, Chábera P, Župčanová A, Hříbek P, Arellano JB, Váchao F, Pšenčík J, Polívka T. Excited state properties of aryl carotenoids. *Phys. Chem. Chem. Phys.* 2010; 12:3112–3120. [PubMed: 20237698]
51. Montano GA, Xin YY, Lin S, Blankenship RE. Carotenoid and bacteriochlorophyll energy transfer in the B808–866 complex from *Chloroflexus aurantiacus*. *J. Phys. Chem. B.* 2004; 108:10607–10611.
52. Balashov SP, Imasheva ES, Boichenko VA, Anton J, Wang JM, Lanyi JK. Xanthorhodopsin: A proton pump with a light-harvesting carotenoid antenna. *Science.* 2005; 309:2061–2064. [PubMed: 16179480]
53. Polívka T, Balashov SP, Chábera P, Imasheva ES, Yartsev A, Sundström V, Lanyi JK. Femtosecond carotenoid to retinal energy transfer in xanthorhodopsin. *Biophys. J.* 2009; 96:2268–2277. [PubMed: 19289053]
54. Debreczeny MP, Wasielewski MR, Shinoda S, Osuka A. Singlet-singlet energy transfer mechanisms in covalently-linked fucoxanthin- and zeaxanthin-pyropheophorbide molecules. *J. Am. Chem. Soc.* 1997; 119:6407–6414.
55. Polívka T, Pellnor M, Melo E, Pascher T, Sundström V, Osuka A, Naqvi KR. Polarity-tuned energy transfer efficiency in artificial light-harvesting antennae containing carbonyl carotenoids peridinin and fucoxanthin. *J. Phys. Chem. C.* 2007; 111:467–476.
56. Macpherson AN, Liddell PA, Kuciauskas D, Tatman D, Gillbro T, Gust D, Moore TA, Moore AL. Ultrafast energy transfer from a carotenoid to a chlorin in a simple artificial photosynthetic antenna. *J. Phys. Chem. B.* 2002; 106:9424–9433.
57. Marino-Ochoa E, Palacios R, Kodis G, Macpherson AN, Gillbro T, Gust D, Moore TA, Moore AL. High-efficiency energy transfer from carotenoids to a phthalocyanine in an artificial photosynthetic antenna. *Photochem. Photobiol.* 2002; 76:116–121. [PubMed: 12126301]
58. Berera R, van Stokkum IHM, Kodis G, Keirstead AE, Pillai S, Herrero C, Palacios RE, Vengris M, van Grondelle R, Gust D, Moore TA, Moore AL, Kennis JTM. Energy transfer, excited-state deactivation, and exciplex formation in artificial carotenoid-phthalocyanine light-harvesting antennae. *J. Phys. Chem. B.* 2007; 111:6868–6877. [PubMed: 17503804]
59. Berera R, Herrero C, van Stokkum IHM, Vengris M, Kodis G, Palacios RE, van Amerongen H, van Grondelle R, Gust D, Moore TA, Moore AL, Kennis JTM. A simple artificial light-harvesting dyad as a model for excess energy dissipation in oxygenic photosynthesis. *Proc. Natl. Acad. Sci. U.S.A.* 2006; 103:5343–5348. [PubMed: 16569703]

Biographies

Tomáš Polívka was born in 1968 in Czech Republic. He received his M.Sc. in Physics (1992) and Ph.D. in Chemical Physics (1996) from Charles University in Prague. He is a Professor of Biophysics at the University of South Bohemia in Czech Republic. Before joining the University of South Bohemia in 2005, he worked at the Department of Chemical Physics of Lund University. His research interests focus on carotenoid photophysics and on functions of carotenoids in various natural and artificial systems.

Harry A. Frank obtained his B.S. degree in Chemistry in 1972 from Memphis State University, his Ph.D. in Chemistry from Boston University in 1977, and subsequently was an NIH postdoctoral fellow with Kenneth Sauer in the Laboratory of Chemical Biodynamics at the University of California, Berkeley. In 1980 he joined the faculty of the Department of Chemistry at the University of Connecticut where he is currently a Board of Trustees Distinguished Professor. He has been a visiting researcher at the Centre Études de Saclay, France, at the University of Glasgow, Scotland, and in 1995 he was a Fulbright Scholar at the University of Leiden, The Netherlands. His research is aimed at understanding the structure and function of carotenoids in biological organisms.





Figure 1. (top left) A three-state model of carotenoid excited states consisting of S_2 (blue) and S_1 (red) states. Relaxation processes are denoted by arrows and corresponding time constants. Internal conversion processes are denoted by blue and red arrows, black arrows denote vibrational relaxation. Blurred lines denotes the other states whose role in energy transfer is less clear: ICT (red), S^* (green), $1^1B_u^-$ (purple) and $3^1A_g^-$ (black). (Top right) Absorption spectra of LH2 (purple), PCP (orange) and LHCII (green). The range of energies of carotenoid S_2 and S_1 transitions are also shown by the orange and red bars beneath the spectra. Dashed line denotes the solar irradiance spectrum emphasizing the importance of carotenoids in light-harvesting. (Bottom) Molecular structures of four important carotenoids. Color coding corresponds to the absorption spectra of the light-harvesting complexes in which they are found.

**Figure 2.**

Structures of four light-harvesting complexes exhibiting energy transfer from carotenoids. (A) LH2 complex of the purple bacterium *Rhodospseudomonas acidophila* having the carotenoid rhodopin glucoside (red) and BChl-*a* (green); (B) LHCII trimer utilizing the carotenoids lutein (red), neoxanthin (yellow) and violaxanthin (orange) in energy transfer to Chl-*a* (green) and Chl-*b* (blue); (C) Peridinin-Chl-*a*-protein from the dinoflagellate, *Amphidinium carterae*. Eight peridinin molecules (red) transfer energy to two Chl-*a* (green); (D) Xanthorhodopsin from the eubacterium *Salinibacter ruber*. The carotenoid salinixanthin (red) transfers energy to the retinal chromophore (green).




Figure 3. Dependence of the S_1 (blue) and S_2 (red) state energies of carotenoids on their number of conjugated double bonds, N . The width of the bands corresponds either to the variability in state energy due to environment or to uncertainty in determining the energy. The green band corresponds to S_2 energies of carbonyl carotenoids. Energies of typical acceptor states are also shown.




Figure 4. Summary of pathways and efficiencies of carotenoid-mediated energy transfer in light-harvesting systems from various sources. Energy is transferred either from the S_2 state of carotenoids (orange) to the Q_x bands of (B)Chl (yellow), or from the carotenoid S_1 state (red) to Q_y bands of BChl (brown) or Chl (green). The S_1 energy of the retinal chromophore in xanthorhodopsin is shown in purple. The height of the rectangles corresponds to the variation in energy of the electronic states in complexes.