

Cholic-2,4-³H acid was prepared from methyl 3-keto-7 α , 12 α -dihydroxy-5 β -cholanoate (4) and conjugated with glycine methyl ester (5). The reaction mixture was purified by column chromatography using silicic acid and a gradient of methanol in chloroform. Fractions containing pure cholyglycine methyl ester were pooled, and reduced to dryness on a rotary evaporator. The residue was dissolved in ethanol: 1 N NaOH (1:1, v/v) and left at room temperature overnight. The pH was then adjusted to neutrality with 5 N HCl; the solution was evaporated to dryness, and the residue was dissolved in chloroform:methanol (2:1, v/v). The final SA was 3.4 mCi/mmole. Glycine-1-¹⁴C methyl ester was conjugated with cholic acid to yield cholyglycine-1-¹⁴C methyl ester of SA 0.5–1 mCi/mmole. The reaction product was purified by column chromatography using silicic acid and a gradient of methanol in chloroform for elution. The radiopurity was assessed by zonal scanning of thin-layer chromatograms. Radiopurity of both compounds was greater than 95%.

Fasting subjects were given 10 μ Ci of cholyglycine-1-¹⁴C and 25 μ Ci of choly-2,4-³H-glycine in a milk shake. The radiation exposure from 10 μ Ci of cholyglycine-1-¹⁴C is calculated as less than 10% of that received from natural and cosmic sources. Samples of duodenal bile were taken during the next 168 hr; 2 ml of bile was removed after the intravenous administration of 37.5 U of cholecystokinin (supplied by Dr. E. Jorpes, Karolinska Institute, Stockholm, Sweden). The samples were collected in 20 ml of ethanol, heated to precipitate protein, filtered, and then dried. The residue was dissolved in 0.5 ml of methanol and stored at 4°C.

Portions of bile were counted in a dioxane-based solvent system, and ³H and ¹⁴C contents were determined by using external standards. The ³H: ¹⁴C ratio became imprecise after 120 hr because of the rapid decrease in ¹⁴C.

A second portion of bile was analyzed by zonal scanning (6) to determine the distribution of radioactivity among individual bile acid conjugates. The disintegrations per minute (dpm) in each 4 mm section were plotted to give a histogram indicating the percentage radioactivity (³H and ¹⁴C) in each component. Reference compounds were chromatographed simultaneously and radioactivity was assigned to cholyglycine, deoxycholyglycine, or conjugates of taurine; cholytaurine and deoxycholytaurine were not separated. These data indicated not only the distribution of radioactivity among bile acid classes but also the ³H: ¹⁴C ratio in cholyglycine.

A third portion of bile was chromatographed on a 20 \times 25 cm silicic acid plate; cholyglycine standards were run on either side of the bile portion. The region of the plate containing cholyglycine was scraped off and eluted with 0.1 ml of methanol, 1 ml of hydrazine hydrate (1 M, pH 9.5), and 1 ml of phosphate buffer (0.1 M, pH 9.5). One-half of the sample was analyzed for bile acid mass using the hydroxysteroid dehydrogenase method (7), and the remaining half was counted for radioactivity to yield the specific activity of cholyglycine in terms of both ³H and ¹⁴C. Samples were run in duplicate; the coefficient of variation was 7%. Because of the rapid decrease of ¹⁴C specific activity in cholyglycine, valid specific activity for the glycine moiety could not be determined for five subjects after 72 hr and for seven subjects after 120 hr.

To determine whether liberated glycine-1-¹⁴C might be absorbed and reincorporated into cholyglycine, one subject (C₆) was given glycine-1-¹⁴C (20 μ Ci) by mouth; bile sam-

ples were collected after 6 and 24 hr and examined by zonal scanning for the presence of cholyglycine-1-¹⁴C.

Bile acid kinetics were calculated according to a first-order kinetic model (8). The equation for the specific activity decay curve of choly-2,4-³H-glycine was calculated by a least-squares computer program; the linearity of the logarithmic specific activity decay was indicated by a correlation coefficient greater than 0.96 for all studies. Choly-2,4-³H-glycine specific activity at time zero was obtained by extrapolation, and the dose of radioactivity administered was divided by this value to give the pool size. The slope of the log specific-activity decay curve is the rate constant or fractional turnover. Cholyglycine-1-¹⁴C specific activities were multiplied by 2.5 to correct for the smaller dose of administered radioactivity. The adjusted specific activities of cholyglycine-1-¹⁴C at 6, 24, 48, and 72 hr, together with the specific activity of cholyglycine at time zero (calculated from decay of choly-2,4-³H-glycine), were used to calculate a specific activity decay curve for the glycine moiety; this curve intercepted the ordinate at the same place as that of choly-2,4-³H-glycine. The half-lives of the choly-2,4-³H and glycine-1-¹⁴C moieties were determined from the equation $t_{1/2} = \log_e 0.5/K$ ($t_{1/2}$ is half-life, and K is daily fractional turnover). The daily syntheses of the choly and glycine moieties of cholyglycine were obtained by multiplying the pool (millimoles) by the rate constant (days⁻¹). In all calculations of glycine kinetics it was assumed that reincorporation of liberated glycine-1-¹⁴C into the glycine moiety of bile acids was negligible (Results).

At 3, 6, 12, 24, 48, 72, 120, and 168 hr after administration of the dose, the subjects exhaled forcibly into duplicate liquid scintillation vials containing 4.0 ml of 1 M Hyamine-ethanol, 1:1 (v/v), with thymolphthalein as indicator (9, 10), until neutralization was indicated by the indicator turning from blue to colorless. Samples were counted in a toluene-based scintillation mixture and converted to dpm by external standardization. Because each counting vial contained 2 mmoles of CO₂, the radioactivity could be expressed as disintegrations/minute per millimole CO₂. The cumulative excretion of ¹⁴C was calculated by multiplying the mean CO₂ specific activity for each time interval by the endogenous production of CO₂ (9 mmoles/kg per hr) (11). The mean specific activity for each period was considered to be the arithmetic average of the specific activity observed at the beginning and end of the period. Total radioactivity was expressed as percentage of the administered dose.

Two 4-day stool samples were collected from each subject. After homogenization, a portion was combusted to ¹⁴CO₂ and ³H₂O using the Packard oxidizer (Packard Instrument Co., Downers Grove, Ill.); in our hands, this instrument gives nearly 100% recovery of ³H and 80–90% recovery of ¹⁴C. Output of ¹⁴C in flatus was not measured. Output of ¹⁴C in urine was not measured, but in other studies with cholyglycine-1-¹⁴C it has been shown that less than 5% of the administered ¹⁴C is excreted in the urine in 72 hr.¹

Because 7-dehydroxylation without deconjugation appeared to occur in several subjects, attempts were made to demonstrate this phenomenon in vitro. Fecal samples from two subjects who showed dehydroxylation without deconjugation and from two subjects who did not were used. A portion of each fecal homogenate was added to thioglycollate broth and incubated for 24 hr; then 0.1 ml of broth was trans-

¹Hofmann, A. F., M. N. Tacker, H. Fromm, P. J. Thomas, and L. H. Smith. Unpublished observations.

ferred to tubes containing 5 ml of thioglycollate broth with 1 μ mole of choyl-2,4- 3 H-glycine-1- 14 C. After anaerobic incubation for 1, 3, 6, and 10 hr, portions of the broth were removed, diluted with 3 vol of ethanol, heated to 100°C, and filtered. The filtrate was dried and dissolved in methanol, and the distribution of radioactivity in bile acid fractions was determined by zonal scanning.

CALCULATIONS OF DECONJUGATION AND EFFICIENCY OF INTESTINAL ABSORPTION

A simple algebraic model for the enterohepatic circulation of bile acids, with a constant pool size, has been described (12). This model is based on a large, recycling pool of bile acids from which a small, constant fraction is lost during each cycle and replaced by concomitant hepatic synthesis in order to maintain a constant pool size. It assumes that the amount of deconjugation and the efficiency of reabsorption are identical during all cycles.

The pool, P, is secreted C times daily into the duodenum. The fraction of the pool which is absorbed as choyl- 3 H-glycine or cholic acid ($f_{abs} \cdot \text{pool}$) is increased by the synthesis of $P(1 - f_{abs})$ per cycle or $C \cdot P(1 - f_{abs})$ daily, in the steady state with constant pool size.

If some bile acid has been deconjugated by bacterial enzymes in the intestinal lumen before returning to the liver, bile acid conjugation equals the sum of bile acid synthesis plus the amount of unconjugated bile acid returning to the liver. The mole fraction of the pool reabsorbed in conjugated form, N_{conj} , does not require conjugation. The unconjugated fraction absorbed during each cycle is equal to $P \cdot f_{abs} (1 - N_{conj})$. Therefore, daily bile acid conjugation equals $C \cdot P [(1 - f_{abs}) + f_{abs} (1 - N_{conj})]$ which can be reduced to $C \cdot P (1 - f_{abs} \cdot N_{conj})$. Since for choyl- 3 H-glycine the daily bile acid conjugation, and hence deconjugation, is equal to the daily synthesis of the glycine moiety of the choyl- 3 H-glycine, it is possible to calculate the percentage of the choyl- 3 H-glycine pool deconjugated/cycle or per day and the percentage of the choyl moiety reabsorbed without deconjugation or 7 α -dehydroxylation.

RESULTS

3 H: 14 C ratio in bile and choyl- 3 H-glycine. The change in isotope ratio with time in whole bile reflects the relative loss of glycine-1- 14 C and choyl-2,4- 3 H from the enterohepatic circulation. The ratio increased above unity in all subjects (Fig. 1a), indicating that the glycine label is lost more rapidly than the steroid label. As will be shown, 3 H in bile was distributed among choyl- 3 H-glycine, deoxychoyl- 3 H-glycine, and taurine conjugates and, accordingly, the numerator of the 3 H: 14 C ratio represents these three bile acids whereas the denominator is predominantly 14 C in choyl- 3 H-glycine.




FIGURE 1 3 H: 14 C ratios. (a), in whole bile; (b), in choyl- 3 H-glycine. C₁ through C₇ represent different subjects.

The 3 H: 14 C ratio in choyl- 3 H-glycine (Fig. 1b) was smaller, in all cases, than in bile; here, the numerator contains only the 3 H from the choyl moiety, whereas the denominator is essentially identical to that in bile. The marked increase in the ratio in choyl- 3 H-glycine indicates absorption of the choyl moiety as such after deconjugation.

Fractional turnover of choyl- 3 H and glycine- 14 C moieties of choyl- 3 H-glycine. The mean fractional turnover of the glycine moiety was about three times that of the choyl moiety (Table I). Accordingly, the calculated daily synthesis of the glycine moiety is more than three times that of the choyl moiety. A plot of the fractional turnovers of the two moieties against each other (Fig. 2) indicates that four subjects had relatively similar turnover rates of steroid and amino acid moieties,




FIGURE 2 Daily fractional turnover of choyl moiety plotted against daily fractional turnover of glycine moiety. These are equal (see *Changes in choyl glycine pool*) to the fraction of choyl glycine pool deconjugated or dehydroxylated per day (per cent) and the fraction of the choyl glycine pool unabsorbed or dehydroxylated per day (per cent) as indicated on the opposite axes. The diagonal is drawn along a line where the daily fractional turnover of the choyl and glycine moieties are equal and indicates that, in all subjects, the daily fractional turnover of the glycine moiety was greater than that of the choyl moiety.

whereas in four others the turnover of the amino acid moiety was much more rapid than that of the steroid moiety.

Distribution of ^3H in biliary bile acids. There was a progressive increase of ^3H in deoxycholyglycine during the first 120 hr, with a reciprocal decrease in the percentage of ^3H in choyl glycine (Fig. 3). The ^3H in taurine conjugates reached a maximum of $11.5 \pm 1.5\%$ at 72 hr.

Heterogeneity of subjects. Different patterns of deconjugation, dehydroxylation, and fractional turnover were seen in the eight subjects studied.

In subject C_1 , rapid deconjugation with considerable reabsorption of the liberated choyl moiety occurred, as evidenced by the rapid increase in the $^3\text{H} : ^{14}\text{C}$ ratio in bile and in choyl glycine (Fig. 4a). That the increase of this ratio above unity is due to loss of liberated glycine- ^{14}C and conservation of choyl- ^3H is shown by comparison of the half-lives of the two moieties: 19 hr for glycine moiety and 59 hr for the choyl moiety (Table I). Little ^3H appeared in deoxycholyglycine in this sub-

TABLE I
Pool Size, Daily Fractional Turnover, and Daily Synthesis in Normal Subjects

Subject	Choyl-glycine pool mmoles	Daily fractional turnover		Half-life		Synthesis	
		Choyl %	Glycine %	Choyl hr	Glycine hr	Choyl mmoles/day	Glycine mmoles/day
C_1	4.95	12	128	138	13	0.60	6.34
C_2	4.07	50	64	33	26	2.04	2.60
C_3	3.02	39	61	43	27	1.18	1.84
C_4	2.20	19	46	86	36	0.42	1.01
C_5	2.18	66	164	25	10	1.44	3.58
C_6	2.75	31	164	54	10	0.85	4.51
C_7	3.02	28	128	60	13	0.85	3.87
C_8	1.83	55	92	30	18	1.01	1.68
Mean	3.00	38	106	59	19	1.05	3.18
SE	0.37	7	17	13	3	0.18	0.62

ject's bile, suggesting either that his intestinal flora did not 7α -dehydroxylate or, contrary to present concepts (1), that his liver rehydroxylated the deoxycholic acid which was absorbed.

In subject C_7 , rapid deconjugation of choyl glycine also occurred, as is shown by the rapid increase in the $^3\text{H} : ^{14}\text{C}$ ratio above unity (Fig. 4b). The conservation of the choyl moiety was greater than that of the glycine moiety, with half-lives of 60 and 13 hr, respectively. However, the $^3\text{H} : ^{14}\text{C}$ ratio increased more markedly in bile than in choyl glycine because of the presence of much ^3H in deoxycholyglycine (67.7% at 168 hr). This subject therefore showed markedly more 7-dehydroxylation than subject C_1 .




FIGURE 3 Time course of distribution of ^3H among conjugated bile acid classes as mean \pm SE for subjects C_1 through C_8 .




FIGURE 4 Cholyglycine metabolism, showing $^3\text{H}:^{14}\text{C}$ ratio in bile and in cholyglycine (left), distribution of ^3H among conjugated bile acid classes (center), and specific activity decay curve of choly and glycine moieties (right). (a), subject C₁; (b), subject C₇; (c), subject C₄.

In a third subject, C₄, less rapid deconjugation was seen (Fig. 4c) with a slow, albeit steady, increase in the $^3\text{H}:^{14}\text{C}$ ratio in bile and in cholyglycine. The half-

life for the glycine moiety was still shorter than that for the choly moiety (36 and 86 hr, respectively), but for both moieties the half-lives were well above the means




FIGURE 5 Time course of distribution of ^3H among conjugated bile acid classes after incubation of choly-2,4- ^3H -glycine with fecal organisms derived from subjects C_1 and C_8 . The appearance of radioactivity in deoxycholyglycine (open circles) indicates 7-dehydroxylation without deconjugation.

for the eight subjects. Almost 42% of the ^3H was in deoxycholyglycine at 120 hr, but this decreased to 22.5% at 168 hr.

7-Dehydroxylation without deconjugation. In vivo. In three subjects, C_1 , C_6 , and C_7 , ^{14}C appeared in deoxycholyglycine, based on chromatography, at specific times during the study (Table II). It was detectable in C_1 from 24 to 120 hr after administration of choly-2,4- ^3H -glycine-1- ^{14}C . In C_6 and C_7 it was present only at 6 and 24 hr. In the five other subjects of this study, no ^{14}C was detected in deoxycholyglycine at any time.

TABLE II
Per Cent of ^{14}C in Deoxycholyglycine Fraction of Bile

Subject	Time interval					
	6	24	48	72	120	168
	hr					
C_1	0	14	20	22	14	0
C_6	25	11	0	0	0	0
C_7	3	7	0	0	0	0
Gb_9^*	1	1	4	10	5	0

* A 55-yr old woman who had had cholecystectomy 9 yr before this study.

TABLE III
Recovery of ^3H and ^{14}C in Stool and Breath in 168 Hr

Subject	^3H	^{14}C	
	Stool	Stool	Breath
	% dose	% dose	
C_1	33	2.0	71
C_2	53	16.9	35
C_3	50	6.0	34
C_4	38	6.9	36
C_5	67	6.3	63
C_6	67	3.5	65
C_7	65	5.0	56
C_8	68	3.3	54
Mean	55	6.2	52
SE	5	1.6	5

After 20 μCi of glycine-1- ^{14}C had been given to one subject (C_8), no cholyglycine-1- ^{14}C could be detected in the bile 6 and 24 hr later. Thus, it seems improbable that any significant reincorporation of glycine-1- ^{14}C occurred when 10 μCi of cholyglycine-1- ^{14}C was administered.

In vitro. Stool samples from subjects C_1 and C_7 (dehydroxylation without deconjugation) and subjects C_3 and C_8 (no glycine label in deoxycholyglycine) were incubated with choly-2,4- ^3H -glycine-1- ^{14}C for 1, 3, 6, and 10 hr. Deoxycholyglycine derived from cholyglycine—that is, labeled with ^{14}C —was found after incubation with stool from C_1 by thin-layer chromatography, but none appeared after incubation of stool from C_7 , who had also shown in vivo 7-dehydroxylation without deconjugation, or from C_3 and C_8 (Fig. 5).




FIGURE 6 Cumulative fecal excretion of ^3H in 168 hr, expressed as percentage of administered dose, calculated from fractional turnover rate of ^3H choly moiety plotted against recovery of ^3H in stool in that period.

Recovery of ^3H and ^{14}C . In stool, the mean ($\pm\text{SE}$) recoveries by 168 hr were $55\pm 5\%$ for ^3H and $6.2\pm 1.6\%$ for ^{14}C (Table III). By contrast, in the breath the recovery of ^{14}C was $52\pm 5\%$. Of the ^{14}C recovered in the breath, $22\pm 32\%$ was recovered in the first 24 hr.

A significant correlation was found between the excretion of ^3H in stool calculated from the fractional turnover rate of the ^3H cholyl moiety and the actual recovery of ^3H in the stool ($r = 0.92$; $P = 0.001$) (Fig. 6). The recovery of ^3H was less than predicted from the daily fractional turnover rate of the cholyl moiety because some cholic acid is 7-dehydroxylated and reabsorbed, thus conserving the steroid moiety as deoxycholic, but not cholic, acid, and because some ^3H is removed from the steroid nucleus and enters body water.

A significant correlation also was found between the daily fractional turnover rate of the glycine moiety and 24-hr $^{14}\text{CO}_2$ excretion in the breath ($r = 0.95$; $P = 0.001$) (Fig. 7). The daily fractional turnover of the glycine moiety equals $1.4 + 4.7\%$ of dose excreted in $^{14}\text{CO}_2$ in 24 hr.

Changes in cholylglycine pool. Daily loss due to 7-dehydroxylation or failure to reabsorb. Loss of the cholyl moiety of cholylglycine can be due to 7-dehydroxylation with or without reabsorption of the resultant deoxycholic acid or deoxycholylglycine or to failure to reabsorb cholic acid or cholylglycine. In the mathematical model described in Methods, the losses due to both of these factors are equal to the daily fractional turnover rate of the cholyl moiety of cholylglycine. Thus, the mean ($\pm\text{SE}$) percentage of the cholylglycine pool lost, for either reason, is $38\pm 7\%$ /day or about $6\pm 1\%$ /cycle for 6 cycles/day.

Daily deconjugation. The percentage, per day, of the cholylglycine pool deconjugated or 7-dehydroxylated without deconjugation is equal to the daily fractional turnover rate of the glycine moiety of cholylglycine in the model. Thus, $106\pm 17\%$ of the pool is deconjugated or 7-dehydroxylated without deconjugation per day or about $18\pm 3\%$ /cycle for 6 cycles/day.

DISCUSSION

Despite considerable variation in the fractional turnover rates of both steroid and amino acid moieties among subjects, glycine degradation (and synthesis) was more rapid than cholic acid loss and 7-dehydroxylation in each subject. Cleavage of the amide bond of cholylglycine is mediated exclusively by bacterial enzymes (13–16), but the site of such bacterial deconjugation was not indicated by this study.

Biotransformation of glycine and cholyl moieties. The ^{14}C present in glycine-1- ^{14}C was rapidly converted to $^{14}\text{CO}_2$ after liberation from cholylglycine-1- ^{14}C ; this conversion may be caused by bacterial or tissue enzymes which mediate an exchange between carboxyl




FIGURE 7 Daily fractional turnover of glycine moiety plotted against $^{14}\text{CO}_2$ excretion in breath in 24 hr as percentage of administered dose. The cumulative excretion of $^{14}\text{CO}_2$ in this period was calculated by multiplying the mean CO_2 specific activity between intervals at which this was measured by the endogenous production of CO_2 (9 mmoles/kg per hr).

carbon and that of bicarbonate in the medium (17, 18). The site of such an enzyme-mediated exchange, however, is not indicated in our study. In contrast to glycine-2- ^{14}C , glycine-1- ^{14}C is known to show little incorporation into tissue proteins (18, 19) and to be rapidly excreted in breath (11, 20)¹; biotransformations other than conversions to $^{14}\text{CO}_2$ were not examined. Metabolism of the nitrogen atom of the glycine moiety has been well characterized (21–23) but varies according to the route of administration (24). The fate of the glycine nitrogen when glycine is attacked by bacteria in the intestinal lumen is unknown.

^3H from the cholyl moiety appeared rapidly in deoxycholylglycine in all eight subjects. However, the rate of bacterial production or of absorption (or both) of deoxycholic acid appeared to differ greatly, since the percentage of ^3H in deoxycholylglycine ranged from 2.6 to 67.4 on the 7th day. No other bile acids containing ^3H were identified.

The rate of 7-dehydroxylation of cholic acid varied greatly among the eight subjects. Like deconjugation, 7-dehydroxylation is caused by intestinal bacterial enzymes (25). The organisms that cause 7-dehydroxylation are strict anaerobes and therefore are unlikely to exist in large numbers in the ileum of healthy per-

sons, where, even distally, aerobic bacteria predominate over anaerobes (26). On the other hand, while deconjugation is caused mainly by anaerobes (15) it may also be caused by aerobic bacteria (27). It is therefore possible that those patients who absorbed large amounts of free cholic acid had deconjugating but not dehydroxylating bacteria in their ileum, while those who formed deoxycholic acid more rapidly had deconjugating bacteria only in a site where 7-dehydroxylating bacteria were also present—namely, the colon.

Kinetics and route of excretion. The radioactivity present in cholyglycine-1-¹⁴C was excreted predominantly in breath, with about 50% of the administered dose recovered in the breath in 1 wk.

The pool size and synthesis rates of the cholyl moiety of cholyglycine agree closely with those reported for cholic acid pool size and synthesis rates (1, 8, 25, 28–30). The latter should be about one-third larger since they include cholyltaurine. To our knowledge, our data on the synthesis of the glycine moiety are the first to be reported.

Dehydroxylation without deconjugation. In four subjects, ¹⁴C appeared in deoxycholyglycine. Since administered glycine-1-¹⁴C itself did not label the cholylglycine pool, in these subjects the labeled deoxycholyglycine must have come from cholyglycine that had been 7-dehydroxylated without deconjugation. The in vitro demonstration of 7-dehydroxylation without deconjugation complements the experiments of Aries and Hill (31) who prepared cell-free extracts of pure strains of enteric bacteria with 7-dehydroxylating activity toward cholic and chenodeoxycholic acids but found that these extracts failed to 7-dehydroxylate cholyglycine or cholyglycine methyl ester. Thus, 7-dehydroxylation without deconjugation has only been observed in vivo and in vitro when a mixed flora was present.

Subject heterogeneity. Our paper presents the first measurements of the extent of bacterial deconjugation of the bile acid pool during enterohepatic cycling in healthy subjects. Striking heterogeneity in the metabolism of cholyglycine was observed with fractional turnover rates of the cholyl moiety ranging from 19 to 66% and of the glycine moiety, from 46 to 164%. Similar heterogeneity existed in the observed biotransformation of the cholyl moiety, with the maximal amount of ³H appearing in deoxycholyglycine ranging from 9 to 67%. Moreover, we observed that there was no apparent correlation between deconjugation (reflected by the fractional turnover rate of the glycine moiety of cholyglycine) and 7-dehydroxylation (reflected by the biotransformation of cholic acid to deoxycholic acid), although both these phenomena are due largely to enteric bacteria. The heterogeneity pre-

sumably reflects differences in intestinal flora as well as in absorptive capacity for free and conjugated bile acids.

Application to disease. The high correlation between the daily fractional turnover rate of the glycine moiety of labeled cholyglycine and ¹⁴CO₂ output suggests that the ¹⁴CO₂ output may be used to predict the turnover of the glycine moiety of cholyglycine. According to our study, the daily fractional turnover of the glycine moiety is about five times the percentage of the dose of cholyglycine-1-¹⁴C excreted as ¹⁴CO₂ in the breath in 24 hr. Measurement of ¹⁴CO₂ production after administration of cholyglycine-1-¹⁴C has recently been shown to be a useful method for detecting patients with increased bile acid deconjugation and is the basis of a simple breath test for altered bile acid metabolism (32, 33).

Patients with bacterial overgrowth or bile acid malabsorption caused by ileal dysfunction have greatly increased turnover of bile acid glycine based on ¹⁴CO₂ measurements after administration of cholyglycine-1-¹⁴C (32, 33). Since such patients have predominantly glycine-conjugated bile acids in bile (34, 35), they must have greatly increased synthesis rates of bile acid glycine. This greatly increased synthesis may be related to the hyperoxaluria which is seen in some of these patients (36) since administration of taurine, which decreases glycine conjugation by increasing conjugation with taurine, also abolishes the hyperoxaluria (37, 38) in some instances.

ACKNOWLEDGMENTS

This investigation was supported in part by Research Grant AM-6908 from the National Institutes of Health, U. S. Public Health Service.

REFERENCES

1. Danielsson, H. 1963. Present status of research on catabolism and excretion of cholesterol. *Adv. Lipid Res.* 1: 335.
2. Norman, A. 1970. Metabolism of glycocholic acid in man. *Scand. J. Gastroenterol.* 5: 231.
3. Garbutt, J. T., R. M. Wilkins, L. Lack, and M. P. Tyor. 1970. Bacterial modification of taurocholate during enterohepatic recirculation in normal man and patients with small intestinal disease. *Gastroenterology.* 59: 553.
4. Hofmann, A. F., P. A. Szczepanik, and P. D. Klein. 1968. Rapid preparation of tritium-labeled bile acids by enolic exchange on basic alumina containing tritiated water. *J. Lipid Res.* 9: 707.
5. Norman, A. 1955. XXXIV. Preparation of conjugated bile acids using mixed carboxylic acid anhydrides: bile acids and steroids. 34. *Ark. Kemi.* 8: 331.
6. Snyder, F., and H. Kimble. 1965. An automatic zonal scraper and sample collector for radioassay of thin-layer chromatograms. *Anal. Biochem.* 11: 510.

7. Stempfel, R. S., Jr., and J. B. Sidbury, Jr. 1964. Studies with hydroxysteroid dehydrogenases. I. A simplified method for the enzymatic estimation of 3- and 17-hydroxysteroids. *J. Clin. Endocrinol. Metab.* **24**: 367.
8. Lindstedt, S. 1957. The turnover of cholic acid in man: bile acids and steroids 51. *Acta Physiol. Scand.* **40**: 1.
9. Abt, A. F., and S. L. von Schuching. 1966. Fat utilization test in disorders of fat metabolism: a new diagnostic method applied to patients suffering with malabsorption syndrome, chronic pancreatitis, and arteriosclerotic cardiovascular disease. *Johns Hopkins Med. J.* **119**: 316.
10. Kaihara, S., and H. N. Wagner, Jr. 1968. Measurement of intestinal fat absorption with carbon-14 labeled tracers. *J. Lab. Clin. Med.* **71**: 400.
11. Winchell, H. S., H. Stahelin, N. Kusubov, B. Slinger, M. Pollycove, and J. H. Lawrence. 1970. Kinetics of CO₂-HCO₃ minus in normal adult males. *J. Nucl. Med.* **11**: 711.
12. Hofmann, A. F. 1966. A physicochemical approach to the intraluminal phase of fat absorption. *Gastroenterology.* **50**: 56.
13. Norman, A., and O. A. Widström. 1964. Hydrolysis of conjugated bile acids by extracellular enzymes present in rat intestinal contents. *Proc. Soc. Exp. Biol. Med.* **117**: 442.
14. Playoust, M. R., and K. J. Isselbacher. 1964. Studies on the transport and metabolism of conjugated bile salts by intestinal mucosa. *J. Clin. Invest.* **43**: 467.
15. Aries, V., and M. J. Hill. 1970. Degradation of steroids by intestinal bacteria. I. Deconjugation of bile salts. *Biochim. Biophys. Acta.* **202**: 526.
16. Norman, A., and R. Grubb. 1955. Hydrolysis of conjugated bile acids by clostridia and enterococci: bile acids and steroids 25. *Acta Pathol. Microbiol. Scand.* **36**: 537.
17. Klein, S. M., and R. D. Sagers. 1966. Glycine metabolism. I. Properties of the system catalyzing the exchange of bicarbonate with the carboxyl group of glycine in *Peptococcus glycinophilus*. *J. Biol. Chem.* **241**: 197.
18. Richert, D. A., R. Amberg, and M. Wilson. 1962. Metabolism of glycine by avian liver. *J. Biol. Chem.* **237**: 99.
19. Tada, K., K. Narisawa, T. Yoshida, T. Konno, Y. Yokoyama, H. Nakagawa, K. Tanno, K. Mochizuki, and T. Arakawa. 1969. Hyperglycinemia: a defect in glycine cleavage reaction. *Tohoku J. Exp. Med.* **98**: 289.
20. Ando, T., W. L. Nyhan, T. Gerritsen, L. Gong, D. C. Heiner, and P. F. Bray. 1968. Metabolism of glycine in the nonketotic form of hyperglycinemia. *Pediat. Res.* **2**: 254.
21. Shemin, D., and D. Rittenberg. 1946. The biological utilization of glycine for the synthesis of the protoporphyrin of hemoglobin. *J. Biol. Chem.* **166**: 621.
22. Shemin, D., and D. Rittenberg. 1946. The life span of the human red blood cell. *J. Biol. Chem.* **166**: 627.
23. Choitz, H. C., D. Kurrie, and O. H. Gaebler. 1963. Utilization of glycine nitrogen at various levels of glycine intake. *J. Nutr.* **80**: 365.
24. Gaebler, O. H., and H. C. Choitz. 1969. Comparative utilization of glycine-N¹⁵ administered orally or parenterally. *Metab. (Clin. Exp.)*, **18**: 416.
25. Nilsson, S., and T. Scherstén. 1969. Importance of bile acids for phospholipid secretion into human hepatic bile. *Gastroenterology.* **57**: 525.
26. Gorbach, S. L. 1971. Intestinal microflora. *Gastroenterology.* **60**: 1110.
27. Yesair, D. W., and P. Himmelfarb. 1970. Hydrolysis of conjugated bile acids by cell-free extracts from aerobic bacteria. *Appl. Microbiol.* **19**: 295.
28. Heaton, K. W., W. I. Austad, L. Lack, and M. P. Tyor. 1968. Enterohepatic circulation of C¹⁴-labeled bile salts in disorders of the distal small bowel. *Gastroenterology.* **55**: 5.
29. Vlahcevic, Z. R., J. R. Miller, J. T. Farrar, and L. Swell. 1971. Kinetics and pool size of primary bile acids in man. *Gastroenterology.* **61**: 85.
30. Danzinger, R. G., A. F. Hofmann, L. J. Schoenfield, and J. L. Thistle. 1971. Altered bile acid metabolism in patients with cholesterol cholelithiasis. *J. Clin. Invest.* **50**: 24a. (Abstr.)
31. Aries, V., and M. J. Hill. 1970. Degradation of steroids by intestinal bacteria. II. Enzymes catalyzing the oxidation-reduction of the 3 α -, 7 α - and 12 α -hydroxyl groups in cholic acid, and the dehydroxylation of the 7-hydroxyl group. *Biochim. Biophys. Acta.* **202**: 535.
32. Fromm, H., and A. F. Hofmann. 1971. Breath test for altered bile acid metabolism. *Lancet.* **2**: 621.
33. Sherr, H. P., Y. Sasaki, A. Newman, J. G. Banwell, H. N. Wagner, Jr., and T. R. Hendrix. 1971. Detection of bacterial deconjugation of bile salts by a convenient breath-analysis technic. *N. Engl. J. Med.* **285**: 656.
34. Garbutt, J. T., K. W. Heaton, L. Lack, and M. P. Tyor. 1969. Increased ratio of glycine- to taurine-conjugated bile salts in patients with ileal disorders. *Gastroenterology.* **56**: 711.
35. McLeod, G. M., and H. S. Wiggins. 1968. Bile-salts in small intestinal contents after ileal resection and in other malabsorption syndromes. *Lancet.* **1**: 873.
36. Hofmann, A. F., P. J. Thomas, L. H. Smith, and J. T. McCall. 1970. Pathogenesis of secondary hyperoxaluria in patients with ileal resection and diarrhea. *Gastroenterology.* **58**: 960. (Abstr.)
37. Dowling, R. H., G. A. Rose, and D. J. Sutor. 1971. Hyperoxaluria and renal calculi in ileal disease. *Lancet.* **1**: 1103.
38. Admirand, W. H., D. L. Earnest, and H. E. Williams. 1971. Hyperoxaluria and bowel disease. *Trans. Ass. Amer. Physicians Philadelphia.* **84**: 307.