

Published in final edited form as:

Tetrahedron Lett. 2009 December 9; 50(49): 6748–6750. doi:10.1016/j.tetlet.2009.09.072.

A Copper-Carbodiimide Approach to the Phomopsin Tripeptide Side Chain

Ning Shangguan and Madeleine Joullie*

Department of Chemistry, University of Pennsylvania, 231 South 34th Street, Philadelphia, PA 19104

Dehydroamino acids are important moieties in biological investigations and are found in many natural products including roquefortine C and E,¹ azinomycins A and B,² AM-toxines and tentoxin.³ Dehydroamino acids introduce conformational rigidity and change the reactivities of peptides. They are also important intermediates in the biosynthesis of other non-proteinogenic amino acids and D-amino acids.

Phomopsins A (**1**) and B (**2**) are natural products isolated from the fungus species *Phomopsis leptostromiformisa* and potent inhibitors of microtubule polymerization (Figure 1).⁴ A significant structural feature of phomopsins A and B is an unsaturated tripeptide side chain, which contains dehydroisoleucine and dehydroaspartate. This side chain was reported to be important for the molecular interaction of **1** with tubulin.⁵ In this report, we describe an efficient and stereoselective synthesis of the phomopsin tripeptide side chain precursor.

Synthesis of the phomopsin side chain requires a stereoselective method to prepare (*E*)-dehydroisoleucine. Many methods are available for the synthesis of dehydroamino acids, and elimination of water from β -hydroxy- α -amino acids is a well established route. Activation of the hydroxyl group for elimination can be achieved by many reagents: DAST/*N,N*-diisopropylethyl amine (DIPEA), tosyl chloride, mesyl chloride, Martin's sulfurane and triphenyl phosphine/diethyl azodicarboxylate (DEAD).⁶ However, these methods are not highly stereoselective for *E/Z* isomers if there is no strong thermodynamic preference. Wandless *et al.* first reported an *anti*-selective two-step cyclic sulfamidate approach to the (*E*)-dehydroisoleucine. Based on this method they reported a total synthesis of phomopsin B in 2007.⁷ Ferreira *et al.* also reported an *anti*-selective Boc anhydride/DMAP elimination method. Under their conditions all amides were Boc-protected.⁸ In addition to elimination approaches, the Horner-Wadsworth-Emmons reaction has also been utilized to form the double bond.⁹ The vinyl amidation method which couples amides and vinyl halides is another useful approach to synthesize dehydroamino acids.¹⁰

We now describe a one-step copper-carbodiimide elimination that provides the required (*E*)-dehydroisoleucine in a highly stereoselective manner. The copper-carbodiimide elimination method was first reported in 1960s,¹¹ Sai and co-workers found that this elimination method could give high *syn*-selective *E/Z* enamides.¹² The utility of the copper-carbodiimide method

© 2009 Elsevier Ltd. All rights reserved.

mjoullie@sas.upenn.edu.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Supplementary data

Supplementary data associated with this article can be found in the online version.

to prepare dehydroamino acids in a natural product was demonstrated in the total synthesis of roquefortine C.¹³

N-Boc-Dehydroproline (**5**) was prepared by a three-step sequence from commercially available pyrrolidine **3** (Scheme 1). Michael addition of methyl cuprate to the commercially available unsaturated ester **6**, at low temperature resulted in a single *anti* addition product **7**.¹⁴ Osmium tetroxide-mediated dihydroxylation gave *syn* diol **8**. (Sharpless asymmetric dihydroxylation was not chosen because an enantiomeric pure compound was not needed since the *syn* elimination would provide a single dehydration product.) Diol **8** was converted to cyclic sulfate **9** and treated with sodium azide to provide β -hydroxy azide **10**. Hydrogenation of azide **10** to amine **11** and subsequent coupling with acid **5** gave amide **12**, the precursor of the dehydro amino acid moiety.

We screened different EDC(1-ethyl-3-(3-dimethylaminopropyl) carbodiimide)-copper mediated dehydration conditions to introduce the unsaturation and found that copper triflate in THF gave the highest yield and the single desired isomer **13** (Table 1, all the entries gave a single isomer **13**). To our knowledge, this is the first example of preparing a trisubstituted enamide by using this method.

The third amino acid fragment, amine **16**, was prepared by a three-step sequence from commercial available (+)-dimethyl tartrate (Scheme 2).¹⁵ The coupling between acid **17** and amine **16** resulted in the formation of an unreactive azlactone **18**.^{7b} To avoid the azlactone formation, the amide nitrogen had to be protected.

Thus, the ethyl ester **13** was converted to allyl ester **19** and the amide nitrogen was protected with Boc to afford compound **20** (Scheme 3). The allyl ester was cleaved under palladium catalyzed conditions to give acid **21** without reducing the double bond in the dehydroproline moiety.¹⁶ A benzyl ester was also tested but the partial reduction of the double bond occurred under hydrogenolysis conditions. Finally, acid **21** was coupled with amine **16** to give tripeptide **22** and Boc deprotection afforded **23** as the side chain precursor in the synthesis of phomopsin.¹⁷ As shown by previous workers, the dehydroaspartate unit in the phomopsin side chain isomerizes readily under basic conditions.¹⁸ Compound **23** will be coupled with the macrocycle part of phomopsins A and B directly and the β -hydroxy group will not be eliminated until the final stage of the synthesis.

A highly stereoselective approach to make the (*E*)-dehydroisoleucine moiety of the phomopsin tripeptide side chain was developed to afford the material for the total syntheses of phomopsins A and B. The copper-carbodiimide method provides an efficient solution to the stereoselective synthesis of dehydroamino acids. The synthesis and evaluation of the biological activities of phomopsins and their analogues will be reported in due courses.

Supplementary Material


Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

Financial support for this research was provided by NIH (CA-40081) and NSF (0515443). Financial Support for the departmental instrumentation was provided by the National Institutes of Health (1S10RR23444-1). We thank Dr. George T. Furst and Dr. Rakesh Kohli of the University of Pennsylvania Spectroscopic Service Center for assistance in securing and interpreting high-field NMR spectra and mass spectra, respectively.


References

1. (a) Ohmomo S, Sato T, Utagawa T, Abe M. *Agric. Biol. Chem* 1975;39:1333. (b) Clark B, Capon RJ, Lacey E, Tennant S, Gill JH. *J. Nat. Prod* 2005;68:1661. [PubMed: 16309319]
2. Yokoi K, Nagaoka K, Nakashima T. *Chem. Pharm. Bulletin* 1986;34:4554.
3. (a) Okuno T, Ishita Y, Sawai K, Matsumoto T. *Chem. Lett* 1974:635. (b) Fulton ND, Bollenbacher K, Templeton GE. *Phytopathology* 1965;55:49–51.
4. Mackay MF, Van Donkelaar A, Culvenor CC. *J. Chem. Soc. Chem. Commun* 1986:1219.
5. Mitra A, Sept D. *Biochemistry* 2004;43:13955. [PubMed: 15518544]
6. (a) Somekh L, Shanzer A. *J. Org. Chem* 1983;48:907. (b) Pattabiraman VR, Stymiest JL, Derksen DJ, Martin N, Vederas JC. *Org. Lett* 2007;9:699. [PubMed: 17286377] (c) Pravdic N, Zissis E, Pokorny M, Fletcher HtG. *Carbohydr. Res* 1974;32:115. (d) Yokokawa F, Shioiri T. *Tetrahedron Lett* 2002;43:8679. (e) Cherney RJ, Wang. L. *J. Org. Chem* 1996;61:2544.
7. (a) Stohlmeyer MM, Tanaka H, Wandless TJ. *J. Am. Chem. Soc* 1999;121:6100. (b) Stohlmeyer, MM. Ph. D Thesis. CA: Stanford University; 2001. p. 1-109. *Investigating biological checkpoints through organic chemistry: synthesis of the phomopsin side-chain and ATR probes.* (c) Grimley JS, Sawayama AM, Tanaka H, Stohlmeyer MM, Woiwode TF, Wandless TJ. *Angew. Chem. Int. Ed* 2007;46:8157.
8. Ferreira PMT, Maia HLS, Monteiro LS. *Tetrahedron Lett* 1998;39:9575.
9. Schmidt U, Griesser H, Leitenberger V, Lieberknecht A, Mangold R, Meyer R, Riedl B. *Synthesis* 1992:487.
10. Jiang L, Job GE, Klapars A, Buchwald SL. *Org. Lett* 2003;5:3667. [PubMed: 14507200]
11. Corey EJ, Andersen NH, Carlson RM, Paust J, Vedejs E, Vlattas I, Winter REK. *J. Am. Chem. Soc* 1968;90:3245. [PubMed: 5649181]
12. Sai H, Ogiku T, Ohmizu H. *Synthesis* 2003:201.
13. Shangguan N, Hehre WJ, Ohlinger WS, Beavers MP, Joullie MM. *J. Am. Chem. Soc* 2008;130:6281. [PubMed: 18412344]
14. Anderson RJ, Corbin VL, Cotterrell G, Cox GR, Henrick CA, Schaub F, Siddall JB. *J. Am. Chem. Soc* 1975;97:1197. [PubMed: 1133379]
15. Reference^{6b} mentioned that their starting material was (–)-dimethyl tartrate but their reported optical rotation data of intermediates agrees with the data of compounds **14–16**. So we believe that (+)-dimethyl tartrate may have been used as the starting material even though the structure of the synthetic target was not affected because the dehydration reaction eliminates the stereocenters at the final stage.
16. Kunz H, Waldmann H. *J. Chem. Soc., Chem. Commun* 1985;10:638.
17. Selected data: compound **13**: ¹H NMR (DMSO-d₆, 70 °C): δ 8.91 (1 H, br), 5.97 (1 H, m), 5.72 (1 H, m), 4.89 (1 H, dd, *J* = 5.1, 2.6 Hz), 4.10 (2 H, m), 4.04 (2 H, q, *J* = 7.2 Hz), 2.37 (2 H, q, *J* = 7.5 Hz), 1.39 (9 H, s), 1.17 (3 H, t, *J* = 7.2 Hz), 1.02 (3 H, t, *J* = 7.5 Hz); ¹³C NMR (DMSO-d₆, 70 °C): δ 167.8, 163.8, 152.7, 144.6, 127.4, 125.7, 121.9, 78.5, 66.9, 59.2, 53.1, 27.6, 26.0, 18.0, 13.4, 12.1; HRMS (ESI) *m/z* calcd for C₁₈H₂₉N₂O₅ (M + H⁺): 353.2076, Found (M + H⁺): 353.2091; IR (cm⁻¹): 3273 (w), 2977(m), 1709 (s), 1520 (w), 1401 (s), 1207 (w), 1105(w); [α]_D²⁵ = –160 (*c* = 1.96, CHCl₃). Compound **23**: ¹H NMR (CDCl₃): δ 9.22 (1 H, br), 6.84 (1 H, d, *J* = 7.8 Hz), 5.92 (2 H, m), 5.19 (1 H, dd, *J* = 7.9, 2.7 Hz), 4.80 (1 H, d, *J* = 2.8 Hz), 4.69 (1 H, m), 4.02 (1 H, m), 3.87 (1 H, m), 3.81 (1 H, s), 3.74 (1 H, s), 2.38 (2 H, q, *J* = 7.6 Hz), 1.74 (3 H, s), 1.10 (3 H, t, *J* = 7.6 Hz); ¹³C NMR (CDCl₃): δ 172.0, 171.7, 170.0, 165.6, 129.2, 127.8, 124.0, 123.6, 71.1, 68.9, 56.2, 54.4, 52.7, 52.5, 26.9, 17.3, 12.8; HRMS (ESI) *m/z* calcd for C₁₇H₂₆N₃O₇Na (M + H⁺): 384.1701, Found (M + Na⁺): 384.1765; IR (cm⁻¹): 3288 (w, br), 2958 (w), 1748 (s), 1661 (s), 1514 (s), 1439 (w), 1223 (m), 1125 (m), 1029 (w); [α]_D¹⁹ = –76 (*c* = 0.21, CHCl₃).
18. Shin C, Obara T, Morita S, Yonezawa Y. *Bull. Chem. Soc. Jpn* 1988;61:3265.





- 1:** phomopsin A, R= Cl
2: phomopsin B, R= H


Figure 1.
Phomopsins A and B.



Scheme 1.
Preparation of the dehydration precursor




Scheme 2.
Preparation of amine **16** and azlactone formation



Scheme 3.
Completion of the side chain

Table 1

Carbodiimide Copper Dehydration.



Conditions	Yield
CuCl ₂ , 80 °C, toluene, overnight	11%
CuCl ₂ , 100 °C, toluene, 2 h	20%
CuI, 100 °C, toluene, 4 h	30%
No copper, toluene, 100 °C, overnight	0
Cu(OTf) ₂ , THF, 45 °C, overnight	42%
Cu(OTf) ₂ , THF, 60 °C, 1.5 h	73%