



Published in final edited form as:

*J Nat Prod.* 2009 September ; 72(9): 1705–1707. doi:10.1021/np900293x.

## Antiproliferative Triterpenoid Saponins of *Dodonaea viscosa* from the Madagascar Dry Forest<sup>1</sup>

Shugeng Cao<sup>†,‡</sup>, Peggy Brodie<sup>†</sup>, Martin Callmänder<sup>§</sup>, Richard Randrianaivo<sup>§</sup>, Jeremi Razafitsalama<sup>§</sup>, Etienne Rakotobe<sup>⊥</sup>, Vincent E. Rasamison<sup>⊥</sup>, Karen TenDyke<sup>||</sup>, Yongchun Shen<sup>||</sup>, Edward M. Suhl<sup>||</sup>, and David G. I. Kingston<sup>\*,†</sup>

Department of Chemistry, M/C 0212, Virginia Polytechnic Institute and State University, Blacksburg, Virginia 24061, Missouri Botanical Garden, P. O. Box 299, St. Louis, Missouri 63166-029 and B.P 3391, Antananarivo, Madagascar, Centre National d'Application des Recherches Pharmaceutiques, B. P. 702, Antananarivo 101, Madagascar, and Eisai Research Institute, 4, Corporate Drive, Andover, Massachusetts 01810

### Abstract

Bioassay-guided fractionation of an EtOH extract obtained from the roots of the Madagascar plant *Dodonaea viscosa* led to the isolation of two new antiproliferative oleanane-type triterpenoid saponins, dodoneasides A and B (**1** and **2**). The structures of these two new compounds were elucidated using 1D and 2D NMR experiments and mass spectrometry. Compounds **1** and **2** showed antiproliferative activity against the A2780 human ovarian cancer cell line with IC<sub>50</sub> values of 0.79 and 0.70 μM, respectively.

In our continuing search for bioactive molecules from the Madagascar rainforests as part of an International Cooperative Biodiversity Group (ICBG) program, we obtained an extract of the roots of *Dodonaea viscosa* (L.) Jacq. (Sapindaceae). This extract, designated MG 3397, showed reproducible cytotoxicity to the A2780 ovarian cancer cell line, with an IC<sub>50</sub> value of 6.0 μg/mL. The extract was selected for bioassay-guided fractionation based on this activity. Previous work on *Dodonaea viscosa* revealed the presence of flavonoids,<sup>2</sup> fatty acids,<sup>3</sup> and cyanolipids.<sup>4</sup> Some *ent*-clerodane diterpenoids were obtained from *D. boroniaefolia*.<sup>5</sup> A southern Brazilian outbreak of acute hepatic insufficiency in which 14 dairy animals died after consumption of *Dodonaea viscosa* has been reported.<sup>6</sup> In this paper, we report the isolation, structure elucidation, and antiproliferative activity of two new triterpenoid saponins (**1** and **2**) obtained from the roots of *Dodonaea viscosa*.

Liquid-liquid partitioning of a portion of an EtOH extract of the roots of *Dodonaea viscosa* into hexane, CH<sub>2</sub>Cl<sub>2</sub> and aqueous MeOH fractions indicated that the CH<sub>2</sub>Cl<sub>2</sub> fraction (326.5 mg) was the most active fraction, with an IC<sub>50</sub> value of 1.0 μg/mL. Purification of the CH<sub>2</sub>Cl<sub>2</sub> fraction using a C<sub>18</sub> open column, followed by preparative C<sub>18</sub> HPLC, led to the isolation of antiproliferative compounds **1** and **2**.

\*To whom correspondence should be addressed. Tel: (540) 231-6570. Fax: (540) 231-3255. dkingston@vt.edu.

<sup>†</sup>Virginia Polytechnic Institute and State University.

<sup>‡</sup>Current address: Department of Biological Chemistry and Molecular Pharmacology & Osher Research Center, Harvard Medical School, Harvard University, 240 Longwood Av., Boston MA 02115

<sup>§</sup>Missouri Botanical Garden

<sup>⊥</sup>Centre National d'Application des Recherches Pharmaceutiques.

<sup>||</sup>Eisai Research Institute

Supporting Information Available: Spectroscopic data, consisting of <sup>1</sup>H NMR spectra of compounds **1** and **2**, are available as Supporting Information. This material is available free of charge via the internet at <http://pubs.acs.org>

Compound **1** was obtained as white amorphous solid. HRFABMS (positive-ion mode) analysis suggested that the molecular formula **1** was C<sub>57</sub>H<sub>88</sub>O<sub>23</sub>. Its 1D NMR spectra revealed seven tertiary methyl groups between  $\delta_{\text{H}}$  0.87 and 1.42 and a double bond with typical <sup>13</sup>C NMR resonances at  $\delta_{\text{C}}$  127.0 and 143.6, indicating an olean-12-ene triterpene derivative since H<sub>3</sub>-27 ( $\delta_{\text{H}}$  1.42, s) showed a <sup>3</sup>J HMBC correlation to C-13 ( $\delta_{\text{C}}$  143.6). The HMBC spectrum (Figure 1) also exhibited correlations between H<sub>3</sub>-23/H<sub>3</sub>-24 ( $\delta_{\text{H}}$  1.09, s/ $\delta_{\text{H}}$  0.87, s) and C-3 ( $\delta_{\text{C}}$  92.3), H<sub>3</sub>-27 ( $\delta_{\text{H}}$  1.42, s) and C-15 ( $\delta_{\text{C}}$  68.5), H<sub>2</sub>-28 ( $\delta_{\text{H}}$  3.00 & 3.20, d,  $J$  = 9.6 Hz) and C-16/C-22 ( $\delta_{\text{C}}$  74.5/ $\delta_{\text{C}}$  73.9), and H<sub>3</sub>-29/H<sub>3</sub>-30 ( $\delta_{\text{H}}$  0.89, s/ $\delta_{\text{H}}$  1.06, s) and C-21 ( $\delta_{\text{C}}$  79.7), indicating that the aglycone was 3,15,16,21,22,28-hexaoxygenated olean-12-ene. Signals for three anomeric protons [ $\delta_{\text{H}}$  5.24 (1H, d,  $J$  = 2.2 Hz, H-1'''), 4.72 (1H, d,  $J$  = 7.7 Hz, H-1''), 4.55 (1H, d,  $J$  = 8.0 Hz, H-1')] were observed in the <sup>1</sup>H NMR spectrum. The <sup>1</sup>H and <sup>13</sup>C NMR data of the sugar moieties were completely assigned on the basis of the <sup>1</sup>H-<sup>1</sup>H COSY, TOCSY, ROESY, HSQC, HSQC-TOCSY and HMBC spectra and by a comparison of their NMR data with those of aesculoside IIe.<sup>7</sup> These three sugar moieties were identified as  $\beta$ -glucuronopyranosyl [GlcA-1'-6' ( $\delta_{\text{H}}/\delta_{\text{C}}$ ): 4.55, d,  $J$  = 8.0 Hz/105.5; 3.78, dd,  $J$  = 8.0, 8.2 Hz/78.1; 3.69, dd,  $J$  = 8.2, 8.2 Hz/86.1; 3.58, dd,  $J$  = 8.2 Hz/72.4; 3.61, d,  $J$  = 8.2 Hz/76.8; 174 (C-6')],  $\beta$ -glucopyranosyl [Glc-1''-6'' ( $\delta_{\text{H}}/\delta_{\text{C}}$ ): 4.72, d,  $J$  = 7.7 Hz/103.7; 3.19, dd,  $J$  = 7.7, 8.8 Hz/76.0; 3.36, dd,  $J$  = 8.8, 8.8 Hz/77.9; 3.09, dd,  $J$  = 8.8, 8.8 Hz/72.0; 3.29, m/77.9; 3.25 & 3.80, m/63.8], and  $\alpha$ -arabinofuranosyl [Ara-1'''-5''' ( $\delta_{\text{H}}/\delta_{\text{C}}$ ): 5.24, d,  $J$  = 2.2 Hz/110.7; 4.14, m/83.4; 3.86, m/78.6; 4.10, m/85.3; 3.64 & 3.76, m/62.8]. The relative stereochemistry of the arabinofuranosyl moiety was assigned based on a comparison of its <sup>13</sup>C NMR chemical shifts with those of the arabinofuranose ring of aesculoside IIe.<sup>7,8</sup> H-1', H-1'' and H-1''' showed <sup>3</sup>J HMBC correlations to C-3, C-2' ( $\delta_{\text{C}}$  78.1) and C-3' ( $\delta_{\text{C}}$  86.1), respectively, which established the connectivities between these sugar moieties and the aglycone. The <sup>1</sup>H NMR spectrum also had signals for two olefinic protons at  $\delta_{\text{H}}$  6.06 (2H, qq,  $J$  = 7.3, 1.4 Hz, H-A3 and H-A3') and four olefinic methyl groups [ $\delta_{\text{H}}$  1.82 (3H, q,  $J$  = 1.4 Hz, H<sub>3</sub>-A5'), 1.84 (3H, q,  $J$  = 1.4 Hz, H<sub>3</sub>-A5), 1.91 (3H, dq,  $J$  = 7.3, 1.4 Hz, H<sub>3</sub>-A4'), 1.91 (3H, dq,  $J$  = 7.3, 1.4 Hz, H<sub>3</sub>-A4)]; these signals were attributed to two angeloyl moieties. The locations of these two angeloyl moieties were determined on the basis of HMBC correlations between C-A1 ( $\delta_{\text{C}}$  169.4) and H-A3/H<sub>3</sub>-A4/H-21 ( $\delta_{\text{H}}$  5.94, d,  $J$  = 10.2 Hz), and C-A1' ( $\delta_{\text{C}}$  169.2) and H-A3'/H<sub>3</sub>-A4'/H-22 ( $\delta_{\text{H}}$  5.61, d,  $J$  = 10.2 Hz). The double bonds in the angeloyl moieties were determined as *E* by the ROESY correlations between H-A3 and H<sub>3</sub>-A5, and H-A3' and H<sub>3</sub>-A5', and also by a comparison of their NMR data with those of floratheasaponin B.<sup>9</sup> A ROESY correlation between H-3 and H-5 indicated the  $\alpha$ -orientation of H-3. A 10.2 Hz coupling constant between H-21 and H-22 was compatible with a 21–22 diaxial orientation of the hydrogens. The  $\beta$ -axial orientation of H-15, and  $\beta$ -equatorial for H-16, were deduced by the ROESY correlations between H-16 and H-28b, and H-15 and H<sub>3</sub>-26 (Figure 2). The above results were confirmed by comparing the <sup>13</sup>C NMR data of the aglycone of **1** with those of the floratheasaponin B (**3**) (Table 1).<sup>8</sup> The identity of these signals confirmed the structure of compound **1** as shown.<sup>10</sup>

Compound **2** was also obtained as a white amorphous solid. Comparison of the NMR data (Table 1) of **1** and **2** in CD<sub>3</sub>OD indicated that there was no substituent at the C-15 position of **2** and that the angeloyl group at the C-21 position of **1** was replaced by an epoxyangeloyl group in **2**. The NMR spectra indicated that the other parts of **2** were identical to those of **1**. The NMR data of the aglycone and the two substituents at both the C-21 and C-22 positions of **2** were compatible with those of 22-angeloyl-21-epoxyangeloylbarringtonenol.<sup>11</sup> Therefore, the structure of **2** was determined as shown.

Compounds **1** and **2** are oleanane-type triterpenoid saponins with a double bond at the 12-position, an OH group at the 16-position, and substituents at the 3-, 21-, and 28-positions, like gummiferaosides A-C.<sup>12</sup> The triterpenoid sapogenin portion of **1** and **2** is similar to 3 $\beta$ ,15 $\alpha$ ,21 $\beta$ ,22 $\alpha$ ,28-pentahydroxy-16 $\alpha$ -angeloyloxy-12-oleanene isolated from *Dodonaea viscosa*.<sup>13</sup>

Compounds **1** and **2** were evaluated for antiproliferative activity against the A2780 human ovarian cancer cell line, and compound **1** was also evaluated in the breast cancer BT-549, prostate cancer DU 145, NSCLC NCI-H460, and colon cancer HCC-2998 cell lines (Table 2). The activities against the A2780 cell line were similar to those shown by gummiferaosides AC, <sup>12</sup> which suggests that the structural features noted are important for their activity. This finding is similar to that of a recent study that showed that acylation with diangeloyl groups at the C-21 and C-22 positions in triterpenoid saponins is essential for cytotoxicity.<sup>14</sup>

## Experimental Section

### General Experimental Procedures

Optical rotations were recorded on a JASCO P-2000 polarimeter. IR and UV spectra were measured on MIDAC M-series FTIR and Shimadzu UV-1201 spectrophotometers, respectively. NMR spectra were obtained on a JEOL Eclipse 500 for <sup>1</sup>H, <sup>13</sup>C, HMQC, and HMBC and an INOVA 400 spectrometer for TOCSY, COSY, ROESY, and HSQC-TOCSY. Chemical shifts are given in  $\delta$  (ppm), and coupling constants are reported in Hz. Mass spectra were obtained on a JEOL JMS-HX-110 instrument, in the positive-ion mode. HPLC was performed on a Shimadzu LC-10AT instrument with a preparative C<sub>18</sub> Varian Dynamax column (8  $\mu$ m, 250  $\times$  21.4 mm) and a semi-preparative C<sub>18</sub> Varian Dynamax column (5  $\mu$ m, 250  $\times$  10 mm).

### Antiproliferative Bioassays

Cytotoxicity measurements were performed at Virginia Polytechnic Institute and State University against the A2780 ovarian cancer cell line, as described previously.<sup>15</sup> The A2780 cell line is a drug-sensitive human ovarian cancer cell line.<sup>16</sup>

### Plant Material

*Dodonaea viscosa* Jacq. (Sapindaceae) was collected on the eastern side of the Montagne des Français in littoral forest on sand at Iovovona, Antsiranana Province, Madagascar, elevation: ca. 5 m, co-ordinates: 12.21.40 S, 49.29.42 E, on July 19, 2005. Its assigned collection number is Randrianaivo et al. 1208. The collection was made from a shrub on the seashore. The genus *Dodonaea* Mill. consists of ca. 50 species, 2 of which occur in Madagascar. One is endemic (*Dodonaea madagascariensis* Radlk.) and the second one *Dodonaea viscosa* has a large distribution throughout the tropics near the sea. Voucher specimens have been deposited at herbaria of the Centre National d'Application des Recherches Pharmaceutiques, Madagascar (CNARP); the Parc Botanique et Zoologique de Tsimbazaza, Madagascar (TAN); the Missouri Botanical Garden, St. Louis, Missouri (MO); and the Muséum National d'Histoires Naturelles, Paris, France (P).


### Extraction and Isolation


Dried roots of *Dodonaea viscosa* (253 g) were ground in a hammer mill, then extracted with EtOH by percolation for 24 h at rt to give the crude extract MG 3397 (8.9 g), of which 2.6 g was made available to Virginia Polytechnic Institute and State University (VPISU). Extract MG 3397 (2 g, IC<sub>50</sub> 6.0  $\mu$ g/mL) was suspended in aqueous MeOH (MeOH-H<sub>2</sub>O, 9:1, 100 mL) and extracted with hexane (3  $\times$  100 mL portions). The aqueous layer was then diluted to 70% MeOH with H<sub>2</sub>O and extracted with CH<sub>2</sub>Cl<sub>2</sub> (3  $\times$  100 mL portions). The CH<sub>2</sub>Cl<sub>2</sub> extract (326.5 mg) was active with an IC<sub>50</sub> 1.0  $\mu$ g/mL, while the hexane extract (153.5 mg) was inactive and the aqueous MeOH extract (1.5 g) was much less active than the CH<sub>2</sub>Cl<sub>2</sub> extract. The CH<sub>2</sub>Cl<sub>2</sub> extract was chromatographed on an open C<sub>18</sub> column (50  $\times$  10 mm) using H<sub>2</sub>O-MeOH (80:20 to 20:80, then 0:100) to yield the three fractions A [40.2 mg (polar, inactive)], B [198.7 mg, IC<sub>50</sub>: 0.5  $\mu$ g/mL], and C [59.7 mg, IC<sub>50</sub>: 4.3  $\mu$ g/mL]. Fraction B furnished 19 subfractions

after HPLC separation on a C<sub>18</sub> column (0-25-30-60-70 min:50-50-70-70-100% MeOH/H<sub>2</sub>O, 10 mL/min). Subfraction 18 yielded compound **1** (*t<sub>R</sub>* 66 min, 5.0 mg). Compound **2** (*t<sub>R</sub>* 21 min, 0.9 mg) was obtained by HPLC of subfraction 16 using C<sub>18</sub> HPLC (0-30-40 min:70-70-100% MeOH/H<sub>2</sub>O, 2 mL/min).

**Dodonaeside A (1)**—white solid;  $[\alpha]_D^{25}$  -44.4 (*c* 0.18, MeOH); UV (MeOH)  $\lambda_{\max}$  (log  $\epsilon$ ) 209 (4.3) nm; IR (film)  $\nu_{\max}$  3389, 1727, 1152, 1033 cm<sup>-1</sup>; <sup>1</sup>H NMR (500 MHz, CD<sub>3</sub>OD) and <sup>13</sup>C NMR (125 MHz, CD<sub>3</sub>OD) see Table 1; HRFABMS *m/z* 1163.5595 (calcd for C<sub>57</sub>H<sub>88</sub>O<sub>23</sub>Na, 1163.5614).

**Dodonaeside B (2)**—white solid;  $[\alpha]_D^{25}$  -80 (*c* 0.08, MeOH); UV (MeOH)  $\lambda_{\max}$  (log  $\epsilon$ ) 207 (4.1) nm; IR (film)  $\nu_{\max}$  3390, 1698, 1076, 1033 cm<sup>-1</sup>; <sup>1</sup>H NMR (500 MHz, CD<sub>3</sub>OD) and <sup>13</sup>C NMR (125 MHz, CD<sub>3</sub>OD) see Table 1; HRFABMS *m/z* 1163.5580 (calcd for C<sub>57</sub>H<sub>88</sub>O<sub>23</sub>Na, 1163.5614).





**3 R =  $\beta$ -D-xylopyranosyl(1- $\rightarrow$ 2)- $\alpha$ -L-arabinopyranosyl**

## Supplementary Material


Refer to Web version on PubMed Central for supplementary material.

## Acknowledgments


This project was supported by the Fogarty International Center, the National Cancer Institute, the National Science Foundation, the National Heart, Lung and Blood Institute, the National Institute of Mental Health, the Office of Dietary Supplements, and the Office of the Director of NIH, under Cooperative Agreement U01 TW000313 with the International Cooperative Biodiversity Groups. This project was also supported by the National Research Initiative of the Cooperative State Research, Education and Extension Service, USDA, Grant #2008-35621-04732. These supports are gratefully acknowledged. We thank Mr. B. Bebout for obtaining the mass spectra and Mr. T. Glass for assistance with the NMR spectra. Field work essential for this project was conducted under a collaborative agreement between the Missouri Botanical Garden and the Parc Botanique et Zoologique de Tsimbazaza and a multilateral agreement between the ICBG partners, including the Centre National d'Applications des Recherches Pharmaceutiques. We gratefully acknowledge courtesies extended by the Government of Madagascar (Ministère des Eaux et Forêts).

## References and Notes

1. Biodiversity Conservation and Drug Discovery in Madagascar, Part 38. For Part 37, see Rasamison VE, Ranaivo-Harimanana L, Cao S, Pan E, Ratovoson F, Randriantafika F, Rakotondrajaona R, Rakotonandrasana S, Andriantsiferana R, Kingston DGI. *Fitoterapia*. 2009 in press.
2. Getie M, Gebre-Mariam T, Rietz R, Neubert RHH. *Pharmazie* 2002;57:320–322. [PubMed: 12061256]
3. Kapur KK, Gupta AS. *J Sci & Ind Res* 1959;18B:528–530.
4. Sherwani MRK, Hasan SQ, Ahmad I, Ahmad F, Osman SM. *Chem & Ind* 1979:523–524.
5. Jefferies PR, Knox JR, Scaf B. *Aust J Chem* 1973;26:2199–2211.
6. Colodel EM, Traverso SD, Seitz AL, Correa A, Oliveira FN, Driemeier D, Gave A. *Vet Human Toxicol* 2003;45:147–148.
7. Zhang Z, Li S. *Phytochemistry* 2007;68:2075–2086. [PubMed: 17599369]
8. The  $^1\text{H}$  NMR chemical shifts of **1** differed from those of the arabinofuranose ring of aesculoside IIe because the latter spectrum was obtained in pyridine- $d_5$ . The  $^1\text{H}$  chemical shifts of carbohydrates are sensitive to solvents, and our  $^1\text{H}$  NMR shifts for the arabinofuranose ring of **1** in MeOH- $d_4$  are intermediate between the values for aesculoside IIe in pyridine- $d_5$  (reference 7) and those of the arabinofuranose ring of annulatophenonoside in acetone- $d_6$ ; Nedialkov PT, Kitanov GM. *Phytochemistry* 2002;59:867–871. [PubMed: 11937168]
9. Yoshikawa M, Morikawa T, Yamamoto K, Kato Y, Nagatomo A, Matsuda H. *J Nat Prod* 2005;68:1360–1365. [PubMed: 16180814]
10. The recorded signals for C-15 and C-16 appear to have been reversed in reference <sup>9</sup>. Our assignment is consistent with the assignment of these signals in similar compounds such as hippocaesculin: Konoshima T, Lee KH. *J Nat Prod* 1986;49:650–6. [PubMed: 3783160]
11. Li ZL, Li X, Li LH, Li N, Yu M, Meng DL. *Planta Med* 2005;71:1068–1070. [PubMed: 16320212]
12. Cao S, Norris A, Miller JS, Ratovoson F, Razafitsalama J, Andriantsiferana R, Rasamison VE, TenDyke K, Suh T, Kingston DGI. *J Nat Prod* 2007;70:361–366. [PubMed: 17263578]
13. Azam A. *Indian J Chem, Sect B* 1993;32B:513–514.
14. Chan PK. *Biochem Pharmacol* 2007;73:341–350. [PubMed: 17092489]
15. Cao S, Brodie PJ, Randrianaivo R, Ratovoson F, Callmander M, Andriantsiferana R, Rasamison VE, Kingston DGI. *J Nat Prod* 2007;70:679–681. [PubMed: 17323994]
16. Louie KG, Behrens BC, Kinsella TJ, Hamilton TC, Grotzinger KR, McKoy WM, Winker MA, Ozols RF. *Cancer Res* 1985;45:2110–2115. [PubMed: 3986765]



**Figure 1.**  
Key HMBC correlations for compound **1**



**Figure 2.**  
Key ROESY correlations for compound 1



Table 1

<sup>1</sup>H and <sup>13</sup>C NMR Data of Compounds **1** - **3**<sup>a,b,c</sup>

position	1		2		3	
	<sup>1</sup> H	<sup>13</sup> C	<sup>1</sup> H	<sup>13</sup> C	<sup>13</sup> C	<sup>13</sup> C
1	1.05; 1.70	40.0	1.10; 1.65	39.9	39.1	
2	1.77; 1.85	27.1	1.77; 1.85	27.0	26.6	
3	3.24	92.3	3.21	92.3	89.6	
4		40.4		40.4	39.6	
5	0.80 br d (11.0)	56.6	0.79 br d (11.0)	56.9	55.7	
6	1.45; 1.55	19.5	1.45; 1.55	19.3	18.9	
7	1.75	37.2	1.75	33.9	36.7	
8		42.3		40.8	41.5	
9	1.62	48.4	1.62	47.7	47.2	
10		37.9		37.7	37.0	
11	1.95	24.8	1.95	24.7	24.0	
12	5.47 br t (3.5)	127.0	5.39 br t (3.5)	125.3	125.5	
13		143.6		142.9	143.7	
14		48.1		41.0	48.5	
15	3.80	68.5	1.65	34.9	73.1	
16	3.84	74.5	3.99	69.7	67.5	
17		48.5		49.0	47.8	
18	2.63	41.6	2.85	42.3	40.9	
19	1.20; 2.54	47.5	1.20; 2.65	47.9	46.9	
20		36.8		37.1	36.4	
21	5.94 d (10.2)	79.7	6.01 d (10.2)	81.9	78.7	
22	5.61 d (10.2)	73.9	5.60 d (10.2)	73.7	73.3	
23	1.09 s	28.2	1.07 s	28.3	28.1	
24	0.87 s	16.9	0.87 s	16.8	16.9	
25	0.99 s	16.2	0.98 s	16.3	15.8	
26	1.02 s	17.9	0.94 s	17.3	17.6	
27	1.42 s	21.0	1.47 s	27.8	21.2	
28	3.00 d (9.6)	63.6	2.92 d (9.6)	64.4	63.1	

position	1		2		3	
	<sup>1</sup> H	<sup>13</sup> C	<sup>1</sup> H	<sup>13</sup> C	<sup>13</sup> C	<sup>13</sup> C
29	0.89 s	29.6	0.90 s	29.8	29.5	29.5
30	1.06 s	20.2	1.11 s	20.2	20.2	20.2
21-angeloyl						
1		169.4		171.34	167.6	167.6
2		129.3		61.2	128.7	128.7
3	6.06 qq (7.3, 1.4)	139.2	3.05 qq (5.5, 1.4)	61.1	138.4	138.4
4	1.91 dq (7.3, 1.4)	16.0	1.15 dq (5.5, 1.4)	13.9	16.0	16.0
5	1.84 q (1.4)	20.9	1.50 br s	19.8	21.0	21.0
22-angeloyl						
1		169.2		169.1	168.2	168.2
2		129.3		128.9	129.1	129.1
3	6.06 qq (7.3, 1.4)	139.1	6.18 qq (7.3, 1.4)	141.4	136.6	136.6
4	1.91 dq (7.3, 1.4)	15.9	2.02 dq (7.3, 1.4)	16.2	15.7	15.7
5	1.82 q (1.4)	20.8	1.86 q (1.4)	21.0	20.6	20.6
3-β-glcA						
1'	4.55 d (8.0)	105.5	4.48 d (8.0)	105.4	105.6	105.6
2'	3.78 dd (8.0, 8.2)	78.1	3.78 dd (8.0, 8.2)	78.1	79.1	79.1
3'	3.69 dd (8.2, 8.2)	86.1	3.69 dd (8.2, 8.2)	86.2	84.0	84.0
4'	3.58 dd (8.2, 8.2)	72.4	3.58 dd (8.2, 8.2)	72.4	71.1	71.1
5'	3.61 d (8.2)	76.8	3.61 d (8.2)	76.8	77.2	77.2
6'		172.0		172.0	172.0	172.0
2'-β-Glc						
1''	4.72 d (7.7)	103.7	4.72 d (7.7)	103.7		
2''	3.19 dd (7.7, 8.8)	76.0	3.19 dd (7.7, 8.8)	75.9		
3''	3.36 dd (8.8, 8.8)	77.9	3.36 dd (8.8, 8.8)	77.8		
4''	3.09 dd (8.8, 8.8)	72.0	3.09 dd (8.8, 8.8)	72.1		
5''	3.29 m	77.9	3.29 m	77.8		
6''	3.80 m	63.8	3.80 m	63.6		
3'-α-Ara(f)						
1'''	5.24 d (2.2)	110.7	5.27 br s	110.7	110.7	110.7
2'''	4.14 m	83.4	4.14 m	83.3	83.3	83.3

position	1		2		3	
	$^1\text{H}$	$^{13}\text{C}$	$^1\text{H}$	$^{13}\text{C}$	$^1\text{H}$	$^{13}\text{C}$
3''	3.86 m	78.6	3.86 m	78.7		
4''	4.10 m	85.3	4.10 m	85.4		
5''	3.64 m, 3.76 m	62.8	3.64 m, 3.76 m	62.8		

<sup>a</sup>  $\delta$  (ppm) 500 MHz for  $^1\text{H}$  and 125 MHz for  $^{13}\text{C}$ ; multiplicities; *J* values (Hz) in parentheses.

<sup>b</sup> The signals of the sugar carbons were assigned by HSQC-TOCSY and  $^{13}\text{C}$  NMR.

<sup>c</sup> In CD<sub>3</sub>OD.

**Table 2**Antiproliferative activity of compounds **1** and **2**.

Cell line	Cancer type	IC <sub>50</sub> (μM)	
		1	2
A2780	ovarian	0.79	0.70
BT-549	breast	>5	NT <sup>a</sup>
DU 145	prostate	>5	NT
NCI-H460	NSCLC	>5	NT
HCC-2998	colon	>5	NT

<sup>a</sup>NT = not tested