

Positive predictive values of $\geq 5\%$ in primary care for cancer: systematic review

Mark Shapley, Gemma Mansell, Joanne L Jordan and Kelvin P Jordan

ABSTRACT

Background

The positive predictive value (PPV) for cancer of symptoms, signs, and non-diagnostic test results of patients routinely consulting a GP (unselected primary care populations) can help to determine when malignancy should be excluded. Comparisons with other illness indicate that a value of 5% or more may be regarded as highly predictive.

Aim

To identify symptoms, signs, and non-diagnostic test results in unselected primary care populations that are highly predictive of cancer.

Design of study

Systematic review.

Setting

Primary care.

Method

Fourteen bibliographic databases were searched, using terms for primary care, cancer, and predictive values. Reference lists of relevant papers were hand-searched. Data were extracted and the quality of each paper was assessed using predefined criteria, and checked by a second reviewer.

Results

Twenty-five studies were identified. PPVs of 5% or more in specific age and sex groups were reported for: rectal bleeding, change in bowel habit, and iron deficiency anaemia and colorectal cancer; haematuria and urological cancer; malignant rectal examination and prostate cancer; haemoptysis and lung cancer; dysphagia and oesophageal cancer; breast lump and breast cancer; and postmenopausal bleeding and gynaecological cancer.

Conclusion

Robust evidence was found for eight symptoms, signs, and non-diagnostic test results as strongly indicative of cancer for specific age and sex groups in unselected primary care populations. These have the potential to improve the early diagnosis of some cancers in primary care by the use of computer warning flags, improved guidelines, audit, and appraisal.

Keywords

cancer; positive predictive values; predictive value of tests; primary care.

INTRODUCTION

When a patient consults a GP, part of the management of the individual involves an assessment of the probability of serious illness including malignancy. One way of expressing this probability is in terms of the positive predictive value (PPV): the proportion of people with the symptom, sign, or test result who develop cancer. Research on medical decision models reveals that probabilities and the valuation of outcomes are distorted by individuals,¹ and thus theories of behaviour based on cognitive assessment of risk are not directly applicable to the consultation.² For instance, in one model of the consultation, the 'evidence' from research that allows the calculation of probabilities of risk and benefit is modified by an individual doctor's knowledge, skills, attitudes, resources, and legal requirements, and an individual patient's ideas, concerns, expectations, beliefs, and values, to produce the final outcome.³ These factors vary widely between individuals.

Decision makers who wish to maximise desirable outcome often adopt models based on probabilities of benefits and harms. Predictive values have been incorporated into guidelines for primary care published by various colleges, academies, and institutes, including the National Institute for Health and Clinical Excellence,⁴ concerning who to investigate or refer for possible malignancy in primary care. A major criticism

M Shapley, DM, FRCGP, GP and senior research fellow; G Mansell, MSc, research assistant; JL Jordan, MSc, MA, research information manager; KP Jordan, PhD, reader in biostatistics, Primary Care Sciences, Keele University.

Address for correspondence

Dr Mark Shapley, Arthritis Research UK Primary Care Centre, Primary Care Sciences, Keele University, Keele, ST5 5BG. E-mail: mark.shapley@northstuffs.nhs.uk

Submitted: 16 June 2010; Editor's response: 29 June 2010; final acceptance: 19 July 2010.

©British Journal of General Practice

This is the full-length article of an abridged version published in print. Cite this article as: *Br J Gen Pract* 2010; DOI: 10.3399/bjgp10X515412.

is that much of the evidence that underpins these guidelines is not derived from those patients who consult a GP during a routine consultation.⁵ This is important, because predictive values are dependent on the prevalence of the disease in the population from which they are derived and thus applied. The prevalence of cancers within populations and their associated PPVs of symptoms, signs, and non-diagnostic test results increase from community to primary care to hospital populations.

The level of the predictive value given in guidelines at which action is required to investigate or not for cancer will depend on the consequences of a delay in the diagnosis of the disease and the physical, psychological, and material cost of investigation. No guidelines for cancer have been published that state the level of risk that necessitates action. NICE guidelines for the primary prevention of cardiovascular disease recommend statin therapy for hyperlipidaemia when the 10-year cardiovascular risk is 20% or more.⁶ This threshold for initiating statin therapy was based on a cost-effectiveness analysis and risk of adverse effects. Studies show a reduction in relative risk of 27% (95% confidence interval [CI] = 14 to 37%) from statin therapy in patients without cardiovascular disease.⁷ This equates to an absolute risk reduction of approximately 5% or a number needed to treat with statins to prevent a cardiovascular event of 20. Statins are a safe group of drugs with a low risk of serious adverse effects at standard doses (less than 1 in 10 000 patient-years for myopathy).⁷ The consequences of delayed cancer diagnosis in terms of morbidity and mortality are comparable to those of cardiovascular disease, and generally the risks of cancer investigation are similar to the risk of statin therapy. For instance, the rate of perforation of the bowel for colonoscopy used in the investigation for colorectal cancer is 1:769 and procedure-related mortality 1:1537.⁸

Fear of cancer may mean it is perceived by patients as a greater risk than cardiovascular disease, even though a greater number of people die of the latter.¹ Emotional reactions to risky situations often diverge from cognitive assessments of those risks.² While the level of risk an individual should take cannot be defined, comparison of risk of cancer to that of cardiovascular disease suggests that if a population consults with a symptom that could be cancer, then the maximum acceptable risk of not excluding cancer is 1 in 20, which equates to a PPV of 5%.

The objective of this systematic review was to identify the symptoms, signs, and non-diagnostic test results in people routinely consulting a GP that have a PPV of 5% or more for cancer and hence mandate action by a GP except in individual patient-centred circumstances.

How this fits in

The positive predictive values of symptoms, signs, and non-diagnostic test results for cancer in primary care are generally low. The level of risk for cancer that a patient and doctor are willing to accept varies. Using systematic review, good evidence has been gathered for eight symptoms, signs, and non-diagnostic test results as being highly predictive of cancer in primary care. Recommendations for clinical practice, audit, education, and research are made.

METHOD

Search strategy

Fourteen electronic databases were searched from their commencement to October 2009 (Box 1), using terms for primary care, cancer, and predictive values. Searches were limited to English language and human studies. The search strategy used in MEDLINE is shown in Box 2.

Reference lists of relevant studies were checked and citations were tracked using electronic databases. Experts in the field were emailed for further information. All the references were downloaded into reference management software, and the duplicates were removed.

Inclusion and exclusion criteria

Primary studies on unselected primary care populations (patients who consult a GP during a routine consultation) that examined symptoms, signs, or tests for a possible cancer were included in the review. The study had to report a PPV for the symptom, sign, or test, or contain the information necessary to calculate it. For final inclusion, the presented or

Box 1. Electronic databases searched.

- MEDLINE (from 1950)
- EMBASE (1980)
- Cumulative Index to Nursing and Allied Health Literature (CINAHL) (1981)
- Web of Science (1970)
- Biosis Previews (1969)
- Ageline (1978)
- Applied Social Sciences Index and Abstracts (1987)
- Biological Sciences Health and Safety Abstracts (1981)
- Social Services Abstracts (1979)
- Sociological Abstracts (1952)
- Academic Search Elite (1975)
- British Nursing Index (2004)
- International Bibliography of the Social Sciences (1919)
- Health Management Information Consortium (1919)

Box 2. Search (example taken from search used for the MEDLINE database).

““GP*” OR exp† FAMILY PRACTICE OR exp. PHYSICIANS, FAMILY OR exp. PRIMARY HEALTH CARE OR (family ADJ2‡ practi*) OR (general ADJ2 practi*) OR (primary ADJ2 care) OR (family ADJ (physician OR doctor*))”

AND

“exp. NEOPLASMS OR neoplasm* OR cancer* OR carcinoma* OR malignan* OR lesion* OR tumo*§r”

AND

“(risk* ADJ3 cancer) OR “PPV” OR (predictive ADJ value*) OR (diagnos* ADJ value*) OR (early ADJ diagnosis) OR (conditional ADJ probabilit*) OR sensitivity OR specificity OR predictor* OR prevalen* OR (time ADJ interval) OR inciden* OR exp “REFERRAL AND CONSULTATION” OR alarm features”

* truncation — picks up various different word endings. † exp. = exploded thesaurus term: picks up all articles which contain the particular terms as a keyword. ‡ ADJ = adjacent: picks up words next to each other in a document (ADJ3 = picks up words which have up to 3 words in between them). § ? = picks up alternative spellings

calculated PPV had to be:

- 5% or more either for the whole study population or for an age/sex stratum; or
- less than 5% but for the same symptom, sign, or test another eligible study gave a point estimate for the PPV of 5% or more.

The latter criterion was necessary in order to calculate a pooled PPV using all the evidence for a symptom, sign, or test and to compare studies with conflicting results. Exclusion criteria for the review were:

- screening studies, as these involve asymptomatic individuals;
- tests in which a histological diagnosis could be made, as clinical judgement on the need for further investigation or referral would be removed (for example, endoscopy with biopsy, skin biopsy);
- metastatic, recurrent, or cancer-therapy studies as these are not early disease;
- studies with 15 cases or fewer, as these are unlikely to produce reliable conclusions;
- studies written in non-English language due to lack of resources for translating papers; and
- studies in healthcare systems that are not comparable to the UK.

Study selection and assessment of methodological quality

One reviewer screened titles and abstracts and retrieved papers for inclusion. A second reviewer checked 150 (7%) of the abstracts and 25% of the full papers to ensure they met the eligibility criteria. All

studies identified in the screening of titles and abstracts that met the eligibility criteria, or for which it was not possible to tell, were retrieved in full. Two authors independently carried out an unblinded assessment of the risk of bias in each eligible study, using the Newcastle-Ottawa Quality Assessment Scales (NOQAS),⁹ for cohort and case-control studies. Studies are rated from 1 to 9 stars in the NOQAS, with 9 stars indicating a high-quality study. Any discrepancies or disagreements were discussed, and if consensus could not be achieved this was resolved by a third reviewer.

Data extraction and analysis

Once the final selection of articles for inclusion had been agreed, two researchers independently extracted data using a standardised data-extraction form. This included the type of study, characteristics of the study participants, the duration of follow-up, and the reported or calculated PPV.

Any subgroups (age and/or sex) reported in the studies that contained fewer than 15 people, or where authors stated that numbers were too low to reasonably calculate a CI and could not be combined with other data, were excluded from the review.

In studies that did not report the PPV, it was calculated as the proportion of those with that symptom, sign, or test who developed cancer. Where 95% CIs for the PPV were not stated in the paper, these were estimated, where possible, using the data presented. Where this was not possible, an attempt was made to contact the authors.

A meta-analysis was performed to obtain a pooled PPV for symptoms reported in four or more studies. The I² statistic was first calculated to assess the heterogeneity of the studies examining a symptom. The I² statistic can be interpreted as the proportion of total variation in study estimates that is due to heterogeneity between studies.¹⁰ For risk factors where studies could be considered homogenous, a meta-analysis using a fixed-effects (no significant interstudy variation) approach was used. Otherwise, a random-effects approach (significant interstudy variation) based on the inverse variance method was used.¹⁰

The analysis was performed using Stata (version 10) and confidence interval analysis software.¹¹

RESULTS

The search resulted in 17 889 unique references. Of these, 25 papers were found to meet all the eligibility criteria (Figure 1).

Sixteen of the included studies were conducted in the UK, two in each of the Netherlands, Belgium, and US, and one in each of Australia, Denmark, and Germany.

Recruitment into the studies was by one of three




Figure 1. Results of search strategy and selection procedure.

processes: retrospective analysis of clinical records, recruitment of patients with symptoms by GPs either as part of routinely collected data or specifically for the study, and postal survey with prospective identification of cancer from GP records.

Studies varied in the method of extraction of medical record data. The source of medical record data was either routinely coded data only, all routinely recorded data including free text, or GP registered onto specific doctor-completed forms. The duration of follow-up when it was reported ranged from no follow-up to 11 years. Five studies reported that cancer was excluded prior to the first consultation.

Nineteen of the studies had a score of ≥ 6 on the NOQAS, indicating good quality. Common flaws were: the cohort was unrepresentative of unselected primary care populations; differences in criteria determining the presence of cancer; incomplete or inadequate blinding of assessment; lack of demonstration that the individual had not had the relevant cancer diagnosed

previously; short duration of follow-up; and loss to follow-up.

Cancer site

Gastrointestinal. For colorectal cancer, 12 studies were identified giving a predictive value of 5% or more, and three further studies were identified giving a value of less than 5% for one of the symptoms or combinations of symptoms. In 11 of the studies, rectal bleeding (definitions varied) had a predictive value of 5% or more in at least one of the age–sex stratifications (Table 1).^{12–22} Eight of these had quality scores of ≥ 6 .

In seven of the 11 studies there was, however, potential selection bias as GPs recruited participants into the study. In three it was stated that participants may not have been representative of all those consulting with the symptom,^{18,20,21} and in two studies it was demonstrated that the effect of this potential bias was not significant.^{16,17} In three additional studies, rectal bleeding had a PPV of less than 5% in all the

Table 1. Studies giving a PPV of ≥5% in one or more strata of rectal bleeding for colorectal cancer.

Study and NOQAS star rating	Type	Data source	Follow-up	Age, years	Sex	Symptom	Stratification with PPV ≥5%			
							Sex	Age, years	Number with symptom	PPV, % (95% CI)
Jones <i>et al</i> , 2007 ¹² (9 stars)	Retrospective cohort	Routinely recorded computer codes	3 years	≥15	All	Rectal bleeding computer codes	M	75–84	633	7.7 (5.8 to 10.1)
								≥85	157	5.1 (2.2 to 9.8)
							F	75–84	930	7.2 (5.6 to 9.1)
Parker <i>et al</i> , 2007 ¹³ (7 stars)	Retrospective cohort	Routinely recorded computer codes	2 years	≥25	All	First ever consultation with rectal bleeding computer code	M+F	75–84	Number with cancer 173	5.5 (4.7 to 6.3)
Lawrenson <i>et al</i> , 2006 ¹⁴ (9 stars)	Retrospective cohort	Routinely recorded computer codes	1 year	40–89	All	Rectal bleeding computer codes	M	60–69	Not given	6.0 ^a
								70–79	Not given	7.7 ^a
								80–89	Not given	9.1 ^a
Wauters <i>et al</i> , 2000 ¹⁵ (6 stars)	Retrospective cohort	Routinely registered	18–30 months	All	All	Cases of rectal bleeding	M+F	60–69	71	11.2 (5.0 to 21.0)
								70–79	66	21.2 (12.0 to 33.0)
								≥80	51	5.8 (1.2 to 16.2)
Du Toit <i>et al</i> , 2006 ¹⁶ (6 stars)	Prospective cohort	Specifically registered	Not given	≥45	All	New onset rectal bleeding	M+F	65–74	63	9.5 (4.4 to 19.3)
								≥75	76	7.9 (3.7 to 16.2)
Heintze <i>et al</i> , 2005 ¹⁷ (7 stars)	Prospective cohort	Specifically registered	Not given	≥15	All	Cases rectal bleeding	M+F	≥50	268	5.6 (3.4 to 9.0)
Ellis and Thompson, 2005 ¹⁸ (6 stars)	Prospective cohort	Specifically registered	18 months	>34	All	Primary complaint rectal bleeding	M+F	≥60	155	5.2 (2.6 to 9.9)
Nørrelund and Nørrelund, 1996 ¹⁹ (3 stars)	Prospective cohort	Specifically registered	22–57 months	≥40	All	First episode or change in rectal bleeding pattern	M+F	≥40	364	14.8 (11.6 to 18.9)
Metcalfe <i>et al</i> , 1996 ²⁰ (5 stars)	Prospective cohort	Specifically registered	Not given	≥40	All	Rectal bleeding of recent onset	M+F	≥40	99	8.1 (4.2 to 15.1)
Mant <i>et al</i> , 1989 ²² (5 stars)	Prospective cohort	Specifically registered	Not given	≥40	All	Rectal bleeding <6 months onset	M+F	≥40	145	11.0 (6.9 to 17.2)
Fijten <i>et al</i> , 1995 ²¹ (6 stars)	Prospective cohort	Specifically registered	Mean 20 months SD 5 months	18–75	All	Overt rectal bleeding reason for encounter or history of rectal blood loss visible to the patient in the previous 3 months	M	18–75	118	5.9 (2.9 to 11.7)
							M+F	60–75	40	20.0 (10.5 to 34.8)

^aNot possible to calculate confidence intervals (author contacted, original data not available). F = female. M = male. NOQAS = Newcastle-Ottawa Quality Assessment Scales. PPV = positive predictive value. SD = standard deviation.

subgroups presented (Table 2).^{23–25} One study¹⁹ provided separate results for new and old symptoms with the former having a PPV of greater than 5% and the latter less than 5%.

A meta-analysis of studies concerning rectal bleeding is shown in Figures 2 and 3. One study was excluded as the original data were unavailable,¹⁴ and one due to ambiguous data.²⁵ Heterogeneity measured by the I² statistic was high (I² = 93% for studies of older adults), and more recent studies (which were of higher quality) appear to give lower PPVs. The pooled PPV for older adults for rectal bleeding was 4.57% (95% CI = 3.68 to 5.46), and exceeded 5% in those patients aged ≥70 years (5.54% 95% CI = 3.91 to 7.17), although the CI crosses 5% indicating some uncertainty (Figure 3).

There were two studies (both high quality) that

identified patients from large electronic databases with a computer code of change in bowel habit in the medical record and used no free text. In a retrospective cohort study, Lawrenson *et al* reported a PPV of more than 5% in men aged ≥60 years for colorectal cancer,¹⁴ while Hamilton and co-workers, in a case-control study, reported PPVs of less than 5% for all age groups (Table 3).²⁴

In two prospective cohort studies,^{18,21} and a randomised controlled trial,²⁶ the symptom of rectal bleeding was combined with other symptoms or test results to give PPVs of 5% or more (Table 4). However, the numbers of symptomatic patients were generally low and CIs wide. Any conclusions drawn from these findings are weak. An additional case-control study gave some combinations of symptoms and signs a

Table 2. Studies giving a PPV of <5% of rectal bleeding for colorectal cancer.

Study and NOQAS star rating	Type	Data	Follow-up	Age, years	Sex	Number with symptom	Symptom	Comments
Hamilton <i>et al</i> , 2009 ²⁴ (8 stars)	Case-control	Routinely recorded computer codes	NA	≥30	M+F	853 cases, 460 controls	Rectal bleeding computer codes	PPV stratified by age and sex but numbers of cases and controls by age and sex not given
Hamilton <i>et al</i> , 2005 ²³ (9 stars)	Case-control	Routinely recorded data including free text	NA	≥40	M+F	148 cases, 73 controls	Rectal bleeding	PPV 2.4% (95% CI = 1.9 to 3.2)
Thompson <i>et al</i> , 1999 ²⁵ (3 stars)	Prospective cohort	Self-reported questionnaire	4–5 years	>16	M+F	197	Rectal bleeding and consulted GP	No cases of rectal cancer; two cases of caecal cancer felt not to cause bleeding but not stated if in consultation cohort
Nørrelund and Nørrelund, 1996 ¹⁹ (3 stars)	Prospective cohort	Specifically registered	22–57 months	≥40	M+F	45	Current bleeding episode similar to previous	PPV 4.4% (95% CI = 1.2 to 14.8)

F = female. M = male. NA = not applicable. NOQAS = Newcastle-Ottawa Quality Assessment Scales. PPV = positive predictive value.

PPV above 5%, but the authors did not calculate CIs for PPVs due to small sample sizes.²³

One study was identified evaluating iron deficiency anaemia as a predictor for upper and lower gastrointestinal cancer,²⁷ one for cancer,²⁸ and one for colorectal cancer only (Table 5).²⁹ The study by Yates *et al* identified 71 patients with cancer from a cohort of 431 people with iron deficiency anaemia.²⁸ Of these, 35 had lower gastrointestinal cancer, 23 non-gastrointestinal cancer, and 13 other gastrointestinal cancer. The PPV for colorectal cancer was above 5% in men aged >20 years with a haemoglobin of ≤12 g/dl, and the PPV was less than 5% in women aged >50 years with a haemoglobin of ≤11 g/dl. Hamilton *et al* used a case-control methodology but felt unable to calculate a PPV for colorectal cancer in men aged 30 to 59 years, as sample sizes were small.²⁹ Only in cohorts aged ≥60 years did the PPV become greater than 5%, although in some strata the PPV was less than 5%. Stellan and Kenwright studied all patients aged >50 years, where they found the PPV of iron deficiency anaemia for colorectal cancer to be greater than 5%.²⁷

Urological. Three high-quality studies examined symptoms of urological cancer (Table 6). One of these studies excluded prostate cancer,¹² one concerned undefined urological cancer,³⁰ and the third only prostate cancer.³¹ Jones *et al* found PPVs in men aged ≥55 years and women aged ≥65 years to be greater than 5% for haematuria.¹² Bruyninckx *et al* found macroscopic haematuria to have a PPV of over 5% in men aged ≥60 years and women aged ≥40 years.³⁰ The study by Hamilton *et al* of prostate cancer only was a case-control design.³¹ They found a 'malignant rectal




Figure 2. PPV of rectal bleeding for colorectal cancer in older adults. Age ranges for studies (years): ≥45 du Toit,¹⁶ Jones,¹² Parker,¹³ ≥60 Ellis,¹⁸ ≥50 Wauters,¹⁵ Heintze,¹⁷ Hamilton (09)²⁴; ≥40 Mant,²² Nørrelund,¹⁹ Metcalf,²⁰ Hamilton (05),²³ 50–75 Fijtjen.²¹ Figures for Hamilton 09²⁴ were estimated from original data sent by the authors.

Figure 3. PPV of rectal bleeding for colorectal cancer age 70–75 years and over. Age ranges for studies (years): ≥75 Du Toit,¹⁶ Jones,¹² Parker,¹³ Hamilton,²⁴ ≥70 Wauters.¹⁵ Figures for Hamilton 09²⁴ were estimated from original data sent by the authors.




Table 3. Studies giving a PPV of $\geq 5\%$ change in bowel habit for colorectal cancer or a predictive value of $< 5\%$ in an equivalent stratum.

Study and NOQAS star rating	Data source	Type	Follow-up	Age, years	Sex	Symptom	Stratification with PPV $\geq 5\%$			
							Sex	Age, years	Number with symptom	PPV, % (95% CI)
Lawrenson <i>et al</i> , 2006 ¹⁴ (9 stars)	Routinely recorded computer codes	Retrospective cohort	1 year	40–89	M+F	Change in bowel habit computer codes	M	60–69	Not given	6.9 ^a
								70–79	Not given	8.5 ^a
								80–89	Not given	7.7 ^a
Hamilton <i>et al</i> , 2009 ²⁴ (8 stars)	Routinely recorded computer codes	Case-control	NA	≥ 30	M+F	Change in bowel habit computer codes	M	60–69	Not given	3.0 (2.1 to 4.2)
								70–79	Not given	4.2 (3.2 to 5.4)
								80–89	Not given	3.9 (2.8 to 5.5)

^aNot possible to calculate confidence intervals (author contacted original data not available). F = female. M = male. NA = not applicable. NOQAS = Newcastle-Ottawa Quality Assessment Scales. PPV = positive predictive value.

examination' to have a PPV of greater than 5% for prostate cancer. Some combinations of symptoms had a PPV above 5% for prostate cancer, but the authors did not calculate CIs for PPVs, because of small sample sizes.

Lung. One study identified haemoptysis as having a PPV of 5% or more for lung cancer,¹² and another study resulted in a PPV for this symptom of less than 5% (Table 7).³² In the latter study, haemoptysis in combination with other symptoms, or recurrent haemoptysis, had high predictive values. However, the authors did not calculate CIs due to small sample sizes. Jones *et al* used only computer codes to identify the symptom,¹² while Hamilton *et al* used codes and free text.³² Both studies were high quality.

Oesophageal. Two studies (one high and one low

quality) were identified for oesophageal cancer (Table 8).^{12,33} No studies were identified that gave a PPV of less than 5% for the symptom of dysphagia.

Breast. Two studies (one high quality) identified a breast lump symptom,^{34,35} and one high-quality study a clinician-palpable breast lump,³⁶ as having a PPV of 5% or more for breast cancer (Table 9). No studies were identified that gave a PPV of less than 5% for this symptom or sign. Two of the studies included all women,^{34,36} even though breast cancer in those aged < 20 years is exceptionally rare.³⁷

Gynaecological. One high-quality retrospective study, using routinely recorded computer codes in the medical records,¹³ identified the first ever consultation for postmenopausal bleeding in women aged 75 to 84 years as having a PPV for relevant cancer of 5.4%

Table 4. Studies giving a PPV of $\geq 5\%$ in one or more strata of rectal bleeding with another symptom for colorectal cancer.

Study and NOQAS star rating	Symptom	Additional symptom, sign, or test result	Sex	Age (years)	Number with symptom	PPV, % (95% CI)
Ellis and Thompson, 2005 ¹⁸ (6 stars)	Primary complaint rectal bleeding	Change in bowel habit	M+F	> 34	119	9.2 (5.2 to 15.8)
		Change in bowel habit (loose and/or frequent)			83	12.1 (6.7 to 20.8)
		Change in bowel habit with no abdominal pain			52	9.6 (4.2 to 20.6)
		No peri-anal symptoms			63	11.1 (5.5 to 21.2)
		Dark blood			31	9.7 (3.3 to 24.9)
		Small volume of blood			187	5.3 (2.9 to 9.6)
Fijten <i>et al</i> , 1995 ²¹ (6 stars)	Overt rectal bleeding reason for encounter or history of rectal blood loss visible to the patient in the previous 3 months	Change in bowel habit more frequent or diarrhoea or variously but not constipation	M+F	18–74	78	9.0 (4.4 to 17.4)
		Blood seen on stool or mixed with only			54	7.4 (2.9 to 17.6)
		Weight loss			42	9.5 (3.8 to 22.1)
		Perianal eczema			17	17.6 (6.2 to 41.0)
		Rectal palpation haemorrhoid			20	10.0 (2.8 to 30.1)
		High ESR			23	8.7 (2.4 to 26.8)
		High WBC			25	12.0 (4.2 to 30.0)
		Haemoccult positive			41	4.9 (1.3 to 16.1)
Leicester <i>et al</i> , 1984 ²⁶ (3 stars)	Any abdominal or bowel complaint	Haemoccult positive	M+F	≥ 40	25	16.0 (6.4 to 34.7)

ESR = erythrocyte sedimentation rate. F = female. M = male. WBC = white cell count. NOQAS = Newcastle-Ottawa Quality Assessment Scales. PPV = positive predictive value.

Table 5. Studies giving a PPV of $\geq 5\%$ in one or more strata of iron deficiency anaemia for colorectal cancer or a predictive value of $< 5\%$ in an equivalent stratum.

Study and NOQAS star rating	Type	Data source	Follow-up	Age, years	Sex	Symptom	Stratification with PPV			
							Sex	Age, years	Number with symptom	PPV, % (95% CI)
Stellon <i>et al</i> , 1997 ²⁷ (6 stars)	Prospective cohort	Specifically registered	Not given	>50	M+F	Hb <12.0 and/or MCV <80 with ferritin ≤ 16	M+F	>50	26	7.7 (2.1 to 24.1)
Yates <i>et al</i> , 2004 ²⁸ (6 stars)	Retrospective cohort	Routinely recorded data	1 year	>20	M	Hb ≤ 12 MCV <hospital normal range, RBC <5.5	M	>20	154	17.5 (12.3 to 24.3)
				>50	F	Hb ≤ 11 MCV <hospital normal range, RBC <5.5	F	>50	277	3.2 (1.7 to 6.1)
Hamilton <i>et al</i> , 2008 ²⁹ (9 stars)	Case-control	Routinely recorded electronic data	NA	≥ 30	M+F	Hb 11.0–11.9 with indicators of iron deficiency	M	60–69	Not given	6.5 (2.0 to 19)
							M	≥ 60	Not given	All age bands and Hb levels PPV >5 (range 5.5 to ≥ 31)
							F	70–79	Not given	All Hb levels PPV >5 (range 5.9 to 10)
							F	≥ 80	Not given	All Hb levels PPV >5 (range 5.7 to 10%)
							F	60–69	Not given	>5 (likelihood ratio)

F = female. Hb = haemoglobin (g dl⁻¹). M = male. MCV = mean corpuscular volume (fL). NA = not applicable. NOQAS = Newcastle-Ottawa Quality Assessment Scales. PPV = positive predictive value. RBC = red blood corpuscles.

Table 6. Studies giving a PPV of $\geq 5\%$ in one or more strata of symptoms for urological cancer.

Study and NOQAS star rating	Cancers	Type	Data	Follow-up	Age, years	Sex	Symptom or sign	Stratification with PPV $\geq 5\%$			
								Sex	Age, years	Number with symptom	PPV, % (95% CI)
Jones <i>et al</i> , 2007 ¹² (9 stars)	Urethra, bladder, ureter and kidney	Retrospective cohort	Routinely recorded computer codes	3 years	≥ 15	M+F	Haematuria computer codes	M	55–64	1104	8.5 (6.9 to 10.3)
									65–74	1517	11.2 (9.7 to 12.9)
									75–84	1198	10.3 (8.6 to 12.1)
									≥ 85	358	9.2 (6.4 to 12.7)
								F	65–74	846	5.9 (4.4 to 7.7)
									75–84	688	6.8 (5.1 to 9.0)
Bruyninckx <i>et al</i> , 2003 ³⁰ (7 stars)	Any of the urological tract	Prospective cohort	Routinely registered	18–30 months	Not given	M+F	Cases macroscopic haematuria	M	≥ 60	Not given	22.1 (15.8 to 30.1)
								F	40–59	Not given	6.4 (1.7 to 18.6)
									≥ 60	Not given	8.3 (3.4 to 17.9)
Hamilton <i>et al</i> , 2006 ³¹ (7 stars)	Prostate	Case-control	Routinely recorded data including free text	NA	≥ 40	M	Rectal examination malignant	M	≥ 40	Cases 5, controls 41	12.0 (5.0 to 37.0)

F = female. M = male. NA = not applicable. NOQAS = Newcastle-Ottawa Quality Assessment Scales. PPV = positive predictive value.

(95% CI = 4.7 to 7.1) (Table 10). No studies were identified that gave a PPV of less than 5% for this symptom in the age group 75 to 84 years.

Other cancers. No other studies were identified with PPVs of over 5% for symptoms, signs, or tests for other sites of cancer.

Table 7. Studies giving a PPV of ≥5% in one or more strata of symptoms for lung cancer or a predictive value of <5% in an equivalent stratum.

Study and NOQAS star rating	Type	Data source	Follow-up	Age, years	Sex	Symptom	Stratification with PPV			
							Sex	Age, years	Number with symptom	PPV, % (95% CI)
Jones <i>et al</i> , 2007 ¹² (9 stars)	Retrospective cohort	Routinely recorded computer codes	3 years	≥15	All	Haemoptysis computer codes	M	55–64	514	8.4 (6.1 to 11.1)
								65–74	552	14.9 (12.0 to 18.1)
								75–84	393	17.1 (13.5 to 21.1)
								≥85	93	20.4 (12.8 to 30.1)
							F	65–74	358	8.4 (5.7 to 11.8)
								75–84	258	10.5 (7.0 to 14.9)
Hamilton <i>et al</i> , 2005 ³² (9 stars)	Case-control	Routinely recorded data including free text	Not applicable	≥40	All	Haemoptysis as a single symptom	M+F	≥40	Cases 50, controls 19	2.4 (1.4 to 4.1)

F = female. M = male. NOQAS = Newcastle-Ottawa Quality Assessment Scales. PPV = positive predictive value.

Table 8. Studies giving a PPV of ≥5% in one or more strata of dysphagia for oesophageal cancer.

Study and NOQAS star rating	Type	Data source	Follow-up	Age, years	Sex	Symptom	Stratification with PPV ≥5%			
							Sex	Age, years	Number with symptom	PPV, % (95% CI)
Jones <i>et al</i> , 2007 ¹² (9 stars)	Retrospective cohort	Routinely recorded computer codes	3 years	≥15	M+F	Dysphagia computer codes	M	55–64	518	6.0 (4.1 to 8.4)
								65–74	576	9.0 (6.8 to 11.7)
								75–84	476	7.1 (5.0 to 9.8)
								≥85	154	9.7 (5.6 to 15.6)
Esfandiyari <i>et al</i> , 2002 ³³ (4 stars)	Prospective cohort	Specifically registered	Not given	Not given	M+F	Dysphagia	M+F	Not given	100	6.0 (2.8 to 12.5)

F = female. M = male. NOQAS = Newcastle-Ottawa Quality Assessment Scales. PPV = positive predictive value.

Table 9. Studies giving a PPV of ≥5% in one or more strata of symptoms for breast cancers.

Study and NOQAS star rating	Type	Data source	Follow-up	Age, years	Sex	Symptom or sign	Stratification with PPV ≥5%			
							Sex	Age, years	Number with symptom	PPV, % (95% CI)
Eberl <i>et al</i> , 2008 ³⁴ (5 stars)	Retrospective cohort	Routinely recorded data	Not given	Not given	F	Breast lump/mass symptom	F	Not given	741	8.1 (6.3 to 10.3)
Barton <i>et al</i> , 1999 ³⁵ (8 stars)	Retrospective cohort	Routinely recorded data	1–11 years	40–69	F	Breast lump/mass symptom	F	40–69	196	10.7 (4.6 to 16.9)
Bywaters 1977 ³⁶ (6 stars)	Retrospective cohort	Routinely recorded data	Not given	All	F	Breast lump/mass sign	F	All	57	24.6 (15.2 to 37.1)

F = female. NOQAS = Newcastle-Ottawa Quality Assessment Scales. PPV = positive predictive value.

Table 10. Studies giving a PPV of ≥5% in one or more strata of symptoms for gynaecological cancer.

Study and NOQAS star rating	Type	Data source	Follow-up	Age, years	Sex	Symptom or sign	Stratification with PPV ≥5%			
							Sex	Age, years	Number with symptom	PPV, % (95% CI)
Parker <i>et al</i> , 2007 ¹³ (7 stars)	Retrospective cohort	Routinely recorded computer codes	2 years	≥40	F	First ever consultation with postmenopausal bleeding computer codes	Female	75–84	856	5.4 (4.7 to 7.1)

F = female. NOQAS = Newcastle-Ottawa Quality Assessment Scales. PPV = positive predictive value.

Table 11. Symptoms, signs and non-diagnostic tests in unselected primary care populations with a PPV of $\geq 5\%$ for cancer for which there is robust evidence.

Symptom	Cancer	Sex	Age, years	Evidence level for PPV of 5% or more in other cohorts
Rectal bleeding computer code or new onset rectal bleeding	Colorectal	M+F	≥ 75	None aged <60 years, aged 60–74 years equivocal
Iron deficiency anaemia Hb <12 g/dl	Colorectal	M	≥ 60	Moderate evidence for gastrointestinal malignancy in men with Hb ≤ 12 g/dl aged >20 years and women with Hb ≤ 11 g/dl aged >50 years
Iron deficiency anaemia Hb <11 g/dl	Colorectal	F	≥ 70	
Iron deficiency anaemia Hb <9 g/dl	Colorectal	F	≥ 60	
Haematuria	Urological	M+F	≥ 60	None aged <40 years, aged 40–60 years equivocal
Rectal examination malignant	Prostate	M	≥ 40	No other evidence for PPV of 5% or more in other cohorts
Haemoptysis computer code	Lung	M	≥ 55	No other evidence for PPV of 5% or more in other cohorts
Haemoptysis computer code	Lung	F	≥ 65	
Dysphagia computer code	Oesophagus	M	≥ 55	No other evidence for PPV of 5% or more in other cohorts
Breast lump or mass	Breast	F	≥ 20	No other evidence for PPV of 5% or more in other cohorts
Postmenopausal bleeding computer code	Gynaecological	F	75–84	No other evidence for PPV of 5% or more in other cohorts

F = female. Hb = haemoglobin. M = male. PPV = positive predictive value.

DISCUSSION

Summary of main findings

The study identified 21 papers that reported a PPV of 5% or more for a symptom, sign, or non-diagnostic test result alone or in combination in unselected primary care populations for cancer, and a further four studies reporting a PPV of less than 5% for the same symptom, sign, or non-diagnostic test result.

Strengths and limitations of the review

The definition of symptom differed between studies. In some, only the computer codes allocated to the consultation in the medical records were used, while others used the whole medical record including free text. The latter method will detect a greater number of patients with symptoms, as the computer code generally represents a summation of the consultation into a diagnosis or term representing patient management. This approach is encouraged in research practices (personal communication, Royal College of General Practitioners' [RCGP] Weekly Returns Service, 2010), with a stated preference for GPs to record diagnostic computer codes rather than symptom codes. It is unusual for a GP to record multiple history or symptom codes for a single problem. The studies that calculate PPVs for computer codes thus give values for patients in whom a GP is unable to make a more specific diagnosis. These studies do not give PPVs for symptoms presented to GPs, unlike studies in which free text is also analysed. Both types of study overestimate the PPV as a consequence of under-recording by GPs, especially when the clinician has a low level of suspicion for cancer. The size of this effect is not known. The PPVs will only remain valid if GP recording and patient consultation behaviour do not change with time.

A significant source of bias in studies using

clinicians to specifically register a patient is selection of those with more severe symptoms and those in whom the clinician has a greater suspicion of malignancy. Such studies are considered to have a low weight in the present review of unselected primary care populations due to this selection bias, and are likely to overestimate the PPV. In the meta-analysis of rectal bleeding, there is a large degree of heterogeneity and a trend for the PPV to be lower in more recent studies. This partially reflects this source of bias and a change in recruitment methods over time from the use of specifically registered patients to routinely recorded data. In studies using computer codes, the actual codes used were not reported and may differ significantly between studies even when the symptom appears to be a single entity (personal communications, W Hamilton and R Jones, 2009).

A recent systematic review and meta-analysis on studies that only involved the recruitment of cases of rectal bleeding by GPs commented on the difficulties of complete identification and follow-up of all incident cases of rectal bleeding in primary care and differences in age cut-off points between studies.³⁸ The authors gave an estimated PPV of over 5% in those aged ≥ 60 years with rectal bleeding.

Several other methodological problems can overestimate the PPV in studies, including the use of cancer registries resulting in the inclusion of cancers not diagnosed through primary care (for example, screen detected, referral within hospital, and admission via accident units), cancers diagnosed as an association and not the cause of symptoms, and recall bias in case-control studies.

Underestimation of the PPV may occur with inadequate follow-up, and the length of follow-up varied between studies from no follow-up to 11 years. The study by Jones *et al* indicates that

some cancers are diagnosed 3 years after initial presentation.¹²

Other potential sources of bias in this review were that it only included English language papers, and that some papers in more specialist journals could not be located. This potential bias is unlikely to affect the results of the review, as, on closer examination of the abstracts, they are unlikely to have matched the study eligibility criteria.

The studies identified that calculated PPVs for combinations of symptoms, signs, or non-diagnostic test results and risk of cancer, involved populations that were too small and/or methods of recruitment that were subject to too high a risk of bias to draw firm conclusions. Three recent systematic reviews on the diagnostic utility of combinations of symptoms, signs, and test results for colorectal cancer showed that evidence in primary care is lacking, and when present only shows modest diagnostic value.³⁸⁻⁴⁰

A potential criticism of this review is that it only included symptoms, signs, and tests with a point estimate PPV of 5% or more. It is likely that it excluded good-quality studies with a point estimate of less than 5%, but in which the 95% CIs included 5%. It is possible that future research may identify additional symptoms, signs, and tests with a point estimate of 5% or more.

The level of risk an individual is prepared to take is specific to that person at that moment, and is dependent on many factors. Twenty-six additional studies were identified with PPVs of other symptoms, signs, and non-diagnostic test results for cancer of less than 5%. A lower level of PPV than that used in this study will identify more symptoms, signs, and non-diagnostic test results for cancer, and additional age and sex groups in unselected primary care populations. Selecting a lower level of risk may diagnose cancer earlier in presenting populations but with the disadvantage that more individuals will undergo negative investigation with physical, psychological, and economic cost from the tests. A maximum level of risk for a population of 5% was chosen for this review, based on stated levels of risk in national guidelines concerning a comparable disease and treatment. The lower the level of risk, the greater the disagreement as to whether or not inaction can be justified.

When communicating with patients, it is preferable to use absolute measures of risk (number needed to investigate) rather than relative measures of risk.⁴¹ In assessing a patient's level of risk using predictive values, it is important to individualise the risk, as the PPV of the population will be modified by the individual's risk factors for the illness. There has been no debate with the public regarding what level of risk is acceptable.¹⁴

Implications for future research and clinical practice

This study found evidence for only nine symptoms, signs, and non-diagnostic test results that have a high predictive value for cancer in patients presenting to GPs during routine consultations. A number of studies were subject to methodological and sample size difficulties, and for one symptom (change in bowel habit and colorectal cancer), high-quality studies using different methodology produced contradictory results. On the basis of good quality studies (with one or more studies on the quality-assessment tool at a level of 6 stars or more and no similar quality contradictory studies), broad consistency with published guidelines,⁴ and the unlikelihood that further research will change the conclusion of studies performed on large databases, it can be concluded that robust evidence exists for a PPV of 5% or more in specific age/sex groups as detailed in Table 11.

The low number of symptoms, signs, and non-diagnostic test results with a PPV of 5% or more was not unexpected, as the difficulties in diagnosing cancer in primary care have been commented on before.⁵ While these symptoms, signs, and non-diagnostic test results occur in the minority of patients with cancer, when they do occur, exclusion of cancer is obligatory unless exceptional circumstances exist. There is some evidence that GPs do not always act upon them.^{26,42,43} A previous effort to trigger GP action concerning one of the tests was unsuccessful,⁴⁴ but this may have been due to the form of trigger used. The identified symptoms, signs, and non-diagnostic test results are broadly consistent with NICE recommendations for referral for suspected cancer. However, NICE generally uses a lower threshold for referral for additional symptoms and tests, and in the case of haemoptysis, even when the computer code is associated with a PPV of 5% or more.

The authors recommend research and development of general practice computer systems to produce effective warning flags when the symptoms, signs, or test results with a PPV of 5% or more from unselected primary care populations are entered for patients within the specified sex and age groups. The management of these patients should be audited and appraised. There is a need to standardise terms and analyses, and for further research on combinations of symptoms, signs, and non-diagnostic test results. There should be more open debate on the level of PPV that triggers a recommendation for referral by a GP.

Funding body

NHS North Staffordshire; NHS Stoke; NHS Executive West Midlands R&D Office; North Staffordshire and Cheshire R&D Consortium.

Ethical approval

Not applicable.

Competing interests

The authors have stated that there are none.

Acknowledgements

We wish to thank Drs Judith Bell and Giri Rajaratnam, Professor Peter Croft and Brian Dudley for commenting on the study design and draft paper.

Discuss this article

Contribute and read comments about this article on the Discussion Forum: <http://www.rcgp.org.uk/bjgp-discuss>

REFERENCES

- Nexoe J, Halvorsen PA, Kistiansen IS. Critiques of the risk concept — valid or not? *Scand J Public Health* 2007; **35**(6): 648–654.
- Loewenstein GF, Weber EU, Hsee CK, Welch N. Risks as feelings. *Psychol Bull* 2001; **127**(2): 267–286.
- Sackett DL, Strauss SE, Richardson WS, et al. *Evidence-based medicine: how to practice and teach EBM*. Edinburgh: Churchill-Livingston, 2000.
- National Institute for Health and Clinical Excellence. *Referral for suspected cancer: a clinical practice guideline. National Clinical Practice Guideline Number 27*. London: National Institute for Health and Clinical Excellence, 2005.
- Mulka O. NICE suspected cancer guidelines. *Br J Gen Pract* 2005; **55**(517): 580–581.
- National Institute for Health and Clinical Excellence. *Cardiovascular risk assessment and the modification of blood lipids for the primary and secondary prevention of cardiovascular disease. National Clinical Practice Guideline Number 67*. London: National Institute for Health and Clinical Excellence, 2009.
- National Prescribing Centre. *Statins: standard doses*. Leeds: National Prescribing Centre, 2009. http://www.npci.org.uk/therapeutics/cardio/cdlipids/resources/pda_Lipids.pdf (accessed 20 Jul 2010).
- Bowles CJA, Leicester R, Romaya C, et al. A prospective study of colonoscopy practice in the UK today: are we adequately prepared for national colorectal screening tomorrow? *Gut* 2004; **53**(2): 277–283.
- Wells GA, Shea B, O'Connell D, et al. *The Newcastle–Ottawa Scale (NOS) for assessing the quality of nonrandomised studies in meta-analyses*. Ottawa: Ottawa Health Research Institute. http://www.ohri.ca/programs/clinical_epidemiology/oxford.htm (accessed 20 Jul 2010).
- Higgins JPT, Thompson SG, Deeks JJ, Altman DG. Measuring inconsistency in meta-analyses. *BMJ* 2003; **327**(7414): 557–560.
- Altman DG, Machin D, Bryant TN, Gardner MJ. *Statistics with confidence*. 2nd edn. London: BMJ Books, 2000.
- Jones R, Latinovic R, Charlton J, Gulliford MC. Alarm symptoms in early diagnosis of cancer in primary care: cohort study using General Practice Research Database. *BMJ* 2007; **334**(7602): 1040.
- Parker C, Hippisley-Cox J, Coupland C, Vinogradova Y. Rectal and postmenopausal bleeding: consultation and referral of patients with and without severe mental health problems. *Br J Gen Pract* 2007; **57**(538): 371–376.
- Lawrenson R, Logie J, Marks C. Risk of colorectal cancer in general practice patients presenting with rectal bleeding, change in bowel habit or anaemia. *Eur J Cancer Care (Engl)* 2006; **15**(3): 267–271.
- Wauters H, Van Casteren V, Buntinx F. Rectal bleeding and colorectal cancer in general practice: diagnostic study. *BMJ* 2000; **321**(7267): 998–999.
- Du Toit J, Hamilton W, Barraclough K. Risk in primary care of colorectal cancer from new onset rectal bleeding: 10 year prospective study. *BMJ* 2006; **333**(7558): 69–70.
- Heintze C, Matysiak-Klose D, Krohn T, et al. Diagnostic work-up of rectal bleeding in general practice. *Br J Gen Pract* 2005; **55**(510): 14–19.
- Ellis BG, Thompson MR. Factors identifying higher risk rectal bleeding in general practice. *Br J Gen Pract* 2005; **55**(521): 949–955.
- Nørrelund N, Nørrelund H. Colorectal cancer and polyps in patients aged 40 years and over who consult a GP with rectal bleeding. *Fam Pract* 1996; **13**(2): 160–165.
- Metcalf JW, Smith J, Jones R, Record CO. Incidence and causes of rectal bleeding in general practice as detected by colonoscopy. *Br J Gen Pract* 1996; **46**(404): 161–164.
- Fijten GH, Starman R, Muris JW, et al. Predictive value of signs and symptoms for colorectal cancer in patients with rectal bleeding in general practice. *Fam Pract* 1995; **12**(3): 279–286.
- Mant A, Bokey EL, Chapuis PH, et al. Rectal bleeding: do other symptoms aid in diagnosis? *Dis Colon Rectum* 1989; **32**(3): 191–196.
- Hamilton W, Round A, Sharp D, Peters TJ. Clinical features of colorectal cancer before diagnosis: a population-based case-control study. *Br J Cancer* 2005; **93**(4): 399–405.
- Hamilton W, Lancashire R, Sharp D, et al. The risk of colorectal cancer with symptoms at different ages and between the sexes: a case-control study. *BMC Med* 2009; **7**: 17.
- Thompson JA, Pond CL, Ellis BG, et al. Rectal bleeding in general and hospital practice; the tip of the iceberg. *Colorectal Dis* 2000; **2**(5): 288–293.
- Leicester RJ, Colinjones DG, Hunt RH, et al. Hemocult testing in general-practice for early diagnosis of colorectal-cancer. *Gut* 1984; **25**(5): A561.
- Stellon AJ, Kenwright SE. Iron deficiency anaemia in general practice: Presentations and investigations. *Br J Clin Pract* 1997; **51**(2): 78–80.
- Yates JM, Logan EC, Stewart RM. Iron deficiency anaemia in general practice: clinical outcomes over three years and factors influencing diagnostic investigations. *Postgrad Med J* 2004; **80**(945): 405–410.
- Hamilton W, Lancashire R, Sharp D, et al. The importance of anaemia in diagnosing colorectal cancer: a case-control study using electronic primary care records. *Br J Cancer* 2008; **98**(2): 323–327.
- Bruyninckx R, Buntinx F, Aertgeerts B, Van Casteren V. The diagnostic value of macroscopic haematuria for the diagnosis of urological cancer in general practice. *Br J Gen Pract* 2003; **53**(486): 31–35.
- Hamilton W, Sharp DJ, Peters TJ, Round AP. Clinical features of prostate cancer before diagnosis: a population-based, case-control study. *Br J Gen Pract* 2006; **56**(531): 756–762.
- Hamilton W, Peters TJ, Round A, Sharp D. What are the clinical features of lung cancer before the diagnosis is made? A population based case-control study. *Thorax* 2005; **60**(12): 1059–1065.
- Esfandyari T, Potter JW, Vaezi MF. Dysphagia: a cost analysis of the diagnostic approach. *Am J Gastroenterol* 2002; **97**(11): 2733–2737.
- Eberl MM, Phillips RL Jr, Lamberts H, et al. Characterizing breast symptoms in family practice. *Ann Fam Med* 2008; **6**(6): 528–533.
- Barton MB, Elmore JG, Fletcher SW. Breast symptoms among women enrolled in a Health Maintenance Organisation: frequency, evaluation, and outcome. *Ann Intern Med* 1999; **130**(8): 651–657.
- Bywaters JL. The incidence and management of female breast disease in a general practice. *J R Coll Gen Pract* 1977; **27**(179): 353–357.
- Office for National Statistics. *Cancer statistics registrations*. http://www.statistics.gov.uk/downloads/theme_health/MB1-37/MB1_37_2006.pdf (accessed 20 Jul 2010).
- Bekkink MO, McCowen C, Falk GA, et al. Diagnostic accuracy systematic review of rectal bleeding in combination with other symptoms, signs and tests in relation to cancer. *Br J Cancer* 2010; **102**(1): 48–58.
- Ford AC, Veldhuyzen van Zanten SJO, Rodgers CC, et al. Diagnostic utility of alarm features for colorectal cancer: Systematic review and meta-analysis. *Gut* 2008; **57**(11): 1545–1552.
- Jellema P, van der Windt DAWM, Bruinvels DJ, et al. Value of symptoms and additional diagnostic tests for colorectal cancer in primary care: systematic review and meta-analysis. *BMJ* 2010; **340**: c1269.
- Gigerenzer G, Edwards A. Simple tools for understanding risk: from innumeracy to insight. *BMJ* 2003; **327**(7417): 741–744.
- Edwards AGK, Robling MR, Wilkinson C, et al. The presentation and management of breast symptoms in general practice in South Wales. *Br J Gen Pract* 1999; **49**(447): 811–812.
- Newton P, Hannay D, Laver R. The presentation and management of female breast symptoms in general practice in Sheffield. *Fam Pract* 1999; **16**(4): 360–365.
- Logan ECM, Yates JM, Stewart RM, et al. Investigation and management of iron deficiency anaemia in general practice: a cluster randomised controlled trial of a simple management prompt. *Postgrad Med J* 2002; **78**(923): 533–537.