

Conserved vertebrate *mir-451* provides a platform for Dicer-independent, Ago2-mediated microRNA biogenesis

Jr-Shiuan Yang^a, Thomas Maurin^a, Nicolas Robine^a, Kasper D. Rasmussen^b, Kate L. Jeffrey^c, Rohit Chandwani^c, Eirini P. Papapetrou^d, Michel Sadelain^d, Dónal O'Carroll^b, and Eric C. Lai^{a,1}

^aDepartment of Developmental Biology, Sloan-Kettering Institute, New York, NY 10065; ^bEuropean Molecular Biology Laboratory, Mouse Biology Unit, 00015 Monterotondo Scalo, Italy; ^cLaboratory for Lymphocyte Signaling, The Rockefeller University, New York, NY 10065; and ^dCenter for Cell Engineering, Molecular Pharmacology and Chemistry Program, Memorial Sloan-Kettering Cancer Center, New York, NY 10065

Edited by Gary Ruvkun, Massachusetts General Hospital, Boston, MA, and approved July 16, 2010 (received for review May 10, 2010)

Canonical animal microRNAs (miRNAs) are generated by sequential cleavage of precursor substrates by the Drosha and Dicer RNase III enzymes. Several variant pathways exploit other RNA metabolic activities to generate functional miRNAs. However, all of these pathways culminate in Dicer cleavage, suggesting that this is a unifying feature of miRNA biogenesis. Here, we show that maturation of miR-451, a functional miRNA that is perfectly conserved among vertebrates, is independent of Dicer. Instead, structure-function and knockdown studies indicate that Drosha generates a short *pre-mir-451* hairpin that is directly cleaved by Ago2 and followed by resection of its 3' terminus. We provide stringent evidence for this model by showing that Dicer knockout cells can generate mature miR-451 but not other miRNAs, whereas Ago2 knockout cells reconstituted with wild-type Ago2, but not Slicer-deficient Ago2, can process miR-451. Finally, we show that the *mir-451* backbone is amenable to reprogramming, permitting vector-driven expression of diverse functional miRNAs in the absence of Dicer. Beyond the demonstration of an alternative strategy to direct gene silencing, these observations open the way for transgenic rescue of Dicer conditional knockouts.

Slicer | gene suppression | miRNA reprogramming

MicroRNAs (miRNAs) are ~22-nt regulatory RNAs that collectively serve essential functions in higher eukaryotes (1). Conserved biogenesis machinery governs the production of canonical miRNAs in invertebrate and vertebrate cells (2). Primary miRNA (pri-miRNA) transcripts containing one or more hairpin structures are first cleaved in the nucleus by the Drosha RNase III enzyme and its dsRNA binding partner DiGeorge syndrome critical region gene 8 (DGCR8)/Pasha to yield ~55–70 nt precursor miRNA (pre-miRNA) hairpins. These are cleaved again in the cytoplasm by the Dicer RNase III enzyme to yield a miRNA/miRNA* duplex, from which one strand matures in a complex with an Argonaute (Ago) protein. The miRNA guides the Ago complex to target transcripts, often bearing 7-nt complements to the 5' end of the miRNA, for destabilization and/or translational inhibition (3). Mammalian genomes encode four Ago-class proteins, of which Ago2 is uniquely capable of directly cleaving highly complementary targets (4, 5), a process popularly termed slicing.

Since the elucidation of this framework, several alternative pathways for miRNA biogenesis have emerged. For example, mirtrons are short hairpin introns that exploit the splicing machinery to generate pre-miRNA hairpins, thereby bypassing Drosha cleavage (6–8). Functional miRNAs can also be generated from certain small nucleolar RNAs (9, 10) and tRNAs (11, 12), presumably through Drosha-independent mechanisms. The viral miRNA precursor miR-M1-7 from murine γ -herpesvirus 68 fuses a tRNA to an miRNA hairpin and is processed by tRNase-Z (13). Endogenous shRNAs (endo-shRNAs) are also independent of the canonical nuclear processing machinery, and their hairpin termini might be defined by RNA polymerase III (11).

Common to these sundry biogenesis pathways is cleavage of an intermediate precursor by Dicer and routing of the mature small

RNA into an Argonaute, suggesting that these are defining features of miRNA-class regulatory RNAs. In this study, we show that the maturation of the highly conserved vertebrate miR-451 bypasses Dicer, and instead requires direct cleavage of its precursor hairpin through Ago2 Slicer activity. We exploit these properties of *mir-451* as a flexible platform for Dicer-independent, vector-mediated expression of miRNAs.

Results

Atypical Conservation and Pattern of Small RNA Reads Derived from *mir-451*. Many animal miRNA loci are clustered, and this property identified a conserved hairpin christened *mir-451*, located 100 bp downstream of *mir-144* (14). As is typical for conserved miRNA genes, *mir-144* exhibits a saddle-shaped pattern of divergence in which its terminal loop exhibits many more nucleotide substitutions than its stems and for which the miRNA* is slightly less constrained than the miRNA (15, 16). In sharp contrast, the terminal loop of *mir-451* is invariant from human to fish, whereas specific nucleotides in the stem are variable (Fig. 1*A* and *B*).

The unusual conservation features of *mir-451* are linked to an atypical pattern of small RNA reads. Its dominantly cloned species extend across the terminal loop into the complementary side of the hairpin (17), and even longer reads were observed. We found these to be reproducible and conserved characteristics of *mir-451*. In addition to dominant 23-nt reads typical of miRNAs, a population of 24- to 30-nt reads extending well into the complementary hairpin arm is produced by *mir-451* in human (18), mouse (19–21), dog (22), and chicken (23) (Fig. 1*C* and [Dataset S1](#)). In contrast to its 3' heterogeneity, the 5' terminus of miR-451 was precisely defined in all species examined ([Dataset S1](#)).

We confirmed the atypical sizes of *mir-451*-derived species with Northern analysis of human K562 cells and murine erythroleukemia (MEL) cells induced with hexamethylene bisacetamide (HMB). By comparison, its partnered gene *mir-144* generated an ~58-nt pre-miRNA and an ~22-nt mature miRNA, as expected for a canonical miRNA gene (Fig. 1*D*). In contrast, a probe antisense to miR-451 detected a series of bands extending past 30 nt (Fig. 1*E*). This was recapitulated by lentiviral transduction of murine *mir-144/451* into uninduced MEL cells or by transfection of a human *mir-144/mir-451* plasmid into HeLa cells. Notably, the >30-nt bands detected by miR-451 probe exhibited variable mobility from ~32 to 42 nt according to gel percentage and temperature (Figs. 1*F* and [Fig. S1](#)). Previous studies noted

Author contributions: J.-S.Y., T.M., and E.C.L. designed research; J.-S.Y., T.M., and N.R. performed research; K.D.R., K.L.J., R.C., E.P.P., and D.O. contributed new reagents/analytical tools; J.-S.Y., T.M., N.R., M.S., and E.C.L. analyzed data; and E.C.L. wrote the paper.

The authors declare no conflict of interest.

This article is a PNAS Direct Submission.

Freely available online through the PNAS open access option.

¹To whom correspondence should be addressed. E-mail: laie@mskcc.org.

This article contains supporting information online at www.pnas.org/lookup/suppl/doi:10.1073/pnas.1006432107/-DCSupplemental.




Fig. 1. Atypical conservation and pattern of small RNAs from *mir-451*. (A) *mir-144/mir-451* region from the University of California Santa Cruz Genome Browser; mature miRNAs (green) are encoded by the bottom strand. *mir-144* exhibits a typical evolutionary pattern with far greater divergence in the terminal loop (triangle) than in mature miRNA or star arm (yellow). *mir-451* is more conserved in its terminal loop than its 3' hairpin arm. (B) Vertebrate *mir-451* alignments illustrate the constrained terminal loop; asterisk marks a variable position in the 3' hairpin arm. (C) Dominant *mir-451* reads are 23 nt (green highlighted box), but larger species up to 30 nt (light green triangle) are observed in multiple species; rare reads are derived from the 3p arm. Data were analyzed from GSM494811 (K562) and GSM433295 (mouse testis); see [Dataset S1](#) for full analysis of these and other *mir-451*-containing libraries. (D) *mir-144* generates typical pre-miRNA hairpin and mature miRNA. (E) *mir-451* generates small RNAs ranging from 22 to 24 nt to >30 nt. Endogenous *mir-144* and *mir-451* were detected in K562 cells or MEL cells induced with HMBA; these miRNAs are not expressed by uninduced MEL or GFP-transduced MEL cells. Ectopic miRNAs were generated by transfection of HeLa cells with *mir-144/451* plasmid or transduction of MEL cells with *mir-144/451* lentivirus. (F) The two upper bands (**) in *mir-451* blots exhibit differential mobility in different acrylamide gels from ~32 to >40 nt. (G) Sensor assays in HeLa cells transfected with *mir-144/451* show repression of *mir-144-3p* and *mir-451* perfect targets and of *mir-451* seed target. Sensor values were normalized against *mir-1-2* construct; SDs from quadruplicate assays are shown.

that short hairpins can exhibit atypical mobility even under denaturing conditions (24), suggesting that these bands correspond to *pre-mir-451* hairpins (Fig. 1B).

Despite its atypical properties, *mir-451* regulates erythroid development and represses endogenous transcripts (25–27). We verified the regulatory activity of *mir-451* using model luciferase constructs containing perfectly complementary sites or bulged *mir-451* target sites. Both targets were repressed by a *mir-451/mir-144* construct relative to a noncognate miRNA expression construct (*mir-1-2*) in HeLa cells, with severalfold greater repression of the perfect targets than the bulged targets (Fig. 1G). Thus, *mir-451* has the regulatory capacity of a typical miRNA.

Structural Requirements for Biogenesis and Function of *mir-451*. To gain insight into the atypical biogenesis of *mir-451*, we investigated structurally variant precursors (Fig. 2A). We first tested the role of the stem basal to the duplex involving mature *mir-451*, which is broadly conserved and resembles the lower stem of ca-

nonical pri-miRNAs. Structurally nonconservative mutations abrogated its ability to repress *mir-451* sensor (Fig. 2B), implicating processing by Drosha/DGCR8. In addition, libraries from pro- and pre-B cells (21) contained reads with heterogeneous 5' ends and precise 3' ends that directly abutted mature *mir-451* ([Dataset S1](#)); such “moR” reads are diagnostic of Drosha processing (28, 29). As well, human blood cells (18), mouse testis (20), and dog lymphocytes (22) express rare RNAs mapping downstream of mature *mir-451* (Fig. 1C and [Dataset S1](#)). Many of these extend to a position constituting a 2-nt 3' overhang to the strict 5' end of *mir-451* (Fig. 2A, black arrowheads), consistent with RNase III cleavage. These are probably not related to *mir-451* biogenesis, because they overlap within the terminal loop of *pre-mir-451*. They might result from breakage within the unstructured loop, perhaps reminiscent of tRNA cleavage within anticodon loops (30). In any case, these data support the existence of an ~42-nt *pre-mir-451* hairpin produced by Drosha/DGCR8 cleavage (Figs. 1C and 2A), as indicated by Northern analysis (Fig. 1F).

mobility compared with wild type. The levels of mature miR-451 paralleled the sensor assays: the double-cleavage site mutant produced almost none, whereas the single mutants produced slightly more miR-451; the lower-stem mutant yielded neither *pre-mir-451* nor shorter matured RNAs. The various *mir-451* mutations did not substantially affect the biogenesis of *mir-144* present on these constructs, which served as an internal control to these experiments (Fig. 2D). Taken together, these tests implied Drosha cleavage and Ago2 slicing in the biogenesis of miR-451.

Drosha-Dependent, Dicer-Independent Biogenesis of miR-451. We followed these structural studies with genetic manipulation of miRNA processing factors by introducing siRNAs to HeLa cells. The knockdown efficiencies were confirmed by qPCR analysis (Fig. S2). Consistent with the inability of the *pri-mir-451* lower-stem mutant to be matured (Fig. 2B), HeLa cells depleted for Drosha or its partner DGCR8 were compromised in their ability to generate *mir-451* intermediates or mature miR-451 (Fig. 3A). We also probed against its “star” arm (that is, the 3p arm complementary to mature miR-451). We expected this to label the *pre-mir-451* hairpin but not the Ago2-cleaved hairpin, because the latter contains <9-nt complementarity to the probe. We observed hybridization to bands of apparent ~35-nt length in 16% gels (labeled pre), which comigrated with the larger bands detected by miR-451 probe (Fig. 3A). This implied that these species contain the full 42-nt *mir-451* hairpin sequence.

The 42-nt Drosha-cleaved *mir-451* hairpin does not possess sufficient duplex to be a Dicer substrate, suggesting that its maturation might bypass Dicer. We performed stringent tests using a viable line of mouse embryonic fibroblasts (MEFs) bearing the floxed allele of *Dicer* lacking exon 22 (32). These cells do not express Dicer protein (Fig. 3B) and are arrested for endogenous miRNA biogenesis at the pre-miRNA stage. Nevertheless, transfection of *mir-144/451* construct into *Dicer*^{-/-} MEFs yielded processed small RNAs (Fig. 3B) with strong repressive capacity, both on perfect and bulged miR-451 sensors (Fig. 3C). These data show a Dicer-independent, miRNA-class regulatory RNA in animal cells.

Ago2 Slicer Activity Is Essential for Maturation of miR-451. Conditional knockout of *mAgo2* in the hematopoietic system reduces the level of many miRNAs (33). However, among miRNAs expressed in bone marrow, miR-451 was uniquely susceptible to loss of *Ago2*. We isolated RNA from wild-type and *Ago2*^{-/-} bone marrow (reconstituted in lethally irradiated recipients from *Ago2*

[fl/fl]; MxCre donors) and analyzed their small RNAs by microarray. Not only was miR-451 the most abundant miRNA in wild-type bone marrow, its nearly complete absence (>400-fold lower) in the *Ago2* mutant made it by far the most highly depleted miRNA (Fig. 4A; see raw and processed data in Dataset 3). We used Northern analysis to confirm the complete loss of intermediate and mature forms of miR-451 in *Ago2*^{-/-} bone marrow, which instead accumulated *pre-mir-451* (Fig. 4B).

The strict Ago2 dependency of miR-451 was consistent with the inefficient maturation of *pri-mir-451* cleavage-site mutants (Fig. 2C). However, these data do not exclude that changes in the erythroid compartment of *Ago2* mutants might contribute to loss of miR-451 (33) nor do they directly address the requirement of Ago2 Slicer activity. We tested this using a panel of *Ago2*^{-/-} MEFs reconstituted with control MigR retrovirus or retroviruses carrying wild-type mAgo2 or catalytically inactive (D669A) mAgo2 (33). Western blots showed equivalent Ago2 expression levels in the reconstituted cells (Fig. 4C Top), and the *mir-451/mir-144* construct was efficiently expressed as assessed by quantitative reverse transcription-polymerase chain reaction across the primary transcript (Fig. S3). However, only *Ago2*^{-/-} cells reconstituted with wild-type Ago2 could mature miR-451 from its hairpin precursor (Fig. 4C Middle).

We examined the state of *mir-451* intermediates in wild-type and catalytic-dead Ago2 proteins by immunoprecipitating Ago2 followed by Northern analysis. As seen in Fig. 4D, wild-type Ago2 contained matured miR-451 species, whereas Ago2[D669A] associated exclusively with the hairpin precursor. Finally, to account for the presence of *pre-mir-451* in the absence of *Ago2*, we checked whether the hairpin was incorporated into other Agos. We observed that Ago1 immunoprecipitated with *pre-mir-451* but not matured miR-451 species (Fig. S4). This confirmed that Ago2-mediated slicing is essential for the biogenesis of miR-451 and cannot be substituted by other Agos. We hypothesize that hairpin cleavage renders its 3' end amenable to resection by ribonuclease (s) to generate 23- to 24-nt mature miR-451 (Fig. 4E).

miR-451 Backbone Confers Dicer-Independent Expression of Other miRNAs. Many studies have used *Dicer* knockouts to assess biological processes that are dependent on miRNAs. For example, maternal and zygotic (MZ) loss of zebrafish *Dicer* impairs clearance of maternal transcripts during the embryonic maternal-zygotic transition (34). This was demonstrably due to miRNAs in the miR-430 family, because injection of synthetic miR-430




Fig. 4. Ago2 Slicer activity mediates miR-451 maturation. (A) Microarray profiling of wild-type and *Ago2*^{-/-} bone marrow revealed that miR-451 is the highest-expressed miRNA in bone marrow and uniquely deficient in the absence of Ago2. (B) Northern blot verifies loss of processed miR-451 intermediates in *Ago2*^{-/-} marrow. (C Top) Western blot verification of *Ago2*^{-/-} MEFs reconstituted with control virus, Ago2-expressing virus, or Ago2[D669A] virus. (Bottom) Transfection of *mir-144/mir-451* construct into this panel of cells shows that Ago2 Slicer function is strictly required for production of mature miR-451. (D Upper) Ago2 proteins were immunoprecipitated from the panel of reconstituted *Ago2*^{-/-} MEFs. (Lower) Associated small RNAs were analyzed by Northern blot and probed for miR-451; Ago2[D669A] cannot mature miR-451. (E) Model for *mir-451* processing.

into *MZ-Dicer* embryos rescued early development. However, transgenic rescue of miRNAs in *Dicer* mutant tissue has not been possible, because miRNA biogenesis generally requires Dicer.

We therefore investigated whether *pri-mir-451* was amenable to reprogramming. This was not necessarily possible, given the strict conservation of miR-451 primary sequence (Fig. 1B). Still, we replaced the human *mir-451* hairpin with a variety of miRNA sequences, while maintaining its secondary structure, and performed initial tests of perfect sensors in HeLa cells. Encouragingly, we observed significant repression by several miRNAs reprogrammed into the *mir-451* backbone (Fig. 5A). With miR-199-3p, the activity of the reprogrammed construct was comparable to its endogenous *pri-miRNA* context, yielding ~7-fold repression compared with a noncognate miRNA (*mir-1-2*). In total, six of seven reprogrammed constructs were active, although they varied in their magnitude of target repression.

We challenged these reprogrammed constructs to function in *Dicer*^{-/-} MEFs, in which canonical miRNA constructs are non-functional (Fig. 5B). We observed robust activity (20- to 50-fold target repression) of all constructs in the absence of Dicer, including ones whose activity was marginal in HeLa cells. Moreover, reprogrammed *mir-451* constructs could strongly repress bulged targets (Fig. 5B), providing evidence for the accurate definition of 5' ends of reprogrammed miRNAs in *Dicer* knockout

cells as with miR-451 (Fig. 3C). Although canonical miRNAs are inactive in these mutant cells, we could perform meaningful normalizations by comparing reprogrammed constructs with a functional reprogrammed 451:miR-23a-3p construct (whose activity was, in turn, normalized to functional 451:miR-199a-3p).

Ago2 can directly use some pre-miRNAs and other long RNAs as guide molecules, and their loading is potentiated in the absence of Dicer (35). Therefore, sensor tests do not necessarily report on Dicer-independent miRNA production. To address this, we analyzed *Dicer*^{-/-} MEFs transfected with reprogrammed *mir-451* constructs and observed accumulation of small RNAs in all cases (Fig. 5C). Some constructs generated a ladder of bands, as with miR-451, whereas others predominantly accumulated a single species. Further analysis of these constructs may provide insight into the nature and efficiency of the resection activity on different Ago2-cleaved hairpin substrates.

Finally, comparing these constructs in *Dicer*^{-/-} MEFs and HeLa cells, we observed several instances where small RNA maturation was apparently increased in the absence of Dicer (Fig. S5). It is possible that lack of Ago2 loading of endogenous Dicer-dependent substrates (i.e., miRNAs and/or siRNAs) might enhance biogenesis and activity of miR-451-based constructs. However, other differences in small RNA pathway status or gene expression could well contribute to the distinct activity of these constructs in




Fig. 5. The *mir-451* backbone confers Dicer-independent expression of other miRNAs. (A) *mir-451* hairpins were reprogrammed with other miRNAs and tested against perfect sensors in HeLa (blue bars). Activities of canonical miRNA constructs are shown for comparison (orange bars); all tests were normalized to *mir-1-2* as a noncognate miRNA (white bars). (B) Canonical miRNA constructs are essentially inactive in *Dicer*^{-/-} MEFs (orange bars), whereas reprogrammed *mir-451* hairpins are highly active (blue bars). We also observed strong repression of bulged sensors (dark blue bars), indicating that the 5' ends of miRNAs from reprogrammed constructs were defined accurately. Data were normalized to sensor activity in the presence of 451:miR-23a-3p as a noncognate functional control miRNA; 451:miR-199a-3p was used as a control for the reprogrammed miR-23a-3p sensor test. Quadruplicate assays were performed, and SDs were plotted. (C) Northern analysis confirms the failure of *Dicer*^{-/-} MEFs to process ectopic canonical miRNAs. In contrast, short RNAs were produced from reprogrammed *mir-451* backbones in *Dicer*^{-/-} MEFs; analysis of untransfected cells reveals the accumulation of some endogenous pre-miRNA species (asterisks).

different cells; thus, additional tests of a competition model for Ago2 loading are necessary. In any case, these data provide strict evidence that the *mir-451* backbone permits diverse, functional, processed miRNAs to be generated in *Dicer* knockout cells.

Discussion

Our studies show that the highly conserved vertebrate *pre-mir-451* hairpin is independent of *Dicer* and instead, is reliant on direct cleavage by Ago2. This differs from its role in generating Ago2-cleaved precursor miRNAs, in which Ago2 was proposed to cleave certain well-paired pre-miRNAs before *Dicer* cleavage, perhaps facilitating removal of passenger-strand products (31). In the case of *mir-451*, slicing of its short pre-miRNA hairpin is absolutely prerequisite for further 3' trimming to generate the mature miRNA and cannot be supported by nonlicing Ago proteins. This adds to other endogenous processes in mammalian cells, including regulation of highly or perfectly complementary targets by certain miRNAs (36) and endogenous siRNAs (11, 19, 37) that may collectively underlie the retention of cleavage activity by vertebrate Ago2.

While our work was under review, the Hannon and Giraldez groups reported similar evidence regarding the Drosha-dependent, *Dicer*-independent, Ago2-mediated biogenesis of miR-451 in mammalian (38) and zebrafish (39) systems. Our biogenesis data are consistent with their data and extend the capacity of this alternative biogenesis pathway for small RNA expression. Beyond the analysis of synthetic *mir-451*-based RNA hairpins (38, 39), we provide broad evidence for flexible reprogramming of *mir-451* using DNA vectors. In particular, we showed efficient processing of an artificial *mir-451*-based construct into miRNA-sized species that could regulate both perfect and bulged targets, even in *Dicer* knockout cells. These data provide proof of principle that the *mir-451* backbone may permit transgenic expression of individual

miRNAs in systemic or conditional murine *Dicer* knockouts (32, 40–42), opening the potential for genetic rescues in the animal.

We have not yet identified a nuclease responsible for resection of the 3' end of Ago2-cleaved *mir-451*. A nuclease is inferred to down-regulate pre-miRNAs after their 3' uridylation by terminal uridylyltransferase 4 (43), but its identity has similarly been elusive. Still, the collected studies suggest that nucleases targeting the 3' ends of small RNA precursors have both negative and positive roles for miRNA biogenesis. We also note that the efficient function of miR-451 and many reprogrammed miRNAs in *Dicer*^{-/-} MEFs indicates that *Dicer* is not essential for substrate loading into Ago2. In light of the discovery that functional *Dicer*-independent “primal RNAs” associate with *Schizosaccharomyces pombe* Ago1 (44), direct loading of RNAs into Argonaute proteins may be an ancient strategy.

Materials and Methods

Primers used for cloning, quantitative polymerase chain reaction, and blotting are listed in [Dataset S2](#). Methods for Northern analysis, cell culture and luciferase sensor assays (6), *Dicer* mutant (32), and Ago2 mutant (33) were previously described. Microarray experiments was performed by LC Sciences. Detailed methods are provided in [SI Materials and Methods](#).

ACKNOWLEDGMENTS. We thank Jidong Liu (Sloan-Kettering Institute, New York) for Ago plasmids, Gunter Meister (Universität Regensburg, Regensburg, Germany) for Ago antibodies, Michael Chen for mapping some of the small RNA data, Katsutomo Okamura for critical comments, and Alexander Tarakhovskiy (The Rockefeller University, New York) for support and discussion. We are grateful to the many laboratories that enabled our studies by depositing their published small RNA datasets into public databases. Work in E.C.L.'s group was supported by the Burroughs Wellcome Fund (1004721), the Alfred Bressler Scholars Fund, the Starr Cancer Consortium (I3-A139), and the National Institutes of Health (R01-GM083300).

- Flynt AS, Lai EC (2008) Biological principles of microRNA-mediated regulation: Shared themes amid diversity. *Nat Rev Genet* 9:831–842.
- Kim VN, Han J, Siomi MC (2009) Biogenesis of small RNAs in animals. *Nat Rev Mol Cell Biol* 10:126–139.
- Filipowicz W, Bhattacharyya SN, Sonenberg N (2008) Mechanisms of post-transcriptional regulation by microRNAs: Are the answers in sight? *Nat Rev Genet* 9:102–114.
- Meister G, et al. (2004) Human Argonaute2 mediates RNA cleavage targeted by miRNAs and siRNAs. *Mol Cell* 15:185–197.
- Liu J, et al. (2004) Argonaute2 is the catalytic engine of mammalian RNAi. *Science* 305:1437–1441.
- Okamura K, Hagen JW, Duan H, Tyler DM, Lai EC (2007) The mirtron pathway generates microRNA-class regulatory RNAs in *Drosophila*. *Cell* 130:89–100.
- Ruby JG, Jan CH, Bartel DP (2007) Intronic microRNA precursors that bypass Drosha processing. *Nature* 448:83–86.
- Berezikov E, Chung WJ, Willis J, Cuppen E, Lai EC (2007) Mammalian mirtron genes. *Mol Cell* 28:328–336.
- Ender C, et al. (2008) A human snoRNA with microRNA-like functions. *Mol Cell* 32:519–528.
- Taft RJ, et al. (2009) Small RNAs derived from snoRNAs. *RNA* 15:1233–1240.
- Babiarz JE, Ruby JG, Wang Y, Bartel DP, Blelloch R (2008) Mouse ES cells express endogenous shRNAs, siRNAs, and other Microprocessor-independent, *Dicer*-dependent small RNAs. *Genes Dev* 22:2773–2785.
- Cole C, et al. (2009) Filtering of deep sequencing data reveals the existence of abundant *Dicer*-dependent small RNAs derived from tRNAs. *RNA* 15:2147–2160.
- Bogerd HP, et al. (2010) A mammalian herpesvirus uses noncanonical expression and processing mechanisms to generate viral microRNAs. *Mol Cell* 37:135–142.
- Altuvia Y, et al. (2005) Clustering and conservation patterns of human microRNAs. *Nucleic Acids Res* 33:2697–2706.
- Lai EC, Tomancak P, Williams RW, Rubin GM (2003) Computational identification of *Drosophila* microRNA genes. *Genome Biol* 4:R42.1–20.
- Berezikov E, et al. (2005) Phylogenetic shadowing and computational identification of human microRNA genes. *Cell* 120:21–24.
- Nelson PT, et al. (2007) A novel monoclonal antibody against human Argonaute proteins reveals unexpected characteristics of miRNAs in human blood cells. *RNA* 13:1787–1792.
- Vaz C, et al. (2010) Analysis of microRNA transcriptome by deep sequencing of small RNA libraries of peripheral blood. *BMC Genomics* 11:288.
- Tam OH, et al. (2008) Pseudogene-derived small interfering RNAs regulate gene expression in mouse oocytes. *Nature* 453:534–538.
- Vagin VV, et al. (2009) Proteomic analysis of murine Piwi proteins reveals a role for arginine methylation in specifying interaction with Tudor family members. *Genes Dev* 23:1749–1762.
- Kuchen S, et al. (2010) Regulation of microRNA expression and abundance during lymphopoiesis. *Immunity* 32:828–839.
- Friedländer MR, et al. (2008) Discovering microRNAs from deep sequencing data using miRDeep. *Nat Biotechnol* 26:407–415.
- Glazov EA, et al. (2008) A microRNA catalog of the developing chicken embryo identified by a deep sequencing approach. *Genome Res* 18:957–964.
- Hirao I, Nishimura Y, Tagawa Y, Watanabe K, Miura K (1992) Extraordinarily stable mini-hairpins: electrophoretic and thermal properties of the various sequence variants of d (CGGAAAGC) and their effect on DNA sequencing. *Nucleic Acids Res* 20:3891–3896.
- Papapetrou EP, Korkola JE, Sadelain M (2010) A genetic strategy for single and combinatorial analysis of miRNA function in mammalian hematopoietic stem cells. *Stem Cells* 28:287–296.
- Dore LC, et al. (2008) A GATA-1-regulated microRNA locus essential for erythropoiesis. *Proc Natl Acad Sci USA* 105:3333–3338.
- Rasmussen KD, et al. (2010) The miR-144/451 locus is required for erythroid homeostasis. *J Exp Med* 207:1351–1358.
- Ruby JG, et al. (2007) Evolution, biogenesis, expression, and target predictions of a substantially expanded set of *Drosophila* microRNAs. *Genome Res* 17:1850–1864.
- Shi W, Hendrix D, Levine M, Haley B (2009) A distinct class of small RNAs arises from pre-miRNA-proximal regions in a simple chordate. *Nat Struct Mol Biol* 16:183–189.
- Thompson DM, Parker R (2009) Stressing out over tRNA cleavage. *Cell* 138:215–219.
- Diederichs S, Haber DA (2007) Dual role for argonautes in microRNA processing and posttranscriptional regulation of microRNA expression. *Cell* 131:1097–1108.
- Yi R, et al. (2006) Morphogenesis in skin is governed by discrete sets of differentially expressed microRNAs. *Nat Genet* 38:356–362.
- O'Carroll D, et al. (2007) A Slicer-independent role for Argonaute 2 in hematopoiesis and the microRNA pathway. *Genes Dev* 21:1999–2004.
- Giraldez AJ, et al. (2005) MicroRNAs regulate brain morphogenesis in zebrafish. *Science* 308:833–838.
- Tan GS, et al. (2009) Expanded RNA-binding activities of mammalian Argonaute 2. *Nucleic Acids Res* 37:7533–7545.
- Yekta S, Shih IH, Bartel DP (2004) MicroRNA-directed cleavage of HOXB8 mRNA. *Science* 304:594–596.
- Watanabe T, et al. (2008) Endogenous siRNAs from naturally formed dsRNAs regulate transcripts in mouse oocytes. *Nature* 453:539–543.
- Cheloufi S, Dos Santos CO, Chong MM, Hannon GJ (2010) A *Dicer*-independent miRNA biogenesis pathway that requires Ago catalysis. *Nature* 465:584–589.
- Cifuentes D, et al. (2010) A novel miRNA processing pathway independent of *Dicer* requires Argonaute2 catalytic activity. *Science* 328:1694–1698.
- Harfe BD, McManus MT, Mansfield JH, Hornstein E, Tabin CJ (2005) The RNaseIII enzyme *Dicer* is required for morphogenesis but not patterning of the vertebrate limb. *Proc Natl Acad Sci USA* 102:10898–10903.
- Muljo SA, et al. (2005) Aberrant T cell differentiation in the absence of *Dicer*. *J Exp Med* 202:261–269.
- Bernstein E, et al. (2003) *Dicer* is essential for mouse development. *Nat Genet* 35:215–217.
- Heo I, et al. (2009) TUT4 in concert with Lin28 suppresses microRNA biogenesis through pre-microRNA uridylation. *Cell* 138:696–708.
- Halic M, Moazed D (2010) *Dicer*-independent primal RNAs trigger RNAi and heterochromatin formation. *Cell* 140:504–516.