

Two Genes on A/J Chromosome 18 Are Associated with Susceptibility to *Staphylococcus aureus* Infection by Combined Microarray and QTL Analyses

Sun-Hee Ahn¹, Hitesh Deshmukh¹, Nicole Johnson², Lindsay G. Cowell², Thomas H. Rude¹, William K. Scott³, Charlotte L. Nelson⁴, Aimee K. Zaas¹, Douglas A. Marchuk⁵, Sehoon Keum⁵, Supaporn Lamlerththorn¹, Batu K. Sharma-Kuinkel¹, Gregory D. Sempowski⁶, Vance G. Fowler Jr.^{1,4*}

1 Department of Medicine, Duke University Medical Center, Durham, North Carolina, United States of America, **2** Department of Biostatistics and Bioinformatics, Duke University Medical Center, Durham, North Carolina, United States of America, **3** Dr. John T. Macdonald Foundation Department of Human Genetics and John P. Hussman Institute for Human Genomics, Miller School of Medicine, University of Miami, Miami, Florida, United States of America, **4** Duke Clinical Research Institute, Durham, North Carolina, United States of America, **5** Department of Molecular Genetics and Microbiology, Duke University Medical Center, Durham, North Carolina, United States of America, **6** Duke Human Vaccine Institute, Durham, North Carolina, United States of America

Abstract

Although it has recently been shown that A/J mice are highly susceptible to *Staphylococcus aureus* sepsis as compared to C57BL/6J, the specific genes responsible for this differential phenotype are unknown. Using chromosome substitution strains (CSS), we found that loci on chromosomes 8, 11, and 18 influence susceptibility to *S. aureus* sepsis in A/J mice. We then used two candidate gene selection strategies to identify genes on these three chromosomes associated with *S. aureus* susceptibility, and targeted genes identified by both gene selection strategies. First, we used whole genome transcription profiling to identify 191 (56 on chr. 8, 100 on chr. 11, and 35 on chr. 18) genes on our three chromosomes of interest that are differentially expressed between *S. aureus*-infected A/J and C57BL/6J. Second, we identified two significant quantitative trait loci (QTL) for survival post-infection on chr. 18 using N₂ backcross mice (F₁ [C18A] × C57BL/6J). Ten genes on chr. 18 (*March3*, *Cep120*, *Chmp1b*, *Dcp2*, *Dtwd2*, *Isoc1*, *Lman1*, *Spire1*, *Tnfaip8*, and *Seh1l*) mapped to the two significant QTL regions and were also identified by the expression array selection strategy. Using real-time PCR, 6 of these 10 genes (*Chmp1b*, *Dtwd2*, *Isoc1*, *Lman1*, *Tnfaip8*, and *Seh1l*) showed significantly different expression levels between *S. aureus*-infected A/J and C57BL/6J. For two (*Tnfaip8* and *Seh1l*) of these 6 genes, siRNA-mediated knockdown of gene expression in *S. aureus*-challenged RAW264.7 macrophages induced significant changes in the cytokine response (IL-1 β and GM-CSF) compared to negative controls. These cytokine response changes were consistent with those seen in *S. aureus*-challenged peritoneal macrophages from CSS 18 mice (which contain A/J chromosome 18 but are otherwise C57BL/6J), but not C57BL/6J mice. These findings suggest that two genes, *Tnfaip8* and *Seh1l*, may contribute to susceptibility to *S. aureus* in A/J mice, and represent promising candidates for human genetic susceptibility studies.

Citation: Ahn S-H, Deshmukh H, Johnson N, Cowell LG, Rude TH, et al. (2010) Two Genes on A/J Chromosome 18 Are Associated with Susceptibility to *Staphylococcus aureus* Infection by Combined Microarray and QTL Analyses. PLoS Pathog 6(9): e1001088. doi:10.1371/journal.ppat.1001088

Editor: Michael R. Wessels, Children's Hospital Boston, United States of America

Received: December 24, 2009; **Accepted:** August 4, 2010; **Published:** September 2, 2010

Copyright: © 2010 Ahn et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: This work was funded by National Institutes of Health research grant R01AI068804 to VGF and R01AI77706 to LGC. Biomarker profiling was performed in the NIAID/NIH-funded Regional Biocontainment Laboratory at Duke University Medical Center (UC6-AI058607). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: Dr. Fowler has served as a consultant for Astellas, Cubist, Inhibitex, Merck, Johnson & Johnson, Leo Pharmaceuticals, NovaDigm, The Medicines Company, Baxter Pharmaceuticals & Biosynexus; has received grant or research support from Astellas, Cubist, Merck, Theravance, Inhibitex, Cerexa, Pfizer, Novartis, Advanced Liquid Logic and National Institute of Health; has received honoraria from Arpida, Astellas, Cubist, Inhibitex, Merck, Pfizer, Targanta, Theravance, Ortho-McNeil, Novartis & Vertex Pharmaceuticals; has served on an advisory committee for Cubist; is employed by Duke University; and has served as a speaker's bureau for Cubist.

* E-mail: fowle003@mc.duke.edu

Introduction

Staphylococcus aureus is an important human pathogen whose clinical spectrum ranges from asymptomatic colonization to endocarditis, shock, and death. Although the importance of genetic factors in determining host susceptibility to *S. aureus* colonization and infection [1–3] is generally accepted, the specific genes responsible for this susceptibility are largely unknown. As with most infectious diseases, the genetics of host susceptibility to *S. aureus* is complex, resulting from variation in multiple genes of small to moderate effect, and from the interaction of these genes with non-genetic factors [4].

Mouse models offer an attractive strategy for investigating complex diseases such as *S. aureus* infections. Abundant breeding, short gestation periods, and the availability of extensive sequence data for inbred strains (<http://mouse.perlegen.com/mouse/index.html> and <http://www.informatics.jax.org/>) facilitate genetic research in mice. In addition, inbred mouse strains demonstrate considerable strain variation in susceptibility to various pathogens [5–11], including *S. aureus* [3]. The determinants of this differential susceptibility to pathogens are probably multigenic, and largely unknown [12].

Among the inbred mouse strains, A/J was recently shown to be highly susceptible to *S. aureus* infection compared with C57BL/6J [3].

Author Summary

Staphylococcus aureus has a wide spectrum of human infection, ranging from asymptomatic nasal carriage to overwhelming sepsis and death. Mouse models offer an attractive strategy for investigating complex diseases such as *S. aureus* infections. A/J mice are highly susceptible to *S. aureus* infection compared with C57BL/6J mice. We showed that genes on chromosomes 8, 11, and 18 in A/J are responsible for susceptibility to *S. aureus* by using chromosome substitution strains (CSS). From the ~4200 genes on these three chromosomes, we identified 191 which were differentially expressed between A/J and C57BL/6J when challenged with *S. aureus*. Next, we identified two significant QTLs on chromosome 18 that are associated with susceptibility to *S. aureus* infection in N₂ backcross mice. Ten genes (*March3*, *Cep120*, *Chmp1b*, *Dcp2*, *Dtwd2*, *Isoc1*, *Lman1*, *Spire1*, *Tnfaip8*, and *Seh1l*) mapped to the two significant QTLs and were differentially expressed between A/J and C57BL/6J. One gene on each QTL, *Tnfaip8* and *Seh1l*, affected expression of cytokines in mouse macrophages exposed to *S. aureus*. These cytokine response patterns were consistent with those seen in *S. aureus*-challenged peritoneal macrophages from CSS 18, but not C57BL/6J. *Tnfaip8* and *Seh1l* are strong candidates for genes influencing susceptibility to *S. aureus* of A/J mice.

By chance, these same genetic backgrounds (A/J and C57BL/6J) were used to construct murine chromosome substitution strains (CSS). CSS were developed to speed the genetic mapping of heritable traits, and were created from a host C57BL/6J strain and a donor A/J inbred strain [13]. Each CSS mouse is homozygous for a specified A/J chromosome but otherwise has a C57BL/6J background. Breeding strategies using these CSS strains greatly enhance the detection of quantitative trait loci (QTLs). Using CSS mice, the impact of a single A/J chromosome can be effectively isolated by eliminating the contribution to variance in phenotype from the other A/J chromosomes. CSS strains have been used to identify QTL for several complex traits, including anxiety [14], diet-induced obesity [15] serum levels of sterols and amino acids [15], testicular cancer [16,17], pubertal timing [18], airway hyperresponsiveness [19], and seizures [20], and provide an ideal means of identifying genetic determinants of murine susceptibility to *S. aureus* sepsis.

In this study, we sought to identify genetic factors associated with susceptibility to *S. aureus* using a multi-step selection process (Figure 1). First, we identified individual chromosomes governing susceptibility to *S. aureus* infection in A/J mice by phenotyping a complete panel of CSS mice. Next, we used peripheral blood mRNA expression to identify genes on these chromosomes that were differentially expressed between susceptible (A/J) and resistant (C57BL/6J) mouse strains in the setting of *S. aureus* infection. Following this, we identified two QTL regions on chromosome 18 that are significantly associated with susceptibility to infection with *S. aureus*. From these two QTL regions, we identified 10 candidate genes that were differentially expressed between A/J and C57BL/6J. We found evidence of biological relevance for two of these genes (*Tnfaip8* and *Seh1l*) using siRNA mediated knockdown and Luminex-based cytokine profiles in *S. aureus*-challenged RAW264.7 mouse macrophages, as well as peritoneal macrophages from CSS18 mice.

Results

Differential susceptibility to *S. aureus* in A/J and C57BL/6J

When mice were injected (intraperitoneally [i.p.]) with the Sanger 476 strain of *S. aureus*, C57BL/6J mice demonstrated a

resistant phenotype (median survival >120 h), whereas A/J mice demonstrated a susceptible phenotype (median survival: 22 h) (Figure 2A). This pattern of differential susceptibility to *S. aureus* between C57BL/6J and A/J mice persisted when the experiments were repeated using both a different *S. aureus* strain (MW2, a methicillin resistant *S. aureus* [MRSA] isolate) (Figure 2B), and when mice were infected by intravenous rather than i.p. route (Figure 2C). A/J mice are known to be deficient in complement factor C5, an important component in neutrophil and macrophage recruitment [21–23]. Thus, the potential impact of C5 deficiency on susceptibility to *S. aureus* infection is important to consider. To ensure that A/J's susceptibility to *S. aureus* is not primarily due to C5 deficiency, we challenged three additional C5 deficient mouse strains (B10.D2/oSnJ, B10.D2-Hc0 H2d H2-T18c/o2SnJ, and NOD/LtJ) with an intraperitoneal injection of *S. aureus* (10⁷ CFU/g). All three additional C5-deficient mice were resistant to *S. aureus* infection (median survival >120 h, Figure 2D), suggesting that factors other than C5 deficiency were responsible for the susceptibility of A/J mice to *S. aureus*.

Genetic susceptibility to *S. aureus* in mice localizes to A/J chromosomes 8, 11, and 18

Since we identified C57BL/6J mice as resistant and A/J mice as susceptible to *S. aureus* infection, we used CSS from these backgrounds to further localize the chromosomal regions responsible for these divergent phenotypes. CSS mice with chromosome 8, 11 or 18 from the susceptible background (A/J) transferred individually into the otherwise resistant background (C57BL/6J) were significantly more susceptible to *S. aureus* infection compared to C57BL/6J mice ($p < 0.005$, log rank test) (Table 1) or other chromosome substituted strains. This suggests that genetic loci mediating susceptibility to *S. aureus* are located on each of these 3 chromosomes.

Rapid mortality in CSS mice with A/J “susceptibility” chromosomes 8, 11, and 18 is associated with increased bacterial load in the peritoneal fluid and kidneys

To further understand the basis of the enhanced susceptibility to *S. aureus* infection, we measured the tissue burden of *S. aureus* in A/J and C57BL/6J mice and the CSS mice 24 hours after infection with *S. aureus*. The animals were euthanized and bacterial load in the kidney and peritoneal fluid was determined. The bacterial load in the peritoneal fluid of susceptible A/J mice was significantly higher than in the resistant C57BL/6J mice ($178.9 \pm 16.3 \times 10^6$ CFU/ml vs. $3.40 \pm 1.55 \times 10^6$ CFU/ml; $P = 0.0001$, F-test; Figure 3A). The bacterial load of the peritoneal fluid was also significantly higher in the CSS mice with A/J chromosome 8 ($119.3 \pm 47.8 \times 10^6$ CFU/ml, $P = 0.0017$, F-test), chromosome 11 ($220 \pm 74.7 \times 10^6$ CFU/ml, $P = 0.016$, F-test) or chromosome 18 ($132.7 \pm 25.7 \times 10^6$ CFU/ml, $P = 0.019$, F-test) compared to C57BL/6J mice and to the rest of the chromosome substituted strains ($1.531 \pm 5.5 \times 10^6$). Similarly, the bacterial load in the kidney of the susceptible A/J mice was significantly higher than in C57BL/6J mice ($111.9 \pm 99.8 \times 10^6$ CFU/g vs. 33.1 ± 18.7 CFU/g; $P = 0.003$, F-test; Figure 3B). Finally, the bacterial count in the kidney of CSS mice with chromosome 8 ($44.8 \pm 27.8 \times 10^6$ CFU/g, $P < 0.001$, F-test), chromosome 11 ($3.5693 \pm 0.1940 \times 10^6$ CFU/g, $P < 0.001$, F-test) and chromosome 18 ($1.3451 \pm 0.2491 \times 10^6$ CFU/g, $P = 0.0013$, F-test) was significantly higher than that in C57BL/6J mice and the remainder of the chromosome substituted strains ($4.75 \pm 86.39 \times 10^3$ CFU/g, $P = 0.016$, F-test). Thus, increased mortality due to *S. aureus* sepsis is associated with a higher bacterial load in the kidneys and peritoneal fluid of susceptible A/




Figure 1. Overall strategy for identifying genes associated with susceptibility to *S. aureus* in A/J mice.
doi:10.1371/journal.ppat.1001088.g001

J, CSS8, CSS11 and CSS18 mice as compared to the resistant C57BL/6J mice and the remainder of the consomic mouse strains (CSS1–CSS7, CSS9–10, CSS12–CSS17, CSS19, CSSX and CSSY). This correlation of susceptibility with bacterial burden provides important insight into the pathogenesis of early death in murine staphylococcal sepsis, as the genes involved in susceptibility of A/J to *S. aureus* sepsis appear to be associated with a relative inability of the host to control infection.

Genetic susceptibility to *S. aureus* in mice is not sex linked and has variable forms of inheritance

We next generated F₁ progeny by crossing the resistant C57BL/6J and susceptible CSS 8, 11, and 18 mice. These crosses result in offspring that, for the relevant chromosome pair (8, 11, or 18), have one C57BL/6J chromosome and one A/J chromosome. Using the same model of infection as described above, we observed no significant

difference in the median survival of each F₁ cohort when the parents were of either sex, indicating that the alleles of the loci mediating susceptibility to *S. aureus* are not imprinted (**Table 2**). F₁ mice from the C57BL/6J×CSS8 (C8A) and C57BL/6J×CSS18 (C18A) crosses retained susceptibility to *S. aureus* infection (median survival <48 h), whereas F₁ mice from C57BL/6J×CSS11 cross (C11A) were resistant (median survival >120 h) to *S. aureus* infection. These results indicate that the individual loci mediating susceptibility to *S. aureus* in A/J mice have different modes of inheritance (i.e., autosomal dominant for loci on chromosomes 8 and 18 and autosomal recessive for loci on chromosome 11). Similar to the homozygous CSS mice, the bacterial load in the kidneys of the susceptible F₁ mice (C8A and C18A) was significantly higher than in the resistant C57BL/6J or C11A mice (data not shown). Taken together, these results provide further evidence that genes governing murine susceptibility to *S. aureus* infection reside on chromosomes 8, 11 and 18.




Figure 2. A/J and C57BL/6J mice exhibit different susceptibility to *S. aureus* that is not primarily due to deficiency in Complement C5. C57BL/6J and A/J were injected via intraperitoneal (i.p.) route (10^7 CFU/g) with *S. aureus* strains Sanger 476 (Figure 2A) and MW2 (Figure 2B) ($n=15$ mice for each strain). (Figure 2C) C57BL/6J and A/J mice were injected via tail vein route (10^7 CFU/g) with Sanger 476 ($n=10$ mice for each strain). Comparison of survival curves was performed by use of log rank test. Values of $P<0.05$ were considered significant. (Figure 2D) Deficiency of Complement C5 is not the primary determinant of susceptibility to *S. aureus* in A/J mice. Three C5-deficient mouse strains (C57BL/10SnJ, B10.D2/oSnJ, B10.D2-Hc0 H2d H2-T18c/o2SnJ, and NOD/LtJ) were injected i.p. (10^7 CFU/g) with Sanger 476 ($n=10$ mice for each strain). doi:10.1371/journal.ppat.1001088.g002

Genes on chromosomes 8, 11, and 18 are differentially expressed between A/J and C57BL/6J mice

Next, to identify genes on A/J chromosomes 8, 11, and 18 that contribute to susceptibility, we compared whole blood genome transcription profiles of A/J and C57BL/6J mice in both an uninfected state and following *S. aureus* infection. A total of 675 genes on chromosomes 8, 11, and 18 were differentially expressed between uninfected A/J and C57BL/6J mice, and 751, 683, 1223 and 639 were differentially expressed at 2, 4, 6, and 12 hours after infection, respectively. However, only 191 of these genes were differentially expressed at all four post-infection time points (Table 3 and Table S1). Of the 191 genes, 37 genes were similarly expressed in uninfected A/J and C57BL/6J mice. The remaining 155 genes were differentially expressed between C57BL/6J and A/J mice in both uninfected and infected states.

QTL for susceptibility to *S. aureus* infection on chromosome 18

To further fine-map the locus (or loci) on chromosome 18 involved in determining susceptibility to *S. aureus* infection, we performed a QTL analysis, generating N_2 backcross mice by mating F_1 mice (C18A) to C57BL/6J mice (Figure S1). A total of 144 N_2 backcross mice were generated and infected with *S. aureus*, and their survival times were measured. No sex-linked suscepti-

bility in N_2 backcross mice was identified (Figure S2). Using J/qtl software, two significant QTLs were identified as linked to survival time after *S. aureus* infection (Figure 4A), one located at 41 to 54 Mb, and one located at 57 to 67Mb. Approximately 317 genes are contained within these two significant intervals.

Next, we identified genes implicated by both our expression array-based and QTL-based selection strategies. A total of 10 genes (*March3*, *Cep120*, *Chmp1b*, *Dcp2*, *Dtwd2*, *Isoc1*, *Lman1*, *Spire1*, *Tnfrsf8*, and *Seh1*) mapping within our two significant QTLs were differentially expressed between *S. aureus* infected A/J and C57BL/6J mice. Four genes (*Dcp2*, *Dtwd2*, *Tnfrsf8*, and *Cep120*) were located within the first significant QTL (41 to 54 Mb) and one gene (*March3*) was located in the suggestive interval for this QTL (56 Mb). The remaining 5 genes (*Isoc1*, *Lman1*, *Chmp1b*, *Spire1*, and *Seh1*) fell within the second significant QTL (57 to 67 Mb). For 6 of these genes (*Dtwd2*, *Tnfrsf8*, *Isoc1*, *Lman1*, *Chmp1b*, and *Seh1*), real-time PCR confirmed significant differences in expression between *S. aureus*-infected A/J and C57BL/6J mice ($P<0.05$, paired two-tailed t-test) (Figure 4B).

Assessing biological relevance: Neutrophil and macrophage function in A/J and C57BL/6J

Because of the central role of neutrophils in the host response to *S. aureus*, we considered whether neutrophil dysfunction might be a

Table 1. Median survival of C57BL/6J, A/J, and Chromosome substituted strain (CSS) mice after intraperitoneal injection with *S. aureus* (10^7 CFU/g).

Strain	Median Survival (d)	Number of Mice Tested (n)	P-Value*
A/J	1	10	0.0005
C57BL/6J	>5	10	1
CSS 1	>5	10	1
CSS 2	>5	10	1
CSS 3	>5	10	1
CSS 4	>5	10	1
CSS 5	>5	10	1
CSS 6	>5	10	1
CSS 7	2.5	10	0.0771
CSS 8	1.5	10	0.0027
CSS 9	>5	10	1
CSS 10	>5	10	1
CSS 11	1	10	0.0023
CSS 12	>5	10	1
CSS 13	>5	10	1
CSS 14	>5	10	1
CSS 15	>5	10	1
CSS 16	>5	10	1
CSS 17	>5	10	1
CSS 18	1	10	0.0023
CSS 19	>5	10	1
CSS X	4	5	0.0805
CSS Y	>5	5	1

CSS mice contain a single A/J chromosome inserted in C57BL/6J background. Thus, CSS 1 contains A/J chromosome 1 inserted in C57BL/6J background, etc. CSS strains 8, 11 and 18 were significantly more susceptible to *S. aureus* infection than C57BL/6J ($P = 0.0027$, 0.0023 , and 0.0023 , respectively).

*Compared to survival for C57BL/6J. Comparison of survival curves was performed by use of log rank test. Values of $P < 0.05$ were considered significant. doi:10.1371/journal.ppat.1001088.t001

primary cause of A/J susceptibility to *S. aureus*. To do this, we assessed ex vivo the bactericidal activity of neutrophils from A/J, C57BL/6J, or CSS18 mice by incubating them with *S. aureus*, and then comparing the number of viable bacteria in the extracellular (unphagocytosed) and intracellular (phagocytosed) fractions of neutrophils between mouse strains. As shown in Figure S3A, the percentage of surviving *S. aureus* was similar in both fractions across all three mouse strains. Thus, no significant strain-specific differences in the function or capacity of neutrophils to kill *S. aureus* were noted. We also found no significant strain-specific differences in bactericidal capacity when we repeated this experiment using peritoneal macrophages from A/J and C57BL/6J mice (Figure S3B). Based on these findings, we considered whether the observed susceptibility of A/J mice to *S. aureus* might be due in part to perturbations in the initial host inflammatory response.

Assessing biological relevance: Cytokine response in A/J and C57BL/6J

To further evaluate the potential importance of our 10 candidate genes in host immune responses to *S. aureus*, we treated mouse macrophage RAW264.7 cells with short interfering RNA (siRNA) oligonucleotides. Knockdown expression of each of the 10 target genes in cells was confirmed by real-time-PCR (Figure S4). The cells were then stimulated with *S. aureus*, and a Luminex-based multiplex cytokine assay was used to evaluate the impact of each genes' individual knockdown expression on the host inflammatory response to *S. aureus*. Cell-culture supernatants from pre- and post-*S. aureus* stimulation conditions were compared (Figure S5). Only TNFAIP8 and SEH1L knockdown cells significantly influenced cytokine levels in *S. aureus*-challenged siRNA-transfected RAW264.7 cells (Figure 5A). Compared to *S. aureus*-challenged negative controls, *S. aureus*-challenged cells with knockdown expression of *Tnfaip8* produced a significantly lower level of IL-1 β , while those with knockdown expression of *Seh1l* produced a significantly higher level of GM-CSF.

To evaluate whether our cytokine results from the RAW264.7 knockdown experiments mimicked what was encountered *in vivo*, we first demonstrated by real time PCR that expression levels of *Tnfaip8* and *Seh1l* are significantly lower in whole blood from A/J mice than from C57BL/6J mice (Figure 4B). Next, we confirmed that the same reduced expression pattern was also encountered in




Figure 3. Bacterial load in peritoneal fluid and kidneys is higher in chromosome substituted strain (CSS) 8, CSS11 and CSS18 mice after *S. aureus* injection. (A) Bacterial load in peritoneal fluid. C57BL/6J, A/J, and CSS mice were injected (i.p.) with Sanger 476 (10^7 CFU/g) and euthanized 24 h later (n = 5 mice for each murine strain). (B) Bacterial load in kidneys. C57BL/6J, A/J, and CSS mice were injected (i.p.) with Sanger 476 (10^7 CFU/g) and euthanized 24 h later (n = 5 mice for each murine strain). P-values were calculated from the F-test. Values of $P < 0.05$ were considered significant. doi:10.1371/journal.ppat.1001088.g003

Table 2. Median survivals of F₁ progeny of Chromosomal Substitution Strains 8, 11, and 18.

F ₁ progeny	Median Survival (d)	Number of Mice Tested (n)
C57BL/6J (M) X A/J (F)*	1	10
A/J (M) X C57BL/6J (F)	1	10
C57BL/6J (M) X CSS8 (F)	2	10
CSS8 (M) X C57BL/6J (F)	2	10
C57BL/6J (M) X CSS18 (F)	2	10
CSS18 (M) X C57BL/6J (F)	2	10
C57BL/6J (M) X CSS11 (F)	5	10
CSS11 (M) X C57BL/6J (F)	5	10

Genetic determinants of susceptibility to *S. aureus* are acquired in a dominant inheritance pattern on chromosomes 8 and 18 and a recessive inheritance pattern on chromosome 11. These data also indicate that the alleles conferring susceptibility to *S. aureus* in A/J are not imprinted.

*M = Male; F = Female.

doi:10.1371/journal.ppat.1001088.t002

peritoneal macrophages (PMΦ) from A/J mice ($54 \pm 13\%$ [*Tnfaip8*] and $70 \pm 10\%$ [*Seh1l*] of C57BL/6J PMΦ mRNA expression, respectively) and CSS18 mice ($42 \pm 3\%$ [*Tnfaip8*] and $78 \pm 5\%$ [*Seh1l*] of C57BL/6J PMΦ mRNA expression, respectively). We then compared cytokine production of PMΦ from C57BL/6J and CSS18 after stimulation with *S. aureus* (Figure S6). We used PMΦ from CSS18 (which contain Chr. 18 from A/J but are otherwise C57BL/6J) in order to eliminate the potential confounding effect on the results of additional unidentified susceptibility genes on Chr. 8 and 11 of A/J mice. As shown in Figure 5B, we encountered the same pattern of IL-1β and GM-CSF in *S. aureus*-stimulated PMΦs from CSS18 mice as we did in the TNFAIP8 and SEH1L knockdown in RAW264.7 cells. Specifically, IL-1β was significantly decreased and GM-CSF was significantly increased in PMΦ of CSS18 compared to that of C57BL/6J ($P = 0.009$ for IL-1β, $P = 0.04$ for GM-CSF; unpaired two tailed t-test).

Presence of SNPs in the QTL regions

All but one (*Dtwd2*) of the 10 differentially expressed genes in the two QTLs on chr. 18 contained SNPs in the region flanking the first and last exon. Such SNPs could potentially lie within transcriptional regulatory sites. A total of 22 genes within the two QTLs on chromosome 18 contained non-synonymous coding SNPs. These genes included *Dcp2*, *Gm5505*, *Ccdc112*, *Pggt1b*, *Comm10*, *Gm5095*, *Gm3790*, *Ftmt*, *Zfp474*, *Fbn2*, *Slc27a6*, *Gm3957*, *Sympo*, *Tcof1*, *Csflr*, *Ppargc1b*, *Il17b*, *Afap11l*, *Sh3tc2*, *Wdr7*, *Alpk2*, *Malt1*, and *5330437102Rik*.

Discussion

Host genetic determinants of susceptibility to *S. aureus* are poorly understood. In the current study, we localized the genetic determinants of *S. aureus* susceptibility in A/J inbred mice to chromosomes 8, 11, and 18. From the ~4200 genes on these three chromosomes, we identified 191 which are differentially expressed between the susceptible A/J and the resistant C57BL/6J inbred mouse strains. We also pursued QTL analysis for chr. 18 to corroborate the findings of our array-based approach with a second strategy. We found two QTLs on chr. 18 over the significance threshold. This suggests that there are at least two genes, one (or more) of which are associated with each peak and which are involved in susceptibility to *S. aureus* infection in A/J mice. In the current report, we have identified two genes, one on each of the two QTL peaks, which are strongly associated with susceptibility to *S. aureus* infection.

We hypothesized that our candidate genes influenced susceptibility to *S. aureus* in A/J mice by regulating macrophage cytokine production. There are several lines of evidence in support of this hypothesis. Macrophages play an important role in initiating host defense to *S. aureus* by secreting cytokines and chemokines [24–26], and impairment of this cytokine production by macrophages results in impaired host inflammatory responses [27,28]. In support of our hypothesis, we found that macrophages from the two mouse lineages did differ significantly in their cytokine response to *S. aureus*. In addition, we found no difference in the bactericidal capacity of either macrophages or neutrophils from A/J and C57BL/6J mice, suggesting that functional deficits within

Table 3. Number of genes on murine chromosomes 8, 11, and 18 that are differentially expressed between A/J and C57BL/6J at all post-infection timepoints following intraperitoneal infection with *S. aureus*.

Chromosome	Genes on Chromosome	Differentially Expressed Genes at All Post-Infection Time Points*
8	1328	56
11	2200	100
18	683	35
Total	4211	191 [†]

*Includes 2 hour, 4 hour, 6 hour, and 12 hour post-infection timepoints.

[†]Includes 37 genes (12 on chromosome 8, 20 on chromosome 11, and 5 on chromosome 18) that are differentially expressed between A/J and C57BL/6J only following infection and 154 genes (44 on chromosome 8, 80 on chromosome 11, and 30 on chromosome 18) that are differentially expressed between A/J and C57BL/6J in both uninfected and post-infection states. See Table S1 for complete list of all 191 genes.

doi:10.1371/journal.ppat.1001088.t003





Figure 4. Ten genes on chromosome 18 were identified by both QTL mapping and expression array for susceptibility to *S. aureus*. (A) Chromosome 18 LOD score plot for susceptibility to *S. aureus* in N_2 backcross mice. Six to eight-week-old backcross mice were injected i.p. with 1×10^7 CFU/g *S. aureus* and observed every 8 h for 5 days. The horizontal dashed lines indicate the thresholds for significant ($P=0.05$) and suggestive ($P=0.63$) LOD score determined by the J/qtl permutation test using 1,000 permuted data sets. The microsatellite markers used for determining genotypes of N_2 backcross mice are presented along the X-axis. The differentially expressed genes are indicated on chromosome 18. Genes identified within significant or suggestive QTL were indicated with * or †, respectively. (B) Quantitative real-time PCR validation of 10 genes (*March3*, *Cep120*, *Chmp1b*, *Dcp2*, *Dtdw2*, *Isoc1*, *Lman1*, *Spire1*, *Tnfaip8*, and *Seh11*) mapping within the two identified QTLs on chr.18 that were also identified by the expression-array selection strategy. Expression values of genes were normalized to 18S rRNA. Six genes (*Dtdw2*, *Tnfaip8*, *Isoc1*, *Lman1*, *Chmp1b*, and *Seh11*) were significantly differentially expressed between A/J and C57BL/6J infected with *S. aureus* in the significant QTLs on chromosome 18. Values of $P<0.05$ were considered significant ($n=5$, paired two-tailed t-test). doi:10.1371/journal.ppat.1001088.g004

A



B




Figure 5. RAW264.7 cells transfected with *Tnfaip8* and *Seh1* siRNA and peritoneal macrophages from C57BL/6J and CSS18 exhibited significant and consistent patterns of change in the cytokine production (IL-1 β and GM-CSF). (A) RAW264.7 cells were treated with negative control, *Tnfaip8* or *Seh1* siRNA and incubated with *S. aureus* for 5 h. (B) Peritoneal macrophages from CSS18 and C57BL/6J were incubated with *S. aureus* for 48 h. Macrophage culture supernatants were collected at two time points (pre- and post- *S. aureus* stimulation) for cytokine/chemokine analysis. IL-1 β and GM-CSF were determined by Luminex-based multiplex cytokine assay. Means and standard deviations of three independent experiments are shown. Values of $P < 0.05$ were considered significant (unpaired two-tailed t-test). doi:10.1371/journal.ppat.1001088.g005

these cells are not the primary cause of the observed differences in susceptibility between the two murine strains. For these reasons, we pursued siRNA-mediated knockdown of our candidate genes in mouse macrophage cells and analyzed the cytokine profiles of knockdown cells exposed to *S. aureus*. As evidenced by real-time PCR, six genes were significantly differentially expressed. More notably, knockdown expression of two of the six genes, *Tnfaip8* and *Seh1*, significantly altered the specific patterns of cytokine production in *S. aureus*-stimulated RAW cells in a manner that was also observed in PM Φ of CSS18, but not C57BL/6J mice. These two genes are prime candidates for influencing susceptibility to *S. aureus*.

Of the 191 genes on chromosomes 8, 11, or 18 that are also differentially expressed between A/J and C57BL/6J mice, 28 have previously been shown to play key roles in the response to various infectious diseases. For example, *Cd209d* has been known to be able to capture Gram-negative bacteria such as *Escherichia coli* and *Salmonella typhimurium* as well as Gram-positive bacteria such as *Streptococcus pneumoniae* [29,30]. *Tbkbp1* may play a role in antiviral innate immunity as part of the TNF/NF κ B interaction pathway [31]. *Map2k3* is induced by various stimuli, including cellular stress, inflammatory cytokines and cell surface receptors, and mediates the activation of p38 mitogen-activated protein kinase

(p38MAPK) [32]. *Grb2* [33], *Prdx2* [34], and *Ppp2cb* [35] are also involved in the MAPKKK pathway. Activation of the MAPKKK cascade confers resistance to both Gram-negative and Gram-positive pathogens, indicating that signaling pathways initiated by various pathogens converge into a MAPKKK cascade [36,37]. Interestingly, almost one-third of these known 28 genes were associated with apoptosis (*Birc5*, *Cln8*, *Cyfp2*, *Psmc3*, *Rffl*, *Rtn4*, *Trp53*, *Tnfaip8*, and *Xaf1*). Although the precise reasons for the high frequency of apoptosis-related genes are not known, it suggests that this pathway may represent a particularly important area for further study. Finally, our strategy also identified a number of novel genes about which little is known. For example, 20 of the 191 differentially expressed genes on chromosomes 8, 11, and 18 were identified entirely by RIKEN cDNA information, and had no known biological function (Table S1).

The TNFAIP8 family of proteins consists of TNFAIP8, TIPE1 (TNFAIP8Like1), TIPE2 and TIPE3. TIPE2 is known as a negative regulator of innate and adaptive immune processes [38]. Woodward et al. showed that TNFAIP8 regulates glucocorticoid-mediated apoptosis of thymocytes [39]. Interestingly, RAW264.7 cells in which TNFAIP8 had been knocked down produced significantly more GM-CSF and significantly less IL-1 β compared to control cells. GM-CSF is a critical Th1 cell-derived cytokine

that mediates pulmonary inflammation *in vivo*, and controls neutrophil and macrophage numbers [40,41]. Interestingly, overexpression of GM-CSF leads to severe inflammation [41]. IL-1 β is a key inflammatory cytokine in orchestrating host defense against *S. aureus*, with mice deficient in this cytokine exhibiting markedly increased susceptibility to *S. aureus* [24,42]. Taken together, *Tnfaip8* is a strong candidate gene governing susceptibility to *S. aureus* through IL-1 β and GM-CSF regulation in mice. *Seh1l*, the other candidate gene, is a part of nuclear pore complex Nup107–160 that was recently found to play a role in embryonic development. Although the infectious diseases-related function of this gene is unknown, its knockdown was also shown to increase GM-CSF production in *S. aureus*-stimulated macrophages. Thus, it may be that SEH1L contributes to susceptibility of A/J to *S. aureus* by increasing, along with TNFAIP8, macrophage production of GM-CSF.

The biological relevance of *Tnfaip8* and *Seh1l* is strengthened by several key observations. First, *Tnfaip8* and *Seh1l* exhibited decreased mRNA expression in both peripheral blood and PM Φ of A/J and CSS18 mice as compared with C57BL/6J mice. Second, of the 10 differentially expressed genes on the significant QTL regions of chromosome 18, only *Tnfaip8* and *Seh1l* induced significant changes in cytokine production from siRNA-inhibited, *S. aureus*-challenged macrophages. Third, the cytokines altered by this siRNA-mediated knockdown of TNFAIP8 and SEH1L are critically important components of host response to *S. aureus* [24,42]. Finally, the cytokine patterns elicited by TNFAIP8 and SEH1L knockdowns were also seen in the peritoneal macrophages of CSS18 mice, but not C57BL/6J mice. Taken together, these data support the importance of our two candidate genes, *Tnfaip8* and *Seh1l*, to the susceptibility to *S. aureus*, both *in vivo* and *in vitro*.

Although regulatory variation can play an important role in various complex traits [43–46], its role in the current study is unresolved. Nine of the 10 differentially expressed candidate genes on our two QTLs contained SNPs in the region flanking the first and last exon. In addition, 22 other genes on our two QTLs contained non-synonymous coding SNPs. Such SNPs could also potentially contribute to strain-specific phenotypes via modification of protein function. Only 10 of these genes (*Fmt*, *Fbn2*, *Synpo*, *Tcof1*, *Csf1r*, *Ppargc1b*, *Il17b*, *Sh3tc2*, *Wdr7*, and *Malt1*) were identified in the map of disease genes described in Online Mendelian Inheritance in Man (OMIM). Of these, three (*Csf1r*, *Il17b*, and *Malt1*) are known to be involved in immune response. For example, *Csf1r* encodes a tyrosine kinase growth factor receptor for colony-stimulating factor-1, a cytokine which controls the production, differentiation, and function of macrophages [47]. IL-17b plays an important role in the pathogenesis of inflammatory arthritis [48] in the animal model. MALT1 is essential for NF-kappa-B activation, cytokine production, and proliferation of primary T and B lymphocytes [49]. Future studies will investigate the role of these genes, if any, on the susceptibility of A/J mice to *S. aureus* infection.

The different susceptibilities of A/J and C57BL/6J mice to *S. aureus* infection were recently confirmed by other investigators [3]. von Kockritz-Blickwede et al. suggested that resistance to *S. aureus* in C57BL/6 mice is critically dependent on an effective and fast recruitment of neutrophils to the site of infection due to different kinetic profiles in expression of KC (*Cxcl1*) and MIP-2 (*Cxcl2*). These two chemokines play a central role in mediating neutrophil recruitment to sites of infection by binding to receptors on the surface of these cells. In support of this hypothesis, we found that *Cxcl1* was significantly differentially expressed between A/J and C57BL/6J mice at 2, 6, and 12 hours after infection with *S. aureus*. Although the gene coding for *Cxcl1* is located on mouse

chromosome 5, CSS5 mice infected with *S. aureus* in the current study retained the resistant phenotype. This finding suggests that genes on mouse chromosome 5 are not directly responsible for A/J's susceptibility, but that the reduced chemokine production observed in *S. aureus*-infected A/J mice [3] may be regulated by one or more genes on chromosomes 8, 11, or 18.

Although C5 deficiency has been shown to contribute to susceptibility to a variety of pathogens, including *Pseudomonas aeruginosa*, *Mycobacterium tuberculosis*, *Listeria monocytogenes*, and *Candida albicans* [22,50–53], its contribution to susceptibility to *S. aureus* is unresolved. For example, Crequetti et al. (1983) showed that C5-deficient mice had impaired lung clearance of *S. aureus* [53], while Toews and Pierce (1984) demonstrated similar lung clearance of *S. aureus* and PMN recruitment in both C5-deficient and sufficient mice [54]. In addition, Easmon and Glynn reported decreased survival of C5-deficient mice compared with C5-sufficient controls after intraperitoneal injection with mucoid *S. aureus* strains [55], while Cunnion et al. observed similar survival in both C5-deficient and sufficient mice challenged via intravenous injection with *S. aureus* [56]. Finally, von Kockritz-Blickwede and colleagues recently demonstrated a protective role of complement C5a in an experimental model of *S. aureus* bacteremia [23]. In the present study, all three of the additional C5-deficient mouse strains we tested were resistant to *S. aureus*. Our consomic experiments also suggest that C5-deficiency is not the primary determinant of susceptibility to *S. aureus* in A/J mice. Although the gene for C5 (Hc) resides on chromosome 2, CSS2 mice (which contain A/J chromosome 2 and thus would be C5 deficient) remained resistant to *S. aureus* infection. Based on this body of evidence, we conclude that factors other than C5-deficiency are likely to be primarily responsible for the susceptibility to *S. aureus* seen in A/J mice.

The current study has limitations. First, linkage scores of the two QTLs on chr. 18 imply that there may be more than one gene on the larger significant peak that plays a role in susceptibility to *S. aureus*. One possible additional susceptibility gene is *Lman1*. In the Figure S5, knockdown of LMAN1 showed dramatically reduced IL-1 α compared to control when stimulated with *S. aureus*. Although it was not statistically significant ($P=0.08$), the quantity of IL-1 α in LMAN1 knockdown cells was below the detection value of standard minimum. Thus, it is possible that testing of a much larger number of samples would reveal a statistical significance of additional candidate genes such as *Lman1* in the larger peak. Second, there may be genes on other chromosomes that contribute to a lesser extent, or that have a strong effect only when acting in conjunction with genes on another chromosome. Third, our strategy will miss genes on chromosomes 8, 11, or 18 that contribute to phenotype at the protein level (e.g., coding SNPs) or by modulating the gene expression patterns of genes on other chromosomes. Fourth, our strategy would not allow detection of polymorphic regulatory sites more than 2kb away from genes, such as those within gene deserts. Finally, we have only completed QTL linkage analysis for chr. 18. Thus, additional work is required, including QTL mapping analysis for A/J chromosomes 8 and 11, defining the pathogenesis of *Tnfaip8* and *Seh1l* using knockout mice, and evaluating the impact of *Tnfaip8* and *Seh1l* on host susceptibility to different pathogen classes. Ultimately, the importance of the identified candidate genes, or their human homologs, will need to be evaluated in patients with *S. aureus* infections.

Despite these limitations, this study makes several key observations. First, using expression array-based strategies, we have identified 191 genes that are differentially expressed between susceptible and resistant mice, and are found on three different chromosomes where loci conferring susceptibility to *S. aureus*

infection have been mapped. Second, we have identified two QTLs on chr. 18 that are linked to survival time after infection with *S. aureus*. Ten differentially expressed genes mapped to the significant- or suggestive threshold of these two QTLs. Of these 10 genes, two (*Tnfaip8* and *Seh1l*) significantly affect the production of IL-1 β and GM-CSF in *S. aureus*-stimulated mouse macrophages in a pattern also seen in peritoneal macrophages from mice containing A/J chromosome 18, thereby supporting a potential role for these two genes in host response to *S. aureus*. These genes in general, and *Tnfaip8* and *Seh1l* in particular, represent promising candidates for the genetic basis of host susceptibility to this serious, common pathogen.

Materials and Methods

Ethics statement

All animal experiments were performed in accordance to NIH guidelines, the Animal Welfare Act, and US federal law. Such experiments were carried out following approval by the Duke University's Institutional Animal Care & Use Committee (IACUC) which has been accredited by the Association for Assessment and Accreditation of Laboratory Animal Care (AAALAC) International. All animals were housed in a centralized and AAALAC-accredited research animal facility that is fully staffed with trained husbandry, technical, and veterinary personnel.

Mouse strains

C57BL/6J and A/J mice were purchased from The Jackson Laboratory (Bar Harbor, ME). Chromosome substitution strains (CSS) were generated by transferring a single, full-length chromosome from the donor strain (A/J) onto the genetic background of the host strain (C57BL/6J) by repeated backcrossing [15]. Male CSS mice (CSS1–19, CSSX and CSSY) were also obtained from the Jackson Laboratory (Bar Harbor, ME).

Preparation of bacterial cells

S. aureus clinical strains, Sanger 476 or MW2 were used in the mortality studies. For preparation of *S. aureus* for injection, an overnight bacterial culture of *S. aureus* was diluted with fresh tryptic soy broth (TSB) and incubated (37°C) with aeration to log-phase growth (optical density at wavelength 600 nm (O.D.600) of ~1.0) [57]. *S. aureus* was harvested by centrifugation, rinsed, and resuspended in saline. To mimic the natural course of *S. aureus* infection in humans, which typically arises from a primary focus of infection and disseminates to other sites, we employed an intraperitoneal (i.p.) route of infection in our animal model [58].

Murine phenotype definitions, injection (*S. aureus*), and specimen procurement

Mouse phenotypes were defined as either **resistant** (survival for >120 hours following *S. aureus* infection) or **susceptible** (survival for <48 hours following *S. aureus* infection). Mice were injected via an intraperitoneal (i.p.) route with 1×10^7 CFU/g *S. aureus* depending on the experiment and observed every 8 h for morbidity. Mice were sacrificed using CO₂ asphyxiation if they appeared moribund. Pain/distress was assessed using a numerical scale for the following characteristics: Appearance (0 = normal; 1 = lack of grooming; 2 = rough hair coat; 3 = abnormal posture); Behavior (0 = normal; 1 = moving slowly; 2 = moving slowly, irregular ambulation; 3 = immobile). A score of 3 indicated significant pain and distress and culminated in the early euthanasia of the animal. Peritoneal lavage was performed as described previously [2]. Kidneys were harvested and either frozen (–80°C) or fixed in 10% buffered formalin, as appropriate, for subsequent

analysis. For experiments involving RNA analysis, blood was collected by intracardiac puncture and stored in RNAlater at –20°C.

Quantifying bacterial load in murine tissues and blood

Kidneys collected from euthanized animals were homogenized in phosphate buffered saline (PBS) and diluted 10 fold serially. Peritoneal lavage (50 μ l) was serially diluted in PBS. The serial dilutions (50 μ l) were plated in Tryptic Soy Agar (TSA) plates and incubated (37°C, overnight) to count the number of colony forming units (CFU) of *S. aureus*.

Generation of F₁ and N₂ progeny

For the current study, susceptible CSS mice with chromosome 8, 11 and 18 from A/J mice on a C57BL/6J background (CSS8, CSS11 and CSS18) were mated with resistant C57BL/6J strain in reciprocal crosses [C57BL/6J male \times CSS8, 11 or 18 female] and (CSS8, 11 or 18 male \times C57BL/6J female) to generate an F₁ population that was heterozygous for the chromosome of interest on an otherwise uniform C57BL/6J genomic background. To generate N₂ backcross mice for QTL linkage analysis on chromosome 18, we bred F₁ mice (C18A) to C57BL/6J (Figure S1).

Microarray

Total RNA was prepared from mouse blood using the Mouse RiboPure Blood RNA isolation kit (Ambion) following the manufacturer's instruction. Globin mRNA was removed from whole blood RNA samples using the GlobinClear kit (Ambion). A total of 30 samples passed the quality criteria of the Agilent Bioanalyzer and were used for microarray analysis. Since the total RNA yield of many samples was low, 1 round of linear amplification was performed for all samples (Ambion MessageAmp Premier). Mouse Genome 430 2.0 Array-Chips were used (Affymetrix). The biotin-labeled cDNA was hybridized to the arrays for 16 hours at 45°C following the manufacturer's instruction. The arrays were then washed and labeled with streptavidinphycoerythrin (strep-PE), and the signal was amplified using biotinylated antistreptavidin followed by another round of staining with strep-PE. Washing and staining were performed on the Affymetrix fluidics station according to the recommended fluidics protocol. Labeled gene chips were scanned using an Affymetrix Genechip Scanner 7G (Affymetrix). This array contains 45,101 probe sets to analyze the expression level of over 39,000 transcripts and variants from over 34,000 well characterized mouse genes. The microarray data have been deposited in the NCBI GEO and are accessible through GEO series accession no. GSE19668.

Analysis of gene expression microarray data

To identify genes for which differential expression between A/J and C57BL/6J mice could contribute to host susceptibility to *S. aureus* infection, we compared the gene expression profiles between uninfected A/J and C57BL/6J mice and between infected A/J and C57BL/6J mice at 2, 4, 6, and 12 hours after infection. We used a multi-step approach that included i) preprocessing to determine probeset expression values, ii) filtering based on probeset annotations, iii) identification of genes with statistically significant differential expression between A/J and C57BL/6J mice after infection, and iv) ranking based on fold change (FC) in gene expression and Gene Ontology (GO) (<http://www.geneontology.org>) annotation.

Preprocessing was conducted using the Robust Multichip Analysis (RMA) [59] implementation in the Bioconductor "affy"

package (<http://www.bioconductor.org/>), with an additional step to account for differences in probe hybridization resulting from single nucleotide polymorphisms (SNP) between A/J and C57BL/6J mice. The additional step is referred to as SNP masking [60] and is applied after background correction and quantile-quantile normalization but prior to the determination of probeset expression values. Genomic locations hybridized by each probe were obtained from the Ensembl database (<http://www.ensembl.org/index.html>), and these genomic locations were compared to the locations of SNPs for which A/J and C57BL/6J mice have different alleles. Probes that hybridize to such SNPs within their target transcripts were excluded from the determination of probeset expression values. There were 7714 probesets for which some of the probes were excluded. In addition, there were four probesets excluded from further analysis because all probes in the set hybridize to SNPs: 1437478_s_at, 1416030_a_at, 1452066_a_at, and 1449635_at. SNP data were obtained from the Perlegen SNP database (<http://mouse.perlegen.com/mouse/index.html>).

We next filtered the probesets based on probeset annotations. We eliminated from further analysis 6041 probesets with “_x_at” in the probeset name, and 2091 probesets with “_s_at” in the probeset name (including 1437478_s_at mentioned above). Probesets with the “_x_at” designation are those that are known to cross-hybridize in an unpredictable manner. Probesets with the “_s_at” designation are those for which multiple different probesets have probes in common.

The remaining 34,706 probesets were analyzed using Analysis of Variance (ANOVA) to determine whether the difference in mean expression levels between A/J and C57BL/6J mice are statistically significant. The following generalized linear model was used:

$$Y_{i,j,k,l} = B_i^1 + B_j^2 + T_k + S * T_{k,l}$$

where B^1 corresponds to IVT batch effects, B^2 to hybridization batch effects, T to time main effects, and $S * T$ to strain-time interaction effects, where the time factor levels correspond to the uninfected state and 2, 4, 6, and 12 hours post-infection). A list of candidate genes for further analysis was compiled by identifying those genes on chromosomes 8, 11, or 18 with $p < 0.05$ for the $S * T$ model coefficient for all post-infection times (i.e. 2, 4, 6, and 12 hours post-infection). There are 191 such genes (**Table S1**). We have not used a generalized family-wise error rate (gFWER), as stringent P -value thresholds have been shown to reduce the reproducibility of lists of differentially expressed genes [61]. We have chosen, therefore, to use non-adjusted p values and to further refine the candidate gene list based on biological rather than statistical criteria, namely differential expression between strains for multiple post-infection timepoints, as well as GO annotations and FC as discussed below. The estimated false discovery rate ranges from 0.09 (6 hours post-infection) to 0.3 (12 hours post-infection) [62].

SNPs allele differences between A/J and C57BL/6J

Single nucleotide polymorphism (SNP) data were obtained from the Mouse Genome Informatics database (MGI) (<http://www.informatics.jax.org/>) for the A/J and C57BL/6J mouse strains [63] for all genes on the candidate gene list. We identified those SNP loci within the gene boundaries or within 2 kilobases of the gene boundaries and which have allele differences between the two strains. We next identified the functional class of each SNP as

assigned in the MGI database. MGI assigns functional classes according to those defined by dbSNP (<http://www.ncbi.nlm.nih.gov/projects/SNP/>). The functional classes are: coding-nonsynonymous, coding-synonymous, intron, locus-region, splice-site, mRNA-UTR, contig-reference, coding, and coding-exception. SNPs in untranscribed regions but within 2 kilobases of the 5' boundary of a gene or within 0.5 kilobases of the 3' boundary of a gene are assigned the locus-region functional class. SNPs that lie within promoter or enhancer sites are therefore most likely assigned the locus-region functional class.

QTL linkage analysis

Polymorphic microsatellite markers for the cross between C57BL/6J and A/J were chosen from a database maintained by Mouse Genomic Informatics (<http://www.informatics.jax.org/>). To cover chromosome 18, nine microsatellite marker were selected with an average inter-marker distance of 5.3 cM. A total of 144 N_2 backcross were generated, all of which were genotyped for each microsatellite marker by PCR amplification and gel electrophoresis. PCR genotyping reactions were used Platinum Taq polymerase (Invitrogen Co.). J/qtl software was used to analyze phenotype and genotype data for association between survival time after infection with *S. aureus* and marker location. Phenotypes were defined as either sensitive or resistant based on the dichotomization of survival data (Survival of less than 1 day is “0” and survival of longer than 1.5 days is “1”, respectively). Because our analyses focused solely upon one chromosome, our significance threshold reflected a chromosome – level analysis [64–66], rather than a genome-wide approach [67]. All linkage analysis results were expressed as LOD scores. LOD score was considered “suggestive” if > 0.41 ($P = 0.63$) and “Significant” if > 1.64 ($P = 0.05$). Threshold values for linkage were determined by a 1,000 permutation test by using J/qtl.

Isolation of bone marrow-derived neutrophils

Bone marrow-derived neutrophils were isolated as described previously [2]. Briefly, bones from the femurs and tibias of 8-week-old male A/J and C57BL/6J mice were removed of all muscles. The bone marrow cells were flushed out with Hank's buffer (without calcium and magnesium) and filtered through a 70- μ m nylon cell strainer to remove cell clumps. Neutrophils were separated from the remaining cells by centrifugation over discontinuous Percoll (Amersham Biosciences) gradients at 500 $\times g$ for 30 min at 4°C, consisting of 52% (vol/vol), 69% (vol/vol), and 78% (vol/vol) Percoll in PBS. Neutrophils from the interface of the 65 and 75% fractions were washed with Hank's buffer (without calcium and magnesium) and cultured (37°C, 10% CO_2) in RPMI 1640 supplemented with 10% fetal calf serum (both from the American Type Culture Collection).

Neutrophil and macrophage bactericidal assay

Killing of *S. aureus* was assayed as described [68] with modification. *S. aureus* cells were opsonized with 10% mouse serum at 37°C for 30 min and washed twice in PBS. In brief, 1 ml of the RPMI 1640 supplemented with 10% fetal calf serum containing neutrophils (2×10^6 per ml) and *S. aureus* (2×10^6 /ml) was placed in the sterile 1.5 ml tubes. The tubes were incubated at 37°C with end-over-end rotation (6 rpm). After 90 min of incubation, the neutrophils were centrifuged at 100 $\times g$ for 5 min at 4°C and washed twice with 1 ml of cold PBS. Supernatants were used for evaluation of the count of extracellular bacteria. The neutrophil pellet was lysed in sterile water (pH 11) to release intracellular bacteria. The each resulting suspension was serially diluted in PBS and bacterial numbers were determined after

plating onto TSB agar. Peritoneal macrophage bactericidal assay was performed in the same manner as neutrophil bactericidal assay except the final step for bacterial counting [69,70]. After 90 min of incubation, the each sample was diluted in sterile water for 5 min to lyse the macrophage, and serially diluted in PBS. The dilutions were plated on TSB agar, and the plates were incubated overnight at 37°C. The amount of killing was determined by plate count of surviving bacteria (extracellular and intracellular bacteria).

Small interfering RNA (siRNA) experiments

To test the role of each candidate gene on cytokine production by host defense cells, we transfected siRNAs into the mouse macrophage cell line RAW264.7. All siRNAs were purchased from Ambion. Cells were transfected by using siPORT *Amine* Transfection Agent in 48-well plate (Ambion), according to the manufacturer's instructions. Transfections were carried out using 30 nM siRNA and 400,000 cells per well. Cells were maintained in DMEM supplemented with 10% FBS. Forty-eight hours after transfection, *S. aureus* Bioparticles (Invitrogen) was added to a final concentration of 10 µg/ml. After 5 h, supernatants (medium) and cells were collected, stored at -70°C for multiplex cytokine assay (described below) and real-time PCR analyses, respectively. RNA was isolated for detecting the level of mRNA expression. The target gene mRNA expression was analyzed by real-time PCR (TaqMan) using 1 µg total RNA from untreated cells or from cells treated with oligonucleotides. The reaction was performed using the High-Capacity cDNA Reverse Transcription kit (Applied Biosystems) and was followed by Taqman real-time PCR. A full list of gene names, siRNA ID numbers, sequences, and Taqman assay ID numbers are shown in Table S2.

Peritoneal macrophage (PMΦ) isolation and stimulation

Resident PMΦs were obtained by lavage of the peritoneal cavity of uninfected mice (CSS18 and C57BL/6J) as described elsewhere [69,71]. Cells were washed once with PBS by centrifugation (300×g, 5 min at 4°C) and resuspended at a concentration of 1×10⁶ cells/ml of RPMI1640 containing 10% FBS. Cells were incubated in a 24-well plate for 2 h and the remaining nonadherent cells were removed by washing with media. The macrophage monolayers were then stimulated to produce cytokines by incubation with 10 µg of *S. aureus* Bioparticles for 48 h at 37°C. Incubation time was determined based on the previous similar studies [72,73]. Culture supernatants for Luminex assay were collected pre- and post- *S. aureus* stimulation.

Measurement of cytokine/chemokine production

Cytokine production was assayed from the collected supernatant of the *S. aureus*-challenged siRNA transfected macrophages and the *S. aureus*-challenged peritoneal macrophages from C57BL/6J and CSS18 using a multiplex cytokine assay kit and Luminex technology (Bio-Rad) available at Duke Human Vaccine Institute. Twenty cytokines were tested: FGF basic, TNFα, IL-1α, IL-1β, IL-2, IL-4, IL-5, IL-6, IL-10, IL-12p40/p70, IL-13, IL-17, IFNγ, IP-10, KC, MIG, MIP-1α, MCP-1, VEGF, and GM-CSF. Of the 20 tested cytokines, 9 (FGF basic, IFNγ, IL-2, IL-4, IL-5, IL-13, IL-17, MIG, and IP-10) were excluded from our analysis due to weak signal in both pre- and post- *S. aureus* stimulated supernatant (5 cytokines); lack of response to *S. aureus* (2 cytokines), and cross-contamination (2 cytokines) (Figure S5). The concentrations for cytokines and chemokines were compared between two groups (negative control vs target gene) with a two-tailed t-test for unpaired samples. Differences were considered to be statistically significant when the *P*-value was smaller than 0.05.

Supporting Information

Figure S1 Breeding scheme used to generate the chromosome 18 backcross mapping. F1 mice (C18A) generated by crossing CSS18 with C57BL/6J. Recombinant N₂ backcross mice made by mating F1 (C18A) mice to the C57BL/6J inbred mouse strain. A total 144 N₂ backcross mice were generated and used genotyped using microsatellite markers and phenotyped for susceptibility to *S. aureus* infection. In the schematic diagram of the genome, chromosomes from C57BL/6J and A/J are indicated by blue- and red-colored boxes, respectively.

Found at: doi:10.1371/journal.ppat.1001088.s001 (1.04 MB TIF)

Figure S2 No sex-linked survival in N₂ backcross for chromosome 18 mice was identified. A total of 144 N₂ backcross mice, generated by crossing C18A with C57BL/6J, were injected i.p. with 1×10⁷ CFU/g *S. aureus* and observed every 8 h.

Found at: doi:10.1371/journal.ppat.1001088.s002 (0.49 MB TIF)

Figure S3 Bactericidal activities of neutrophils and peritoneal macrophages isolated from A/J and C57BL/6J mice are indistinguishable. (A) Neutrophils (2×10⁶) from bone marrow of A/J, C57BL/6J, or CSS18 mice were incubated with 2×10⁶ CFU of *S. aureus* with end-over-end rotation for 90 min at 37°C. Neutrophils were centrifuged at 100×g for 5 min at 4°C and washed twice with PBS. Supernatants were used for evaluation of the count of extracellular viable bacteria. Neutrophil pellets were disrupted with sterile water (pH 11) to release and quantify intracellular surviving bacteria. Bars represent the mean ± SD of three individual mice. Values of *P*<0.05 were considered significant (unpaired two-tailed t-test). (B) Peritoneal macrophages (2×10⁶) from C57BL/6J and A/J mice were incubated with 2×10⁶ CFU of *S. aureus* with end-over-end rotation for 90 min at 37°C. The each sample was diluted in sterile water to release and quantify surviving bacteria. Bars represent the mean ± SD of three individual mice. Values of *P*<0.05 were considered significant (unpaired two-tailed t-test).

Found at: doi:10.1371/journal.ppat.1001088.s003 (0.83 MB TIF)

Figure S4 Knockdown efficiencies of the 10 target genes in RAW264.7 cells by real-time PCR. mRNA expression levels of each gene in wild-type (WT) and knockdown (KD) RAW264.7 macrophage were determined by real-time PCR. Real-time PCR measurements are normalized to GAPDH mRNA.

Found at: doi:10.1371/journal.ppat.1001088.s004 (1.22 MB TIF)

Figure S5 Cytokine/chemokine profiles from the knockdowns of the 10 target genes in RAW264.7 cells pre- and post- *S. aureus* stimulation. RAW264.7 macrophage cells were transfected with siRNA for each candidate gene or negative control siRNA (30nM). At 48 h after transfection, the levels of cytokines/chemokines in *S. aureus*-stimulated cell culture supernatants were measured by Luminex-based multiplex cytokine assay. The horizontal dashed lines indicate the minimum detectable concentration. Means and standard deviations of three independent experiments are shown. Values of *P*<0.05 were considered significant (unpaired two-tailed t-test).

Found at: doi:10.1371/journal.ppat.1001088.s005 (6.96 MB TIF)

Figure S6 Cytokine/chemokine profiles of peritoneal macrophage from Chromosomal Substitution Strain 18 (CSS18) and C57BL/6J mice pre- and post- *S. aureus* stimulation. CSS18 mice, which contain A/J chromosome 18 but are otherwise genetically C57BL/6J, were used to minimize potential confounding effects of genetic susceptibility factors on A/J chromosomes 8 and 11. The peritoneal macrophage cells were isolated from C57BL/6J and CSS18 and cultured in RPMI1640 containing 10% FBS for 2 h.

The remaining nonadherent cells were removed by washing with media and the macrophage monolayers were then stimulated to produce cytokine by incubation with 10 µg of *S. aureus* Bioparticles for 48 h at 37°C. The levels of cytokines/chemokines in cell culture supernatants at both pre- and post-stimulated conditions were measured by Luminex-based multiplex cytokine assay. The horizontal dashed lines indicate the minimum detectable concentration. Means and standard deviations of three independent experiments are shown. Values of $P < 0.05$ were considered significant (unpaired two-tailed t-test).

Found at: doi:10.1371/journal.ppat.1001088.s006 (3.52 MB TIF)

Table S1 Genes differentially expressed between A/J and C57BL/6J mice at 2, 4, 6, and 12 hours after infection with *S. aureus*. The gene symbol for each of the 191 differentially expressed genes is shown, along with its gene name, chromosome location, human homologue, and fold change and P -value for differential expression at 2 hours post infection. Fold change is shown as the mean expression level in A/J mice divided by the mean expression level in C57BL/6J mice. For genes with multiple probesets, the fold change and P -value are shown for the probeset with largest fold change at 2 hours post infection. All gene information was taken from the Mouse Genome Informatics (MGI) (<http://www.informatics.jax.org/>) and National Center for Biotechnology Information (NCBI) (<http://www.ncbi.nlm.nih.gov/>) databases. * Genes do not have SNP differences identified between A/J and C57BL/6J.

Found at: doi:10.1371/journal.ppat.1001088.s007 (9.73 MB TIF)

Table S2 List of siRNAs and Taqman assay used in this study. Found at: doi:10.1371/journal.ppat.1001088.s008 (2.08 MB TIF)

Acknowledgments

We thank Nikki Walter (Oregon Health & Science University) for critical comments and providing R code for the SNP mask procedure, and Joseph Lucas for guidance on the conduct of the microarray data analysis.

This work has been presented in part at the 47th Infectious Disease Society of America Annual Meeting, Philadelphia, USA, 2009; Abstract 1335.

Author Contributions

Conceived and designed the experiments: SHA HD NJ LGC WKS AKZ DAM SK SL BKSK VGF. Performed the experiments: SHA HD THR. Analyzed the data: SHA HD NJ LGC CLN GDS VGF. Wrote the paper: SHA HD LGC VGF.

References

- Fournier B, Philpott DJ (2005) Recognition of *Staphylococcus aureus* by the innate immune system. *Clin Microbiol Rev* 18: 521–540.
- Deshmukh HS, Hamburger JB, Ahn SH, McCafferty DG, Yang SR, et al. (2009) Critical role of NOD2 in regulating the immune response to *Staphylococcus aureus*. *Infect Immun* 77: 1376–1382.
- von Kockritz-Blickwede M, Rohde M, Oehmcke S, Miller LS, Cheung AL, et al. (2008) Immunological mechanisms underlying the genetic predisposition to severe *Staphylococcus aureus* infection in the mouse model. *Am J Pathol* 173: 1657–1668.
- Hill PC, Wong CG, Voss LM, Taylor SL, Pottumarthy S, et al. (2001) Prospective study of 125 cases of *Staphylococcus aureus* bacteremia in children in New Zealand. *Pediatr Infect Dis J* 20: 868–873.
- Bullock AE, Slobe BS, Vazquez V, Collins AC (1997) Inbred mouse strains differ in the regulation of startle and prepulse inhibition of the startle response. *Behav Neurosci* 111: 1353–1360.
- Logue SF, Paylor R, Wehner JM (1997) Hippocampal lesions cause learning deficits in inbred mice in the Morris water maze and conditioned-fear task. *Behav Neurosci* 111: 104–113.
- Paylor R, Crawley JN (1997) Inbred strain differences in prepulse inhibition of the mouse startle response. *Psychopharmacology (Berl)* 132: 169–180.
- Ralph RJ, Paulus MP, Geyer MA (2001) Strain-specific effects of amphetamine on prepulse inhibition and patterns of locomotor behavior in mice. *J Pharmacol Exp Ther* 298: 148–155.
- Varty GB, Walters N, Cohen-Williams M, Carey GJ (2001) Comparison of apomorphine, amphetamine and dizocilpine disruptions of prepulse inhibition in inbred and outbred mice strains. *Eur J Pharmacol* 424: 27–36.
- Willott JF, Tanner L, O'Steen J, Johnson KR, Bogue MA, et al. (2003) Acoustic startle and prepulse inhibition in 40 inbred strains of mice. *Behav Neurosci* 117: 716–727.
- Zaas AK, Liao G, Chien JW, Weinberg C, Shore D, et al. (2008) Plasminogen alleles influence susceptibility to invasive aspergillosis. *PLoS Genet* 4: e1000101.
- Marquis JF, Gros P (2008) Genetic analysis of resistance to infections in mice: A/J meets C57BL/6J. *Curr Top Microbiol Immunol* 321: 27–57.
- Nadeau JH, Singer JB, Matin A, Lander ES (2000) Analysing complex genetic traits with chromosome substitution strains. *Nat Genet* 24: 221–225.
- Singer JB, Hill AE, Nadeau JH, Lander ES (2005) Mapping quantitative trait loci for anxiety in chromosome substitution strains of mice. *Genetics* 169: 855–862.
- Singer JB, Hill AE, Burrage LC, Olszens KR, Song J, et al. (2004) Genetic dissection of complex traits with chromosome substitution strains of mice. *Science* 304: 445–448.
- Matin A, Collin GB, Asada Y, Varnum D, Nadeau JH (1999) Susceptibility to testicular germ-cell tumours in a 129.MOLF-Chr 19 chromosome substitution strain. *Nat Genet* 23: 237–240.
- Youngren KK, Nadeau JH, Matin A (2003) Testicular cancer susceptibility in the 129.MOLF-Chr19 mouse strain: additive effects, gene interactions and epigenetic modifications. *Hum Mol Genet* 12: 389–398.
- Krewson TD, Supelak PJ, Hill AE, Singer JB, Lander ES, et al. (2004) Chromosomes 6 and 13 harbor genes that regulate pubertal timing in mouse chromosome substitution strains. *Endocrinology* 145: 4447–4451.
- Ackerman KG, Huang H, Grasmann H, Puma C, Singer JB, et al. (2005) Interacting genetic loci cause airway hyperresponsiveness. *Physiol Genomics* 21: 105–111.
- Winawer MR, Kuperman R, Niethammer M, Sherman S, Rabinowitz D, et al. (2007) Use of chromosome substitution strains to identify seizure susceptibility loci in mice. *Mamm Genome* 18: 23–31.
- Gerard C, Gerard NP (1994) C5A anaphylatoxin and its seven transmembrane-segment receptor. *Annu Rev Immunol* 12: 775–808.
- Mullick A, Leon Z, Min-Oo G, Berghout J, Lo R, et al. (2006) Cardiac failure in C5-deficient A/J mice after *Candida albicans* infection. *Infect Immun* 74: 4439–4451.
- von Kockritz-Blickwede M, Konrad S, Foster S, Gessner JE, Medina E (2009) Protective role of complement C5a in an experimental model of *Staphylococcus aureus* bacteremia. *J Innate Immun* 2: 87–92.
- Shimada T, Park BG, Wolf AJ, Brikos C, Goodridge HS, et al. (2010) *Staphylococcus aureus* evades lysozyme-based peptidoglycan digestion that links phagocytosis, inflammasome activation, and IL-1 β secretion. *Cell Host Microbe* 7: 38–49.
- Lowy FD (1998) *Staphylococcus aureus* infections. *N Engl J Med* 339: 520–532.
- Bubeck Wardenburg J, Williams WA, Missiakas D (2006) Host defenses against *Staphylococcus aureus* infection require recognition of bacterial lipoproteins. *Proc Natl Acad Sci U S A* 103: 13831–13836.
- Smith AM, Rahman FZ, Hayee B, Graham SJ, Marks DJ, et al. (2009) Disordered macrophage cytokine secretion underlies impaired acute inflammation and bacterial clearance in Crohn's disease. *J Exp Med* 206: 1883–1897.
- Cailhier JF, Sawatzky DA, Kipari T, Houlberg K, Walbaum D, et al. (2006) Resident pleural macrophages are key orchestrators of neutrophil recruitment in pleural inflammation. *Am J Respir Crit Care Med* 173: 540–547.
- Takahara K, Yashima Y, Omatsu Y, Yoshida H, Kimura Y, et al. (2004) Functional comparison of the mouse DC-SIGN, SIGNR1, SIGNR3 and Langerin, C-type lectins. *Int Immunol* 16: 819–829.
- Kang YS, Kim JY, Bruening SA, Pack M, Charalambous A, et al. (2004) The C-type lectin SIGN-R1 mediates uptake of the capsular polysaccharide of *Streptococcus pneumoniae* in the marginal zone of mouse spleen. *Proc Natl Acad Sci U S A* 101: 215–220.
- Bouwmeester T, Bauch A, Ruffner H, Angrand PO, Bergamini G, et al. (2004) A physical and functional map of the human TNF- α /NF- κ B signal transduction pathway. *Nat Cell Biol* 6: 97–105.
- Raingaud J, Whitmarsh AJ, Barrett T, Derjard B, Davis RJ (1996) MKK3- and MKK6-regulated gene expression is mediated by the p38 mitogen-activated protein kinase signal transduction pathway. *Mol Cell Biol* 16: 1247–1255.
- Zhang S, Weinheimer C, Courtois M, Kovacs A, Zhang CE, et al. (2003) The role of the Grb2-p38 MAPK signaling pathway in cardiac hypertrophy and fibrosis. *J Clin Invest* 111: 833–841.
- Kamata H, Honda S, Maeda S, Chang L, Hirata H, et al. (2005) Reactive oxygen species promote TNF α -induced death and sustained JNK activation by inhibiting MAP kinase phosphatases. *Cell* 120: 649–661.
- Arachchige Don AS, Dallapiazza RF, Benmin DA, Brake T, Cowan CE, et al. (2006) Cyclin G2 is a centrosome-associated nucleocytoplasmic shuttling protein that influences microtubule stability and induces a p53-dependent cell cycle arrest. *Exp Cell Res* 312: 4181–4204.

36. Asai T, Tena G, Plotnikova J, Willmann MR, Chiu WL, et al. (2002) MAP kinase signalling cascade in Arabidopsis innate immunity. *Nature* 415: 977–983.
37. Lu HT, Yang DD, Wysk M, Gatti E, Mellman I, et al. (1999) Defective IL-12 production in mitogen-activated protein (MAP) kinase kinase 3 (Mkk3)-deficient mice. *Embo J* 18: 1845–1857.
38. Sun H, Gong S, Carmody RJ, Hilliard A, Li L, et al. (2008) TIPE2, a negative regulator of innate and adaptive immunity that maintains immune homeostasis. *Cell* 133: 415–426.
39. Woodward MJ, de Boer J, Heidorn S, Hubank M, Kioussis D, et al. (2010) Tnfai β 8 is an essential gene for the regulation of glucocorticoid-mediated apoptosis of thymocytes. *Cell Death Differ* 17: 316–323.
40. Hamilton JA, Stanley ER, Burgess AW, Shadduck RK (1980) Stimulation of macrophage plasminogen activator activity by colony-stimulating factors. *J Cell Physiol* 103: 435–445.
41. Shi Y, Liu CH, Roberts AI, Das J, Xu G, et al. (2006) Granulocyte-macrophage colony-stimulating factor (GM-CSF) and T-cell responses: what we do and don't know. *Cell Res* 16: 126–133.
42. Miller LS, Pietras EM, Uricchio LH, Hirano K, Rao S, et al. (2007) Inflammation-mediated production of IL-1 β is required for neutrophil recruitment against *Staphylococcus aureus* in vivo. *J Immunol* 179: 6933–6942.
43. Mackay TF (2001) The genetic architecture of quantitative traits. *Annu Rev Genet* 35: 303–339.
44. Doebley J (1992) Mapping the genes that made maize. *Trends Genet* 8: 302–307.
45. Stam LF, Laurie CC (1996) Molecular dissection of a major gene effect on a quantitative trait: the level of alcohol dehydrogenase expression in *Drosophila melanogaster*. *Genetics* 144: 1559–1564.
46. Lemon WJ, Swinton CH, Wang M, Berbari N, Wang Y, et al. (2003) Single nucleotide polymorphism (SNP) analysis of mouse pulmonary adenoma susceptibility loci 1–4 for identification of candidate genes. *J Med Genet* 40: e36.
47. Glenn G, van der Geer P (2008) Toll-like receptors stimulate regulated intramembrane proteolysis of the CSF-1 receptor through Erk activation. *FEBS Lett* 582: 911–915.
48. Yamaguchi Y, Fujio K, Shoda H, Okamoto A, Tsuno NH, et al. (2007) IL-17B and IL-17C are associated with TNF- α production and contribute to the exacerbation of inflammatory arthritis. *J Immunol* 179: 7128–7136.
49. Ruefli-Brasse AA, French DM, Dixit VM (2003) Regulation of NF- κ B-dependent lymphocyte activation and development by paracaspase. *Science* 302: 1581–1584.
50. Jagannath C, Hoffmann H, Sepulveda E, Actor JK, Wetsel RA, et al. (2000) Hypersusceptibility of A/J mice to tuberculosis is in part due to a deficiency of the fifth complement component (C5). *Scand J Immunol* 52: 369–379.
51. Gervais F, Desforges C, Skamene E (1989) The C5-sufficient A/J congenic mouse strain. Inflammatory response and resistance to *Listeria monocytogenes*. *J Immunol* 142: 2057–2060.
52. Cerquetti MC, Sordelli DO, Bellanti JA, Hooke AM (1986) Lung defenses against *Pseudomonas aeruginosa* in C5-deficient mice with different genetic backgrounds. *Infect Immun* 52: 853–857.
53. Cerquetti MC, Sordelli DO, Ortegón RA, Bellanti JA (1983) Impaired lung defenses against *Staphylococcus aureus* in mice with hereditary deficiency of the fifth component of complement. *Infect Immun* 41: 1071–1076.
54. Toews GB, Pierce AK (1984) The fifth component of complement is not required for the clearance of *Staphylococcus aureus*. *Am Rev Respir Dis* 129: 597–601.
55. Easmon CS, Glynn AA (1976) Comparison of subcutaneous and intraperitoneal staphylococcal infections in normal and complement-deficient mice. *Infect Immun* 13: 399–406.
56. Cunnion KM, Benjamin DK, Jr., Hester CG, Frank MM (2004) Role of complement receptors 1 and 2 (CD35 and CD21), C3, C4, and C5 in survival by mice of *Staphylococcus aureus* bacteremia. *J Lab Clin Med* 143: 358–365.
57. Rice KC, Firek BA, Nelson JB, Yang SJ, Patton TG, et al. (2003) The *Staphylococcus aureus* cidAB operon: evaluation of its role in regulation of murein hydrolase activity and penicillin tolerance. *J Bacteriol* 185: 2635–2643.
58. Thakker M, Park JS, Carey V, Lee JC (1998) *Staphylococcus aureus* serotype 5 capsular polysaccharide is antiphagocytic and enhances bacterial virulence in a murine bacteremia model. *Infect Immun* 66: 5183–5189.
59. Irizarry RA, Hobbs B, Collin F, Beazer-Barclay YD, Antonellis KJ, et al. (2003) Exploration, normalization, and summaries of high density oligonucleotide array probe level data. *Biostatistics* 4: 249–264.
60. Walter NA, McWeeney SK, Peters ST, Belknap JK, Hitzemann R, et al. (2007) SNPs matter: impact on detection of differential expression. *Nat Methods* 4: 679–680.
61. Shi L, Jones WD, Jensen RV, Harris SC, Perkins RG, et al. (2008) The balance of reproducibility, sensitivity, and specificity of lists of differentially expressed genes in microarray studies. *BMC Bioinformatics* 9 Suppl 9: S10.
62. Storey JD, Tibshirani R (2003) Statistical significance for genomewide studies. *Proc Natl Acad Sci U S A* 100: 9440–9445.
63. Frazer KA, Eskin E, Kang HM, Bogue MA, Hinds DA, et al. (2007) A sequence-based variation map of 8.27 million SNPs in inbred mouse strains. *Nature* 448: 1050–1053.
64. Petryshen TL, Kirby A, Hammer RP, Jr., Purcell S, O'Leary SB, et al. (2005) Two quantitative trait loci for prepulse inhibition of startle identified on mouse chromosome 16 using chromosome substitution strains. *Genetics* 171: 1895–1904.
65. Churchill GA, Doerge RW (1994) Empirical threshold values for quantitative trait mapping. *Genetics* 138: 963–971.
66. Prows DR, Hafertepen AP, Winterberg AV, Gibbons WJ, Jr., Wesselkamper SC, et al. (2008) Reciprocal congenic lines of mice capture the aliq1 effect on acute lung injury survival time. *Am J Respir Cell Mol Biol* 38: 68–77.
67. Lander E, Kruglyak L (1995) Genetic dissection of complex traits: guidelines for interpreting and reporting linkage results. *Nat Genet* 11: 241–247.
68. Clarke TB, Davis KM, Lysenko ES, Zhou AY, Yu Y, et al. (2010) Recognition of peptidoglycan from the microbiota by Nod1 enhances systemic innate immunity. *Nat Med* 16: 228–231.
69. Leijh PC, van Zwet TL, ter Kuile MN, van Furth R (1984) Effect of thioglycolate on phagocytic and microbicidal activities of peritoneal macrophages. *Infect Immun* 46: 448–452.
70. Wilson KR, Napper JM, Denvir J, Sollars VE, Yu HD (2007) Defect in early lung defence against *Pseudomonas aeruginosa* in DBA/2 mice is associated with acute inflammatory lung injury and reduced bactericidal activity in naive macrophages. *Microbiology* 153: 968–979.
71. van Furth R, Diesselhoff-den Dulk MM (1980) Method to prove investigation of particles by macrophages with light microscopy. *Scand J Immunol* 12: 265–269.
72. Zellweger R, Zhu XH, Wichmann MW, Ayala A, DeMaso CM, et al. (1996) Prolactin administration following hemorrhagic shock improves macrophage cytokine release capacity and decreases mortality from subsequent sepsis. *J Immunol* 157: 5748–5754.
73. Whitaker SM, Colmenares M, Pestana KG, McMahon-Pratt D (2008) *Leishmania pifanoi* proteoglycolipid complex P8 induces macrophage cytokine production through Toll-like receptor 4. *Infect Immun* 76: 2149–2156.