

Concentrations, Viability, and Distribution of *Cryptosporidium* Genotypes in Lagoons of Swine Facilities in the Southern Piedmont and in Coastal Plain Watersheds of Georgia[∇]

Michael B. Jenkins,^{1*} Janice L. Liotta,² Araceli Lucio-Forster,² and Dwight D. Bowman²

USDA-Agriculture Research Service, J. Phil Campbell, Sr., Natural Resource Conservation Center, Watkinsville, Georgia,¹ and Department of Microbiology and Immunology, College of Veterinary Medicine, Cornell University, Ithaca, New York²

Received 17 February 2010/Accepted 5 July 2010

Waste lagoons of swine operations are a source of *Cryptosporidium* oocysts. Few studies, however, have reported on oocyst concentrations in swine waste lagoons; none have reported on oocyst viability status, nor has there been a systematic assessment of species/genotype distributions across different types of swine facilities. Ten swine waste lagoons associated with farrowing, nursery, finishing, and gestation operations were each sampled once a month for a year. Oocysts were extracted from triplicate 900-ml effluent samples, enumerated by microscopy, and assessed for viability by dye exclusion/vital stain assay. DNA was extracted from processed samples, and 18S ribosomal DNA (rDNA) genes were amplified by PCR and sequenced for species and genotype identification. Oocysts were observed at each sampling time at each lagoon. Annual mean concentrations of total oocysts and viable oocysts ranged between 24 and 51 and between 0.6 and 12 oocysts ml⁻¹ effluent, respectively. The species and genotype distributions were dominated (95 to 100%) by *Cryptosporidium suis* and *Cryptosporidium* pig genotype II, the latter of which was found at eight of the lagoons. The lagoon at the gestation facility was dominated by *Cryptosporidium muris* (90%), and one farrowing facility showed a mix of pig genotypes, *Cryptosporidium muris*, and various genotypes of *C. parvum*. The zoonotic *C. parvum* bovine genotype was observed five times out of 407 18S rDNA sequences analyzed. Our results indicate that pigs can have mixed *Cryptosporidium* infections, but infection with *C. suis* is likely to be dominant.

Over the last few decades, pork production in North America has undergone significant growth and centralization into large concentrated swine (*Sus scrofa*) operations with more animals on fewer farms (18). A consequence of the increase in numbers of swine per facility is a concomitant increased concentration of swine waste. Present housing facilities for swine are designed to collect feces and urine in wastewater lagoons, in which the waste undergoes anaerobic transformations. One of several public health concerns over swine lagoons is the potential presence of infectious bacteria, viruses, and protozoa (4). Because of the notoriety given to swine waste lagoon spills in the coastal flood plain of North Carolina that were associated with a series of hurricanes in 1998 and 1999 (21), large-scale swine operations have become a focus of environmental and public health concerns.

The cause of the massive outbreak of cryptosporidiosis in Milwaukee, WI, in 1993 was afterwards determined to be *Cryptosporidium hominis*, the human genotype of *C. parvum* and an obligate parasite of humans (33, 44). At the time, however, it was thought to be caused by *C. parvum* (22). Because of this initial misidentification of the cryptosporidial source of the outbreak, the connection between *C. parvum* and large-scale confined livestock operations has become a focused area of research. Although manure-associated outbreaks of *C. parvum*

have implicated bovine sources, a Canadian study found that the prevalence of *Cryptosporidium* in swine lagoons was greater than that in dairy liquid manure (9). Olson et al. (24) also reported the presence of *Cryptosporidium* oocysts of undetermined genotype at four of six hog operations in Canada. Atwill et al. (2) observed *C. parvum* oocysts in feces of feral pigs. Hutchison et al. (13) observed *C. parvum* oocysts of undetermined genotype in 5 and 13% of fresh and stored fecal samples, respectively, from pigs of undeclared age. Guselle et al. (10) followed the course of a naturally occurring *C. parvum* infection in 33 weaned pigs. Following the protocol of the genetic analysis of Morgan et al. (23), Guselle et al. (10) identified this *C. parvum* genotype as being adapted to pigs. At the time, the zoonotic potential of this *C. parvum* pig-adapted genotype was considered uncertain (23).

Recently, two genotypes of *Cryptosporidium* have been recognized as host adapted to swine: *Cryptosporidium suis* (formerly *Cryptosporidium* pig genotype I) and *Cryptosporidium* pig genotype II (28, 29). Xiao et al. (37) reported on an immunocompromised person who was infected with a *Cryptosporidium* pig genotype and thus implicated *Cryptosporidium* from swine as potentially zoonotic and a public health concern. Before molecular methods were developed to differentiate pig genotypes of *Cryptosporidium* from other species, *C. parvum* was thought to infect 152 species of mammals and consist of several cryptic species (6). An extensive survey of swine effluent from swine finishing operations in Ireland indicated a prevalence of both *C. suis* and *Cryptosporidium* pig genotype II (39). Hamnes et al. (11) reported prevalence of both *C. suis* and *Cryptosporidium* pig genotype II in feces of suckling pigs across Norway

* Corresponding author. Mailing address: USDA-ARS, J. Phil Campbell, Sr., Natural Resource Conservation Center, Watkinsville, GA 30677. Phone: (706) 769-5631. Fax: (706) 769-8962. E-mail: michael.jenkins@ars.usda.gov.

[∇] Published ahead of print on 16 July 2010.




FIG. 1. Flow diagram of sampling, processing, and analysis protocol. SSU, small subunit.

and thus implicated farrowing operations as sources of this parasite.

Other than the prevalence of *Cryptosporidium* in feces of young pigs and effluent lagoons of older pigs in finishing operations, little comprehensive data on oocyst concentrations, viability of oocysts, and distributions of *Cryptosporidium* species and genotypes have been reported. No systematic study of swine lagoon effluents from large-scale facilities has been reported for the four separate stages of swine development, (i) breeding and gestation, (ii) farrowing (parturition), (iii) nursery (in which weaned piglets are kept until 8 to 9 weeks of age), and (iv) finishing (in which 8- to 9-week-old pigs are kept to market weight). The objective of this investigation was to determine for 1 year the frequencies, concentrations, viability statuses, and distributions of *Cryptosporidium* species and genotypes in lagoons associated with the four types of swine operations in the Southern Piedmont and in coastal plain watersheds of Georgia.

MATERIALS AND METHODS

Sampling sites. Ten swine lagoons were sampled. Seven lagoons were located in the Southern Piedmont and three were located in the coastal plain of Georgia. Three lagoons were on farrowing operations, four were on finishing operations, one was on a nursery operation, one was on a gestation operation, and one was on a lagoon that received effluent from a farrowing and nursery operation (designated Far/Nur). Each lagoon was sampled once a month from June 2007 to May 2008.

Lagoon sampling protocol. The sampling protocol that was developed considered all safety aspects. Since small boats were launched on lagoons to take samples, all personnel were trained in small boat safety. Three 1-liter surface samples were taken at three different designated locations across each lagoon, one of which was taken near the effluent source pipe. The samples were contained in one-liter Nalgene bottles. Samples were stored on ice while in transit to the laboratory.

The percentage of solids in each lagoon effluent was determined. Ten- to 50-ml subsamples were filtered through tared Whatman no. 2 filter paper. The filter papers were then weighed before and after drying in an oven at 105°C for 24 h. Percentages of solids were determined by dividing the difference between the weight of the dried filter and the tared filter weight by the volume of sample filtered. After subsampling for percentages of solids, the samples were sent on ice by Federal Express overnight to Cornell University for oocyst extraction and further analysis as depicted in Fig. 1. Enumerations and viability assays were performed at the ARS laboratory in Georgia, and BLAST analyses and sequence determinations were performed at Cornell University.

Sample preparation. Of the three effluent samples collected from each of the 10 lagoons, 900 ml of each was filtered through U.S. sieves (no. 200 and 500, with pore sizes of 75 and 25 μm , respectively; Seedburo Equipment Company), and the filtrate was collected in five 250-ml conical tubes. These tubes were centrifuged at 1,500 $\times g$ for 10 min. The supernatant was gently vacuumed off, and the

pellets were combined. The centrifugation, vacuuming, and combining of pellets continued until the total volume fit into a 50-ml tube. The 50-ml tube was centrifuged at 1,500 $\times g$ for 10 min. The resulting pellet was <10 ml. Particularly dense lagoon samples were first floated with 1.3-specific gravity (sp. gr.) sugar before filtration in 50-ml centrifuge tubes and centrifuged at 1,000 $\times g$ for 15 min. The top 15 ml was collected and poured over U.S. sieves no. 200 and 500. The resulting filtrate was processed as stated above for less-dense samples.

To further purify any *Cryptosporidium* oocysts present in the sieved and centrifuged processed pellet, the pellet was placed on a discontinuous sugar gradient of 1.05, 1.08, and 1.10 sp. gr. and concentrated by ultracentrifugation (55,000 $\times g$) in a swinging-bucket rotor for 15 min at 8°C. Oocysts were extracted in and just above the 1.08-sp. gr. layer. After the oocysts were extracted, deionized water was added to bring the volume to three times the extracted volume to reduce the specific gravity to 1.0. This tube was centrifuged at 1,500 $\times g$ for 10 min and the supernatant discarded by vacuuming. The pellet was washed into a smaller centrifuge tube. The centrifugation, vacuuming, and placing into a smaller centrifuge tube continued until the total volume was reduced to 1 ml. This 1-ml suspension was divided in half; one half was used for enumeration and viability assessment, and the other was used for genotyping.

Dye exclusion/vital stain assay. The details of this assay have been described previously (1, 3, 14). Briefly, 10 μl of 4',6-diamidino-2-phenylindole (DAPI; 2 mg ml^{-1} in methanol) and 10 μl of propidium iodide (PI; 1 mg ml^{-1} in 0.1 \times phosphate-buffered saline, pH 7.2) were added to a 100- μl suspension of extracted oocysts in a 1.5-ml microcentrifuge tube. After incubation of the suspension in the dark at room temperature for 1 h, 10 μl of Merifluor fluorescent antibody (Meridian Bioscience Inc., Cincinnati, OH) was added and mixed, and the suspension was incubated in the dark at room temperature for 0.5 h (1, 14). Similar to the result reported by Anguish and Ghiorse (1), oocysts were initially identified with the fluorescent antibody. Further details of its applicability as an indicator of viability or potential for animal infectivity and thus the capability of reproduction under appropriate conditions have been previously discussed (27). As explained in the reply to Robertson et al. (27), to assay the viability of oocysts extracted from an environmental matrix, such as soil, the dye permeability assay is treated explicitly as an indirect assessment of potential infectivity. As such, this dye permeability method (14) was adapted to the U.S. EPA's method 1622 in a study assessing viability of oocysts extracted from surface waters (31). All phases of dye permeability and impermeability have been illustrated in color micrographs (1) and exemplified in images of nonviable oocysts extracted from lagoon effluent (Fig. 2). Viable oocysts were the sum of DAPI-negative (DAPI⁻), PI-negative (PI⁻) oocysts and DAPI-positive (DAPI⁺) PI⁻ oocysts; DAPI⁺ PI⁺ oocysts were considered inactivated. Empty oocyst shells were also counted as inactivated in this assay. The number of viable oocysts was determined as the percentage of total oocysts enumerated.

Microscopy and quantification of oocysts. For enumeration and determination of the viability status of oocysts from each processed sample, duplicate 10- μl aliquots of a stained oocyst suspension were each placed on an agar-coated microscope slide (1% Noble agar air dried on an ethanol-cleaned slide) and each mounted with a 22-mm² coverslip. The slides were examined with a Leica DMR fluorescent microscope with a 100 \times /1.40-0.7 (magnification/numerical aperture) oil PL APO differential interference contrast (DIC) objective and 10 \times eyepieces and with filters to observe oocysts stained with the fluorescent antibody and the two nucleic acid dyes DAPI and PI. Images of each microscopic field were captured with SimplePCI imaging software (Compix Inc., Sewickley, PA) and




FIG. 2. Image of oocysts from the lagoon at the nursery is a combination of images of the immunofluorescent stain and DAPI and PI dyes. The oocyst near the center is DAPI⁺ PI⁻ and is considered viable and potentially infective. The oocysts at 11 and 5 o'clock have excysted and are not viable; the oocysts at 12 and 6 o'clock are DAPI⁺ PI⁺ and are not viable. Bar with internal subdivisions, 5.085 μm .

examined for oocysts. Twenty random fields (out of $>10^5$ possible fields) were brought into view, and oocysts, if present, were counted and their viabilities were assessed for each duplicate subsample of each replicate processed effluent sample. Total counts were determined by the following equation: $\text{cells ml}^{-1} = [(N \times A_c)/(A_f \times V_u)] \times (V_c/V_s)$ (where N is the mean cell count, A_c is the area under the coverslip, A_f is the area of the microscopic field of view, V_u is the volume of sample under the coverslip, V_c is the volume of final oocyst concentrate, and V_s is the volume of effluent from which the oocysts were concentrated).

Genotyping. From the sample preparation, a 0.5-ml portion was used for genotyping as published by Xiao et al. (36, 38). In brief, the genotyping procedure consisted of immunomagnetic separation (Dynabeads anti-*Cryptosporidium* kit; Invitrogen Dynal), with the resulting bead-oocyst complex being used directly in the DNA extraction procedure (QIAamp DNA minikit; Qiagen). Several volumes of DNA, 1, 3, 4, and 6 μl , were used in the PCR. Primers used in the analysis are as delineated by Xiao et al. (36). Positive PCR products were identified by electrophoresis on a 2% agarose gel. In the interest of time, a limited number of positive PCR products were used for the restriction fragment length polymorphism (RFLP) protocol. Instead, all amplified PCR products, including those that were identified by RFLP, were sequenced (Applied Biosystems automated 3730xl DNA analyzer; Biotechnology Resource Center, Cornell University). The resulting sequences were analyzed using the Basic Local Alignment Search Tool (National Center for Biotechnology Information).

Data analysis. Because data like those for the raw oocyst counts have been shown to fit a Poisson distribution (16), the count data were transformed for statistical analysis with the equation $X' = (X + 0.5)^{1/2}$ (43) and transformed means were determined. The transformed means were then back transformed into numbers of oocysts ml^{-1} of effluent for presentation. To make the data behave as if they were normally distributed, these oocyst densities were transformed into natural log numbers to conduct an analysis of variance with lagoons as fixed effects and replicate samples as a random effect in analysis of variance while comparing means for significant differences (at a P value of ≤ 0.05) with Proc Mixed, version 9.2 (30). Significant differences were assigned to tests of transformed means, which are presented in untransformed units of concentration. Analysis of variance was also conducted on the data for percent solids with Proc Mixed (30). Regression relationships between concentrations of total oocysts, viable oocysts, and percent solids from lagoon effluent were analyzed with Proc Reg (30). A chi-square analysis was also performed on the frequency distribution of observed *Cryptosporidium* genotypes with Proc Freq (30), and the logistic regression procedure of SAS (30) was performed on normalized (frequency of genotype/total number of genotypes per lagoon) frequency data.

RESULTS

Concentration and viability assessment. *Cryptosporidium* oocysts were recovered from every lagoon and on each sampling date. Mean oocyst concentrations ranged between 11.0 and 215.7 ml^{-1} of lagoon effluent for finishing facilities, be-




FIG. 3. Annual mean concentrations of total oocysts (A) and viable oocysts (B) at each swine facility. Gest, gestation; Far, farrowing; Nur, nursery; Fin, finishing. Different letters above the bars indicate differences at a P value of ≤ 0.05 .

tween 11.0 and 210.8 ml^{-1} lagoon effluent for the farrowing facilities, between 11.0 and 354.1 ml^{-1} for the nursery facilities, and between 11.0 and 126.8 ml^{-1} lagoon effluent for the one gestation facility sampled. The annual mean concentration of oocysts over the 12 months of sampling by the lagoon indicated that the nursery had a greater concentration of oocysts than three of the finishing facilities, one farrowing facility, and the gestation facility (Fig. 3A). The annual mean concentration of viable oocysts in lagoon effluent was greater at the nursery than at the lagoons of the other facilities (Fig. 3B). Of the mean total annual oocysts at the nursery, 24.2% were observed to be viable; the percentages of viable oocysts among mean annual total oocysts for the other swine facilities ranged between 2.4 and 4.4%. Regression analysis indicated no correlation between total oocyst concentrations and viable oocyst concentrations. The annual mean percentage of lagoon effluent solids was significantly greater for the lagoon receiving effluent from the gestation facility than the other lagoons; differences in annual mean percentages of solids between the other lagoons were not observed (data not shown). Regression analysis indicated no correlation between percentages of solids and total oocyst concentrations.




FIG. 4. RFLP patterns of the dominant *Cryptosporidium* spp. from the different swine lagoons. Lane 1, 50-bp ladder; lane 2, Nur, *C. suis*; lane 3, Nur, *C. suis*; lane 4, *C. parvum* bovine genotype (control); lane 5, Nur, *C. suis*; lane 6, Far3, pig genotype II; lane 7, *C. parvum* bovine genotype (control); lane 8, unsequenced isolate; lane 9, Far/Nur, *C. suis*; lane 10, Far/Nur, *C. suis*; lane 11, Fin4, *C. suis*; lane 12, Fin4, pig genotype II.

Distribution of species and genotypes. Nine species and genotypes of *Cryptosporidium* oocysts were observed in the effluent of the 10 lagoons sampled over the 12-month sampling period: *C. suis*, *Cryptosporidium* pig genotype II, *C. muris*, *C. parvum* rat genotypes, *C. parvum* mouse genotypes, *C. parvum* bovine genotypes, *Cryptosporidium meleagridis*, *Cryptosporidium felis*, and one unknown *C. parvum* genotype. Genetic analysis indicated that *C. suis* and pig genotype II (Fig. 4) were observed in nearly all of the lagoons. The distributions of *Cryptosporidium* oocyst species and genotypes (Table 1) were dominated (73.9%) by *C. suis*, which was observed in every lagoon except for the lagoon at the gestation facility. *Cryptosporidium* pig genotype II was observed in lagoons of eight of the 10 facilities; *C. muris* was dominant in the lagoon of the gestation facility and was observed in one finishing facility and one farrowing facility lagoon, as well as in the lagoon receiving effluent from both a nursery and farrowing facility. Of the less

TABLE 1. Distribution of *Cryptosporidium* genotypes among the 10 swine lagoons examined^a

Facility	No. (%) of isolates of indicated species/genotype			
	<i>C. suis</i>	Pig genotype II	<i>C. muris</i>	Other
Fin1	54 (94.7)	3 (5.3)	0	0
Fin2	41 (100)	0	0	0
Fin3	44 (81.5)	4 (7.4)	6 (11.1)	0
Fin4	21 (63.6)	12 (36.4)	0	0
Far1	9 (26.5)	2 (5.9)	5 (14.7)	18 (52.9)
Far2	33 (93.9)	0	0	2 (6.1)
Far3	17 (85.0)	2 (15.0)	0	0
Nur	42 (80.8)	7 (13.5)	0	1 (5.7)
Far/Nur	40 (90.9)	2 (4.5)	1 (2.3)	1 (2.3)
Gestation	0	3 (7.5)	36 (90.0)	1 (2.5)
Total	301 (73.9)	35 (8.6)	48 (11.8)	23 (5.6)

^a Gest, gestation; Far, farrowing; Nur, nursery; Fin, finishing.

TABLE 2. Results of chi-square analysis of the four sets of *Cryptosporidium* species and genotypes across the 10 locations^a

Analysis type	df	Chi-square	Probability
Location by CrypType	27	480.458	<0.0001
FacType by CrypType	9	328.266	<0.0001
Location by CrypType (<i>C. suis</i> /other) ^b	9	202.809	<0.0001
FacType by CrypType (<i>C. suis</i> /other) ^c	3	139.490	<0.0001
Location by CrypType (<i>C. suis</i> /other, without Far1/Gest) ^d	7	30.805	<0.0001
FacType by CrypType (<i>C. suis</i> /other, without Gest) ^e	2	0.0706	0.9653

^a Results of chi-square analysis of the four sets of *Cryptosporidium* species and genotypes (CrypType; i.e., *C. suis*, *Cryptosporidium* pig genotype II, *C. muris*, and the other less frequently observed species and genotypes [as delineated in Table 1]) across the 10 locations and by type of facility (FacType).

^b Analysis of the distribution of *C. suis* and all other species and genotypes across locations.

^c Analysis of the distribution of *C. suis* and all other species and genotypes across facilities.

^d Analysis of the distribution of *C. suis* and other species and genotypes across locations with Far1 and the gestation facility (Gest) removed.

^e Analysis of the distribution of *C. suis* and other species and genotypes across facilities with only Gest removed.

frequently observed species and genotypes of *Cryptosporidium* oocysts, the *C. parvum* rat genotype (GenBank accession number FJ205699) was observed eight times, *C. parvum* mouse genotypes (GenBank accession numbers AF112571, AF108863, and EU553589) were observed seven times, a *C. parvum* bovine genotype (GenBank accession number AF308600) was observed two times, and *C. meleagridis* (GenBank accession number AF112574) was observed once over the 12 sampling times at the farrowing facility designated Far1. Two *C. parvum* bovine genotypes (GenBank accession numbers DQ656354 and AF308600) were observed in the lagoon of the farrowing facility designated Far2. An unknown *C. parvum* genotype (GenBank accession number EF489037) (45) was observed in the nursery lagoon, and a *C. parvum* bovine genotype (GenBank accession number AF308600) was observed in the lagoon receiving effluent from both a nursery and farrowing facility. *Cryptosporidium felis* was observed once in the lagoon effluent of the gestation facility. The viability status was unknown for the oocysts from which the DNA was extracted for genetic analysis.

For the purpose of performing a chi-square analysis, all of the less frequently observed genotypes of *C. parvum*, *Cryptosporidium meleagridis*, and *Cryptosporidium felis* and the one unknown *C. parvum* genotype were considered other genotypes (Table 1). The chi-square analysis (Table 2) indicated that the observations were not similarly distributed across the 10 swine operations ($P < 0.0001$). When the facilities were analyzed by production type (finishing, farrowing, nursery, and gestation), the distributions of the species and genotypes of *Cryptosporidium* oocysts were still not similar among the facility types ($P < 0.001$) and were likely skewed by there being only one gestation facility in which no *C. suis* was observed. When the chi-square analysis was run with *C. suis* and the other species and genotypes were truncated into a single set, the distributions were still not similar among production facilities ($P < 0.0001$). If data for the gestation facility and farrowing facility Far1 were removed from the data set, there was still

TABLE 3. Odds ratio estimates of the distribution of *Cryptosporidium* genotypes/species across the 10 swine lagoons^a

Genotype/species	Odds ratio ^b	95% Wald confidence interval
<i>C. suis</i>	71.616***	8.830–531.355
Pig genotype II	2.382	0.487–9.049
<i>C. muris</i>	0.766**	0.168–4.504

^a The distribution of each of the shown genotypes/species is compared to the distributions of the other less frequently observed species and genotypes, such as the zoonotic bovine *C. parvum* type II, as well as others.

^b **, significant at a *P* of ≤0.01; ***, significant at a *P* of ≤0.001.

significant variation among the remaining eight operations when testing for the proportion of *C. suis* in positive samples (*P* < 0.0001). However, when the facility types excluding the gestation facility were analyzed, no significant variation in the proportions of *C. suis* was observed (*P* = 0.9653).

Results of a logistic regression analysis (Table 3) indicated that the odds of observing *C. suis* in the lagoon effluent of the swine facilities sampled would be 71.6 times greater than the odds of observing *C. parvum* genotypes. Although not statistically significant, the odds of observing *Cryptosporidium* pig genotype II would be 2.4 times the odds of observing *C. parvum* genotypes; the odds of observing *C. muris* would be 0.8 times the odds of observing *C. parvum* genotypes.

DISCUSSION

Viability assessment and distribution of *Cryptosporidium* genotypes. *Cryptosporidium* oocysts were detected and quantified in all the effluent lagoons on each sampling occasion. In contrast to our results, previous prevalence studies on the distribution of *Cryptosporidium* oocysts in swine effluent lagoons have reported observation frequencies less than 100%. Reinoso and Becares (26) reported detecting *Cryptosporidium* oocysts in 53% of the effluent samples taken from 13 intensive swine operations in Spain. Hutchison et al. (13) reported detecting oocysts in only 3 of 58 swine effluent lagoons sampled in the United Kingdom. Cote et al. (5) reported detecting oocysts in only one operation out of 32 finishing swine operations sampled in Quebec, Canada. Xiao et al. (39) reported detecting *Cryptosporidium* in effluent from 14 of 33 finishing facilities in Ireland. The difference between our frequency of observations and those of Reinoso and Becares (26), Hutchison et al. (13), and Xiao et al. (39) may be attributed to differences in sampling scheme, sample size, and methods of oocyst extraction and detection. The concentrations that we have reported (Fig. 3) are in the ranges that Reinoso et al. (26) and Hutchison et al. (13) reported, 16 to 233 and 140 to 310 oocysts ml⁻¹ effluent, respectively. Assessment of viability was not reported in any of these previous studies.

Our observation that the nursery operation, which has two houses and a 0.3-ha lagoon, compared to the finishing operations with four houses and 0.6-ha lagoons, had the greatest concentration of oocysts is in agreement with the observation of Vitovec et al. (34). They reported that the greatest prevalence of *Cryptosporidium* oocysts in porcine fecal samples was associated with postweaning piglets between 6 and 8 weeks old.

The greatest proportion of viable oocysts was also associated with the nursery operation (Fig. 3). The nursery's greater frequency of viable oocysts may be attributable to the greater influx and load of oocysts in the effluent lagoon. However, the observation that the viability assessment for the lagoon receiving effluent from a farrowing and nursery operation was not different from those of the other lagoons implies that our observation of the elevated oocyst viability at the nursery operation may be anomalous. Apart from the nursery operation we sampled, the mean concentration of viable oocysts in the other lagoons was less than 10% of the mean total. Our results indicate that there appears to be a continuous input of oocysts into the lagoons from their respective housing facilities. Based on infection prevalences of 16, 31, and 100% in sows, piglets, and postweaning piglets (20), respectively, we can infer that the apparent inflow of oocysts into the lagoon indicates that some of the pigs in the houses were shedding oocysts in their feces at any given time. The apparent die-off of oocysts may, in part, be attributed to known ammonia emissions from lagoons (12) and deleterious effects of ammonia on oocyst viability (15).

Our determination of the mean concentration of oocysts between 25,000 and 500,000 oocysts dm⁻³ is indicative of their persistence in effluent lagoons for all four types of swine operations and expands on the observations of Xiao et al. (39) that oocysts of unknown viability appeared to persist in swine effluent from finishing operations in Ireland. We observed that a small fraction of the oocysts appeared to be viable and potentially infective. Thus, an application of swine effluent to spray fields would include a load of potentially infective *Cryptosporidium* oocysts. Given that 500 m³ of swine effluent may be applied to a spray field as a source of N for plant growth (32) and given that the concentration range of viable oocysts for any of the lagoons in our study was between 1,000 and 12,000 dm⁻³, the load of viable oocysts applied to a spray field could range between 5 × 10⁸ and 6 × 10⁹ viable oocysts.

The distribution of *Cryptosporidium* oocysts among the lagoons of the swine facilities sampled indicated that the frequency of observing *C. suis* would be significantly greater than that for *C. parvum* genotypes, at a ratio of >10:1. Combining *C. suis* and *Cryptosporidium* pig genotype II further increases the ratio of swine-adapted genotypes to *C. parvum* genotypes. Whereas Langkjaer et al. (19) reported that pig genotype II was most prevalent for weaners, we observed that *C. suis* dominated the nursery lagoon. Except in the gestation facility, the ratio of *C. suis* to *Cryptosporidium* pig genotype II favored the former and ranged from 20:1 to 7:4. The yearlong sampling of the lagoon receiving effluent from the gestation facility proved to be anomalous: 90% of the samples were identified as *C. muris*, no *C. suis* was observed, a small percentage of *Cryptosporidium* pig genotype II was identified, and *C. felis* was observed once. *Cryptosporidium felis* has been known to infect cows and both immunocompromised and immunocompetent humans (7). Because this observation of *C. felis* was associated with an effluent lagoon (which could average a volume of 20,000 m³) into which fecal waste from gestating sows was collected, its presence appears to implicate swine as a source, which would be, to the best of our knowledge, a first.

The observation of the *C. parvum* rat genotype at farrowing facility Far1 may be a first for swine facilities in the United States. This *C. parvum* rat genotype matched one that Feng et

al. (8) reported isolating from a wastewater treatment plant in Shanghai, China, and suggests a worldwide distribution of this genotype. The source of this rat genotype was likely a known population of rats inhabiting the houses. The *C. meleagridis* isolate from Far1 matched one isolated from a turkey (36). In addition to *C. muris* being observed in the lagoon at Far1, several isolates of a *C. parvum* mouse genotype that matched those identified in three different studies were observed (23, 25, 36). The source of this genotype was likely a population of mice inhabiting the operation. An unknown "other" *Cryptosporidium* genotype observed at the nursery matched an isolate observed in Ireland (45) and suggests an isolate with a continental distribution.

Our observations over the yearlong period of sampling the lagoons of the 10 swine facilities support previous observations that pigs can be infected with *C. suis*, *Cryptosporidium* pig genotype II, and *C. parvum* (29) as well as *C. muris* (39, 45). We emphasize that the percentages of *C. parvum* genotypes observed ranged from 0 to 2% except for that at farrowing facility Far1. This observation is similar to the prevalence study that Zintl et al. (45) reported. The low frequency of *C. parvum* genotypes that we observed also supports the prevalence study that Johnson et al. (17) reported, in which they maintained that domestic pigs do not appear to pose a significant public health risk.

Because the morphologies of *C. suis* and *C. parvum* oocysts are indistinguishable (29), observations by microscopy of these oocysts in swine feces and lagoon effluent may have been assumed to be *C. parvum*. The reports in the literature that pigs and lagoons of swine facilities may be a significant source of *C. parvum* (2, 9, 13, 24) may actually refer to *C. suis*. The weaner pigs infected with the *C. parvum* pig genotype that Guselle et al. (10) reported were likely infected by *C. suis*, as Ryan et al. (29) later described. As Zintl et al. (45) observed, mixed infections in pigs can be common, and our data support this view. *Cryptosporidium suis* is, however, likely to be the dominant genotype, and the frequency of observing *C. parvum* genotypes in pigs has been low (42). Based on recent studies, the zoonotic potential of *C. suis* is not great. Most human cases of cryptosporidiosis in the United States and worldwide (over 90%) are associated with *C. hominis* and *C. parvum*. A smaller proportion of cases have been associated with *C. meleagridis*, *C. felis*, and *C. canis* (40, 41). Although *C. suis* and *C. suis*-like organisms have been associated with sporadic cases of cryptosporidiosis, these reports number in the single digits (35, 40). *Cryptosporidium suis* appears not to be an important cause of zoonotic disease in humans; however, immunocompromised individuals, known to be at greater risk of infection with *Cryptosporidium* spp., should take special precautions to prevent exposure to these parasites.

ACKNOWLEDGMENTS

We express our appreciation to Gregory Surratt, Michael Thornton, and Shaheen Humayoun for their expert technical assistance.

This project was supported in part by the USDA-CSREES-National Research Initiative Competitive Grants Program.

REFERENCES

1. Anguish, L. J., and W. C. Ghiorse. 1997. Computer-assisted laser scanning and video microscopy for analysis of *Cryptosporidium parvum* oocysts in soil, sediment, and feces. *Appl. Environ. Microbiol.* **63**:724–733.
2. Atwill, E. R., R. A. Sweitzer, M. D. G. C. Pereira, I. A. Gardner, D. Van Vuren, and W. M. Boyce. 1997. Prevalence of and associated risk factors for shedding *Cryptosporidium parvum* oocysts and *Giardia* cysts within feral pig populations in California. *Appl. Environ. Microbiol.* **63**:3946–3949.
3. Campbell, A. T., L. J. Robertson, and H. V. Smith. 1992. Viability of *Cryptosporidium parvum* oocysts: correlation of in vitro excystation with inclusion or exclusion of fluorogenic vital dyes. *Appl. Environ. Microbiol.* **58**:3488–3493.
4. Cole, D., L. Todd, and S. Wing. 2000. Concentrated swine feeding operations and public health: a review of occupational and community health effects. *Environ. Health Perspect.* **108**:685–699.
5. Cote, C., A. Villeneuve, L. Lessard, and S. Quessy. 2006. Fate of pathogenic and nonpathogenic microorganisms during storage of liquid hog manure in Quebec. *Livestock Sci.* **102**:204–210.
6. Fayer, R., U. Morgan, and S. J. Upton. 2000. Epidemiology of *Cryptosporidium*: transmission, detection, and identification. *Int. J. Parasitol.* **30**:1305–1322.
7. Fayer, R., M. Santin, J. M. Trout, and J. P. Dubey. 2006. Detection of *Cryptosporidium felis* and *Giardia duodenalis* assemblage F in a cat colony. *Vet. Parasitol.* **140**:44–53.
8. Feng, Y., N. Li, L. Duan, and L. Xiao. 2009. *Cryptosporidium* genotype and subtype distribution in raw wastewater in Shanghai, China: evidence for possible unique *Cryptosporidium hominis* transmission. *J. Clin. Microbiol.* **47**:153–157.
9. Guan, T. Y., and R. A. Holley. 2003. Pathogen survival in swine manure environments and transmission of human enteric illness—a review. *J. Environ. Qual.* **32**:383–392.
10. Guselle, N. J., A. J. Applebee, and M. E. Olson. 2003. Biology of *Cryptosporidium parvum* in pigs: from weaning to market. *Vet. Parasitol.* **113**:7–18.
11. Hamnes, I. S., B. K. Gjerde, T. Forberg, and L. J. Robertson. 2007. Occurrence of *Cryptosporidium* and *Giardia* in suckling piglets in Norway. *Vet. Parasitol.* **144**:222–233.
12. Harper, L. H., R. R. Sharpe, T. B. Parkin, A. D. Visscher, O. van Cleemput, and F. M. Byers. 2002. Nitrogen cycling through swine production systems: ammonia, dinitrogen, and nitrous oxide emissions. *J. Environ. Qual.* **33**:1189–1201.
13. Hutchison, M. L., L. D. Walters, S. M. Avery, B. A. Synge, and A. Moore. 2004. Levels of zoonotic agents in British livestock manures. *Lett. Appl. Microbiol.* **39**:207–214.
14. Jenkins, M. B., L. J. Anguish, D. D. Bowman, M. J. Walker, and W. C. Ghiorse. 1997. Assessment of a dye permeability assay for determination of inactivation rates of *Cryptosporidium parvum* oocysts. *Appl. Environ. Microbiol.* **63**:3844–3850.
15. Jenkins, M. B., D. D. Bowman, and W. C. Ghiorse. 1998. Inactivation of *Cryptosporidium parvum* oocysts by ammonia. *Appl. Environ. Microbiol.* **64**:784–788.
16. Jenkins, M. B., A. J. Franzluebbers, and S. B. Humayoun. 2006. Assessing short-term responses of prokaryotic communities in bulk and rhizosphere soils to tall fescue endophyte infection. *Plant Soil* **289**:309–320.
17. Johnson, J., R. Buddle, S. Reid, A. Armson, and U. M. Ryan. 2008. Prevalence of *Cryptosporidium* genotypes in pre- and post-weaned pigs in Australia. *Exp. Parasitol.* **119**:418–421.
18. Key, N., and W. McBride. 2007. The changing economics of U.S. hog production. ERR-52. United States Department of Agriculture, Economic Research Service, Washington, DC. <http://www.ers.usda.gov/Publications/ERR52/>. Accessed 10 June 2010.
19. Langkjaer, R. B., H. Vigre, H. L. Enemark, and C. Maddox-Hyttel. 2007. Molecular and phylogenetic characterization of *Cryptosporidium* and *Giardia* from pigs and cattle in Denmark. *Parasitology* **134**:339–350.
20. Maddox-Hyttel, C., R. B. Langkjaer, H. L. Enemark, and H. Vigre. 2006. *Cryptosporidium* and *Giardia* in different age groups of cattle and pigs—occurrence and management associated risk factors. *Vet. Parasitol.* **141**:48–59.
21. Mallin, M. A. 2000. Impacts of industrial animal production on rivers and estuaries. *Am. Sci.* **88**:26–37.
22. McDonald, A. C., W. R. MacKenzie, D. G. Addiss, M. S. Gradus, G. Linke, E. Zembrowski, M. R. Hurd, M. J. Arrowood, P. J. Lammie, and J. W. Priest. 2001. *Cryptosporidium parvum*-specific antibody responses among children residing in Milwaukee during the 1993 waterborne outbreak. *J. Infect. Dis.* **183**:1373–1379.
23. Morgan, U. M., L. Xiao, R. Fayer, T. K. Graczik, A. A. Lal, P. Deplazes, and R. C. A. Thompson. 1999. Phylogenetic analysis of *Cryptosporidium* isolates from captive reptiles using 18S rDNA sequence data and random amplified polymorphic DNA analysis. *J. Parasitol.* **85**:525–530.
24. Olson, M. E., C. L. Thorlakson, L. Deselliers, D. W. Morck, and T. A. McAllister. 1997. *Giardia* and *Cryptosporidium* in Canadian farm animals. *Vet. Parasitol.* **68**:375–381.
25. Pedraza-Diaz, S., L. M. Ortega-Mora, B. A. Carrion, V. Navarro, and M. Gomez-Bautista. 2009. Molecular characterization of *Cryptosporidium* isolates from pet reptiles. *Vet. Parasitol.* **160**:204–210.
26. Reinoso, R., and E. Becares. 2008. The occurrence of intestinal parasites in

- swine slurry and their removal in activated sludge plants. *Bioresour. Technol.* **99**:6661–6665.
27. **Robertson, L. J., A. T. Campbell, and H. V. Smith.** 1998. Viability of *Cryptosporidium parvum* oocysts: assessment by the dye permeability assay. *Appl. Environ. Microbiol.* **64**:3544–3545. (Letter.)
 28. **Ryan, U. M., B. Samarasinghe, C. Read, J. R. Buddle, I. D. Robertson, and R. C. A. Thompson.** 2003. Identification of a novel *Cryptosporidium* genotype in pigs. *Appl. Environ. Microbiol.* **69**:3970–3974.
 29. **Ryan, U. M., P. Monis, H. L. Enemark, I. Sulaiman, B. Samarasinghe, C. Read, R. Buddle, I. Robertson, L. Zhou, R. C. A. Thompson, and L. Xiao.** 2004. *Cryptosporidium suis* n. sp. (Apicomplexa: Cryptosporidiidae) in pigs (*Sus scrofa*). *J. Parasitol.* **90**:769–773.
 30. **SAS Institute.** 2004. SAS/STAT 9.1 user's guide. SAS Institute, Cary, NC.
 31. **Simmons, O. D., III, M. D. Sobsey, C. D. Heaney, F. W. Schaefer III, and D. S. Franczy.** 2001. Concentration and detection of *Cryptosporidium* oocysts in surface water samples by method 1622 using ultrafiltration and capsule filtration. *Appl. Environ. Microbiol.* **67**:1123–1127.
 32. **Sistani, K. R., M. R. McLaughlin, and G. E. Brink.** 2008. Soil nutrient evaluation from swine effluent application to five forage-system practices. *Nutr. Cycl. Agroecosyst.* **82**:265–271.
 33. **Sulaiman, I. M., L. Xiao, C. Yang, L. Escalante, A. Moore, C. B. Beard, M. J. Arrowood, and A. A. Lal.** 1998. Differentiating human from animal isolates of *Cryptosporidium parvum*. *Emerg. Infect. Dis.* **4**:681–685.
 34. **Vitovec, J., K. Hamadejova, L. Landova, M. Kvac, D. Kvetonova, and B. Sak.** 2006. Prevalence and pathogenicity of *Cryptosporidium suis* in pre- and post-weaned pigs. *J. Vet. Med. B* **53**:239–243.
 35. **Wang, J. C., L. X. Zhang, C. S. Ning, F. C. Jian, and M. F. Sun.** 2007. Identification and phylogenetic analysis of a *Cryptosporidium* isolate from a patient admitted in hospital. *Chin. J. Zoon.* **23**:997–1000.
 36. **Xiao, L., U. M. Morgan, J. Limor, A. Escalante, M. Arrowood, W. Shulaw, R. C. A. Thompson, R. Fayer, and A. A. Lal.** 1999. Genetic diversity within *Cryptosporidium parvum* and related *Cryptosporidium* species. *Appl. Environ. Microbiol.* **65**:3386–3391.
 37. **Xiao, L., C. Bern, M. Arrowood, I. Sulaiman, L. Zhou, V. Kawai, A. Vivar, A. A. Lal, and R. H. Gilman.** 2002. Identification of the *Cryptosporidium* pig genotype in a human patient. *J. Infect. Dis.* **185**:1845–1848.
 38. **Xiao, L. L., A. A. Lal, and J. Jiang.** 2004. Detection and differentiation of *Cryptosporidium* oocysts in water by PCR-RFLP. *Methods Mol. Biol.* **268**:163–176.
 39. **Xiao, L., J. E. Moore, U. Ukoh, W. Gatei, C. J. Lowery, T. M. Murphy, J. S. G. Dooley, B. C. Miller, P. J. Rooney, and J. R. Rao.** 2006. Prevalence and identity of *Cryptosporidium* spp. in pig slurry. *Appl. Environ. Microbiol.* **72**:4461–4463.
 40. **Xiao, L. L., and Y. Y. Feng.** 2008. Zoonotic cryptosporidiosis. *FEMS Immunol. Med. Microbiol.* **52**:309–323.
 41. **Xiao, L.** 2009. Molecular epidemiology of human cryptosporidiosis in developing countries, p. 25–30. *In* M. G. Ortega-Pierres, S. Caccio, R. Fayer, T. Mank, and R. C. A. Thompson (ed.), *Giardia and Cryptosporidium: from molecules to disease*. CABI, Wallingford, United Kingdom.
 42. **Xiao, L.** 2010. Molecular epidemiology of cryptosporidiosis: an update. *Exp. Parasitol.* **124**:80–89.
 43. **Zar, J. H.** 1999. *Biostatistical analysis*. Prentice-Hall, Upper Saddle River, NJ.
 44. **Zhou, A., A. Singh, J. Jiang, and L. Xiao.** 2003. Molecular surveillance of *Cryptosporidium* spp. in raw waste water in Milwaukee: implications for understanding outbreak occurrence and transmission dynamics. *J. Clin. Microbiol.* **41**:5254–5257.
 45. **Zintl, A., D. Neville, D. Maguire, S. Fanning, G. Mulcahy, H. V. Smith, and T. De Waal.** 2007. Prevalence of *Cryptosporidium* species in intensively farmed pigs in Ireland. *Parasitology* **134**:1575–1582.