

**Kavitha Marapakala,^a A. Abdul
 Ajees,^a Jie Qin,^a Banumathi
 Sankaran^b and Barry P. Rosen^{a,b,*}**

^aDepartment of Cellular Biology and
 Pharmacology, Florida International University,
 Herbert Wertheim College of Medicine, Miami,
 Florida, USA, and ^bBerkeley Center for Structural
 Biology, Lawrence Berkeley Laboratory,
 1 Cyclotron Road, BLDG 6R2100, Berkeley,
 CA 94720, USA

Correspondence e-mail: brosen@fiu.edu

Received 26 June 2010

Accepted 13 July 2010

Crystallization and preliminary X-ray crystallographic analysis of the ArsM arsenic(III) *S*-adenosylmethionine methyltransferase

Arsenic is the most ubiquitous environmental toxin and carcinogen and consequently ranks first on the Environmental Protection Agency's Superfund Priority List of Hazardous Substances. It is introduced primarily from geochemical sources and is acted on biologically, creating an arsenic biogeochemical cycle. A common biotransformation is methylation to monomethylated, dimethylated and trimethylated species. Methylation is catalyzed by the ArsM (or AS3MT) arsenic(III) *S*-adenosylmethionine methyltransferase, an enzyme (EC 2.1.1.137) that is found in members of every kingdom from bacteria to humans. ArsM from the thermophilic alga *Cyanidioschyzon* sp. 5508 was expressed, purified and crystallized. Crystals were obtained by the hanging-drop vapor-diffusion method. The crystals belonged to the monoclinic space group *C*2, with unit-cell parameters $a = 84.85$, $b = 46.89$, $c = 100.35$ Å, $\beta = 114.25^\circ$ and one molecule in the asymmetric unit. Diffraction data were collected at the Advanced Light Source and were processed to a resolution of 1.76 Å.

1. Introduction

Arsenic is the most ubiquitous environmental toxin and carcinogen and consequently ranks first on the Environmental Protection Agency's Superfund Priority List of Hazardous Substances (<http://www.atsdr.cdc.gov/cercla/07list.html>). It is introduced primarily from geochemical sources and is acted on biologically, creating an arsenic biogeochemical cycle. Members of every kingdom methylate arsenite in a series of alternating oxidative methylations and reductions, producing the pentavalent species methylarsenate (MAs^V), dimethylarsenate (DMAs^V) and trimethylarsine oxide (TMAs^VO) and the trivalent species MAs^{III}, DMAs^{III} and TMAs^{III} (Challenger, 1951). Humans and other mammals methylate arsenite to species such as DMAs^V, which is excreted in the urine (Thomas *et al.*, 2004). The mammalian liver enzyme that catalyzes transfer of the methyl group of *S*-adenosylmethionine to As^{III} has been called AS3MT (Styblo *et al.*, 2002). It is not clear whether methylation in humans is a detoxification process. Because the more toxic trivalent intermediates MAs^{III} and DMAs^{III} are formed during liver biotransformation of inorganic arsenate and arsenite, methylation may activate inorganic arsenic to more carcinogenic species.

In contrast, it has been clearly demonstrated that microbial methylation detoxifies arsenic. Genes for orthologues of the human AS3MT are found in bacteria, archaea, fungi and lower plants. These genes have been termed *arsM* and their protein product has been termed ArsM (arsenite *S*-adenosylmethyltransferase; Qin *et al.*, 2006). Since ArsMs are widespread in every kingdom, we have hypothesized that this biotransformation has a significant impact on the global arsenic cycle (Qin *et al.*, 2009). ArsM (accession No. ACN39191) from the Yellowstone acidothermoacidophilic eukaryotic red alga *Cyanidioschyzon* sp. 5508 is a 400-residue enzyme [411 residues in the cloned gene after the addition of a His tag (AAALEHHHHHH) at the C-terminus] with a mass of 44 980 Da that methylates As^{III} to a final product of volatile TMAs^{III}. When the algal *arsM* gene was expressed in an arsenic-hypersensitive strain of *Escherichia coli*, it conferred As^{III} resistance in *E. coli* in the absence of any other arsenic-resistance genes, demonstrating that methylation



is sufficient to detoxify arsenic. Purified ArsM is a thermostable enzyme with a temperature optimum of 333–343 K.

Here, we report the crystallization of two active derivatives of the 400-residue ArsM7 from *Cyanidioschyzon* sp. 5508, one lacking the C-terminal 28 residues (ArsM7A) and the other lacking these residues plus the N-terminal 31 residues (ArsM7B). ArsM7A crystallized in the monoclinic space group *C2*, with unit-cell parameters $a = 84.85$, $b = 46.89$, $c = 100.35$ Å, $\beta = 114.25^\circ$ and one molecule in the asymmetric unit. Diffraction data were collected at the Advanced Light Source and processed to a resolution of 1.76 Å.

2. Materials and methods

2.1. Construction of *arsM* genes, ArsM expression and purification

Attempts to crystallize full-length ArsM were unsuccessful. From alignment of putative ArsM orthologues, it appears that the algal ArsM has N-termini and C-termini that are not conserved in other members of the group. For this reason, the genes for two catalytically active derivatives lacking the C-terminal 28 residues or both the C-terminal 28 residues and the N-terminal 31 residues were constructed from the sequence of the full-length *arsM7* gene (Qin *et al.*, 2009).

For construction of a plasmid with the *arsM7A* deletion, a PCR fragment was amplified from the full-length *arsM7* gene using the forward primer 5'-GATATA**CCATGGCGTGCAGCTGTGCGTCTGG**-3' (*NcoI* site in bold) and the reverse primer 5'-**TGCGGCCGCTTCACAGACAAGCTGC**-3' (*NotI* site in bold) and then cloned into vector plasmid pET28a(+) (Novagen) as an *NcoI/NotI* digest, generating plasmid pET28-arsM7A, in which the *arsM7A* gene is under the control of the T7 promoter. This construct encodes an *arsM7* gene lacking the last 28 residues but with the last two residues (Cys-Glu) changed to Ser-Gly; the sequence for the same histidine tag was added at the 3'-end. The 383-residue ArsM7A protein has a molecular mass of 42 228 Da.

While the location of the C-terminal truncation was decided by inspection of a multiple alignment, the decision for truncation from




Figure 1
ArsM7A was purified by Ni-NTA chromatography as described in Qin *et al.* (2009). The SDS-PAGE profile of ArsM7A is shown. Lane 1 contains molecular-weight markers (labeled in kDa). Lanes 2–5 are sequential fractions eluted with an imidazole gradient. Fractions 4 and 5 were used for crystallization trials. The arrow indicates the position of ArsM7A.

the N-terminus was made following limited trypsin digestion of full-length ArsM7. Limited trypsin digestion was performed at room temperature in 50 mM KHPO₄ pH 7.4 buffer and 2 mg ml⁻¹ protein. The ArsM7:trypsin ratio was 500:1(w:w). Digestion was initiated by the addition of *N*-*p*-tosyl-L-phenylalanine chloromethyl ketone-treated trypsin (Sigma–Aldrich, St Louis, Missouri, USA). Proteolysis was terminated at times between 10 and 300 min by the addition of 0.2% trifluoroacetic acid and was analyzed by 12% SDS-PAGE (Laemmli, 1970) with Coomassie Blue staining. During this entire period, a fragment of approximately 38 kDa remained resistant to trypsin digestion. Gel slices were excised from the trypsin-resistant band, cut into 1 mm³ cubes and transferred into Eppendorf tubes. The bands were subjected to in-gel trypsin digestion and the peptides were analyzed by mass spectrometry by the Proteomics Facility Core of the Institute of Environmental Health Sciences at Wayne State University, Detroit, Michigan, USA. From this analysis, the sites of limited trypsin digestion were determined to be Lys31 at the N-terminal end and Lys375 at the C-terminal end, indicating a trypsin-resistant core of 344 residues with a mass of 38 185 Da. Since ArsM7A was already truncated at Val370, the gene for an N-terminally truncated derivative was constructed from the *arsM7A* gene. In this construct, the first 31 residues of the sequence were deleted and the codons for two residues (Met-Ala) were added to the 5'-end, encoding a protein of 354 residues with a mass of 39 223 Da. Plasmid pET28-arsM7A was used as a template for PCR amplification using the forward primer 5'-GATATA**CCATGGAAACGCTTCAGTCAAG**-3' (*NcoI* site in bold) and the reverse primer 5'-**TGCGGCCGCTTCACAGACAAGCTGC**-3' (*NotI* site in bold). The PCR product was cloned into pET28a(+) as an *NcoI/NotI* digest, generating plasmid pET28-arsM7B encoding the doubly N- and C-terminally truncated ArsM. All genes were verified by DNA sequencing. All ArsM proteins were purified as described in Qin *et al.* (2009).

2.2. Crystallization

Proteins were exchanged into a buffer consisting of 50 mM MOPS pH 7.0 containing 0.5 M NaCl and 10 mM DTT. Initially, sitting-drop crystallization experiments were set up with 3.0 µl 15 mg ml⁻¹ protein solution in 96-well plates (Corning 3785) at 293 K using a variety of crystal screens from Hampton Research (Aliso Viejo, California, USA), Emerald BioSystems Inc. (Bainbridge Island, Washington,




Figure 2
Crystals of ArsM7A were grown in hanging drops as described in §2. The crystal indicated by the arrow has approximate dimensions of 0.4 × 0.3 × 0.2 mm.




Figure 3
A 1° oscillation image collected from an Arsm7A crystal at the Lawrence Berkeley National Laboratory Advanced Light Source at 100 K. The edge of the oscillation image corresponds to 1.76 Å resolution.

USA) and Jena Bioscience GmbH (Jena, Germany). Drops consisting of 1.5 µl protein solution mixed with 1.5 µl reservoir solution were equilibrated against 100 µl reservoir solution. Crystals were obtained under condition No. 18 of Wizard II from Emerald BioSystems [20.0% (w/v) polyethylene glycol 3000, 0.1 M Tris–HCl pH 7.0 containing 0.2 M calcium acetate]. Optimization of the initial crystal conditions were performed by microseeding for Arsm7A. Initial crystals of Arsm7A were transferred to an Eppendorf tube containing Seed Beads (Hampton Research) and 50 µl reservoir solution. The sample was vortexed for 90 s and three serial dilutions (1:10, 1:100 and 1:1000) were made before seeding. Drops were prepared by mixing 2.5 µl 15 mg ml⁻¹ Arsm7A solution with 2.5 µl seed stock consisting of 20% polyethylene glycol 3000, 0.2 M calcium acetate, 0.1 M Tris–HCl pH 7.0. Diffraction-quality crystals were obtained from 1:100 dilution drops. The Arsm7B construct produced suitable crystals without microseeding by mixing 2.5 µl 15 mg ml⁻¹ Arsm7B solution with 2.5 µl reservoir solution consisting of 18% polyethylene glycol 3350, 0.2 M calcium acetate and 0.1 M Tris–HCl pH 7.0. The hanging drops were equilibrated against 0.5 ml well solution and crystals were obtained at 293 K using Linbro 24-well plates (HR3-110) from Hampton Research.

2.3. Data collection

Crystals were transferred to a cryoprotectant solution (25% polyethylene glycol 3350, 0.2 M calcium acetate, 0.1 M Tris–HCl pH 7.0 and 10% glycerol) and flash-cooled in liquid nitrogen. A native data set was recorded on beamline 5.02 of the Lawrence Berkeley National Laboratory Advanced Light Source. Data were collected using an ADSC Quantum 315r (3 × 3 CCD array) detector at 100 K under a liquid-nitrogen stream. Data obtained from 143 images using a crystal-to-detector distance of 250 mm, 1° oscillation per image and an exposure time per image of 2 s were indexed, integrated and scaled using *HKL-2000* (Otwinowski & Minor, 1997).

Table 1
Data-collection and processing statistics.

Values in parentheses are for the highest resolution shell.

	Arsm7A	Arsm7B
Wavelength (Å)	0.9795	0.9795
Space group	C2	C2
Unit-cell parameters (Å, °)	$a = 84.85, b = 46.89,$ $c = 100.35, \beta = 114.25$	$a = 84.75, b = 47.20,$ $c = 101.68, \beta = 115.67$
Resolution range (Å)	45.78–1.76 (1.82–1.76)	50.78–1.78 (1.84–1.78)
No. of observations	99708	153634
No. of unique reflections	34898 (3092)	34951 (3450)
Completeness (%)	96.7 (86.2)	99.9 (99.5)
Redundancy	2.9 (2.7)	4.4 (3.5)
$\langle I \rangle / \langle \sigma(I) \rangle$	17.51 (2.47)	37.06 (5.83)
$R_{\text{merge}}^{\dagger}$ (%)	5.3 (35.2)	3.4 (19.1)

$\dagger R_{\text{merge}} = \frac{\sum_{hkl} \sum_i |I_i(hkl) - \langle I(hkl) \rangle|}{\sum_{hkl} \sum_i I_i(hkl)}$, where $I_i(hkl)$ is the intensity of the i th measurement of the reflection with Miller indices hkl and $\langle I(hkl) \rangle$ is the mean intensity of that reflection.

3. Results and discussion

Crystallization trials were attempted with native Arsm7, but this protein did not yield crystals. The active derivatives Arsm7A and Arsm7B were purified (Fig. 1) and diffraction-quality crystals were obtained with both. Initial crystals of Arsm7A diffracted to only 4 Å resolution, but microseeding using small crushed crystals of Arsm7A as seeds was more successful. Crystals appeared within 1 d and reached maximum dimensions of 0.4 × 0.3 × 0.2 mm within 2–3 d (Fig. 2). Crystals kept for longer than 3 d began to disintegrate in the drop, so newly obtained crystals were used for diffraction. Complete data sets were collected to resolutions of 1.76 and 1.78 Å using single crystals of Arsm7A (Fig. 3) and Arsm7B, respectively. The data-collection statistics for Arsm7A and Arsm7B are shown in Table 1. The asymmetric unit contains a single molecule of Arsm7A; the crystal volume per unit molecular weight, V_M , was calculated to be 2.22 Å³ Da⁻¹, corresponding to a solvent content of 44.7% (Matthews, 1968). No structure of a homologous protein is available for molecular-replacement phasing, so crystals with heavy metals or of selenomethionine-labeled protein will be obtained for structure determination.

This study was supported in part by US National Institutes of Health Grant GM55425. The Berkeley Center for Structural Biology is supported in part by the National Institutes of Health, National Institute of General Medical Sciences and the Howard Hughes Medical Institute. The Advanced Light Source is supported by the Director, Office of Science, Office of Basic Energy Sciences of the US Department of Energy under Contract No. DE-AC02-05CH11231.

References

- Challenger, F. (1951). *Adv. Enzymol. Relat. Subj. Biochem.* **12**, 429–491.
- Laemmli, U. K. (1970). *Nature (London)*, **227**, 680–685.
- Matthews, B. W. (1968). *J. Mol. Biol.* **33**, 491–497.
- Otwinowski, Z. & Minor, W. (1997). *Methods Enzymol.* **276**, 307–326.
- Qin, J., Lehr, C. R., Yuan, C., Le, X. C., McDermott, T. R. & Rosen, B. P. (2009). *Proc. Natl Acad. Sci. USA*, **106**, 5213–5217.
- Qin, J., Rosen, B. P., Zhang, Y., Wang, G., Franke, S. & Rensing, C. (2006). *Proc. Natl Acad. Sci. USA*, **103**, 2075–2080.
- Styblo, M., Drobna, Z., Jaspers, I., Lin, S. & Thomas, D. J. (2002). *Environ. Health Perspect.* **110**, Suppl. 5, 767–771.
- Thomas, D. J., Waters, S. B. & Styblo, M. (2004). *Toxicol. Appl. Pharmacol.* **198**, 319–326.