

Population genetic relationships between *Casearia sylvestris* (Salicaceae) varieties occurring sympatrically and allopatrically in different ecosystems in south-east Brazil

Marcelo Mattos Cavallari^{1,2,3,6,*}, Marcos Aparecido Gimenes^{1,4}, Claire Billot³, Roseli Buzanelli Torres⁵, Maria Imaculada Zucchi⁶, Alberto Jose Cavalheiro⁷ and Jean-Marc Bouvet²

¹Departamento de Genética, Instituto de Biociências, UNESP - Universidade Estadual Paulista 'Julio de Mesquita Filho' (UNESP), Distrito de Rubião Junior s/n, Botucatu, 18618-000, SP, Brazil, ²Unité Propre de Recherche de Génétique Forestière, CIRAD, Campus International de Baillarguet TA10/C, 34398 Montpellier Cedex 5, France, ³Mixed Unit of Research 'Plant Development and Adaptation', CIRAD, Avenue d'Agropolis TA A-96/03, F-34398 Montpellier Cedex 5, France, ⁴Embrapa Recursos Genéticos e Biotecnologia, Av. W5 Norte, CP 02372, Brasília, 70770-900, DF, Brazil, ⁵Núcleo de Pesquisa e Desenvolvimento do Jardim Botânico, ⁶Centro de Pesquisa e Desenvolvimento em Recursos Genéticos Vegetais, Instituto Agrônomo – IAC, Av. Barão de Itapura, 1481, Campinas, 13020-902, SP, Brasil and ⁷Instituto de Química, UNESP, Rua Prof. Francisco Degni s/n, 14800-900, Araraquara, SP, Brazil

*For correspondence. E-mail mmcavall@gmail.com

Received: 24 February 2009 Returned for revision: 3 November 2009 Accepted: 18 June 2010 Published electronically: 10 August 2010

- **Background and Aims** Species delimitation can be problematic, and recently diverged taxa are sometimes viewed as the extremes of a species' continuum in response to environmental conditions. Using population genetic approaches, this study assessed the relationship between two *Casearia sylvestris* (Salicaceae) varieties, which occur sympatrically and allopatrically in the landscape of south-east Brazil, where intermediate types are also found.
- **Methods** In total, 376 individuals from nine populations in four different ecosystems were sampled, and nine microsatellite markers were used to assess the relative effects of the ecosystems and varieties on the distribution of genetic diversity among populations of this species.
- **Key Results** As a by-product of this study, several PCR products with more than two alleles were observed. The possibility that extra bands represent non-specific amplification or PCR artefacts was discarded by sequencing a sample of these bands. We suggest that (partial) genome duplication in *C. sylvestris* most probably explains this phenomenon, which may be a key factor in the differentiation of the two taxa, as it was markedly more frequent in one of the varieties. AMOVA indicated that approx. 22 % of the total genetic diversity was found between the two varieties. Bayesian analysis identified varieties and ecosystems as evolutionary units, rather than the individual populations sampled.
- **Conclusions** The results are in agreement with field observations and support the recognition of two varieties, as well as documenting the occurrence of hybridization between them.

Key words: Atlantic Forest, *Casearia sylvestris*, Cerrado, ecotones, hybrid zone, microsatellites, population genetic structure, SSR, sympatry.

INTRODUCTION

Ecology is of fundamental importance in speciation (Orr and Smith, 1998; Schluter, 2001; Wiens, 2004; Gegeer and Burns, 2007), as incipient species occur in different ecological locations, and local adaptation leads to evolutionary divergence (Wiens, 2004). Such divergence may arise through the evolution of ecotypes (Abbott and Comes, 2007), which are distinct forms of a species adapted to different environmental conditions or habitats (Turesson, 1922). However, reproductive isolation between recently diverged taxa may be incomplete, hindering the delimitation of the boundaries among them (Rieseberg and Willis, 2007). In plants, phenotypic variation does not necessarily assort into discrete categories, and, for this reason, and due to interspecific hybridization, the definition of a species has been a major impediment to botanical studies of speciation (Rieseberg and Willis, 2007).

Casearia sylvestris Sw. (Salicaceae – APG II, 2003; Flacourtiaceae – Cronquist, 1981) is a shrub-to-tree species that is widespread in Central and South America, from Mexico to Argentina and Uruguay (Sleumer, 1980). It presents hermaphrodite flowers that are attractive to insects, mainly diptera, while its fruits are utilized by birds (Torres and Ramos, 2007). Two varieties, proposed by Sleumer (1980), are traditionally recognized: *C. sylvestris* var. *sylvestris*, which inhabits humid, dense forests, and *C. sylvestris* var. *lingua*, which is restricted to open, xeric habitats (Sleumer, 1980). The latter variety has also been recognized at the level of species [*Casearia lingua* (Cambess.) Eichler, Saint-Hilaire, 1829], and presents several traits related to drought and fire stress, including smaller, coriaceous leaves and shrubby habit (Silva, 1996). Despite differences in morphology and habitat preference, a gradation of intermediate forms can be found, making it

difficult to delimit the taxa (Sleumer, 1980; Torres and Yamamoto, 1986; Silva *et al.*, 2006). For this reason, in a recent review of Flacourtiaceae in São Paulo State, Brazil, Torres and Ramos (2007) did not recognize infraspecific taxa in *C. sylvestris*. Due to a lack of studies on the genetic structure of this species, whether the two taxa are independent biological units or represent the genetic continuum of a single species in response to different environmental conditions remains unresolved.

Molecular methods for assaying genetic diversity provide numerous tools to estimate relationships among natural populations of closely related organisms, including detecting species at shallow levels of evolutionary divergence (Drummond and Hamilton, 2007). In this context, microsatellite markers have proved to be powerful tools to solve biological problems, and are largely used to track the biological history of populations (Chambers and MacAvoy, 2000), providing a better understanding of species boundaries and of interspecific hybridization (e.g. Lexer *et al.*, 2005; Drummond and Hamilton, 2007).

The present study characterizes the genetic relationships among populations of two closely related taxa, *C. sylvestris* var. *sylvestris* and *C. sylvestris* var. *lingua*, which occur sympatrically and allopatrically in different ecosystems in south-east Brazil. Microsatellite markers are used to assess the relative effects of the ecosystems and varieties in the distribution of genetic diversity in this species, thus helping to clarify the genetic relationships between its varieties. We hypothesized that if the varieties are in fact divergent lineages rather than the continuum of a single taxon, distinct populations of the same variety would tend to be more closely related than sympatric populations of different varieties. In other words, geographical distribution of sampling sites would not be the major factor determining the distribution of genetic diversity.

MATERIALS AND METHODS

Sampling

Although *Casearia sylvestris* var. *sylvestris* is widespread throughout the neotropics, *C. sylvestris* var. *lingua* has a more restricted distribution and occurs in open areas in eastern and south-eastern South America, more specifically in the Brazilian Cerrado and Caatinga, usually in poor soils and at low altitudes (Sleumer, 1980). This study focuses on south-east Brazil, more specifically the State of São Paulo, which is a mosaic of ecosystems where the two varieties occur sympatrically and allopatrically.

We sampled nine locations (Table 1, Fig. 1) representing four different ecosystems [Atlantic Forest *sensu stricto* (*s.s.*), semi-deciduous Atlantic forest, Cerrado and ecotones). The Atlantic Forest *s.s.* is a dense, humid, evergreen forest that occurs in the mountains along the coast, while the semi-deciduous Atlantic Forest occurs in more continental areas and has a well-characterized dry season. The Cerrado is characterized by shrub savannas under fire regime occurring on acid, aluminium-rich soils, with a severe dry season (Ratter *et al.*, 1997). Contact zones (ecotones) between the semi-deciduous Atlantic Forest and the Cerrado are found throughout the landscape.

Sampling sites differed in size, isolation from other natural areas and conservation status (Table 1), with Carlos Botelho (Atlantic Forest *s.s.*) and Morro do Diabo (semi-deciduous Atlantic Forest) being the largest and least disturbed localities. Ilha Anchieta (Atlantic Forest *s.s.*) is an oceanic island separated from the mainland by about 500 m. The population in Botucatu (Cerrado) is particularly small and disturbed: until 1970 it was a cattle pasture, and since then natural regeneration of Cerrado has been taking place. Although the site covers 33 ha, *C. sylvestris* is only found in an area of approx. 1 ha.

TABLE 1. Sampling of *Casearia sylvestris*: ecosystems sampled, name of sampling sites and their characteristics, and number of individuals of each variety and of dubious morphology sampled in each location and ecosystem

Sampling site and ecosystem	Coordinates	Altitude (m)	Area (ha)	Dry season	Site characteristics	var. <i>sylvestris</i>	var. <i>lingua</i>	Dubious	Total
<i>Atlantic Forest s.s.</i>						122	–	–	122
Ilha Anchieta	23°32'S, 45°03'W	100	828	Absent	Oceanic island/moderate disturb	32	–	–	32
Santa Virginia	23°20'S, 45°08'W	900	17 000	Absent	Part of a larger continuous forest/low disturbance	39	–	–	39
Carlos Botelho	24°02'S, 47°57'W	800	37 000	Absent	Part of a larger continuous forest/low disturbance	51	–	–	51
<i>Semi-deciduous Atlantic Forest</i>						87	–	–	87
Morro do Diabo	22°30'S, 52°19'W	450	34 000	Moderate	Large fragment isolated/low disturbance	54	–	–	54
Caetetus	22°23'S, 49°42'W	600	2178	Moderate	Medium fragment isolated/medium disturbance	33	–	–	33
<i>Ecotones</i>						48	10	34	92
Mogi-Guaçu	22°15'S, 47°11'W	650	980	Moderate	Isolated/disturbed	16	6	25	47
Porto Ferreira	21°51'S, 47°25'W	560	612	Moderate	Isolated/disturbed	32	4	9	45
<i>Cerrado</i>						4	68	3	72
Assis	22°34'S, 50°24'W	550	1170	Severe	Isolated/disturbed	4	39	3	43
Botucatu	22°56'S, 48°27'W	850	33	Severe	Very small fragment isolated and disturbed	–	29	–	29
Total						261	78	37	376




FIG. 1. Map showing (approximately) the original ecosystem distribution in Sao Paulo state, south-east Brazil, and populations of *Casearia sylvestris* sampled. AS = Assis, BT = Botucatu, CB = Carlos Botelho, CT = Caetetus, IA = Ilha Anchieta, MD = Morro do Diabo, MG = Mogi-Guaçu, PF = Porto Ferreira, SV = Santa Virginia.

The average distance between sampling sites is 320 km, varying from 25 km (Ilha Anchieta – Santa Virginia) to 745 km (Ilha Anchieta – Morro do Diabo). The number of individuals sampled in each site is given in Table 1. Leaf samples were dried in sealed plastic bags containing silica gel and stored at -20°C prior to DNA extraction.

Classifying individuals at the level of variety

For each individual sampled, a voucher was deposited in the IAC (Instituto Agrônomo de Campinas) Herbarium, Campinas, Brazil. Morphological traits were used to assign the individuals to a variety according to Sleumer (1980): *var. lingua* presents more coriaceous, oval leaves, with prominent reticulation and a slender petiole. We also considered cymbiform leaves to be characteristic of *var. lingua*, based on the field and taxonomic experience of R.B.T. on the genus *Casearia*. Individuals were classified as *lingua*, *sylvestris* or dubious.

DNA isolation

DNA was isolated from leaf samples according to the acid protocol of Csaikl et al. (1998). We added 2% (w/v) sodium bisulfite to the extraction buffer to avoid oxidation, particularly observed in *var. sylvestris* samples.

Markers, PCR and electrophoresis conditions, and allele scoring

Ten microsatellite markers specifically designed for *C. sylvestris* (Cavallari et al., 2008) were tested. Locus Csy08 presented null alleles according to MICRO-CHECKER software (Van Oosterhout et al., 2004) and was

not retained for subsequent analysis. The remaining nine loci were polymorphic and were thus used in this study. PCR conditions followed Cavallari et al. (2008). The amplified fragments were analysed at 700 and 800 nm by electrophoresis on an IR-DNA analyser (LI-COR 4200 sequencer, www.licor.com) at the Montpellier Languedoc-Roussillon Genopole genotyping platform. Allele scoring was performed with the SAGA Generation 2 v.3.2 (LI-COR) software.

Verifying the microsatellite identity of the bands

We observed several PCR products with more than two alleles (see Results), which is not expected for microsatellite markers in diploid species (the ploidy of *C. sylvestris* is not known). A new set of PCR reactions was performed with 90 randomly chosen individuals (for all primer pairs), and the same results were obtained. To follow up, we randomly chose three individuals and two primer pairs (Csy04 and Csy09) to perform a more detailed study. For this study, PCR products were run in a 3.8% agarose gel and detected bands were excised from the gel under UV light. Gel slices containing the bands were passed through a purification column (Wisard SV Gel and PCR Clean-Up System, Promega) and the purified products were ligated into pGEM-T Easy vector (Promega, www.promega.com) and used to transform DH5 α competent cells. Five positive colonies (using blue/white β -galactosidase selection) from each Petri dish (i.e. representing each of the bands detected in the gel) were sequenced using the SP6 primer and BigDye terminator version 3.1 Cycle Sequencing Kit (Applied Biosystems, www.appliedbiosystems.com) and run on a 3100 DNA Analyser (Applied Biosystems). The sequences obtained were examined for the presence of the microsatellite repeats and their flanking regions.

Statistical analysis

Genetic diversity. Due to the detection of several PCR products with more than two alleles (see Results), correct allele frequencies could not be calculated, and most of the analyses that are usually performed with co-dominant markers could not be performed (H_E , H_O , F_{IS} , etc.). Descriptive analyses of genetic diversity were thus restricted to those that were completely independent of allele frequencies, i.e. by estimating the number of alleles per locus and allele richness (El Mousadik and Petit, 1996). This enabled us to describe and compare the genetic diversity of sampling sites, ecosystems and varieties without making assumptions on the ploidy level of the species. Allele richness in particular has been considered to be one of the most relevant criteria for measuring diversity, especially in the context of genetic conservation (reviewed by Petit *et al.*, 1998). To perform these analyses, PCR products with more than two alleles (which correspond to only 3.4 % of the total PCR products – see Results) were removed from the data set; this did not distort the results as all the alleles that were excluded were present in other individuals of the same population (data not shown). Analyses were performed using FSTAT software (Goudet, 1995). Sampling sites, ecosystems and varieties were compared to get an idea of their relative genetic diversity. For each variety, we also determined the size range of alleles, the number of private alleles and the frequency of the most common allele. To this end, all morphologically dubious individuals were excluded.

In addition, we calculated Shannon's information index to estimate the degree of genetic diversity within each sampling site and ecosystem using POPGENE software v.1.31 (Yeh *et al.*, 1997). This was done by coding alleles as dominant marker bands (binary coding), considering all PCR products, including those with more than two alleles.

Population genetic structure. The usual F -statistics performed with co-dominant markers could not be carried out due to the lack of allele frequencies. Based on the presence/absence of bands (binary coding, considering all PCR products), population genetic structure was inferred by analysis of molecular variance (AMOVA) with ARLEQUIN v.2.0 software (Schneider *et al.*, 2000), estimating Φ -statistics (Excoffier *et al.*, 1992), which are analogous to Wright's hierarchical fixation indices under the island model of gene flow (Wright, 1951). However, it should be borne in mind that the alleles from a given microsatellite locus are not independent samples and, for this reason, P values obtained in the AMOVAs may present some bias. Two independent AMOVAs were performed, which hierarchically partitioned genetic diversity by (1) ecosystems/sampling sites/individuals and (2) varieties/sampling sites/individuals.

Population genetic structure was also revealed using the Bayesian approach of Pritchard *et al.* (2000), implemented in the software STRUCTURE. The latest versions (2.2.3 and 2.3.2) of the software include the algorithm of Falush *et al.* (2007), which can handle genotypic ambiguity (where it is not possible to identify the exact genotypes in heterozygotes, e.g. tetraploid individuals displaying two or three alleles). As we observed individuals with up to four bands in some loci, STRUCTURE was run considering all the individuals as tetraploids. Following the user's guide for the software, the four

rows of each individual at each locus (representing the four alleles) were filled up with the alleles observed (one, two, three or four alleles). The integer for missing data was used to complete the infile, and the option RECESSIVEALLELES was set to 1. The program thus considers that the four alleles are co-dominant to each other, and that there is genotypic ambiguity. In this way, we incorporated all alleles observed in the sample in the study of population genetic structure.

STRUCTURE 2.3.2 was used to check the ancestry of individuals within populations, ecosystems and varieties. The optimal number of clusters (K) was determined by varying the value of K from 1 to 10, and five runs of each K value were performed without any prior information using the admixture model and correlated alleles frequencies, with 200 000 generations sampled as burn-in and 500 000 generations sampled in the Monte Carlo Markov chain (MCMC). The *ad hoc* statistics ΔK (Evanno *et al.*, 2005) and $\text{Pr}(X|K)$, given by $\ln P(D)$ (Pritchard *et al.*, 2000), were used as predictors of the real number of clusters. At $K = 2$, the clusters that formed agreed closely with the morphological classification of the individuals (see Results), and, for this reason, it was possible to use the admixture proportions (Q) of each individual to reveal admixed individuals between the two varieties and to give an idea of the extent of hybridization in each population. This was possible because the admixture model estimates the proportion of each individual's genome that originated from each of the K inferred clusters (Pritchard *et al.*, 2000). Although the threshold adopted here is somewhat arbitrary, Vähä and Primmer (2006) and Lepais *et al.* (2009) demonstrated through simulation studies that a threshold of 10 % is probably the best choice to correctly classify purebreds and hybrids in such situations. Following these authors, we considered as admixed between the two varieties all individuals presenting a Q value of between 0.1 and 0.9. The STRUCTURE results were displayed graphically by the software DISTRUCT (Rosenberg, 2004).

RESULTS

Classification of individuals into varieties and their distribution among ecosystems

For the majority of individuals sampled, classification at the variety level was not problematic. Key morphological features were generally clear, leading to immediate recognition of the variety. As a general rule, all individuals sampled in the Cerrado were easily recognized as var. *lingua*, and all individuals sampled in the Atlantic Forest (both *s.s.* and semi-deciduous) were undoubtedly var. *sylvestris*. We found the two varieties occurring sympatrically in the ecotones, where individuals of dubious morphology were also observed. The population of Assis (Cerrado) also included the two varieties and intermediate morphs, with var. *lingua* restricted to Cerrado *s.s.* (shrub savannas) areas with var. *sylvestris* and the intermediate morphs restricted to 'Cerradão' areas (dense arrangements of Cerrado trees). The number of individuals of each variety sampled in each ecosystem/sampling site is given in Table 1.

Detection of PCR products with more than two alleles and study of band identity

Several PCR products with more than two alleles were detected (see below). Sequencing of some of these PCR products (three individuals each for primer pair Csy04 and Csy09) revealed that all bands contained the expected microsatellite repeats (CT_n for Csy04 and AG_n for Csy09; Cavallari *et al.*, 2008) located between the same flanking regions. This result shows that the additional bands observed are true microsatellite alleles, and not PCR artefacts.

PCR products with more than two alleles were observed in all but one locus (Csy18), with locus Csy04 being the most affected (12 % of PCR products). Considering all loci, multiple alleles were observed in 3.4 % of the PCR products, and detected in 83 (22 %) of the 376 individuals analysed. In most of them (64 individuals), the phenomenon was restricted to only one locus, although some individuals had more than two alleles at two, three or four loci. We detected many more cases of PCR products with three alleles (106) than with four alleles (only four). Of the 78 var. *lingua* individuals genotyped, 53 (68 %) presented more than two alleles in at least one locus. PCR products with more than two alleles were observed in only 7.7 % (20 of 261 individuals) of var. *sylvestris* samples and in 27 % (ten of 37 individuals) of the morphologically dubious individuals. Possible explanations for the existence of more than two alleles per locus are provided in the Discussion.

Genetic diversity

The number of alleles sampled and the allele richness of each sampling site, ecosystem and variety are presented in Table 2. The mean number of alleles per locus per sampling site was 8.87, ranging from 3.1 (Botucatu) to 12.2 (Porto Ferreira). Botucatu displayed a very particular genetic composition: in addition to the extremely low number of alleles per locus (ranging from one to six alleles, whereas in other sampling sites the number ranged from three to 21) and low total allele richness (19.89, whereas the mean total allele richness was 37.89), in six of the nine loci no homozygotes were observed. In the remaining sampling sites, homozygotes were observed at all loci.

The remaining sampling sites displayed very similar patterns of diversity (Table 2), with the mean number of alleles per locus per sampling site ranging from 8.88 (Mogi-Guaçu, Ilhan Anchieta) to 12.2 (Porto Ferreira), and mean allele richness per locus ranging from 4.12 (Santa Virginia) to 5.15 (Porto Ferreira).

As a whole, the ecotones presented the highest number of alleles per locus and the highest allele richness. When the minor variety (var. *lingua*) in these sites was excluded from the data set, these numbers decreased and Atlantic Forest *s.s.* and semi-deciduous Atlantic Forest appear as the most genetically diverse ecosystems (Table 2).

The allele richness observed in each variety differed significantly (Table 2). Of the total number of alleles detected, 64 (42.1 %) were unique to var. *sylvestris* and eight (5.25 %) were unique to var. *lingua* (Table 3), implying that only 52.6 % of the bands are shared by the two varieties (dubious

TABLE 2. Mean number of alleles sampled per locus, total number of alleles sampled, mean allele richness per locus and total allele richness of each ecosystem, sampling site and variety studied

Ecosystems/sampling sites/varieties	Number of alleles sampled		Allele richness*	
	Mean	Total	Mean	Total
<i>Atlantic Forest s.s.</i>	12.56	113	9.97	89.77
Ilha Anchieta	8.89	80	4.33	38.94
Santa Virginia	9.33	84	4.12	37.05
Carlos Botelho	9.00	81	4.19	37.71
<i>Semi-deciduous Atlantic Forest</i>	11.67	105	9.95	89.52
Morro do Diabo	10.00	90	4.59	41.34
Caetetus	9.44	85	4.67	42.01
<i>Cerrado (combined)</i>	9.56	86	8.06	72.56
Assis (combined)	9.00	81	4.55	40.97
Botucatu	3.11	28	2.21	19.89
<i>Cerrado (only major variety)</i>	8.67	78	7.53	67.80
Assis (only major variety)	8.11	73	4.26	38.31
Botucatu	3.11	28	2.21	19.89
<i>Ecotones (combined)</i>	13.22	119	10.48	94.32
Mogi-Guaçu (combined)	8.89	80	4.08	36.76
Porto Ferreira (combined)	12.22	110	5.15	46.35
<i>Ecotones (only major variety)</i>	10.67	96	9.84	88.53
Mogi-Guaçu (only major variety)	5.00	45	3.51	31.55
Porto Ferreira (only major variety)	9.78	88	4.83	43.46
var. <i>sylvestris</i>	15.56	140	11.36	102.21
var. <i>lingua</i>	9.44	85	7.76	69.82
unknown var.	10.11	91	9.73	87.60

* Allele richness for sampling sites based on minimum sample size of five diploid individuals; for ecosystems based on minimum sample size of 33 diploid individuals; and for varieties based on minimum sample size of 30 diploid individuals.

individuals were not taken into consideration). The size range of the alleles, the most common allele and its frequency, and the number of private alleles for each locus and each variety are shown in Table 3. The varieties differed significantly in the presence and frequency of alleles, especially at loci Csy04, Csy06, Csy11, Csy14 and Csy18 (Table 3).

Values obtained for Shannon's information index (data not shown) were similar for all sampling sites, ranging from 0.16 (Santa Virginia) to 0.21 (Porto Ferreira). The only discrepancy was observed for Botucatu (0.05). The total diversity observed in all populations was 0.23. Ecosystems and varieties showed similar Shannon's information index values ranging from 0.17 (Cerrado) to 0.20 (ecotones). Varieties presented similar levels of genetic diversity (Shannon's information index approx. 0.21 in each variety).

Population genetic structure

The first hierarchical AMOVA (Table 4) revealed that approx. 13 % of the total molecular genetic diversity was attributable to differences among ecosystems. The second AMOVA revealed that differences between the varieties accounted for approx. 22 % of the total genetic diversity sampled. Estimated Φ -statistics were shown to be significant at the 1 % level ($P < 0.01$) in both AMOVAs, but this result should be interpreted with caution due to the lack of independence between alleles of the same locus. In both AMOVAs, the

TABLE 3. Comparison between varieties of *C. sylvestris*: size range of alleles, most common allele and number of private alleles detected in each locus studied

Locus	Range size of alleles			Most common allele*		Number of private alleles	
	var. <i>sylvestris</i>	var. <i>lingua</i>	Total	var. <i>sylvestris</i>	var. <i>lingua</i>	var. <i>sylvestris</i>	var. <i>lingua</i>
Csy_04	115–167	113–159	113–167	145 (0.49)	115 (0.28)	9	5
Csy_06	276–322	276–318	276–326	280 (0.13)	280 (0.60)	14	0
Csy_07	246–276	246–276	244–276	246 (0.45)	246 (0.60)	6	0
Csy_09	185–209	191–202	185–209	191 (0.60)	199 (0.47)	5	0
Csy_11	140–182	142–182	140–186	162 (0.23)	172 (0.31)	3	2
Csy_14	218–244	218–242	218–244	230 (0.66)	226 (0.37)	6	0
Csy_15	251–285	251–287	251–287	271 (0.24)	263 (0.22)	7	1
Csy_16	272–286	276–282	270–286	278 (0.67)	278 (0.55)	3	0
Csy_18	271–313	273–297	271–313	287 (0.26)	273 (0.90)	11	0
Total						64	8

*Allele frequency in parentheses.

TABLE 4. Results of AMOVA

Source of variation	d.f.	SSD	Percentage total variance	Φ -statistics
AMOVA ecosystems				
Ecosystems	3	492.505	0.1281	$\Phi_{CT} = 0.13$
Sampling sites within ecosystems	5	251.099	0.1146	$\Phi_{SC} = 0.11$
Individuals within sampling sites	367	2587.465	0.7573	$1 - \Phi_{ST} = 0.76$
Total	375	3331.069		
AMOVA varieties				
Varieties	1	331.316	0.2199	$\Phi_{CT} = 0.22$
Sampling sites within varieties	10	455.916	0.1385	$\Phi_{SC} = 0.14$
Individuals within sampling sites	327	2197.718	0.6416	$1 - \Phi_{ST} = 0.64$
Total	338	2984.95		

d.f., degrees of freedom; SSD, sum of squared deviations.

greater part of the genetic diversity sampled was within populations, but diversity was greater when individuals of the same variety were grouped together in the same sampling sites (first AMOVA, 76 %) than when they were separated into different populations (second AMOVA, 64 %).

Bayesian analysis performed using STRUCTURE with K from 1 to 10 provided either two or three clusters ($K = 2$ and $K = 3$; Fig. 2). Figure 3 shows clustering results for $K = 2$ and $K = 3$. At $K = 2$ (Fig. 3A), clustering closely matched our morphological classification. When the threshold of 10 % was used, individuals with an admixture coefficient (Q) below 0.1 or above 0.9 can be considered as not admixed (or as ‘pure grey’ and ‘pure black’, respectively, in Fig. 3A). Of the 69 ‘pure black’ individuals, 65 (94.2 %) were morphologically classified as var. *lingua* (underlined by a black bar in Fig. 3). In the same way, of the 287 ‘pure grey’ individuals, 261 (91 %) were morphologically classified as var. *sylvestris* (underlined by a grey bar in Fig. 3). There were no individuals with var. *lingua* morphology in the ‘grey’ cluster, and vice versa. Individuals of intermediate morphology (underlined by a white bar in Fig. 3) were revealed to have different degrees of admixture, ranging from ‘pure black’ to ‘pure grey’.

Once we confirmed that this clustering agreed closely with our morphological classification, the admixture coefficient for each individual (Q) was used to reveal putatively admixed individuals between the two varieties. Thirty of 376 individuals (7.9 %) presented a value of Q of between 0.1 and 0.9, and can be considered as admixed between the two varieties. Only seven putatively admixed individuals out of 238 (i.e. 2.9 %) were found in the populations of the Atlantic Forest, semi-deciduous Forest and Cerrado (only Botucatu). In the mixed populations (Assis, Mogi-Guaçu and Porto Ferreira), 23 of 138 individuals (16.6 %) can be considered as putatively admixed between the two varieties, and this proportion increased to 19.5 % (18 of 92) when only the ecotones (Mogi-Guaçu and Porto Ferreira) were taken in account.

At $K = 3$, clustering agreed closely with the ecosystems sampled (Fig. 3B), showing the ability of the analysis to detect biologically meaningful clusters.

DISCUSSION

Genome duplication in *C. sylvestris*

The unusual microsatellite banding pattern obtained suggests the occurrence of genome duplication in *C. sylvestris*, although it was not possible to distinguish between chromosome duplication (polyploidy, aneuploidy) and small-scale duplication (only some genes duplicated) with the tools used.

We observed individuals with up to four alleles in three loci, and with up to three alleles in eight of the nine loci studied. Moreover, in Botucatu, which had a low number of alleles in all loci, six of the nine loci showed fixed heterozygous profiles. This pattern is similar to that obtained by Segarra-Moragues *et al.* (2004) in *Borderea pyrinaica* (Dioscoreaceae): they observed individuals with up to four bands in 11 of 17 microsatellite loci they screened, whereas two loci showed fixed heterozygous profiles. They interpreted the fixed heterozygous profile of these loci as the amplification of two duplicate loci that are fixed for one allele each, and based on overall patterns observed they suggest that this species is allotetraploid.

One of the obvious approaches to solve this question is cytogenetics, which will be done in the near future. Cytogenetic




FIG. 2. Plots of $\ln P(D)$ (Pritchard *et al.*, 2000) and ΔK (Evanno *et al.*, 2005) for each K obtained from the STRUCTURE analysis.




FIG. 3. Graphical representation of results of STRUCTURE obtained for two (A) and three (B) clusters (K). Each individual is represented by a vertical line divided into K coloured fragments proportional to its membership in the corresponding genetic cluster. Q denotes the proportion of admixture for each individual and the threshold of 10 % is represented by a dashed white line in (A). Dashed black lines separate individuals from different populations and ecosystems, which are indicated at the top. The thin bar under the figures indicates the morphological classification of individuals (black = var. *lingua* morphology, grey = var. *sylvestris* morphology, white = individuals with dubious morphology).

studies on *Casearia* are very rare in the literature. The *Index to Plant Chromosome Numbers* (available at <http://mobot.mobot.org/W3T/Search/ipcn.html>), which contains data from published indices from 1979 onward, lists only one publication on *Casearia* chromosome numbers ($n = 21$ for *C. elliptica* Tul.; Bir *et al.*, 1980). However, in Salicaceae, aneuploids have been reported in *Populus* (Bradshaw and Stettler, 1993; Cervera *et al.*, 2001; Yin *et al.*, 2004), while *Salix* species present various ploidy levels (Leskinen and Alström-Rapaport, 1999).

Molecular and genetic studies in plants have shown that autopolyploidy is much more common than traditionally assumed (Soltis *et al.*, 2003). A large number of angiosperm families, including Salicaceae, which were traditionally considered to be diploids, are probably the products of ancient

polyploid events (Soltis and Soltis, 2000). These ancient polyploids exhibit extra copies of some of their genes above the level that one would expect for diploid plants (Gottlieb, 1982). For example, *Populus trichocarpa* has 8000 duplicated genes (Tuskan *et al.*, 2006). The complete sequencing of the *P. trichocarpa* genome and the comparison of *Populus* and *Salix* orthologous genes revealed that the common ancestor of these two genera was polyploid (Tuskan *et al.*, 2006). Thus, we may find that *C. sylvestris*, as a member of Salicaceae, shares this history and also presents duplicated genes.

Duplicated genes and genomes can undergo divergent evolution and evolve new functions (Soltis and Soltis, 2000; Cui *et al.*, 2006), facilitating the colonization of unstable habitats (Lawton-Rauh, 2003). It is important to note that in the

present study, the great majority of individuals with gene (or chromosome) duplication were found in the Cerrado (var. *lingua*), an unstable, harsh environment under fire regime.

Genetic diversity

The primer pairs utilized in this work were developed from *C. sylvestris* var. *sylvestris* and tested for transferability for other *Casearia* species, with no positive results (Cavallari et al., 2008). In the present work, these markers successfully amplified the DNA of the two varieties of *C. sylvestris*, but a high percentage of private alleles (47.35 %) was observed, and the most common allele and its frequency differed markedly between the two varieties at most loci. When studying the boundaries between two sympatric varieties of *Lupinus microcarpus* (Leguminosae), Drummond and Hamilton (2007) observed a similar pattern of allele frequency distribution, revealing a mix of shared and private alleles. According to these authors, this result is consistent with shared ancestral polymorphism and recent divergence between species.

The low genetic diversity observed in Botucatu is congruent with the recent (after 1970) foundation of the population. The distribution of individuals (restricted to 1 ha) also suggests that few founders arrived in the locality (which has a total area of 33 ha). Although fragmentation and a reduction in population size are expected to decrease gene diversity (Young et al., 1996), apart from Botucatu, observed gene diversity was not positively correlated with fragment size or conservation status. Sampling sites harbouring the two varieties, although relatively isolated and disturbed, presented high values of Shannon's information index and allele richness. The ecotones as a whole had the highest values of Shannon's index and allele richness, higher than the main values observed for ecosystems. Our sampling strategy (two varieties occurring sympatrically and allopatrically) and the high percentage of bands restricted to one or the other variety may explain these trends. Indeed, excluding from the data set the minor variety occurring in these sampling sites/ecosystem resulted in decreased diversity (although this was not the case of Porto Ferreira, which even without considering the minor variety still displayed the highest genetic diversity). These results are in accordance with recent research on a wide range of taxa, which suggests that environmental gradients are important in diversification and speciation, and may deserve special attention (Smith et al., 2001).

Population genetic structure: main components

The population structure of *C. sylvestris* in this region can be described in terms of 'varieties' and 'ecosystems', which were shown to be significant factors in the distribution of genetic diversity. This was predicted from field observations and corroborated by statistical analyses, especially by the Bayesian analysis.

Clustering by STRUCTURE at $K = 2$ (Fig. 3A) and at $K = 3$ (Fig. 3B) grouped the individuals in a way that is highly concordant with varieties and ecosystems, respectively. The agreement between clusters and varieties indicates a non-random distribution of alleles and allele frequencies between them, the same applying to the ecosystems. As

ΔK (Fig. 2) provided poor statistical support for $K > 3$, the subdivision of the species' genetic diversity into varieties and ecosystems was very strong, with each variety or ecosystem recognized as a discrete unit, although this was not the case for individual sampling sites.

According to Minder and Widmer (2008), 'good' species are expected to be genetically distinct from each other, with geographically distant populations of the same species being genetically more closely related to each other than to geographically proximate populations of the other species. This pattern was clearly supported by our Bayesian analysis. In the Bayesian analysis, the clusters obtained agreed closely with our morphological classification. Thus, in addition to being morphologically recognizable, the two varieties also possess significant differences at neutral (microsatellite) loci, suggesting little connectivity between the two taxa. Other results that also support divergence between them are the marked differences in the proportion of individuals with evidence of partial genome duplication in each variety, the high proportion of private alleles in each variety (47.4 %) and results of the AMOVA showing that approx. 22 % of the total genetic diversity sampled is attributable to genetic differences between varieties. The results of the AMOVA are similar to those observed between *Croton alabamensis* varieties separated by more than 1000 km (Van Ee et al., 2006) and between two species of *Iliamna* (Malvaceae) (Slotta and Porter, 2006), and, although there is no absolute measure of genetic differentiation for determining species separation (Morgan-Richards and Wolff, 1999), our results indicate strong differentiation between the two taxa.

The partitioning of the genetic diversity of the species according to the ecosystem was supported by the clustering at $K = 3$ (Fig. 3B). Individuals of var. *sylvestris* were separated into two clusters: one composed of individuals from the Atlantic Forest *s.s.*, and the other of individuals from the remaining ecosystems. This is reasonable as the semi-deciduous Atlantic Forest and the Cerrado are connected by ecotones, allowing more connectivity between its individuals than between individuals from the Atlantic Forest *s.s.*, which is more isolated. Genetic divergence between Atlantic Forest *s.s.* and the semi-deciduous Atlantic Forest (both harbouring exclusively var. *sylvestris*) may be explained by genetic drift, as they are all isolated fragments.

Putative hybrid zones

The literature on *C. sylvestris* refers to the existence of morphologically intermediate individuals (Sleumer, 1980; Torres and Ramos, 2007) without referring to their habitat preference or co-occurrence with 'pure forms'. We observed that intermediate forms occur exclusively in the presence of both of the 'pure' varieties. The sympatric occurrence of the three forms (var. *sylvestris*, var. *lingua* and intermediate forms) suggests that morphologically intermediate individuals are the product of hybridization, rather than of phenotypic responses to environmental intermediacy. For instance, we visited several open Cerrado-like areas within the Atlantic Forest, where although the trees had a somewhat different morphology due to full exposure to the sun, they could undoubtedly be classified as pure var. *sylvestris* individuals.

Of course, to disentangle phenotypic plasticity and genetic divergence leading to divergent morphology, other approaches should also be used (e.g. reciprocal transplants, quantitative genetic analyses of morphological and adaptive traits to separate genetic effects from environmental effects), and in this sense, our results are preliminary.

In addition, the literature on *C. sylvestris* refers to the existence of two varieties inhabiting different environments (var. *sylvestris* in dense forests and var. *lingua* in open areas) (Sleumer, 1980; Silva *et al.*, 2006), and, to our knowledge, sympatry of these forms has never been directly reported. Our fieldwork allowed us to observe sympatry of the two varieties: in the ecotones, individuals of each variety were found side by side.

As expected, populations harbouring the two varieties contained a greater number of admixed individuals displaying a wide range of admixture proportions (revealed by Q values), suggesting the coexistence of many hybrid generations and backcrosses. According to these observations, these regions can be considered putative hybrid zones, a pattern extensively reported in the literature at the species level (e.g. Fritsche and Kaltz, 2000; Lexer *et al.*, 2005; Valbuena-Carabaña *et al.*, 2005; Remington and Robichaux, 2007).

The study of plant species hybrid zones in the Cerrado/Atlantic Forest geographical region is in its infancy. A putative hybrid zone was studied by Lacerda *et al.* (2002) and Goulart *et al.* (2005), in ecotonal regions where the Fabaceae species *Plathymenia reticulata* (from Cerrado) and *P. foliolosa* (from the Atlantic Forest) meet. These authors evoked the 'refugium theory' (Haffer, 1969) to explain the origin of the two species: changes in vegetation cover and in the distribution of plant species during climatic changes in the Pleistocene would have restricted some populations to dry areas, which may have evolved into *P. reticulata*, whereas populations of warmer and humid refuges evolved into *P. foliolosa*. This hypothesis may be applicable to *C. sylvestris* populations in the same region, leading to the differentiation between populations from humid refuges (which may have evolved into var. *sylvestris*) and xeric areas (which may have evolved into var. *lingua*). It is reasonable to propose that the ecological differences between these habitats (e.g. different pollinators, habitat preference of pollinators and seed dispersers, different pluviometric regimes leading to flowering asynchrony) today represent strong barriers to gene exchange between varieties, thus helping to maintain separation of the taxa.

It would be worth checking if there are preferential crossing patterns or partial reproductive barriers between the two taxa in the ecotones. In plant species, many closely related taxa differ in flowering time to minimize niche competition (Friberg *et al.*, 2008). During sampling, we observed that pure parental forms persist sympatrically with intermediate forms. Our field observations indicate the absence of flowering synchrony between the two varieties (in Assis, Mogi-Guaçu and Porto Ferreira, we noted that only individuals of var. *lingua* were flowering), but there is no statistical robustness in this observation, and more detailed studies are necessary.

In summary, our field observations and experimental results suggest that the two *C. sylvestris* varieties are relatively independent biological units, rather than representing the genetic continuum of a single species in response to different

environmental conditions. We also documented the existence of putative hybrid zones between these two varieties. To corroborate these findings, additional studies with chloroplast DNA microsatellites, cpDNA sequences and rDNA sequences will be performed in the near future. It is important to stress that our sampling was restricted to the geographical region of south-east Brazil, which harbours a particular assortment of ecosystems. It is possible that the genetic relationships between the two varieties of *C. sylvestris* differ in central northern Brazil, where they also may occur sympatrically.

ACKNOWLEDGEMENTS

We thank the staff of the Conservation Units visited, especially Dr Giselda Durigan (Assis) and Dr Léo Zimback (Botucatu), Instituto Florestal de São Paulo, for the information they provided on site history and distribution of *C. sylvestris*, Eduardo Moretti for help with fieldwork, Dr Mateus Mondin (ESALQ/USP) for critical review of part of the manuscript and Dr Aluana G. Abreu (IAC) for insightful discussion. This work was partially supported by the Fundação de Amparo à Pesquisa do Estado de São Paulo (FAPESP). M.M.C. received a PhD fellowship from the Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq) and from the Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES). Part of this work was carried out using the resources of the CBSU from Cornell University, which is partially funded by Microsoft Corporation.

LITERATURE CITED

- Abbott RJ, Comes HP. 2007. Blowin' in the wind – the transition from ecotype to species. *New Phytologist* **175**: 197–200.
- APG II. 2003. An update of the Angiosperm Phylogeny Group classification for the orders and families of flowering plants: APG II. *Botanical Journal of the Linnean Society* **141**: 399–436.
- Bir SS, Gill BS, Bedi YS, Singhal VK. 1980. Evolutionary status of the woody taxa of Garhwal Himalaya. In: Khosla PK, ed. *Improvement of forest biomass*. Solan: Society of Tree Scientists, 81–96.
- Bradshaw HD, Stettler RF. 1993. Molecular genetics of growth and development in *Populus*. I. Triploidy in hybrid poplars. *Theoretical and Applied Genetics* **86**: 301–307.
- Cavallari MM, Billot C, Bouvet J-M *et al.* 2008. Isolation and characterization of microsatellite markers for *Casearia sylvestris* Sw. (Salicaceae), a neotropical medicinal tree. *Molecular Ecology Resources* **8**: 802–804.
- Cervera M-T, Storme V, Ivens B *et al.* 2001. Dense genetic linkage maps of three *Populus* species. *Populus deltoides*, *P. nigra* and *P. trichocarpa* based on AFLP and microsatellite markers. *Genetics* **158**: 787–809.
- Chambers GK, MacAvoy ES. 2000. Microsatellites: consensus and controversy. *Comparative Biochemistry and Physiology* **126**: 455–476.
- Cronquist A. 1981. *An integrated system of classification of flowering plants*. New York: Columbia University Press.
- Csaikl UM, Bastian H, Brettschneider R *et al.* 1998. Comparative analysis of different DNA extraction protocols: a fast, universal maxi-preparation of high quality plant DNA for genetic evaluation and phylogenetic studies. *Plant Molecular Biology Reporter* **16**: 69–86.
- Cui L, Wall PK, Leebens-Mack JH *et al.* 2006. Widespread genome duplications throughout the history of flowering plants. *Genome Research* **16**: 738–749.
- Drummond CS, Hamilton MB. 2007. Hierarchical components of genetic variation at a species boundary: population structure in two sympatric varieties of *Lupinus microcarpus*. (Leguminosae). *Molecular Ecology* **16**: 753–769.
- El Mousadik A, Petit RJ. 1996. High level of genetic differentiation for allelic richness among populations of the argan tree [*Argania*

- spinosa* (L.) Skeels] endemic to Morocco. *Theoretical and Applied Genetics* **92**: 832–839.
- Evanno G, Regnaut S, Goudet J. 2005.** Detecting the number of clusters of individuals using the software structure: a simulation study. *Molecular Ecology* **14**: 2611–2620.
- Excoffier L, Smouse PE, Quattro JM. 1992.** Analysis of molecular variance inferred from metric distances among DNA haplotypes – application to human mitochondrial – DNA restriction data. *Genetics* **131**: 479–491.
- Falush D, Stephens M, Pritchard JK. 2007.** Inference of population structure using multilocus genotype data: dominant markers and null alleles. *Molecular Ecology Notes* **7**: 574–578.
- Friberg M, Bergman M, Kullberg J, Wahlberg N, Wiklund C. 2008.** Niche separation in space and time between two sympatric sister species—a case of ecological pleiotropy. *Evolutionary Ecology* **22**: 1–18.
- Fritsche F, Kaltz O. 2000.** Is the *Prunella* (Lamiaceae) hybrid zone structured by an environmental gradient? Evidence from a reciprocal transplant experiment. *American Journal of Botany* **87**: 995–1003.
- Gegear RJ, Burns JG. 2007.** The birds, the bees, and the virtual flowers: can pollinator behavior drive ecological speciation in flowering plants? *American Naturalist* **170**: 551–556.
- Gottlieb LD. 1982.** Conservation and duplication of isozymes in plants. *Science* **216**: 373–380.
- Goudet J. 1995.** FSTAT version 1.2: a computer program to calculate F-statistics. *Journal of Heredity* **86**: 485–486.
- Goulart MF, Lemos Filho JP, Lovato MB. 2005.** Phenological variation within and among populations of *Plathymania reticulata* in Brazilian Cerrado, the Atlantic Forest and transitional sites. *Annals of Botany* **96**: 445–455.
- Haffer J. 1969.** Speciation in Amazonian forest birds. *Science* **165**: 131–137.
- Lacerda DR, Lemos Filho JP, Acedo MDP, Lovato MB. 2002.** Molecular differentiation of two vicariant neotropical tree species, *Plathymania foliolosa* and *P. reticulata*. (Mimosoideae), inferred using RAPD markers. *Plant Systematics and Evolution* **235**: 67–77.
- Lawton-Rauh A. 2003.** Evolutionary dynamics of duplicated genes in plants. *Molecular Phylogenetics and Evolution* **29**: 396–409.
- Lepais O, Petit RJ, Guichoux E et al. 2009.** Species relative abundance and direction of introgression in oaks. *Molecular Ecology* **18**: 2228–2242.
- Leskinen E, Alström-Rapaport C. 1999.** Molecular phylogeny of Salicaceae and closely related Flacourtiaceae: evidence from 5-8S, ITS1, and ITS2 of the rDNA. *Plant Systematics and Evolution* **215**: 209–227.
- Lexer C, Fay MF, Joseph JA, Nica MS, Heinze B. 2005.** Barrier to gene flow between two ecologically divergent *Populus* species, *P. alba* (white poplar) and *P. tremula* (European aspen): the role of ecology and life history in gene introgression. *Molecular Ecology* **14**: 1045–1057.
- Minder AM, Widmer A. 2008.** A population genomic analysis of species boundaries: neutral processes, adaptive divergence and introgression between two hybridizing plant species. *Molecular Ecology* **17**: 1552–1563.
- Morgan-Richards M, Wolff K. 1999.** Genetic structure and differentiation of *Plantago major* reveals a pair of sympatric sister species. *Molecular Ecology* **8**: 1027–1036.
- Orr MR, Smith TB. 1998.** Ecology and speciation. *Trends in Ecology and Evolution* **13**: 502–506.
- Petit RJ, El Mousadik A, Pons O. 1998.** Identifying populations for conservation on the basis of genetic markers. *Conservation Biology* **12**: 844–855.
- Pritchard JK, Stephens M, Donnelly P. 2000.** Inference of population structure using multilocus genotype data. *Genetics* **155**: 945–959.
- Ratter JA, Riveiro JF, Bridgewater S. 1997.** The Brazilian cerrado vegetation and threats to its biodiversity. *Annals of Botany* **80**: 223–230.
- Remington DL, Robichaux RH. 2007.** Influences of gene flow on adaptive speciation in the *Dubautia arborea*–*D. ciliolata* complex. *Molecular Ecology* **16**: 4014–4027.
- Rieseberg LH, Willis JH. 2007.** Plant speciation. *Science* **317**: 910–914.
- Rosenberg NA. 2004.** Distruct: a program for the graphical display of population structure. *Molecular Ecology Notes* **4**: 137–138.
- Saint-Hilaire A. 1829.** *Casearia lingua*. Cambess. (Eichler). *Flora brasiliae meridionalis* **2**: 236–236.
- Schluter D. 2001.** Ecology and the origin of species. *Trends in Ecology and Evolution* **16**: 372–380.
- Schneider S, Roessli D, Excoffier L. 2000.** *ARLEQUIN: a software for population data analysis*, version 2.0. Geneva: University of Geneva.
- Segarra-Moragues JG, Palop-Esteban M, Gonzalez-Candelas F, Catalan P. 2004.** Characterization of seven (CTT)_n microsatellite loci in the Pyrenean endemic *Borderea pyrenaica* (Dioscoreaceae): remarks on ploidy level and hybrid origin assessed through allozymes and microsatellite analyses. *Journal of Heredity* **95**: 177–183.
- Silva JF. 1996.** Biodiversity and stability in tropical savannas. In: Solbrig OT, Medina E, Silva JF, eds. *Biodiversity and savanna ecosystem processes: a global perspective*. Berlin: Springer-Verlag, 161–171.
- Silva MAS, Ming LC, Pereira AMS, Bertoni BW, Batistini AP, Pereira PS. 2006.** Phytochemical and genetic variability of *Casearia sylvestris* Sw. from São Paulo State Atlantic Forest and Cerrado populations. *Revista Brasileira de Plantas Medicinadas* **8**: 159–166.
- Sleumer HO. 1980.** *Flora Neotropica Monograph n.22. FLACOURTIACEAE*. New York: The New York Botanical Garden.
- Slotta TAB, Porter DM. 2006.** Genetic variation within and between *Iliamna corei* and *I. remota* (Malvaceae): implications for species delimitation. *Botanical Journal of the Linnean Society* **151**: 345–354.
- Smith TB, Kark S, Schneider CJ, Wayne RK, Moritz C. 2001.** Biodiversity hotspots and beyond: the need for preserving environmental transitions. *Trends in Ecology and Evolution* **16**: 431.
- Soltis DE, Soltis PS, Tate JA. 2003.** Advances in the study of polyploidy since plant speciation. *New Phytologist* **161**: 173–191.
- Soltis PS, Soltis DE. 2000.** The role of genetic and genomic attributes in the success of polyploids. *Proceedings of the National Academy of Sciences* **97**: 7051–7057.
- Torres RB, Ramos R. 2007.** Flacourtiaceae. In: Wanderley MGL, Shepherd GJ, Melhem TSA, Giulietti AM, eds. *Flora Fanerogâmica do Estado de São Paulo*. São Paulo: Instituto de Botânica, 201–223.
- Torres RB, Yamamoto K. 1986.** Taxonomia das espécies de *Casearia* Jacq. (Flacourtiaceae) do estado de São Paulo. *Revista Brasileira de Botânica* **9**: 239–258.
- Turesson G. 1922.** The genotypical response of the plant species to the habitat. *Hereditas* **3**: 211–350.
- Tuskan GA, DiFazio S, Jansson S et al. 2006.** The genome of black cottonwood, *Populus trichocarpa*. Torr. and Gray. *Science* **313**: 1596–1604.
- Vähä J-P, Primmer CR. 2006.** Efficiency of model-based Bayesian methods for detecting hybrid individuals under different hybridization scenarios and with different numbers of loci. *Molecular Ecology* **15**: 63–72.
- Valbuena-Carabaña M, González-Martínez SC, Sork VL et al. 2005.** Gene flow and hybridization in a mixed oak forest [*Quercus pyrenaica* Willd. and *Q. petraea* (Matts.) Liebl.] in central Spain. *Heredity* **95**: 457–465.
- Van Ee BW, Jelinski N, Berry PE, Hipp AL. 2006.** Phylogeny and biogeography of *Croton alabamensis* (Euphorbiaceae), a rare shrub from Texas and Alabama, using DNA sequence and AFLP data. *Molecular Ecology* **15**: 2735–2751.
- Van Oosterhout C, Hutchinson WF, Wills DP, Shipley P. 2004.** MICRO-CHECKER: software for identifying and correcting genotyping errors in microsatellite data. *Molecular Ecology Notes* **4**: 535–538.
- Wiens J. 2004.** Speciation and ecology revisited: phylogenetic niche conservatism and the origin of species. *Evolution* **58**: 193–197.
- Wright S. 1951.** The genetical structure of populations. *Annals of Eugenics* **15**: 395–420.
- Yeh FC, Yang RC, Boyle TBJ, Ye ZH, Mao JX. 1997.** *POPGENE, the user-friendly shareware for population genetic analysis*. Edmonton: Molecular Biology and Biotechnology Centre, University of Alberta.
- Yin TM, DiFazio SP, Gunter LE, Riemenschneider D, Tuskan GA. 2004.** Large-scale heterospecific segregation distortion in *Populus* revealed by a dense genetic map. *Theoretical and Applied Genetics* **109**: 451–463.
- Young A, Boyle T, Brown T. 1996.** The population genetic consequences of habitat fragmentation for plants. *Trends in Ecology and Evolution* **11**: 413–418.