



Published in final edited form as:

*ChemMedChem*. 2010 August 2; 5(8): 1221–1225. doi:10.1002/cmdc.201000186.

## HTS and Rational Drug Design [RDD] to Generate a Class of 5-HT<sub>2C</sub>-Selective Ligands for Possible Use in Schizophrenia

**Alan P. Kozikowski,**

Drug Discovery Program, Department of Medicinal Chemistry and Pharmacognosy, College of Pharmacy University of Illinois at Chicago, Illinois 60612-7230 (USA)

**Sung Jin Cho,**

Drug Discovery Program, Department of Medicinal Chemistry and Pharmacognosy, College of Pharmacy University of Illinois at Chicago, Illinois 60612-7230 (USA)

**Niels H. Jensen,**

Departments of Pharmacology and Psychiatry, Comprehensive Cancer Center, Center for Neurobiology, Division of Medicinal Chemistry and Natural Products and NIMH Psychoactive Drug Screening Program, University of North Carolina Medical School, Chapel Hill CB 7365, North Carolina 27599 (USA)

**John A. Allen,**

Departments of Pharmacology and Psychiatry, Comprehensive Cancer Center, Center for Neurobiology, Division of Medicinal Chemistry and Natural Products and NIMH Psychoactive Drug Screening Program, University of North Carolina Medical School, Chapel Hill CB 7365, North Carolina 27599 (USA)

**Andreas M. Svennebring,** and

Drug Discovery Program, Department of Medicinal Chemistry and Pharmacognosy, College of Pharmacy University of Illinois at Chicago, Illinois 60612-7230 (USA)

**Bryan L. Roth**

Departments of Pharmacology and Psychiatry, Comprehensive Cancer Center, Center for Neurobiology, Division of Medicinal Chemistry and Natural Products and NIMH Psychoactive Drug Screening Program, University of North Carolina Medical School, Chapel Hill CB 7365, North Carolina 27599 (USA)

### Keywords

CNS; Drug Discovery; 5-HT<sub>2C</sub>; Medicinal Chemistry; Schizophrenia

---

The 5-hydroxytryptamine (**1**, 5-HT) 2C receptor (5-HT<sub>2C</sub>), a prominent central serotonin receptor subtype, is widely distributed throughout the central nervous system (CNS) and is thought to play a role in regulating a wide variety of behavioral processes such as mood, appetite, and sexual behavior.[1-4] The 5-HT<sub>2A</sub> receptor mediates the hallucinogenic activity of drugs such as lysergic acid diethylamide (LSD) and is a major target for treating schizophrenia, insomnia and other disorders.[5-8] The 5-HT<sub>2B</sub> receptor mediates the potentially lethal valvulopathic side effects of several compounds that were used as prescription drugs.[9,10]

---

Fax: +1 312-996-7107 kozikowa@uic.edu.

Supporting information for this article is available on the WWW under <http://www.chemmedchem.org> or from the author.

5-HT<sub>2C</sub> agonists have demonstrated efficacy in preclinical models of depression, obesity, addiction, and psychosis.[11-13] Targeting the 5-HT<sub>2C</sub> receptor thus appears to offer a promising means for developing novel therapeutics for the treatment of CNS related disorders. However, as this receptor is homologous to the two other family members, 5-HT<sub>2A</sub> and 5-HT<sub>2B</sub>, [14] it is essential that 5-HT<sub>2C</sub> agonists being developed for clinical use show little if any activity at these subtypes.[15] To date, several 5-HT<sub>2C</sub> agonists have shown efficacy in preclinical animal models (Figure 1), [16-18] and are currently undergoing human trials. [16] In particular, one of the most advanced 5-HT<sub>2C</sub> ligands is Lorcaserin (**2**) which is being developed by Arena Pharmaceuticals and which has been demonstrated in two Phase III trials to be an orally active, antiobesity medication.[19]

Based upon the identification of tranlycypromine as the initial hit from an HTS campaign employing a library of FDA approved drugs, we undertook a structural optimization campaign that led to a potent, but moderately selective agonist **8** with 120- and 14-fold selectivity over 5-HT<sub>2A</sub> and 5-HT<sub>2B</sub>, respectively (EC<sub>50</sub> = 585, 65, and 4.8 nM at the 2A, 2B, and 2C subtypes, respectively). Compound **8** (10–60 mg/kg) was also demonstrated to exhibit moderate antidepressant-like effects in a commonly used behavioral assay.[18]

However, because compound **8** fails to exhibit sufficient selectivity over the 5-HT<sub>2B</sub> receptor, further optimization was required to identify a potential clinical candidate. Currently, the degree of selectivity that is actually needed to avoid side effects is unknown, however, this is a question that can ultimately be addressed only by studies in humans. Thus, we explored additional, selected modifications of these *trans*-2-phenylcyclopropylmethylamine analogues in order to improve upon this subtype selectivity issue. Our efforts to modify the 5-HT<sub>2C</sub> agonist **8** led to the discovery of several new drug candidates with an increased subtype selectivity including dual 5-HT<sub>2B</sub> antagonism / 5-HT<sub>2C</sub> agonism in the functional assays, and which were thus found suitable for in vivo testing as detailed herein.

By stepwise structural modifications of the *trans*-(2-aryl)cyclopropylmethylamine aromatic moiety, we found that potent, 5-HT<sub>2C</sub>-selective compounds could be produced using 5-hydroxy or 5-fluoro substituted systems with a medium sized 2-alkyloxy group. The 2-cyclopropylmethoxy-5-fluoro substituted derivative **12** was synthesized according to the steps shown in Scheme 1. Thus, the starting compound **9** was prepared by employing a standard sequence of reactions as previously reported.[18] The amino group of the phenolic derivative **9** was protected using Boc-anhydride. The *N*-Boc protected derivative **10** was then alkylated with cyclopropylmethyl bromide followed by subsequent deprotection to provide the racemic compound **12**.

To synthesize the 2-cyclopropylmethoxy-5-hydroxy substituted derivative **18**, the 2-cyclopropylmethoxy intermediate **16** was prepared in a straightforward manner through a sequence of selective protection and alkylation steps.[20] Next, the pivaloyl and Boc protecting groups were removed sequentially to afford the final product **18** (Scheme 2).

To prepare the optically pure enantiomers of **12** (Scheme 3), we first carried out a chiral separation of the *N*-Boc protected fluoro derivative **11** using a Chiralpack AD column. Under isocratic conditions (4% isopropanol in hexane), the individual enantiomers could be conveniently separated in pure form (> 99%). Due to the high resolution, stacked injections could be employed in order to increase throughput. The choice of intermediate **11** on which to carry out the chiral separation was based on the ease of the separation and the ready cleavage of the resulting enantiomers under acidic conditions.

The resulting enantiomers (+)-**11** and (–)-**11** were then converted individually to (–)- and (+)-*trans*-[2-(2-cyclopropylmethoxy-5-fluorophenyl)cyclopropyl]-methylamine hydrochlorides ((–)-**12** and (+)-**12**), respectively, using the same method as described above for the racemate.

For preparation of the pure enantiomers of compound **18**, intermediate **17** was used for the chiral separation.

The functional activity of these two sets of compounds was determined by measuring  $G_{\alpha q}$  mediated intracellular calcium mobilization in HEK-293 cells stably expressing the human 5-HT<sub>2A</sub>, human 5-HT<sub>2B</sub>, and human 5-HT<sub>2C</sub> (INI) receptors.[21] The results are summarized in Table 1. In the first round of the functional assays, racemic **12** was found not to activate either the 5-HT<sub>2A</sub> or 5-HT<sub>2B</sub> receptors and to have an EC<sub>50</sub> of 19 nM at the 5-HT<sub>2C</sub> receptor. The more active enantiomer (+)-**12** also showed the expected selectivity profile in these functional assays. In contrast, the less active isomer (–)-**12** had an EC<sub>50</sub> of 918 nM in the 5-HT<sub>2C</sub> assay. For comparison purposes we also tested the 5-HT<sub>2C</sub> ligand **2**, which is in Phase III clinical trials for obesity. In our hands this compound has a low nM potency at the 5-HT<sub>2C</sub> receptor, however, it is also fairly active at the 5-HT<sub>2B</sub> subtype, with an EC<sub>50</sub> of 85 nM and an E<sub>max</sub> of 93%. As such, there may be some risk for possible valvulopathic side effects induced by this drug. Compound (+)-**12** has an even lower selectivity margin towards the 5HT<sub>2B</sub>-subtype, but due to its low maximal activity, 21%, we believe this will be a very minor concern, especially in view of the 2B-antagonist activity described below. Three other known 5-HT<sub>2C</sub> agonists Vabicaserin (**3**) and the Way compounds **4** and **5** along with 5-HT (**1**) are also included in Table 1 for reference purposes.

In order to further characterize the pharmacology of the lead compounds, both (+)-**12** and (+)-**18**, which show minimal 5-HT<sub>2B</sub> activation, were tested for their ability to function as 5-HT<sub>2B</sub> antagonists. Compounds (+)-**12** and (+)-**18** were found to shift the concentration curve of 5-HT in calcium flux experiments rightwards without depressing the maximal 5-HT response, indicating fast binding kinetics (Figure 2). Schild analyses yielded pA<sub>2</sub> (± SEM) values of 5.50 ± 0.06, 5.79 ± 0.07, and 5.92 ± 0.02, respectively (n = 3). These results show that the tested compounds act as moderate potency, full antagonists at the 5-HT<sub>2B</sub> receptor.

On the basis of their promising in vitro pharmacology, the active enantiomers (+)-**12** and (+)-**18** were selected for further in vivo studies. Specifically, we examined the ability of these compounds to normalize disrupted prepulse inhibition (PPI) in phencyclidine (PCP) treated animals.[22] The ability to normalize the effects of PCP (an NMDA receptor antagonist)-induced disruption of PPI in mice is a well accepted model of measuring atypical antipsychotic activity [23,24] and has been used previously to characterize the antipsychotic activity of 5-HT<sub>2C</sub> agonists.[25] We have successfully used an identical PPI testing paradigm previously to demonstrate clozapine, a classic atypical antipsychotic, robustly normalizes PCP-disrupted PPI.[26] We tested the ability of compounds (+)-**12** and (+)-**18** as well as **3** as the reference compound, to normalize PCP effects on PPI (Figure 3). As is apparent from the accompanying figures, both (+)-**12** and (+)-**18** are able to normalize PCP-disrupted PPI at doses of 10 mg/kg (and are comparable in activity to **3**), while at the 5 mg/kg dose level, (+)-**12** and **3** are still effective.

Because of its robust activity at normalizing PCP-disrupted PPI, we carried out cytochrome P-450 screening, metabolic stability studies, and hERG assays on (+)-**12** in order to qualify it as a possible candidate for further development. In the recombinant CYP inhibition test, (+)-**12** showed relatively low inhibition against CYP2C9 (19.6%), CYP2D6 (27.0%), and CYP3A4 (21.7% and 25.1% using midazolam and testosterone as the substrates, respectively, at 10 μM (Table 2). Furthermore, compound (+)-**12** had an acceptable microsomal stability in a 1h assay (90.5% left in human microsomes and 66% in rat, respectively) compared with the control drug, verapamil (39.6% using human microsomes). Additionally, no hERG inhibition was detected for (+)-**12**, (–)-**12**, or **18**, nor for the comparison compounds **2**, **3**, and **5**. However, **4** caused very modest inhibition (3.6 mM, 97%), and the standard cisapride gave full inhibition (35 nM, 100%).

In conclusion, our optimization efforts led to the discovery of the highly selective 5-HT<sub>2C</sub> agonists (±)-**12** and (±)-**18**, and to the identification of (+)-**12** and (+)-**18** as the more active enantiomers. Compounds (+)-**12** and (+)-**18** were as effective as the known reference compound **3** in the PCP-PPI animal model of antipsychotic drug activity. Compound **3** was recently in Phase II clinical trials for the treatment of acute exacerbations of schizophrenia. Compound (+)-**12** has an acceptable DMPK profile while showing no adverse side effects in animals. Compound (+)-**12** represents a structurally simple 5-HT<sub>2C</sub> partial agonist of excellent potency and high selectivity. The structural relationship of **12** with the marketed monamine oxidase inhibitor tranylcypromine is also noteworthy, and when combined with the information now in hand, bodes well for the further development of this class of molecules. The study of (+)-**12** in other animal disease models including cocaine addiction is underway.

## Supplementary Material

Refer to Web version on PubMed Central for supplementary material.


## Acknowledgments

This work was supported by NIH grants R01 DA022317 (A.P.K.) and R01 MH61887, N01 MH80032, and U19 MH82441 (B.L.R.). S.J.C. was supported by the Korea Research Foundation Grant funded by the Korean Government (KRF-2007-357-C00071) and the NIDA fellowship (2006–2007). J.A.A. was supported by fellowship NICHD T32HD040127 and the UNC Neurodevelopmental Disorders Research Center. We thank Drs. Arsen Gaysin and Rong He for technical assistance. We thank Barbara Caldarone for reviewing the article and providing comments.


## References

1. Chou-Green JM, Holscher TD, Dallman MF, Akana SF. *Physiol. Behav* 2003;78:641. [PubMed: 12782219]
2. Rocha BA, Goulding EH, O'Dell LE, Mead AN, Coufal NG, Parsons LH, Tecott LH. *J. Neurosci* 2002;22:10039. [PubMed: 12427861]
3. Tecott LH, Sun LM, Akana SF, Strack AM, Lowenstein DH, Dallman MF, Julius D. *Nature* 1995;374:542. [PubMed: 7700379]
4. Millan MJ, Peglion JL, Perrin-Monneyron GSL. *Eur. J. Pharmacol* 1997;325:9. [PubMed: 9151932]
5. Nichols DE. *Pharmacol. Ther* 2004;101:131. [PubMed: 14761703]
6. Nilsson BM. *J. Med. Chem* 2006;49:4023. [PubMed: 16821762]
7. Lacivita E, Leopoldo M. *Curr. Top. Med. Chem* 2006;6:1927. [PubMed: 17017967]
8. Ahmad S, Ngu K, Miller KJ, Wu G, Hung CP, Malmstrom S, Zhang G, O'Tanyi E, Keim WJ, Cullen MJ, Rohrbach KW, Thomas M, Ung T, Qu Q, Gan J, Narayanan R, Pellemounter MA, Robl JA. *Bioorg. Med. Chem. Lett* 20:1128. [PubMed: 20022752]
9. Rothman RB, Baumann MH, Savage JE, Rauser L, McBride A, Hufeisen SJ, Roth BL. *Circulation* 2000;102:2836. [PubMed: 11104741]
10. Roth BL. *N. Engl. J Med* 2007;356:6. [PubMed: 17202450]
11. Dunlop J, Marquis KL, Lim HK, Leung L, Kao J, Cheesman C, Rosenzweig-Lipson S. *CNS Drug. Rev* 2006;12:167. [PubMed: 17227285]
12. Burbassi S, Cervo L. *Psychopharmacology (Berl)* 2008;196:15. [PubMed: 17899022]
13. Miller KJ. *Mol. Interv* 2005;5:282. [PubMed: 16249524]
14. Kroeze WK, Kristiansen K, Roth BL. *Curr. Top. Med. Chem* 2002;2:507. [PubMed: 12052191]
15. Berger M, Gray JA, Roth BL. *Annu. Rev. Med* 2009;60:355. [PubMed: 19630576]
16. Wacker DA, Miller KJ. *Curr. Opin. Drug Discovery Dev* 2008;11:438.
17. Booth RG, Fang L, Huang Y, Wilczynski A, Sivendran S. *Eur. J. Pharmacol* 2009;615:1. [PubMed: 19397907]
18. Cho JS, Jensen HN, Kurome T, Kadari S, Manzano LM, Malberg EJ, Caldarone B, Roth LB, Kozikowski PA. *J. Med. Chem* 2009;52:1885. [PubMed: 19284718]


19. Wolfson W. *Chem. Biol* 2008;15:1139. [PubMed: 19022173]
20. Yao Q. *Angew. Chem., Int. Ed* 2000;39:3896.
21. Jensen NH, Rodriguiz RM, Caron MG, Wetsel WC, Rothman RB, Roth BL. *Neuropsychopharmacology* 2008;33:2303. [PubMed: 18059438]
22. Swerdlow NR, Braff DL, Taaid NM, Geyer MA. *Arch. Gen. Psychiatry* 1994;51:139. [PubMed: 8297213]
23. Geyer MA, Ellenbroek B. *Prog. Neuro-Psychopharmacol. Biol. Psychiatry* 2003;27:1071.
24. Geyer MA, Krebs-Thomson K, Braff DL, Swerdlow NR. *Psychopharmacology (Berlin, Ger.)* 2001;156:117.
25. Marquis LK, Sabb LA, Logue FS, Brennan AJ, Piesla JM, Comery AT, Grauer MS, Ashby RC, Nguyen QH, Dawson AL, Barrett EJ, Stack G, Meltzer YH, Harrison LB, Rosenzweig-Lipson S. J. *Pharmacol. Exp. Ther* 2007;320:486. [PubMed: 17038512]
26. Abbas IA, Yadav NP, Yao W-D, Arbuckle IM, Grant GNS, Caron GM, Roth LB. *J. Neurosci* 2009;29:7124. [PubMed: 19494135]




**Figure 1.**  
Several recently developed 5-HT<sub>2C</sub> agonists.




**Figure 2.** 5-HT<sub>2B</sub> antagonism (Schild Plot) of compounds (+)-**18** (A) and (+)-**12** (B). (pA<sub>2</sub> (± SEM): 5.79 ± 0.07 and 5.92 ± 0.02 (n = 3)).




**Figure 3.** Compounds (+)-**12** and (+)-**18** normalize PCP-disrupted PPI in mice. Mice were administered (ip) saline or 6 mg/kg PCP and immediately tested for PPI of the acoustic startle response. (A) and (B). 30 min pre-treatments of mice with (+)-**12** or (+)-**18** effectively normalize PCP-disruption of PPI. (C). Compound **3**, a positive reference control compound, also normalizes PCP-disrupted PPI. \*,  $p < 0.05$  versus saline treated animals. Data are the mean  $\pm$  SEM.  $N = 9, 10$  or  $7$  animals in each testing group for (+)-**12**, (+)-**18** or **3**, respectively.

**Scheme 1.**

Reagents and conditions: (a) Boc<sub>2</sub>O, triethylamine, CH<sub>2</sub>Cl<sub>2</sub>, 0 °C to rt, 5 h. (b) (bromomethyl)cyclopropane, K<sub>2</sub>CO<sub>3</sub>, DMF, 60 °C, 20 h. (c) 2M HCl, rt, 48 h.

**Scheme 2.**

Reagents and conditions: (a) (i)  $\text{BBr}_3$ ,  $\text{CH}_2\text{Cl}_2$ ,  $-78^\circ\text{C}$  to rt, 6 h; (ii)  $\text{Boc}_2\text{O}$ , triethylamine,  $\text{CH}_2\text{Cl}_2$ ,  $0^\circ\text{C}$  to rt, 1 h. (b)  $\text{Piv-Cl}$ ,  $\text{CH}_2\text{Cl}_2$ , triethylamine,  $0^\circ\text{C}$  to rt, 6 h. (c) (bromomethyl) cyclopropane,  $\text{K}_2\text{CO}_3$ , DMF,  $60^\circ\text{C}$ , 20 h. (d)  $\text{NaOtBu}$ , MeOH, rt, 1 h. (e) 2 M HCl, rt, 48 h.

**Scheme 3.**

Reagents and conditions: (a) 2M HCl, rt, 48 h.

Table 1

Functional Activity and Selectivity of Racemic **12** and **18**, their Enantiomers, and Comparison Compounds at the Human 5-HT<sub>2A</sub>, 5-HT<sub>2B</sub>, and 5-HT<sub>2C</sub> Receptors in Calcium Flux Assays Using Stably Transfected HEK-293 Cells.

Compound <sup>[a]</sup>	5-HT <sub>2A</sub>			5-HT <sub>2B</sub>			5-HT <sub>2C</sub>			Selectivity	
	EC <sub>50</sub> ± SEM (nM)	%MAX <sup>[b]</sup> ± SEM (nM)	N <sup>[c]</sup>	EC <sub>50</sub> ± SEM (nM)	%MAX <sup>[b]</sup> ± SEM (nM)	N <sup>[c]</sup>	EC <sub>50</sub> ± SEM (nM)	%MAX <sup>[b]</sup> ± SEM (nM)	N <sup>[c]</sup>	2A/2C	2B/2C
<b>1</b> (5-HT)	6.9 ± 0.42	100%	15	0.7 ± 0.02	100%	16	0.1 ± 0.01	100%	16	63	6.7
(+)- <b>12</b>	NA	7% ± 1.8%	3	NA	6% ± 0.6%	3	19 ± 3.5	68% ± 5.2%	3	-	-
(+)- <b>18</b>	761 ± 160	11% ± 1.9%	3	NA	2% ± 0.9%	3	9.9 ± 1.7	68% ± 4.5%	3	77	-
<b>2</b> (Lorcaserin)	264 ± 31	24% ± 0.9%	6	85 ± 7.0	93% ± 1.2%	6	2.1 ± 0.29	99% ± 1.0%	7	123	40
<b>3</b> (Vabicaserin)	NA	2% ± 0.1%	6	NA	5% ± 1.0%	6	6.0 ± 1.0	95% ± 0.8%	7	-	-
<b>4</b> (WAY-161503)	76 ± 9.1	80% ± 1.0%	3	15 ± 1.6	92% ± 1.3%	3	1.1 ± 0.16	97% ± 1.6%	4	70	14
<b>5</b> (WAY 629)	NA	1% ± 0.2%	6	NA	6% ± 1.3%	6	286 ± 40	80% ± 1.1%	7	-	-
<b>1</b> (5-HT)	6.1 ± 0.38	100%	9	0.8 ± 0.08	100%	9	0.1 ± 0.0	100%	9	62	7.7
(+)- <b>12</b>	894 ± 91	29 ± 1.6%	3	289 ± 34	21% ± 4.9%	3	21 ± 2.2	71% ± 4.4%	3	42	14
(-)- <b>12</b>	NA	0% ± 0.0%	3	NA	4% ± 2.7%	3	918 ± 83	35% ± 2.4%	3	-	-
(+)- <b>18</b>	372 ± 131	18% ± 3.0%	3	NA	6% ± 2.8%	3	9.3 ± 0.05	70% ± 5.4%	3	40	-
(-)- <b>18</b>	NA	3% ± 0.6%	3	NA	0% ± 0.5%	3	361 ± 21	51% ± 2.4%	3	-	-
<b>2</b> (Lorcaserin)	136 ± 24	31% ± 0.5%	3	50 ± 10	86% ± 2.3%	3	1.7 ± 0.09	94% ± 0.8%	3	79	29
<b>3</b> (Vabicaserin)	NA	7% ± 0.1%	3	47 ± 6.4	15% ± 3.3%	3	8.0 ± 1.6	88% ± 1.8%	3	-	5.8
<b>4</b> (WAY-161503)	61 ± 8.3	79% ± 4.5%	3	16 ± 3.2	86% ± 4.7%	3	1.5 ± 0.18	94% ± 3.6%	3	40	11
<b>5</b> (WAY 629)	NA	7% ± 2.0%	3	> 5 μM	17% ± 2.7%	3	451 ± 52	82% ± 3.5%	3	-	-

<sup>[a]</sup> Tested in two independent screening campaigns using different cell lines / passages; direct comparisons of the potencies and efficacies are only valid within the bounds of each particular table section.

<sup>[b]</sup> Percent of maximal activation by 5-HT; activation at 10 μM for compounds without EC<sub>50</sub> value.

<sup>[c]</sup> n: Number of concentration curves from ≥ 2 (typically ≥ 3) independent experiments. NA: E<sub>max</sub> ≤ 12%. In contrast to binding affinities, the potencies in functional assays can vary strongly depending on cell type, receptor expression level, and passage number.