

Influence of Apolipoprotein E Polymorphism on Apolipoprotein B-100 Metabolism in Normolipemic Subjects

Thomas Demant, Dorothy Bedford, Christopher J. Packard, and James Shepherd
Institute of Clinical Biochemistry, Glasgow Royal Infirmary, Glasgow G4 OSF, Scotland

Abstract

This study examined apolipoprotein (apo) B metabolism in normolipemic subjects homozygous for the apo E₂ (n = 4), apo E₃ (n = 5), or apo E₄ (n = 5) phenotype. Radioiodinated very low density lipoprotein (VLDL₁) (ultracentrifuge flotation rate [S_T] 60–400) and VLDL₂ (S_T 20–60) were injected into volunteers and the conversion of apo B was followed through intermediate density lipoprotein (IDL) to low density lipoprotein (LDL). Subjects homozygous for E₃ converted ~ 50% of VLDL₂ to LDL, the remainder being lost by direct catabolism. Those with the E₂ phenotype produced less VLDL₁ but converted more of it to VLDL₂ (compared to E₃ subjects). They displayed a characteristic dyslipidemia with the presence of slowly catabolized VLDL₁ and VLDL₂ remnants. LDL levels were low owing to increased direct catabolism of VLDL₂ and IDL and a reduced efficiency of delipidation; only 25% of VLDL₂ apo B was directed to LDL production. In contrast, E₄ subjects converted more VLDL₂ apo B to LDL than E₃ subjects. About 70% of VLDL₂ apo B was found in LDL; direct catabolism of VLDL and IDL was reduced as was the fractional catabolic rate of LDL (0.2 vs. 0.26 in E₃ subjects). These changes in the VLDL → IDL → LDL metabolic cascade can in part be explained by alterations in hepatic LDL receptors with E₂ subjects having higher and E₄ subjects lower activities than those in E₃ homozygotes. (*J. Clin. Invest.* 1991. 88:1490–1501.) **Key words:** catabolism • modeling • kinetics • synthesis • very low density lipoprotein

Introduction

In humans the gene locus for apolipoprotein E (apo E) is polymorphic with three alleles (E₂, E₃, and E₄) being present in all populations studied so far (1–3). The commonest variant is E₃ for which ~ 70% of Caucasians are homozygous. It codes for a protein of 299 amino acid residues, which although minor in terms of plasma concentration, has an important role in regulating lipoprotein metabolism by virtue of its ability to act as a ligand for specialized lipoprotein receptors. Individuals who possess the E₂ allele synthesize an apo E in which the arginine residue at position 158 is replaced with cysteine as the result of a C → T point mutation. Those with E₄ in contrast generate a

product with an arginine at residue 112 compared with cysteine in apo E₃. These mutations in the protein sequence cause charge changes which are readily detected by isoelectric focusing of very low density lipoproteins (VLDL) apoproteins (4) or by isoelectric focusing of plasma followed by detection with immunoblotting (5, 6). Structure–function studies of apo E have revealed that a domain covering amino acid 140–160 is of critical importance in receptor interaction. The E₂ mutation occurs in this region and in vitro studies have demonstrated that the resulting protein has impaired receptor-binding properties (4, 7).

Epidemiological surveys have demonstrated that the apo E polymorphism has a significant impact on plasma cholesterol and apolipoprotein B (apo B) levels (3). Compared to apo E₃ homozygotes, those who are homozygous for the E₂ allele exhibit plasma cholesterol levels that are 10–15% lower while the opposite holds for those who are E₄ homozygotes. Examination of the lipoprotein profile in the various apo E phenotypes has revealed that the gradation in plasma cholesterol from E₂ to E₃ to E₄ homozygotes is due to an increase in the level of low density lipoprotein (LDL). However, E₂ homozygotes have higher VLDL cholesterol levels and display a dyslipidemia that is qualitatively similar to that seen in type III hyperlipidemia. The latter is usually associated with E₂ homozygosity and inheritance of another as yet unidentified mutation that causes an elevation of plasma lipid levels (8). A number of studies have examined the metabolic changes underlying the altered lipoprotein profiles associated with the apo E polymorphism. Chylomicron clearance is reduced in subjects possessing the E₂ allele, whereas E₄ heterozygotes and homozygotes show accelerated clearance (9, 10). Metabolic studies of LDL apo B have shown that type III and, to a lesser extent, E₂ homozygous subjects degrade LDL faster than normal (11, 12). Previous studies from this laboratory have examined VLDL, intermediate density lipoprotein (IDL), and LDL apo B metabolism in a number of genetically determined dyslipidemias. In particular, we found in patients with homozygous familial hypercholesterolemia that the LDL receptor was important not only in LDL catabolism but also in the processing of apo B-containing lipoproteins throughout much of the delipidation cascade from VLDL to LDL (13). Since apo E is important for lipoprotein receptor interaction, we surmised that structural and functional variation in this protein would modulate the metabolism of the lipoproteins within the entire ultracentrifuge flotation rate (S_T)¹ 0–400 spectrum. To investigate this, we conducted VLDL turnover studies in groups of individuals homozygous for the three E variants. The results demonstrate the profound impact that this polymorphism has on apo B metabolism.

Dr. Demant's current address is Institut für Klinische Chemie, Klinikum Grosshadern, D-8000 Munich 70, Federal Republic of Germany.

Address reprint requests to Dr. Demant.

Received for publication 2 February 1991 and in revised form 27 June 1991.

J. Clin. Invest.

© The American Society for Clinical Investigation, Inc.

0021-9738/91/11/1490/12 \$2.00

Volume 88, November 1991, 1490–1501

1. *Abbreviations used in this paper:* FCR, fractional catabolic rate; FH, familial hypercholesterolemia; S_T, ultracentrifuge flotation rate; TMU, tetramethylurea.

Methods

Subjects. The study participants were selected from individuals who attended a coronary screening program in health centers in Glasgow. The Institute of Biochemistry received samples for cholesterol assay as part of the screen and those in which the level lay in the range of 5.0–6.2 mmol/liter (195–240 mg/dl) were subjected to apo E phenotyping using the isoelectric focusing/immunoblotting technique described below. Subjects who were found to be homozygous for the presence of apo E₂ (n = 4), apo E₃ (n = 5), or apo E₄ (n = 5) were recruited for the study. For ease of presentation throughout this report subjects are referred to as the E₂, E₃, or E₄ group to denote phenotypically determined homozygosity for the E₂, E₃, or E₄ alleles (14). They were screened for cardiometabolic, renal, endocrine, hepatic, or metabolic disease by routine clinical examination and biochemical testing. Age distribution and body weight index were similar in the three study groups. Each individual's dietary intake was recorded over a 7-d period and this was analyzed for the percent contribution of calories derived from protein, carbohydrates and fat (15).² Although daily energy intake (kilo-calorie per day) varied considerably, approximately in proportion to

differences in body frame, the composition of food was virtually the same in the three groups.

Plasma lipid and lipoprotein levels of study participants given in Table I were measured according to the Lipid Research Clinics protocol (16).

Apo E phenotyping. Apo E isoforms were detected as described by Menzel et al. (5) and Havekes et al. (6). 10 µl of plasma were delipidated, redissolved in 6 M urea containing 0.1 M Tris, pH 10.0, 5% β-mercaptoethanol, and 1% sodium decyl sulfate and applied to a vertical polyacrylamide slab gel of 5% (wt/vol) acrylamide in 8 M urea containing 1% ampholyte (pH range 4–6). The cathode buffer was 0.2 M NaOH and the anode buffer 0.1 M H₃PO₄. Isoelectric focusing was carried out at 3 W per gel for 16 h. Transfer by electrophoresis from the acrylamide slab gels to nitrocellulose membranes was performed as described by Towbin et al. (17). The transfer buffer contained 0.2 M glycine, 0.025 M Tris, and 20% methanol and the current applied was 0.4 A for 3 h. Apo E isoforms bound to the membrane were visualized by immunostaining using a monoclonal or polyclonal apo E-specific antibody and an appropriate IgG-binding second antibody linked to horseradish peroxidase.

Lipoprotein isolation and labeling. The methods for preparation of tracer VLDL subfractions VLDL₁ (S_r 60–400) and VLDL₂ (S_r 20–60), have been described in detail in previous publications (18). Briefly, 250 ml of plasma was removed by plasmapheresis from subjects who had been fasted overnight and total VLDL of *d* < 1.006 g/ml (S_r 20–400)

2. Intake was analyzed by using the computer program MICRODIET (Department of Computer Sciences, Salford University, Salford, UK).

Table I. Plasma Lipid and Lipoprotein Levels in E₂, E₃, and E₄ Subjects

Subject	Sex	Age yr	Weight kg	Total triglyceride	Total cholesterol	VLDL cholesterol	LDL cholesterol	HDL cholesterol	VLDL cholesterol/plasma triglyceride ratio
				mmol/liter					
E_{2/2}									
1	M	47	74.0	1.60±0.79	4.88±0.13	1.03±0.64	2.43±0.43	1.42±0.19	0.64
2	F	39	47.0	1.80±0.27	4.85±0.35	1.07±0.10	2.45±0.26	1.33±0.08	0.59
3	F	45	63.0	1.85±0.27	6.14±0.83	1.41±0.47	3.25±0.49	1.48±0.23	0.76
4	F	57	53.0	1.64±0.08	5.96±0.76	1.25±0.24	3.25±0.59	1.48±0.21	0.76
Mean±SD	mmol/liter (mg/dl)			1.72±0.10 (151±9)	5.46±0.60 (211±23)	1.19±0.15* (46±6)	2.84±0.40 (110±15)	1.43±0.06 (55±2)	0.69±0.07* (0.30±0.03)
E_{3/3}									
5	M	36	71.0	1.71±0.38	4.96±0.54	0.79±0.21	3.14±0.35	1.01±0.13	0.46
6	M	43	86.0	2.55±0.71	6.23±0.79	1.02±0.25	4.09±0.50	1.36±0.19	0.40
7	M	46	89.0	1.09±0.13	5.76±0.22	0.61±0.14	3.96±0.17	1.19±0.08	0.56
8	F	40	71.0	0.95±0.06	4.97±0.32	0.35±0.08	2.76±0.17	1.86±0.12	0.37
9	F	44	76.0	2.17±0.29	5.63±0.35	0.97±0.21	3.74±0.14	1.20±0.05	0.45
Mean±SD	mmol/liter (mg/dl)			1.69±0.56 (148±49)	5.51±0.49 (213±19)	0.75±0.25 (29±10)	3.54±0.51 (137±20)	1.32±0.29 (51±11)	0.45±0.06 (0.20±0.03)
E_{4/4}									
10	M	29	85.0	2.00±0.75	6.55±1.00	1.08±0.54	4.37±0.83	1.10±0.22	0.54
11	F	41	62.0	1.49±0.28	6.74±0.35	0.61±0.23	4.66±0.41	1.39±0.19	0.41
12	M	45	79.0	2.01±0.59	6.19±0.34	0.99±0.50	3.96±0.20	1.23±0.14	0.49
13	F	29	50.0	0.90±0.27	5.66±0.40	0.59±0.28	3.35±0.49	1.73±0.09	0.66
14	F	36	48.0	0.93±0.18	5.33±0.46	0.32±0.14	3.45±0.44	1.56±0.11	0.34
Mean±SD	mmol/liter (mg/dl)			1.47±0.49 (129±43)	6.09±0.53 (236±21)	0.72±0.30 (28±12)	3.96±0.51 (153±20)	1.40±0.23 (54±9)	0.49±0.11 (0.22±0.05)

* Significantly different from E₃ group *P* < 0.05; † *P* < 0.01 by Student's *t* test.

isolated by centrifugation for 18 h at 36,000 rpm and 10°C in a Ti60 rotor (Beckman Instruments, Inc., Palo Alto, CA). The supernatant was harvested by suction and used for preparation of the subfractions. The VLDL solution was adjusted to a density of 1.118 kg/liter by the addition of NaCl (0.34 g per 2 ml of solution) and layered in a SW40 rotor tube (Beckman Instruments, Inc.). A six-step gradient from 1.0988–1.0588 kg/liter was constructed above the sample and centrifugation carried out at 23°C to separate sequentially VLDL₁ (1 h 38 min, 39,000 rpm) and VLDL₂ (15 h 41 min; 18,500 rpm). The subfractions were labeled with ¹³¹I and ¹²⁵I, respectively, by a modification of the iodine monochloride method (19) and sterilized by filtration through an 0.45 micron filter (Millipore, Molsheim, France).

Turnover protocol. Subjects were fasted from 8 p.m. and injected with autologous trace-labeled ¹³¹I-VLDL₁ and ¹²⁵I-VLDL₂ at 8.00 a.m. on the following day. The tracers (50 μCi of each iodine isotope) were administered within 72 h of blood being withdrawn for their preparation. During the first day of the turnover to minimize chylomicron production the volunteers were fasted until 6.00 p.m. Plasma samples were obtained at frequent intervals over the 72 h after injection and then daily in the fasting state for 14 d. The apo B-containing lipoproteins VLDL₁ (S_r 60–400), VLDL₂ (S_r 20–60), IDL (S_r 12–20), and LDL (S_r 0–12) were prepared by a modification (13) of the cumulative gradient ultracentrifugation (20). From these apo B was isolated by tetramethylurea (TMU) precipitation (21) and its specific activity was determined by radioactivity counting and protein determination (22).

The apo B pool circulating with each of these lipoprotein fractions was determined by replicate analyses of plasma samples collected intermittently throughout the turnover study. Correction was made for possible B protein loss during TMU precipitation by comparing the apo B recovered at the end of the procedure with the values calculated as the difference between total and TMU-soluble (apo E and C) protein contents (21). The composition of each fraction was determined by assay of total and esterified cholesterol, triglyceride, phospholipid, and protein (23). Protein determinations were performed with the addition of sodium dodecyl sulfate to the Biuret reagent (22).

Subjects received potassium iodate in tablet form (180 mg/d) for 3 d before and 1 mo after injection to block thyroidal sequestration of radioactive iodine released by catabolism.

Kinetic analysis. The radioactivity associated with the apo B protein present in each lipoprotein fraction was calculated from the apo B specific activities and the individual pool sizes. These were expressed as a percentage of the total apo B radioactivity (i.e., VLDL₁ plus VLDL₂, IDL, and LDL) present in the plasma 10 min after injection and the resulting values were used to construct decay curves which were analyzed by the SAAM 30 (24) multicompartmental modeling program. The metabolic model employed (Fig. 1) was basically the same as de-




Figure 1. Multicompartmental model for apo B metabolism in VLDL₁ (S_r 60–400), VLDL₂ (S_r 20–60), IDL (S_r 12–20), and LDL (S_r 0–12). U₁, U₅, U₇, and U₁₂ represent de novo input of apo B into VLDL₁, VLDL₂, and IDL. Synthesis into the LDL density interval was calculated as the difference between the absolute catabolic rate of apo B in this fraction (observed mass × overall FCR) and the input from VLDL₂ and IDL.

scribed elsewhere (13) with the addition of a second subcompartment (compartment 12) in VLDL₁ to account for the second exponential seen in E₂ homozygotes. Its main features are as follows: (a) apo B synthetic input occurs at the level of VLDL₁, VLDL₂, and, where indicated, IDL and LDL; (b) VLDL is delipidated in a stepwise manner following the concept of Berman et al. (25); (c) slowly catabolized remnant subpopulations are present in VLDL₂ (compartment 6) and IDL (compartment 9); (d) parallel processing pathways exist leading from VLDL₂ through IDL to LDL.

Rate constants were determined and in combination with B protein pool sizes were used to calculate flux rates and steady-state synthetic inputs. The rate constants, fluxes, and apo B masses were compared in the E₂, E₃, and E₄ homozygotes by Student's *t* test and Mann-Whitney nonparametric test.

Ethical consideration. All subjects participating in the study gave informed consent. The study met the requirements of the Ethical Committee of the Glasgow Royal Infirmary.

Results

The subjects for this study were selected from coronary screening clinic attendees whose plasma cholesterol at the initial visit fell in the range 5.0–6.2 mmol/liter. Mean values for total plasma cholesterol and triglyceride (Table I) in the three groups did not differ significantly from each other or from the population means (± 1 SD) for these lipids of 5.8 ± 1.2 and 1.8 ± 1.4 mmol/liter, respectively. Using this selection process we attempted to minimize the effect that gross perturbations in plasma lipid levels would have on apo B kinetics. The distribution of cholesterol in the lipoprotein fractions, however, was characteristically different in the groups. VLDL cholesterol was significantly higher and LDL cholesterol lower in E₂ vs. E₃ homozygotes. Likewise, the VLDL cholesterol/plasma triglyceride ratio was increased in the E₂ individuals. In this small series there was no significant difference in plasma lipoprotein levels between E₃ and E₄ homozygotes, although LDL cholesterol was moderately elevated in the latter as might be expected (3). HDL cholesterol was the same in all three groups. These perturbations in lipid levels were reflected in the distribution of apo B in the four lipoprotein fractions prepared by cumulative ultracentrifugation (Table II). Total apo B concentration, which was calculated as the sum of the apoprotein levels observed in VLDL₁, VLDL₂, IDL, and LDL, was markedly decreased in E₂ homozygotes. This was due to a much reduced LDL apo B level, which was approximately a third of that seen in E₃ subjects. VLDL₂ apo B in contrast was elevated in the E₂ compared with the E₃ group, whereas VLDL₁ and IDL apo B concentrations were similar. Individuals in the E₄ group had VLDL₁, VLDL₂, and IDL apo B levels that were close to those seen in E₃ subjects, although there was a tendency for LDL apo B to be higher in E₄ and the IDL/LDL ratio differed across all three groups (E₂ 0.63 ± 0.15 , E₃ 0.16 ± 0.016 , E₂ vs. E₃ $P < 0.05$ by *t* test; E₄ 0.13 ± 0.019 , E₃ vs. E₄ $P < 0.05$ by *t* test).

The composition of the four apo B-containing lipoprotein fractions is given in Table III. Compared with E₃ individuals, those with E₂ showed a high unesterified cholesterol content in VLDL₁ and an enrichment of cholesteryl ester at the expense of triglyceride in VLDL₂. The percentage of free cholesterol in LDL was decreased in E₂ homozygotes, and in this lipoprotein fraction there was also a graded decrease in triglyceride content comparing E₂ with E₃ and E₄.

Apo B kinetic studies. The metabolic behavior of apo B in VLDL₁, VLDL₂, IDL, and LDL after simultaneous injection

Table II. Apo B Concentrations in E₂, E₃, and E₄ Subjects

Subject	VLDL ₁	VLDL ₂	IDL	LDL	Total Apo B
	mg/dl				
E _{2/2}					
1	1.3	7.6	7.1	17.0	33
2	2.8	9.5	13.4	21.3	47
3	3.8	7.6	16.8	24.8	53
4	2.6	9.5	15.1	19.8	47
Mean±SD	2.6±0.9	8.6±1.0*	13.1±3.7	20.7±3.2*	45±8*
E _{3/3}					
5	3.3	6.3	8.7	57.8	76
6	2.4	7.3	13.8	86.6	110
7	3.7	5.8	11.3	76.3	97
8	1.8	2.6	8.1	50.5	63
9	4.9	7.5	12.6	66.0	91
Mean±SD	3.2±1.1	5.9±2.0	10.9±2.2	67.4±14.4	87±18
E _{4/4}					
10	4.1	7.5	13.6	82.8	108
11	1.8	4.6	13.3	103.3	123
12	4.6	5.0	8.1	58.3	76
13	1.3	2.9	7.7	64.1	76
14	0.2	3.5	10.1	88.1	102
Mean±SD	2.4±1.7	4.7±1.6	10.6±2.5	79.3±16.4	97±18

* Significantly different from E₃ group, *P* < 0.02 by Student's *t* test.

of VLDL₁ and VLDL₂ tracers is shown in Fig. 2. In order to examine intergroup differences more closely, the observed data at each time point were averaged within group and are presented as a mean and standard error for the E₂, E₃, and E₄ subjects. Individual decay curves and apo B masses were used

in multicompartmental modeling to generate the kinetic parameters given in Tables IV and V.

VLDL₁ apo B radioactivity was cleared rapidly from the plasma of E₃ and E₄ homozygotes (Fig. 2 *a*). The decay curve was monoexponential and only ~ 1% remained at the last time point on the first day, 14 h after injection. Clearance of this lipoprotein was slower in E₂ subjects with ~ 10% remaining at 14 h. The decay curve in the last group was biexponential containing a slow component which accounted for up to 10% of the apo B present in the VLDL₁ tracer. This feature necessitated the addition to the model of a second compartment (compartment 12) in VLDL₁ (Fig. 1). VLDL₁ apo B appeared in VLDL₂ with a peak radioactivity ~ 5 h after injection in all subjects (Fig. 2 *b*). Transfer of VLDL₁ to VLDL₂ was greater and catabolism of apo B from VLDL₂ was slower in E₂ compared with E₃ and E₄ subjects. The latter was true whether label in this fraction was introduced directly as a VLDL₂ tracer or derived from VLDL₁.

The metabolism of IDL derived from both tracers (Fig. 2 *c*) was similar in all three groups. Radioactivity peaked at 8–10 h and decayed in a multiexponential fashion, slightly faster in E₃ than in E₂ or E₄ subjects. Substantial differences however were observed in the metabolic behavior of LDL apo B. Peak radioactivity values of 20% for the VLDL₁ apo B tracer and ~ 35% for the VLDL₂ apo B tracer were observed in LDL for E₃ and E₄ homozygotes. In contrast, < 5% and ~ 12% of apo B radioactivity derived from the VLDL₁ and VLDL₂ tracers, respectively, was seen in the LDL fraction of E₂ subjects. The decay curve for LDL apo B in E₃ homozygotes appeared to decline more sharply than that for the E₄ group, particularly over the first 5 d of the turnover.

There was little difference in VLDL₁ apo B pool size in the three groups (Table IV *a*) in concordance with their similar plasma triglyceride levels (Table I). Apo B in this flotation interval was synthesized at about 800 mg/d and cleared at a rate

Table III. Composition of Apo B-containing Lipoproteins in E₂, E₃, and E₄ Subjects

	Free cholesterol	Cholesteryl ester	Triglyceride	Phospholipids	Protein
	g/100 g				
VLDL ₁					
E _{2/2}	4.7±0.5**	19.9±3.9	50.9±4.2	17.3±1.2	7.2±1.2
E _{3/3}	1.5±1.8	16.2±3.2	57.4±4.1	15.4±2.6	9.4±2.0
E _{4/4}	3.6±1.7	14.4±5.8	57.0±4.3	17.7±2.7	7.5±1.0
VLDL ₂					
E _{2/2}	8.6±1.4	30.5±3.7 [‡]	27.5±5.2 [§]	21.5±1.4	12.0±1.4
E _{3/3}	6.4±1.9	22.4±4.3	36.9±2.9	20.4±2.3	14.1±1.4
E _{4/4}	7.0±2.2	22.6±2.1	34.8±2.8	21.2±0.8	14.5±0.9
IDL					
E _{2/2}	9.5±1.6	38.3±2.2	11.6±2.8	23.7±1.0	17.0±0.8
E _{3/3}	8.9±3.0	35.6±4.7	14.3±1.7	22.1±1.7	19.0±1.7
E _{4/4}	9.4±2.5	37.0±1.7	11.4±2.0	23.2±1.2	19.0±1.4
LDL					
E _{2/2}	8.4±1.6 [‡]	38.2±1.9	7.9±1.2 [‡]	22.8±0.7	22.8±0.9
E _{3/3}	11.6±1.6	37.0±2.0	6.0±0.9	22.0±0.5	23.4±1.4
E _{4/4}	11.2±1.2	36.5±0.7	4.4±0.6 [§]	21.9±1.7	26.0±2.4

* Mean±1 SD. Significantly different from E₃ group, [‡] *P* < 0.05, [§] *P* < 0.02 by Student's *t* test.




Figure 2. Apo B radioactivity decay curves in (a) VLDL₁, (b) VLDL₂, (c) IDL, and (d) LDL (opposite page) in E₂, E₃, and E₄ subjects. Tracers of ¹³¹I-VLDL₁ and ¹²⁵I-VLDL₂ were injected simultaneously and their metabolism was followed through IDL to LDL. The data were calculated for individual subjects and are here presented as a mean ± SE for each group: (■) E₂, (◆) E₃, (▲) E₄.

of 10 pools/d in E₃ subjects. It had two metabolic fates, on average two-thirds were catabolized directly from plasma whereas the remainder was lipolyzed to VLDL₂. The results in E₄ homozygotes did not differ significantly from those seen in the E₃ group. However, E₂ homozygotes exhibited low apo B synthetic rates and greatly reduced direct catabolism of VLDL₁ (Table IV a, Fig. 2 b). It is noteworthy that the VLDL₁ to VLDL₂ transfer rate was not affected by apo E phenotype. Approximately 10% of the VLDL₁ apo B mass in E₂ subjects was placed in the slowly metabolized second compartment [M (12), Table V a]. Material from this pool was cleared directly from the plasma or delipidated to VLDL₂. The mathematical model (Fig. 1) was constructed so that apo B from remnant compartment 12 in VLDL₁, when delipidated, appeared in the VLDL₂ remnant pool (compartment 6).

Apo E phenotype appeared not to influence the rate of input of apo B into VLDL₂ whether the material was derived from direct synthesis or from delipidation of VLDL₁ (Table IV b). The overall pool size of VLDL₂ apo B did not differ in the three groups although the remnant compartment (compartment 6) was specifically increased in the E₂ subjects ($P < 0.02$

by Mann-Whitney test, Table V c). This was apparently due to increased diversion of apo B from the delipidation chain into the slowly metabolized compartment {cf. $L(6, 2)/[L(4, 2) + L(6, 2)]$ in E₂ vs. E₃ and E₄ subjects in Table V b} as well as a contribution of material derived from VLDL₁ remnant catabolism [by $L(6, 12)$, Table V a]. Direct catabolism of VLDL₂ apo B was reduced (although not significantly, $P = 0.11$) in E₂ homozygotes compared to E₃ subjects (Table IV b) while on-going delipidation to IDL was the same in all three groups. In no subject was there a requirement for de novo apo B synthesis at the level of IDL (Table IV c); the plasma pool of IDL apo B which was ~ 300–500 mg in all subjects was derived entirely from VLDL₂ catabolism. The metabolic fate of this fraction differed according to apo E phenotype. The transfer rate of IDL to LDL was reduced significantly by 66% in the E₂ group compared to E₃ and E₄ homozygotes. In contrast, direct IDL apo B catabolism was increased in E₂ subjects (Tables IV c and V d). This redirection was found to be a feature of both parallel pathways of apo B delipidation [Table V d $L(10, 7)$, $L(11, 8)$]. The differences in E₄ vs. E₃ subjects were not as dramatic. There was a tendency for reduced direct IDL apo B catabolism in E₄ ho-




Figure 2 (Continued)

mozygotes (Table IV *c*). This can be seen in the individual fractional rates of delipidation and catabolism for compartments 7 and 8 in Table V *d*. When the proportion of VLDL₂ turnover in milligrams per day directed towards LDL apo B was calculated (i.e., LDL production/VLDL₂ synthesis, Tables IV *b* and *d*), there was a clear gradation across the phenotypes. In E₂ homozygotes 23±8% of VLDL₂ apo B turnover was used to make LDL. This was significantly different from the 50±13% in E₃ subjects ($P < 0.02$ by Mann-Whitney test). Homozygotes for E₄, however, converted significantly more VLDL₂ apo B to LDL (70±14%, $P < 0.05$) compared to the E₃ group.

The main cause of the lower LDL apo B plasma concentration in E₂ subjects was reduced synthesis from VLDL (Table IV *d*). Both LDL subcompartments (compartments 10 and 11) were reduced in mass compared to E₃ (Table V *e*). There was little direct synthesis of apo B in the LDL flotation interval of the E₂ group and the catabolic rate for the apoprotein was similar to that seen in E₃ homozygotes. In E₄ homozygotes the overall fractional catabolic rate (FCR) of LDL apo B was reduced by 23% (Table IV *d*) owing to a specific decrease in the elimination constant for the faster catabolized compartment 10 (Table V *e*). In the majority of E₂ and E₃ individuals, L(0,

10) exceeded L(0, 11) by ~ 63%, whereas in the E₄ group the rate of LDL apo B catabolism from these two compartments was virtually identical. This is in keeping with the flat nature of the LDL apo B decay curve in E₄ subjects (Fig. 2 *d*).

Discussion

The polymorphism in the gene coding for apo E is the most important inherited trait modulating plasma cholesterol levels in the normal population (3). This study was designed to investigate the perturbations in apo B metabolism associated with homozygosity for the three common phenotypes, E₂, E₃, and E₄. Subjects were located during the screening of several thousand individuals for coronary risk factors. A decision was taken to select volunteers from those whose total plasma cholesterol levels on initial presentation were in a range (5.0–6.2 mmol/liter) that straddled the mean population value. By this approach we hoped to minimize any effect that gross differences in plasma lipid levels might have on apo B kinetics. Mean (±1 SD) cholesterol levels in subjects homozygous for apo E₂, apo E₃, and apo E₄ in the whole screened population were 5.28±0.71, 5.51±0.72, and 5.93±1.18 mmol/liter, respectively. There was no significant difference in plasma chole-

Table IV. Apo B Metabolism in E₂, E₃, and E₄ Normolipidemic Subjects

a. VLDL ₁					c. IDL					
	Synthesis	Plasma pool	Direct catabolism	Transfer to VLDL ₂		Direct synthesis	Flux from VLDL ₂	Plasma pool	Direct catabolism	Transfer to LDL
	mg/d	mg	pools/d			mg/d	mg	pools/d		
E_{2/2}					E_{2/2}					
1	140	37	1.24	2.54	1	0	455	253	1.55	0.24
2	396	54	2.57	4.76	2	0	370	330	0.70	0.42
3	374	85	0.00	4.39	3	0	510	467	0.77	0.32
4	338	50	1.92	4.84	4	0	334	309	0.73	0.34
Median	356**	52	1.58**	4.58	Median	0	413	320	0.75	0.33**
E_{3/3}					E_{3/3}					
5	625	67	4.17	5.16	5	0	621	241	0.76	1.80
6	522	52	5.75	4.49	6	0	495	408	0.18	1.02
7	1,288	97	10.79	2.48	7	0	517	358	0.26	1.17
8	1,120	106	6.91	3.66	8	0	385	351	0.33	0.76
9	819	47	14.73	2.71	9	0	189	174	0.37	0.68
Median	819	67	6.91	3.66	Median	0	495	351	0.33	1.02
E_{4/4}					E_{4/4}					
10	894	107	5.36	2.99	10	0	593	442	0.41	0.94
11	474	105	2.53	1.98	11	0	387	272	0.33	1.10
12	515	45	7.56	3.88	12	0	305	324	0.01	0.93
13	294	12	16.80	7.68	13	0	235	171	0.17	1.25
14	146	16	0.00	9.12	14	0	370	209	0.12	1.65
Median	474	45	5.36	3.88	Median	0	370	272	0.17	1.10
b. VLDL ₂					d. LDL					
	Direct synthesis	Flux from VLDL ₁	Plasma pool	Direct catabolism	Transfer to IDL and LDL		Direct [§] synthesis	Flux from IDL and VLDL ₂	Plasma pool	LDL FCR
	mg/d	mg	mg	pools/d			mg/d	mg	pools/d	
E_{2/2}					E_{2/2}					
1	373	94	224	0.06	2.03	1	10	61	507	0.14
2	238	257	237	0.42	1.67	2	14	165	460	0.39
3	333	374	215	0.84	2.46	3	12	164	650	0.27
4	234	242	214	0.65	1.56	4	2	106	450	0.24
Median	284	250	220	0.54	1.85	Median	11**	135 [‡]	484 [‡]	0.26
E_{3/3}					E_{3/3}					
5	301	346	160	0.18	3.88	5	0	435	1,433	0.30
6	761	229	231	2.14	2.14	6	233	418	2,505	0.26
7	500	241	199	0.92	2.81	7	10	460	2,044	0.23
8	280	388	218	1.15	1.93	8	153	301	1,620	0.28
9	203	127	69	2.14	2.74	9	183	119	1,160	0.26
Median	301	241	199	1.15	2.74	Median	153	418	1,620	0.26
E_{4/4}					E_{4/4}					
10	273	320	249	0.00	2.38	10	83	415	2,490	0.20
11	180	208	170	0.01	2.28	11	69	299	1,750	0.21
12	480	175	120	2.89	2.54	12	58	305	2,588	0.14
13	173	92	57	0.46	4.14	13	41	214	1,274	0.20
14	332	146	79	1.15	4.90	14	65	360	1,770	0.24
Median	273	175	120	0.46	2.54	Median	65	304	1,770	0.20 [‡]

* Differences between groups were assessed by the Mann-Whitney test. Significantly different from E₃ group, † P < 0.02. § Direct synthesis in LDL was calculated as the difference between the total absolute catabolic rate (observed mass × overall FCR) and the input from VLDL and IDL.

terol and triglyceride in the three groups of this study (Table I), although the E₂ homozygotes displayed the characteristic dyslipidemia associated with this phenotype, i.e., cholesterol-

enriched VLDL and low LDL levels. Apo B levels in this group were low (Table II) despite the similarity in total plasma cholesterol in E₂ and E₃ subjects owing to the higher cholesterol to

apo B ratios in VLDL₁ and VLDL₂ (Table III) and higher concentrations of VLDL₂ and IDL particles in the circulation (Table II) of the former group.

Kinetics studies using tracers of radioiodinated VLDL₁ and VLDL₂ revealed marked differences in the metabolic behavior of apo B in E₂ and E₃ homozygotes and more subtle distinctions between the E₃ and E₄ groups. For comparative purposes, E₃ is taken as the norm since it represents the commonest variant in the population. The decay curves obtained for the four lipoprotein fractions had the general features observed in previous investigations (13, 18). Only a minor modification to the model (compartment 12, Fig. 1) was required to accommodate VLDL₁ apo B kinetics in the E₂ group. This model developed for normal and hypertriglyceridemic subjects has been used successfully to explain apo B kinetics in other genetically determined dyslipidemic conditions such as homozygous familial hypercholesterolemia (FH) and hepatic lipase deficiency (13, 26).

Apo B metabolism in E₃ homozygotes. In E₃ subjects most apo B entered the system at the level of VLDL₁. This material had two fates, catabolism and transfer to VLDL₂ by delipidation. Direct removal of VLDL₁ apo B from the circulation has been noted in virtually all subjects examined with this technique. The pathway was previously found to be present in normal and homozygous FH subjects (13) and its activity was not affected by a chemical modification of the VLDL₁ tracer designed to prevent the interaction of apo B with LDL receptors (27). These findings appear to eliminate the LDL receptor as the agent responsible for this catabolic route and at present the mechanism of this process is unknown. Previous work indicates that lipoprotein lipase is the agent responsible for the VLDL₁ to VLDL₂ conversion. Tracer studies show delayed catabolism of large VLDL in lipoprotein lipase-deficient subjects (28), whereas VLDL₁ transfer to VLDL₂ occurs at a normal rate in hepatic lipase (26) and LDL receptor (13) deficiency. When the enzyme is deficient (i.e. in type I hyperlipidemia [29]) or inhibited in its action (30), large triglyceride-rich VLDL accumulate. Apo B entering VLDL₂ by direct synthesis and by lipolysis of VLDL₁ in E₃ subjects was rapidly and efficiently converted to IDL and LDL (Table IV b); a trivial amount entered the VLDL₂ remnant pool (compartment 6, Table V c). About half of the apo B passing through the delipidation cascade from VLDL₂ to LDL was lost by direct catabolism probably via LDL receptors. These removal mechanisms are inhibited in homozygous FH (13) and blocked by chemical modification of apo B (27). LDL apo B was divided into two metabolically distinct pools to account for the observation that apo B derived from VLDL₂ appeared and was cleared more rapidly than material derived from VLDL₁ (13, 18).

Apo B metabolism in E₂ homozygotes. The metabolism of VLDL₁ apo B differed in E₂ compared with E₃ subjects in three aspects. First, the decay curve was biexponential probably as a result of the presence of chylomicron remnants. It has been shown that subjects homozygous or heterozygous for the apo E₂ phenotype clear chylomicrons slowly and therefore remnants are likely to be present even after a 12-h fast (9, 10). Remnant accumulation is a hallmark of type III hyperlipidemia, and we noted that the VLDL₁ apo B curves observed in the present E₂ group were qualitatively similar to those found previously in type III subjects (18). The slowly metabolized species in the VLDL₁ fraction in the present study was esti-

mated at 10% of the apo B mass but direct quantitation of B48 vs. B100 in the tracer (a measure of chylomicron remnant content) was not performed. Secondly, the rate of VLDL₁ apo B synthesis was significantly reduced in comparison to the E₃ subjects. The reason for this is not clear although it is possible that impaired delivery of chylomicron lipid to the liver in E₂ homozygotes causes that organ subsequently to secrete less triglyceride-rich VLDL. Thirdly, there was less direct catabolism of VLDL₁ apo B. This pathway accounted for 25% of VLDL₁ clearance compared with 65% in E₃ subjects (Table IV a). As mentioned above this unknown mechanism of apo B catabolism operates in most normal and hyperlipidemic subjects. Its decreased activity in E₂ homozygotes indicates that functional apo E may be required for it to proceed efficiently and raises the possibility that a receptor may be involved. It is unlikely that the classical LDL receptor has a substantial role in this regard since the pathway works as normal in FH (13). There is evidence that the recently discovered LDL receptor-related protein has apo E-binding properties (31) and is a candidate agent for mediating chylomicron remnant removal. If very large triglyceride-rich VLDL were catabolized in a similar fashion to chylomicrons, then this would explain the decreased removal in E₂ homozygotes. Indeed, *in vitro* apo E₂ shows impairment in binding to the LDL receptor-related protein (32).

Apo B synthesis into VLDL₂ whether *de novo* or from VLDL₁ occurred at the same rate in E₂ and E₃ subjects (Table IV b). More apo B passing through the delipidation chain in E₂ homozygotes was diverted to remnants and the expansion of this presumably cholesteryl ester-rich population of particles in VLDL₂ to 13% of total apo B mass (compared to < 2% in E₃ subjects, Table V c) explains the abnormal composition of this lipoprotein fraction in the E₂ group (Table II). Surprisingly the catabolic rate of VLDL₂ remnants [L(9, 6) and L(0, 6), Table V] was not consistently reduced in our E₂ subjects. IDL formation and clearance occurred at approximately the same rate in E₂ and E₃ subjects. However, the metabolic fate of apo B differed greatly between them with a 66% reduction in the rate of LDL formation in the former. This difference was the principal cause of reduced LDL levels in E₂ homozygotes. The data in Tables IV and V indicate that in E₂ subjects the impaired IDL to LDL conversion is balanced by increased catabolism of the intermediate fraction: an observation that suggests that functional apo E is not required for effective IDL clearance. Direct removal of IDL probably occurs via LDL receptors since it is inhibited by 1,2-cyclohexanedione modification of the lipoprotein (27) and is depressed in homozygous FH (13). In fact, there was no inhibition of VLDL₂, IDL or LDL clearance in our E₂ subjects suggesting either that it is apo B that acts as the ligand for receptors in these lipoproteins fractions (as is found *in vitro* [33, 34]) or that an increase in receptor activity more than compensates for the reduced affinity of apo E₂ for the receptor. The reason why direct IDL catabolism is favored over conversion to LDL in E₂ homozygotes is unknown. The phenomenon was recorded previously in our study of the effects of bezafibrate on apo B metabolism in type III hyperlipidemic subjects (35). These were apo E₂ homozygotes with elevated lipid levels that were corrected by bezafibrate therapy. The mean on-therapy concentrations of plasma cholesterol, triglyceride, VLDL, LDL, and HDL cholesterol were 5.87, 2.12, 1.63, 2.79, and 1.46 mmol/liter, respectively; values close to those seen in our normolipemic E₂ group (Table I). Apo B kinetics in the type III

Table V. Computed Masses and Rate Constants in E_2 , E_3 , and E_4 Subjects

a. VL DL_1	L(0,1)*	L(2,1)	L(0,12)	L(6,12)	M(1)	M(12)	U(1) [§]	U(12)			
	d^{-1}				<i>mg</i>						
$E_{2/2}$											
1	1.60	3.23	0.69	0.33	27	10	130	10			
2	2.78	5.36	0.00	0.95	48	6	390	6			
3	0.00	4.61	0.19	0.00	81	4	373	1			
4	1.94	5.27	0.00	1.78	46	4	331	7			
Median	1.77	4.94	0.10	0.64	47	5	352	6.5			
$E_{3/3}$											
5	4.17	5.16	—	—	67	—	625	—			
6	5.75	4.49	—	—	51	—	522	—			
7	10.79	2.48	—	—	97	—	1288	—			
8	6.91	3.66	—	—	106	—	1120	—			
9	14.73	2.71	—	—	47	—	819	—			
Median	6.91	3.66	—	—	67	—	819	—			
$E_{4/4}$											
10	5.36	2.99	—	—	107	—	894	—			
11	2.53	1.98	—	—	105	—	474	—			
12	7.56	3.88	—	—	45	—	515	—			
13	16.80	7.68	—	—	12	—	294	—			
14	0.00	9.12	—	—	16	—	146	—			
Median	5.36	3.88	—	—	45	—	474	—			
b. VL DL_2	L(4,2)	L(6,2)	L(9,6)	L(0,6)	L(0,4)	L(8,4)	L(11,4)	L(9,4)	L(0,5)	L(7,5)	L(10,5)
	d^{-1}										
$E_{2/2}$											
1	8.40	0.67	0.03	0.36	0.00	0.70	0.00	0.30	0.00	3.77	0.00
2	8.40	0.33	0.13	0.30	0.00	1.23	0.14	0.02	10.97	17.38	0.00
3	8.40	0.65	0.68	0.00	1.99	1.70	0.14	0.00	0.00	7.75	0.00
4	7.48	0.24	0.40	0.00	0.18	1.04	0.00	0.30	10.14	11.13	0.00
Median	8.40	0.49	0.27	0.15	0.09	1.14	0.07	0.01	5.07	9.44	0.00
$E_{3/3}$											
5	6.22	0.03	0.56	0.00	0.00	5.76	0.00	0.90	0.56	5.45	0.00
6	7.16	0.02	0.41	0.00	1.92	2.22	0.00	0.60	2.72	2.42	0.00
7	6.96	0.13	0.76	0.00	0.00	3.14	0.68	0.00	1.87	3.23	0.00
8	8.92	0.09	0.31	0.32	1.63	2.19	0.08	0.00	1.24	2.37	0.39
9	7.03	0.10	0.00	0.90	0.33	10.35	0.00	1.26	3.65	1.56	0.00
Median	7.03	0.02	0.41	0.00	0.33	3.14	0.00	0.60	1.87	2.37	0.00
$E_{4/4}$											
10	8.40	0.03	0.38	0.00	0.00	2.44	0.00	0.00	0.00	3.54	0.00
11	3.89	0.01	0.00	0.28	0.00	1.84	0.00	0.44	0.00	7.51	0.00
12	8.64	0.40	0.34	0.39	1.94	0.87	0.06	0.00	7.12	7.87	0.00
13	8.59	0.07	0.26	0.14	0.78	1.64	0.02	0.28	0.02	14.40	0.00
14	8.10	0.04	0.00	0.52	2.56	0.51	0.47	0.58	0.00	13.26	0.00
Median	8.40	0.04	0.26	0.28	0.78	1.64 [§]	0.02	0.28	0.00	7.87	0.00
c. VL DL_2	M(2)	M(4)	M(6)	M(5)	U(5)						
	<i>mg</i>										
$E_{2/2}$											
1	10	81	35	99	373						
2	29	177	23	8	238						
3	41	91	41	43	333						
4	31	153	18	11	234						
Median	30	122	29	27	284						
$E_{3/3}$											
5	56	52	3	50	301						
6	32	48	3	148	761						
7	34	62	6	98	500						
8	43	99	6	70	280						
9	18	11	2	39	203						
Median	34	52	3	70	301						
$E_{4/4}$											
10	38	172	3	77	273						
11	53	91	2	24	180						
12	19	58	10	32	480						
13	10	32	2	12	173						
14	18	35	2	25	332						
Median	18	58	2	25	273						

Table V. (Continued)

d. IDL	L(10,7)	L(0,7)	L(0,8)	L(11,8)	L(0,9)	M(7)	M(8)	M(9)
			d^{-1}				<i>mg</i>	
E_{2/2}								
1	1.89	0.34	4.97	0.42	0.34	167	10	75
2	0.06	1.27	1.24	0.00	0.16	110	176	44
3	0.77	0.60	0.91	0.04	0.43	242	161	64
4	0.26	0.54	3.23	0.60	0.46	150	41	118
Median	0.52	0.57 [‡]	2.24	0.23 [‡]	0.39	159	101	69
E_{3/3}								
5	0.00	4.09	1.65	2.04	0.52	66	81	94
6	0.00	1.39	0.80	1.10	0.32	256	56	95
7	0.00	1.58	0.61	0.70	0.37	199	148	11
8	0.00	1.44	0.49	0.44	0.38	114	231	5
9	0.00	0.80	1.21	1.39	0.24	76	42	56
Median	0.00	1.44	0.80	1.10	0.37	114	81	56
E_{4/4}								
10	0.00	1.53	0.69	0.55	0.19	178	260	5
11	0.00	2.03	0.58	1.41	0.39	89	84	103
12	0.00	1.11	0.00	0.63	0.20	225	81	18
13	0.00	2.55	0.37	0.66	0.20	70	52	49
14	0.00	2.69	0.30	0.74	0.29	123	17	69
Median	0.00	2.03	0.37	0.66	0.20	123	81	49
e. LDL		L(0,10)		L(0,11)		M(10)		M(11)
			d^{-1}				<i>mg</i>	
E_{2/2}								
1		0.14		0.15		391		30
2		0.41		0.29		341		88
3		0.32		0.13		447		154
4		0.28		0.17		296		145
Median		0.30		0.16		365		117
E_{3/3}								
5		0.42		0.21		640		793
6		0.30		0.15		1,206		417
7		0.28		0.16		1,124		920
8		0.32		0.23		606		485
9		0.28		0.25		219		231
Median		0.30		0.21		606		485
E_{4/4}								
10		0.20		0.18		1,136		782
11		0.22		0.20		819		592
12		0.14		0.15		1,830		353
13		0.21		0.17		828		199
14		0.24		0.22		1,406		132
Median		0.21 [†]		0.18		1,136		353

* Rate constants, L, k (destination, source) and masses, M(). ‡ U() represents de novo synthesis of apo B into a compartment. † Significantly different from E₃ group, P < 0.05, †† P < 0.02, ††† P < 0.01. Difference between groups was assessed by the Mann-Whitney test.

Table VI. Comparison of Apo B Turnover in Normolipemic E₂ Homozygotes and Type III Hyperlipidemic Subjects before and during Bezafibrate Therapy

Subjects	VLDL ₁			VLDL ₂			IDL			LDL		
	Total production	Plasma concn.	FCR	Total production	Plasma concn.	FCR	Total production	Plasma concn.	FCR	Total production	Plasma concn.	FCR
	<i>mg/d</i>	<i>mg/dl</i>	<i>pool/d</i>	<i>mg/d</i>	<i>mg/dl</i>	<i>pools/d</i>	<i>mg/d</i>	<i>mg/dl</i>	<i>pools/d</i>	<i>mg/d</i>	<i>mg/dl</i>	<i>pools/d</i>
Type III* (n = 6)	885	15.9	2.0	1125	30.8	1.2	630	13.7	1.3	439	21	0.21
Type III on bezafibrate	385	2.9	4.5	783	14.6	1.7	568	14.0	1.2	138	17	0.33
E _{2/2} normolipemic	356	2.6	6.2	534	8.6	2.3	413	13.1	1.1	135	21	0.26

* Data on type III hyperlipidemic subjects were taken from reference 35.

subjects before and during bezafibrate treatment are presented in Table VI. There is a remarkable similarity between treated type III and normolipemic E₂ subjects in both the apo B content of lipoprotein fractions and metabolic behavior indicating that bezafibrate removes the hyperlipidemia component from the type III pattern but does not correct the abnormality owing to E₂ genetic variant. Interestingly, before drug therapy the type III patients converted 76% of IDL to LDL (similar to the E₃ subjects in Table IV), whereas on treatment this fell to 26%, i.e., close to the value seen in normolipemic E₂ subjects. Thus bezafibrate had two effects; it reduced VLDL synthesis correcting the hyperlipidaemia (but not the dysbetalipoproteinemia) and altered the nature of the IDL to LDL metabolic link from that seen in E₃ to that seen in E₂ subjects.

There was no increase in the fractional clearance rate of LDL apo B in E₂ homozygotes in the present study (Table IV *d*). This contrasts with an earlier preliminary report by Gregg et al. (12) in which a small increase in LDL FCR was observed in two normolipemic E₂ homozygotes. However, further work by the same authors in an apo E-deficient patient demonstrated a normal catabolic rate for autologous LDL apo B (36, 37). The low LDL apo B level observed in that condition was due to reduced synthesis at 20% of normal and in general LDL kinetics in apo E deficiency were similar to those reported here for E₂ subjects (Table IV *d*).

Apo B metabolism in E₄ homozygotes. Apo B synthesis in VLDL₁ was lower in E₄ than E₃ subjects (Table IV *a*) but the difference was not significant. The circulating mass of apo B and clearance rate of the fraction were similar in the two groups. Likewise, VLDL₂ apo B synthesis, pool size, and overall catabolic rate were similar in E₃ and E₄ subjects as was the turnover of IDL (Table IV). However, calculation of the amount of apo B undergoing direct catabolism from VLDL₂ and IDL compared to that channeled down the delipidation cascade revealed that E₄ homozygotes exhibited a relative decrease in direct removal and, per milligram of apo B entering VLDL₂, directed more towards LDL production (Table IV). Thus, despite the fact the E₄ subjects synthesized apo B in VLDL (VLDL₁ plus VLDL₂) at only two-thirds of the rate seen in E₃ homozygotes (Table IV *a* and *b*) they had apo B levels that were at least as high as those seen in the latter group (Table II). LDL levels in E₄ subjects were further increased by the low FCR seen for apo B in this flotation interval (Tables IV *d* and *V e*). This was due to a decrease in the rate of apo B degradation from the larger of the two LDL compartments (compartment 10, Table V *e*) with the result that both LDL pools were cleared at the same rate. This is consistent with a reduced receptor-mediated LDL clearance in E₄ subjects. In fact, the concentration and distribution of apo B-containing lipoproteins in E₄ homozygotes appears to be the result of suppressed receptor-mediated catabolism of VLDL₂, IDL, and LDL. It is likely that the low synthetic rate for apo B observed in our E₄ group was a result of patient selection. E₄ homozygotes with an apo B synthetic rate similar to that seen in the E₃ group (800 mg/d) would be predicted to have a plasma apo B level of about 150 mg/dl and a plasma cholesterol in excess of 7.0 mmol/liter.

Influence of apo E on apo B metabolism. It is clear from the above that variation in the apo E gene has a profound impact on apo B metabolism throughout the S_r 0–400 lipoprotein spectrum. This information can be integrated with other studies on cholesterol metabolism (38) and chylomicron kinetics (9, 10) to generate an overall picture of the influence of the

polymorphism which refines and in places corrects the model previously suggested by Davignon et al. (3). Cholesterol absorption from the gut is reportedly higher in E₄ vs. E₃ vs. E₂ subjects. This together with differential clearance rate of chylomicron remnants (E₄ > E₃ > E₂; references 9 and 10) will affect the delivery of intestinal (including dietary) cholesterol to the liver. The result is a predicted decrease in the hepatic sterol pool in E₂ vs. E₃ and E₄ subjects which gives rise to the observed graded difference (E₂ > E₃ > E₄) in the rates of cholesterol and bile acid production (38, 39).

According to current concepts changes in the liver cholesterol pool will alter hepatic LDL receptor activity. The latter is predicted to be higher in E₂ compared to E₃ subjects and they in turn will be higher than E₄ subjects. Since we have found previously that LDL receptors play a role in VLDL₂, IDL, and LDL metabolism (13, 27), this mechanism explains why E₂ homozygotes exhibit more and E₄ homozygotes less direct catabolism of VLDL₂ and IDL than E₃ subjects. In this scenario LDL removal should be increased in E₂ homozygotes but this was not observed in our group (Table IV *d*) or in an apo E-deficient patient (37) possibly because the small amount of LDL that is present is abnormal, having a conformation of apo B that fails to recognize receptors efficiently. Evidence to support this hypothesis comes from studies of the kinetic behavior of LDL from apo E-deficient or E₂ homozygous subjects in normals; it is cleared more slowly than autologous LDL (12, 37). In contrast, normal LDL given to an apo E-deficient patient is cleared rapidly as would be the case if LDL receptors were up-regulated (37). The redirection of the metabolic fate of IDL from conversion to LDL to direct catabolism in E₂ homozygotes may be further influenced by a requirement for functional apo E for lipolysis to proceed efficiently. In vitro studies suggest that VLDL from E₂ subjects is not a good substrate for lipase and lipolysis of this lipoprotein does not lead to the formation of LDL (40). Addition of apo E₃ enhances the reaction and LDL is formed. Retarded delipidation would provide a mechanism for the formation of VLDL₂ remnants and the very low conversion of IDL to LDL in our E₂ subjects. This integrated model, which is now modified and considerably strengthened by the availability of detailed apo B kinetic data, can be used to help explain why individuals with varying apo E phenotypes respond differently to diet and drug therapies (41, 42).

Acknowledgments

The authors thank Patricia Price for her excellent secretarial help.

This work was supported by grants from the British Heart Foundation (87/6 and 89/109). Dr. Demant was the recipient of a scholarship from Stiftung Volkswagen, Hanover, Federal Republic of Germany.

References

1. Utermann, G. 1975. Isolation and partial characterization of an arginine-rich apolipoprotein from human plasma very low density lipoproteins: apolipoprotein E. *Hoppe-Seyler's Z. Physiol. Chem.* 356:1113–1121.
2. Zannis, V. I., and J. L. Breslow. 1981. Human very low density lipoprotein apolipoprotein E isoprotein polymorphism is explained by genetic variation and post-translational modification. *Biochemistry.* 20:1033–1041.
3. Davignon, J., R. E. Gregg, and C. F. Sing. 1988. Apolipoprotein E polymorphism and atherosclerosis. *Arteriosclerosis.* 8:1–21.
4. Mahley, R. W., T. L. Innerarity, S. C. Rall, and K. H. Weisgraber. 1984. Plasma lipoproteins: apolipoprotein structure and function. *J. Lipid Res.* 25:1277–1294.

5. Menzel, H. J., and G. Utermann. 1986. Apolipoprotein E phenotyping from serum by Western blotting. *Electrophoresis*. 7:492-495.
6. Havekes, L. M., P. de Knijff, U. Beisiegel, J. Havinga, M. Smit, and E. Klases. 1987. A rapid micro-method for apolipoprotein E phenotyping directly in serum. *J. Lipid Res.* 28:455-463.
7. Weisgraber, K. H., T. L. Innerarity, and R. W. Mahley. 1982. Abnormal lipoprotein receptor-binding activity of the human E apolipoprotein due to cysteine-arginine interchange at a single site. *J. Biol. Chem.* 257:2518-2521.
8. Utermann, G., M. Hees, and A. Steinmetz. 1977. Polymorphism of apolipoprotein E and occurrence of dysbetalipoproteinaemia in man. *Nature (Lond.)*. 269:604-607.
9. Weintraub, M. S., S. Eisenberg, and J. L. Breslow. 1987. Dietary fat clearance in normal subjects is regulated by genetic variation in apolipoprotein E. *J. Clin. Invest.* 80:1571-1577.
10. Brenninkmeijer, B. J., P. M. J. Stuyt, P. N. M. Demacker, A. F. H. Stalenhoef, and A. van't Larr. 1987. Catabolism of chylomicron remnants in normolipidaemic subjects in relation to the apoprotein E phenotype. *J. Lipid Res.* 28:361-370.
11. Turner, P. R., C. Cortese, R. Wootton, C. Marenah, N. E. Miller, and B. Lewis. 1985. Plasma apolipoprotein B metabolism in familial type III dysbetalipoproteinaemia. *Eur. J. Clin. Invest.* 15:100-112.
12. Gregg, R. E., L. A. Zech, C. Gabelli, D. Stark, D. Wilson, and H. B. Brewer, Jr. 1984. LDL metabolism in normolipidemic apoE₂ homozygotes. *Circulation*. 70:II-312. (Abstr.)
13. James, R. W., B. Martin, D. Pometta, J. C. Fruchart, P. Duriez, P. Puchois, J. P. Farreux, A. Tacquet, T. Demant, R. J. Clegg, et al. 1989. Apolipoprotein B metabolism in homozygous familial hypercholesterolemia. *J. Lipid Res.* 30:159-169.
14. Zannis, V. I., J. L. Breslow, G. Utermann, R. W. Mahley, K. H. Weisgraber, R. J. Havel, J. L. Goldstein, M. S. Brown, G. Schonfeld, W. R. Hazzard, et al. 1982. Proposed nomenclature of apoE isoproteins, apoE genotypes, and phenotypes. *J. Lipid Res.* 23:911-914.
15. Paul, A. A., and D. A. T. Southgate, editors. 1978. McCance and Widdowson's: The Composition of Food. Her Majesty's Stationery Office, London.
16. Lipid Research Clinics Program Manual of Laboratory Operations. 1975. DHEW Publications, No. (NIH) 75-268. Government Printing Office, Washington, DC.
17. Towbin, H., T. Staehelin, and J. Gordon. 1979. Electrophoretic transfer of proteins from polyacrylamide gels to nitrocellulose sheets: procedure and some applications. *Proc. Natl. Acad. Sci. USA*. 76:4350-4354.
18. Packard, C. J., A. Munro, A. R. Lorimer, A. M. Gotto, and J. Shepherd. 1984. Metabolism of apolipoprotein B in large triglyceride-rich very low density lipoproteins of normal and hypertriglyceridemic subjects. *J. Clin. Invest.* 74:2178-2192.
19. Bilheimer, D. W., S. Eisenberg, and R. I. Levy. 1982. The metabolism of very low density lipoprotein proteins. I. Preliminary in vitro and in vivo observations. *Biochim. Biophys. Acta*. 260:212-221.
20. Lindgren, F. T., C. L. Jensen, and F. T. Hatch. 1972. The isolation and quantitative analysis of serum lipoproteins. In *Blood Lipids and Lipoproteins: Quantitation, Composition and Metabolism*. G. J. Nelson, editor. Wiley-Interscience, New York. 221-245.
21. Kane, J. P., T. Sata, R. L. Hamilton, and R. J. Havel. 1975. Apoprotein composition of very low density lipoproteins of human serum. *J. Clin. Invest.* 56:1622-1634.
22. Lowry, O. H., N. J. Rosenbrough, A. L. Farr, and R. J. Randall. 1951. Protein measurement with the folin phenol reagent. *J. Biol. Chem.* 193:265-275.
23. Shepherd, J., C. J. Packard, S. M. Grundy, D. Yeshurun, A. M. Gotto, and O. D. Taunton. 1980. Effects of saturated and polyunsaturated fat diets on the chemical composition and metabolism of low density lipoproteins in man. *J. Lipid Res.* 21:91-99.
24. Berman, M., and M. F. Weiss. 1974. SAAM Manual. U. S. Public Health Service Publication No. 1703. U. S. Government Printing Office, Washington, DC.
25. Berman, M., M. Hall III, R. I. Levy, S. Eisenberg, D. W. Bilheimer, R. D. Phair, and R. M. Goebel. 1978. Metabolism of apoB and apoC lipoproteins in man: kinetic studies in normal and hyperlipoproteinemic subjects. *J. Lipid Res.* 19:38-56.
26. Demant, T., L. A. Carlson, L. Holmquist, F. Karpe, P. Nilsson-Ehle, C. J. Packard, and J. Shepherd. 1988. Lipoprotein metabolism in hepatic lipase deficiency: studies on the turnover of apolipoprotein B and on the effect of hepatic lipase on high density lipoprotein. *J. Lipid Res.* 29:1603-1611.
27. Packard, C. J., D. E. Boag, R. Clegg, D. Bedford, and J. Shepherd. 1985. Effects of 1,2-cyclohexanedione modification on the metabolism of very low density lipoprotein apolipoprotein B: potential role of receptors in intermediate density lipoprotein catabolism. *J. Lipid Res.* 26:1058-1066.
28. Stalenhoef, A. F., M. J. Malloy, J. P. Kane, and R. J. Havel. 1984. Metabolism of apolipoproteins B48 and B100 of triglyceride rich lipoproteins in normal and lipoprotein lipase deficient humans. *Proc. Natl. Acad. Sci. USA*. 81:1837-1843.
29. Nikkila, E. A. 1983. Familial lipoprotein lipase deficiency and related disorders of chylomicron metabolism. In *Disorders of Lipoprotein and Lipid Metabolism*. J. B. Stanbury, J. B. Wyngaarden, D. S. Fredrickson, J. L. Goldstein, and M. S. Brown, editors. McGraw-Hill Book Co., Inc., New York. 622-642.
30. Goldberg, I. J., N. A. Le, H. N. Ginsberg, R. M. Krauss, and F. T. Lindgren. 1988. Lipoprotein metabolism during acute inhibition of lipoprotein lipase in the cynomolgus monkey. *J. Clin. Invest.* 81:561-568.
31. Herz, J., U. Hamann, S. Rogne, O. Myklebost, H. Gausepohl, and K. K. Stanley. 1988. Surface location and high affinity for calcium of a 500 kD liver membrane protein closely related to the LDL-receptor suggest a physiological role as lipoprotein receptor. *EMBO (Eur. Mol. Biol. Organ.) J.* 7:4119-4127.
32. Kowal, R., J. Herz, K. H. Weisgraber, R. W. Mahley, M. S. Brown, and J. L. Goldstein. 1990. Opposing effects of apolipoproteins E and C on lipoprotein binding to low density lipoprotein receptor-related protein. *J. Biol. Chem.* 265:10771-10779.
33. Bradley, W., S.-L. Hwang, J. B. Karlin, A. H. Y. Lin, S. C. Prasad, A. M. Gotto, and S. H. Gianturco. 1984. Low density lipoprotein receptor-binding determinants switch from apolipoprotein E to apolipoprotein B during conversion of hypertriglyceridaemic very low density lipoprotein to low density lipoproteins. *J. Biol. Chem.* 259:14728-14735.
34. Krul, E. S., M. J. Tikkanen, T. G. Cole, J. M. Davie, and G. Schonfeld. 1985. Roles of apolipoprotein B and E in the cellular binding of very low density lipoproteins. *J. Clin. Invest.* 75:361-369.
35. Packard, C. J., R. J. Clegg, M. H. Dominiczak, A. R. Lorimer, and J. Shepherd. 1986. Effects of bezafibrate on apolipoprotein B metabolism in type III hyperlipoproteinemic subjects. *J. Lipid Res.* 27:930-938.
36. Schaefer, E. J., R. E. Gregg, G. Ghiselli, T. M. Forte, J. M. Ordovas, L. A. Zech, and M. B. Brewer. 1986. Familial apolipoprotein E deficiency. *J. Clin. Invest.* 78:1206-1219.
37. Gabelli, C., R. E. Gregg, L. A. Zech, E. Manzato, and H. B. Brewer, Jr. 1986. Abnormal low density lipoprotein metabolism in apolipoprotein E deficiency. *J. Lipid Res.* 27:326-333.
38. Kesaniemi, Y. A., C. Ehnholm, and T. A. Miettinen. 1987. Intestinal cholesterol absorption efficiency in man is related to apoprotein E phenotype. *J. Clin. Invest.* 80:578-581.
39. Angelin, B., L. Holmquist, B. Leijd, and K. Einarsson. 1990. Bile acid metabolism in familial dysbetalipoproteinaemia: studies in subjects with the apolipoprotein E₂ phenotype. *Eur. J. Clin. Invest.* 20:143-149.
40. Ehnholm, C., R. W. Mahley, D. A. Chappell, K. H. Weisgraber, and J. L. Witztum. 1984. Role of apolipoprotein E in the lipolytic conversion of β -very low density lipoproteins to low density lipoproteins in type III hyperlipoproteinemia. *Proc. Natl. Acad. Sci. USA*. 81:5566-5570.
41. Eto, M., T. Sato, K. Watanabe, Y. Iwashima, and I. Makino. 1990. Effects of probucol on plasma lipids and lipoproteins in familial hypercholesterolaemic patients with and without apolipoprotein E₄. *Atherosclerosis*. 84:49-53.
42. Tikkanen, M. J., J. K. Huttunen, C. Ehnholm, and P. Pietinen. 1990. Apolipoprotein E₄ homozygosity predisposes to serum cholesterol elevation during high fat diet. *Arteriosclerosis*. 10:285-288.