

An *in vivo* RNAi assay identifies major genetic and cellular requirements for primary piRNA biogenesis in *Drosophila*

Daniel Olivieri¹, Martina M Sykora¹,
Ravi Sachidanandam², Karl Mechtler^{1,3}
and Julius Brennecke^{1,*}

¹Institute of Molecular Biotechnology of the Austrian Academy of Sciences (IMBA), Vienna, Austria, ²Department of Genetics and Genomic Sciences, Mount Sinai School of Medicine, New York, NY, USA and ³Institute of Molecular Pathology (IMP), Vienna, Austria

In *Drosophila*, PIWI proteins and bound PIWI-interacting RNAs (piRNAs) form the core of a small RNA-mediated defense system against selfish genetic elements. Within germline cells, piRNAs are processed from piRNA clusters and transposons to be loaded into Piwi/Aubergine/AGO3 and a subset of piRNAs undergoes target-dependent amplification. In contrast, gonadal somatic support cells express only Piwi, lack signs of piRNA amplification and exhibit primary piRNA biogenesis from piRNA clusters. Neither piRNA processing/loading nor Piwi-mediated target silencing is understood at the genetic, cellular or molecular level. We developed an *in vivo* RNAi assay for the somatic piRNA pathway and identified the RNA helicase Armitage, the Tudor domain containing RNA helicase Yb and the putative nuclease Zucchini as essential factors for primary piRNA biogenesis. Lack of any of these proteins leads to transposon de-silencing, to a collapse in piRNA levels and to a failure in Piwi-nuclear accumulation. We show that Armitage and Yb interact physically and co-localize in cytoplasmic Yb bodies, which flank P bodies. Loss of Zucchini leads to an accumulation of Piwi and Armitage in Yb bodies, indicating that Yb bodies are sites of primary piRNA biogenesis.

The EMBO Journal (2010) 29, 3301–3317. doi:10.1038/emboj.2010.212; Published online 3 September 2010

Subject Categories: RNA; development

Keywords: *Drosophila*; piRNA pathway; Piwi; transposon silencing

Introduction

Selfish genetic elements such as transposons populate all eukaryotic genomes. Though in a few cases the host benefits from transposons, they overall have a negative impact on the host's reproductive fitness because of their mutagenic character (Slotkin and Martienssen, 2007).

Genetic studies on *Drosophila melanogaster* illustrate the threat emanating from active transposons. Here, uncontrolled

activity of a single transposable element (e.g. the *I*-element, a retro-element or the *P*-element, a DNA-element) leads to defects in gametogenesis and sterility (Rubin *et al*, 1982; Bucheton *et al*, 1984). Given that the *D. melanogaster* genome harbours over a 100 different transposon families (Bergman *et al*, 2006), many of which are still active, a strong selective pressure to silence transposons must exist.

After the discovery of small RNA-silencing pathways, it has become clear that this regulatory mechanism is at the root of transposon control in animals and that different lineages deploy the basic principle of small RNA pathways in different ways to guarantee specific and efficient silencing of selfish genetic elements (reviewed in Girard and Hannon, 2008; Malone and Hannon, 2009).

Although transposon control is important for all cells, their silencing is of pivotal importance in the germline, the only cell lineage that passes its genetic information onto the next generation. Indeed, multi-cellular animals possess a unique small RNA pathway targeted towards silencing selfish genetic elements in their gonads. At the centre of this pathway is a subclass of Argonaute proteins, the so-called PIWI proteins, complexed with 23–30 nt long PIWI-interacting RNAs (piRNAs). Mutations in PIWI proteins result in strong de-repression of transposable elements and lead to widespread defects in gametogenesis and sterility (reviewed in Klattenhoff and Theurkauf, 2008; Malone and Hannon, 2009).

The piRNA pathway is best understood in *Drosophila*, where insight from three research areas has accumulated. First, studies on transposon biology have uncovered important concepts of the host control system and have also identified a number of essential genetic loci involved (Prud'homme *et al*, 1995; Aravin *et al*, 2001; Jensen *et al*, 2002; Desset *et al*, 2003; Ronsseray *et al*, 2003; Sarot *et al*, 2004; Pelisson *et al*, 2007). Second, sterility and egg patterning screens have uncovered more than a dozen factors that turned out to be piRNA pathway members (Schupbach and Wieschaus, 1991; Clegg *et al*, 1997; Lin and Spradling, 1997; Cook *et al*, 2004; Chen *et al*, 2007; Pane *et al*, 2007). And finally, research centred on small RNA pathways and bioinformatics analysis of piRNA populations has allowed combining the diverse genetic findings into a coherent model of transposon control in *Drosophila* (Aravin *et al*, 2003; Vagin *et al*, 2006; Brennecke *et al*, 2007; Gunawardane *et al*, 2007). According to this, PIWI proteins are loaded with piRNAs in a Dicer and presumably double-stranded RNA (dsRNA)-independent manner (Vagin *et al*, 2006). Most piRNAs originate from transposon- and other repeat regions in the genome (Aravin *et al*, 2003; Saito *et al*, 2006; Vagin *et al*, 2006; Brennecke *et al*, 2007; Gunawardane *et al*, 2007). A hallmark of the piRNA pathway is that discrete genomic loci (piRNA clusters) that harbour a diverse collection of transposon and repeat fragments are major sources of piRNAs (Brennecke

*Corresponding author. Institute of Molecular Biotechnology of the Austrian Academy of Sciences, IMBA, Dr Bohrgasse 3, Vienna 1030, Austria. Tel.: +43 179 044 4805; Fax: +43 17 904 4110; E-mail: julius.brennecke@imba.oew.ac.at

Received: 22 July 2010; accepted: 10 August 2010; published online: 3 September 2010

et al, 2007). Another unique feature of the piRNA pathway is the involvement of PIWI proteins in a target-dependent amplification loop (Brennecke *et al*, 2007; Gunawardane *et al*, 2007). In this so-called ping-pong cycle, the PIWI proteins Aubergine and AGO3 cleave reciprocally sense (from active elements) and antisense (from piRNA clusters) transcripts, respectively. It is postulated that each cleavage event triggers the production of a novel piRNA from the cleaved RNA. Thus, the ping-pong cycle leads to a preferential amplification of silencing competent piRNAs. Gonad-specific activity of the piRNA pathway, repeat enriched piRNA clusters and signatures of the ping-pong cycle are all conserved features in vertebrates (reviewed in Malone and Hannon, 2009).

Major open questions are how the cell distinguishes transcripts from piRNA clusters, transposons and endogenous genes and how piRNA biogenesis and loading into PIWI proteins is controlled. Equally unclear is how PIWI-piRNA complexes silence the array of selfish genetic elements in the genome. Various studies indicate that post-transcriptional control through target slicing and degradation and transcriptional control through guiding DNA and/or chromatin modifications both have an active function (Carmell *et al*, 2007; Klenov *et al*, 2007; Kuramochi-Miyagawa *et al*, 2008; Lim *et al*, 2009).

In *Drosophila*, germline cells and somatic support cells of male and female gonads possess a functional piRNA pathway. However, in somatic support cells, only one of the three PIWI family proteins (Piwi) is expressed and a simplified pathway (the somatic piRNA pathway) lacking the ping-pong cycle is active (Sarot *et al*, 2004; Pelisson *et al*, 2007; Lau *et al*, 2009; Li *et al*, 2009; Malone *et al*, 2009; Saito *et al*, 2009). Deep sequencing of small RNA populations indicates that single-stranded RNAs from a subset of piRNA clusters and to a lower extent also from over a 1000 genes are processed into piRNAs (Lau *et al*, 2009; Malone *et al*, 2009; Robine *et al*, 2009). The most prominent piRNA cluster in somatic support cells is the genetically identified *flamenco* or *COM* locus (Prud'homme *et al*, 1995; Desset *et al*, 2003). It contains an exceptionally high density of transposon fragments, nearly all of which belong to the *gypsy* family of retro-elements. As nearly all fragments are oriented antisense to the transcription direction, piRNA processing from *flamenco* yields almost exclusively antisense piRNAs (Pelisson *et al*, 2007; Brennecke *et al*, 2007; Malone *et al*, 2009). Many *gypsy* family transposons expressed in somatic support cells encode functional *gag*, *pol* and *env* genes. In the absence of efficient silencing, they form viral particles that invade the neighbouring oocyte potentially through cellular transport vesicles (Pelisson *et al*, 1994; Chalvet *et al*, 1999; Leblanc *et al*, 2000; Brassat *et al*, 2006).

The somatic cells of the *Drosophila* gonad are the only described cell type with an active piRNA pathway that lacks the ping-pong cycle. These cells are, therefore, ideally suited for genetic and biochemical approaches towards elucidating the core concepts of the piRNA pathway. Here, we describe a genetic assay that allows for testing any gene of interest for its involvement in the somatic piRNA pathway. We use it to identify the three evolutionarily conserved proteins Armitage (Armi), Zucchini (Zuc) and Yb (King *et al*, 2001; Cook *et al*, 2004; Pane *et al*, 2007) as essential components of primary piRNA biogenesis in *Drosophila* and link the Yb body, a

previously identified cellular structure (Szakmary *et al*, 2009) to piRNA biogenesis.

Results

An in vivo assay for the identification of novel piRNA pathway components

To facilitate the identification of novel somatic piRNA pathway members, we constructed a tester fly line that could be combined with the Vienna *Drosophila* RNAi collection (VDRC) (Dietzl *et al*, 2007). The VDRC library consists of Gal4-inducible UAST lines allowing expression of dsRNA hairpins against >90% of the *Drosophila* genes. The tester line combines a GAL4 driver specific for somatic support cells with a sensor capable of monitoring the integrity of the somatic piRNA pathway.

We identified the previously reported *traffic jam* GAL4 line (*tj*-GAL4) as an optimal driver for RNAi in somatic support cells (Tanentzapf *et al*, 2007). GAL4 expression mirrored the restricted expression of the host gene *tj* to gonadal somatic support cells when crossed to UAST-GFP (Figure 1A) (Li *et al*, 2003). GFP signal was only detected in somatic support cells of the ovary including somatic stem cells in the germarium. To determine the tissue specificity of *tj*-GAL4, we crossed the line to UAST-Luciferase and measured reporter activity in whole females, dissected ovaries and carcasses (flies with manually removed ovaries). The *tj*-GAL4 driver induced Luciferase signal ~50-fold and on average, 94% of this signal originated from ovaries (Figure 1B).

Crosses to a broad spectrum of RNAi lines targeting essential genes resulted in missing or rudimentary ovaries with no impact on viability (Supplementary Table S1). This is significant, as it has been reported that a number of strong follicle cell GAL4 drivers were incapable of inducing efficient RNAi (Zhu and Stein, 2004). Importantly, *tj*-GAL4-driven RNAi against *piwi*, the only undisputed member of the somatic piRNA pathway, resulted in ovaries that phenocopied *piwi* mutant ovaries (Figure 1C) (Cox *et al*, 2000). A known target of the somatic piRNA pathway, the retroviral *gypsy* transposon (Sarot *et al*, 2004), was de-repressed in these ovaries with a strong accumulation of the *gypsy* Env protein at the follicle cell cortex (Figure 1D). No staining by the *gypsy* Env antibody was detectable in wild-type ovaries and staining was restricted to patches of cells in which *piwi* RNAi was induced clonally (Supplementary Figure S1). Quantitative RT-PCR analysis further indicated that besides *gypsy*, the retro-elements *ZAM* and *Tabor*, two additional targets of the somatic piRNA pathway (Desset *et al*, 2003; Li *et al*, 2009; Malone *et al*, 2009), are strongly de-silenced in *piwi* RNAi crosses compared with *aubergine* RNAi crosses (*aub* shows no detectable expression in follicle cells) (Figure 1E). Notably, RNA levels of the telomeric retro-elements *HeT-A* and *TART*, which are sensitive to defects in the germline piRNA pathway (Vagin *et al*, 2006; Klattenhoff *et al*, 2009), were unaffected. These results are in line with the abundance of piRNAs against *gypsy*, *ZAM* and *Tabor* and the low level of piRNAs against *HeT-A* and *TART* in cultured ovarian somatic cells (OSCs), which are derived from somatic support cells (Supplementary Figure S2) (Niki *et al*, 2006; Lau *et al*, 2009).

We combined the *tj*-GAL4 driver with a *gypsy*-lacZ reporter that in flies carrying a restrictive allele of the *flamenco* cluster is fully silenced by the somatic piRNA pathway (Sarot *et al*,





Figure 1 An *in vivo* RNAi-based assay for the somatic piRNA pathway. **(A)** Cartoon depicting the two major cell types in the ovary with somatic cells in green and germline cells in beige. The *tj-GAL4* driver is expressed at all developmental stages in somatic support cells (labelled with Tj) as judged by a cross to UAS-GFP. **(B)** Relative Luciferase reporter activity is shown from flies of the indicated genotypes (left) and from ovaries and carcasses of *tj-GAL4* > UAS-Luc flies. **(C)** Morphology of ovaries from flies expressing dsRNA against *piwi* and *aub* through *tj-GAL4* compared with ovaries from *piwi* mutants. **(D)** Immuno-staining of the retroviral *gypsy* Env protein (green) in ovaries with *tj-GAL4* induced Piwi knockdown. The lower panels show a zoom onto the follicular epithelium. Actin (red) marks the cell cortex. **(E)** Shown are changes in steady-state RNA levels ($n = 3$) of the endogenous retro-elements *ZAM* and *Tabor* upon Piwi (red) or Aubergine (black) knockdown in comparison with *tj-GAL4* / + flies. The germline-controlled *HeT-A* and *TART* elements serve as controls. **(F)** Shown is the *gypsy-lacZ* reporter in restrictive and permissive *flamenco* backgrounds (left) and in a restrictive background upon RNAi against *aub* and *piwi* (two independent hairpins). β -Gal staining (blue) is strongest in the columnar follicle epithelium of stages 9–10 egg chambers.



gypsy sensor with *tj*-GAL-4 driven RNAi against indicated gene (restrictive *flam.* background)

Figure 2 Comprehensive analysis of the involvement of piRNA pathway members in the somatic pathway. Shown are β -Gal stainings as readout for *gypsy* silencing for genetically identified piRNA pathway genes in which a VDR line was available using *gypsy*-lacZ and *tj*-GAL4 in a restrictive *flamenco* background.

2004). Lack of the somatic piRNA pathway or homozygosis for a permissive *flamenco* allele strongly de-silences β -Gal expression (Figure 1F) (Sarot *et al*, 2004). When the *tj*-GAL4; *gypsy*-lacZ line was crossed to an *aub*-RNAi line, no β -Gal signal was detectable. In contrast, crosses to two independent *piwi* RNAi lines resulted in strong β -Gal activity (Figure 1F). Thus, given the near genome-wide collection of RNAi lines at the VDR, almost any gene can be tested for its involvement in the somatic piRNA pathway with a single genetic cross.

Armitage and Zucchini are essential components of the somatic piRNA pathway

Over a dozen genes have been implicated in the *Drosophila* piRNA pathway (Boswell and Mahowald, 1985; Gillespie and Berg, 1995; Cox *et al*, 1998; Harris and Macdonald, 2001; Cook *et al*, 2004; Vagin *et al*, 2004; Anne *et al*, 2007; Chen *et al*, 2007; Horwich *et al*, 2007; Lim and Kai, 2007; Pane *et al*, 2007; Saito *et al*, 2007; Klattenhoff *et al*, 2009; Li *et al*, 2009; Malone *et al*, 2009; Patil and Kai, 2010). To test their involvement in the somatic piRNA pathway, we

crossed the *tj*-GAL4; *gypsy*-lacZ tester strain to RNAi lines for all candidates if a VDR line was available. Most candidates are seemingly not essential components of the somatic piRNA pathway (Figure 2). Consistent with this, loss of *spindle-E*, *squash*, *aubergine*, *rhino* or *vasa* does not affect *flamenco*-derived piRNA levels, indicating their specific involvement in the germline-specific ping-pong amplification cycle (Klattenhoff *et al*, 2009; Malone *et al*, 2009). In addition, loss of the piRNA methyl-transferase Hen1 has been reported to impact transposon silencing only very mildly (Horwich *et al*, 2007). It is of interest to note that lack of symmetrically methylated Arginines (mediated by knocking down the arginine methyl-transferase *capsuleen* or its cofactor *valois*) did not lead to detectable sensor activation. Thus, this post-translational modification of PIWI proteins (Kirino *et al*, 2009; Nishida *et al*, 2009; Vagin *et al*, 2009) has no or only a modulatory function in the somatic piRNA pathway, although we cannot exclude the involvement of other arginine methyl transferases.

In contrast, the *gypsy* sensor was strongly de-silenced in the cross to the *zucchini* (*zuc*) RNAi line (Figure 2). This

is in agreement with published findings that loss of *Zuc* affects *flamenco*-derived piRNA levels as well as those derived from the *tj*-3'UTR, a gene expressed in somatic support cells only (Malone *et al*, 2009; Robine *et al*, 2009; Saito *et al*, 2009).

To our surprise, knockdown of *armitage* (*armi*) led to a robust de-silencing of the sensor (Figure 2). This contradicts earlier findings that *armi* mutations specifically affect the germline piRNA pathway (Malone *et al*, 2009): in *armi*[1]/*armi*[72.1] mutants, Piwi-bound piRNAs and nuclear localization of Piwi were only lost in germline cells. Furthermore, *flamenco*-derived piRNA levels were unchanged in those mutants. In support of the previous genetic results, the *gypsy-lacZ* sensor was not de-silenced in follicle cells of *armi*[1]/*armi*[72.1] ovaries (Figure 3A). In these ovaries, we observed weak β -Gal activity in nurse cells, indicating a slight activity of the *gypsy* promoter and an *Armi*-dependent silencing mechanism in ovarian germline cells.

The *armi*[1] and *armi*[72.1] alleles are not null alleles as noted previously (Cook *et al*, 2004). Nevertheless, they cause a severe reduction of *Armi* protein in germline cells. No difference in immuno-staining, however, has been detected in somatic follicle cells of these mutants, which has been interpreted as antibody cross-reactivity (Cook *et al*, 2004). The following observations, however, indicate that *Armi* is expressed in follicle cells and that it is an essential component of the somatic piRNA pathway. We first verified our knockdown result with an independent RNAi line (Figure 3A). Second, knockdown of *Armi* led to a strong de-silencing of the *ZAM* and *Tabor* retro-elements with no impact on *HeT-A* and *TART* (Figure 3B). Similar results were obtained from flies expressing the *zuc* RNAi construct. We next verified robust *Armi* expression in somatic support cells by three independent means: first, a transgenic fly line expressing *Armi* as a C-terminal GFP fusion under its own promoter showed robust GFP signal in germline and somatic cells of the ovary, similar to the published immuno-stainings (Figure 3D) (Cook *et al*, 2004). Second, the follicle cell staining of *Armi* was specifically lost in clones of cells expressing the *armi* RNAi construct (Figure 3E). Third, western analysis showed that ovary lysate contains two *Armi* isoforms that appear as a duplet and that both isoforms are also present in a lysate from OSCs (cultured somatic cells) (Figure 3C). To test for the specificity of the western signal, we generated a genomic deletion that removes the entire *armi* locus including two downstream genes (*armi*[Δ 1]). Homozygous *armi*[Δ 1] flies are viable, contain malformed ovaries and do not lay eggs. Extracts from these ovaries lack both *Armi* isoforms (Figure 3C).

This indicates that the two published alleles *armi*[1] and *armi*[72.1] are not null alleles at the protein level. Indeed, both *Armi* isoforms were detected by western analysis in *armi*[1]/*armi*[72.1] flies albeit at ratios that resemble that found in OSCs (Figure 3C). Furthermore, *Armi* protein was detectable in follicle cells only of *armi*[1]/*armi*[Δ 1] and *armi*[72.1]/*armi*[Δ 1] mutant ovaries (Figure 3D). We, therefore, suspected that the *armi* locus contains two promoters and, therefore, gives rise to two distinct mRNAs encoding the two observed protein isoforms. Indeed, two mRNA isoforms have been observed by northern analysis with only the larger one being severely affected by the published alleles (Cook *et al*, 2004). We performed 5'RACE analysis on OSC RNA and

identified two distinct mRNA 5' ends with one corresponding to the Flybase annotated transcription start site (TSS) (Figure 3F). The second TSS is a few nucleotides downstream of the annotated ATG start codon and, therefore, is predicted to generate an ~4 kDa smaller isoform. In this model, the insertion site of the P-element in *armi*[1] and the small deletion in the *armi*[72.1] allele, which was derived from *armi*[1] by imprecise excision are expected to affect only the longer isoform. Although the 5'RACE analysis indicates that both promoters are active in follicle cells, a western analysis from lysate obtained from 0 to 0.5 h old embryos identified only the larger isoform, indicating that the downstream promoter is not active in germline cells (Figure 3C). Our results, therefore, offer a consistent explanation for the discrepancy of the published literature with our result that *Armi* is an essential component of the somatic piRNA pathway.

***Yb* co-localizes with *Armitage* in cytoplasmic foci and is essential for the somatic piRNA pathway**

Both GFP tagged and endogenous *Armi* localized to the cytoplasm of follicle cells and OSCs with a strong enrichment in 1–3 peri-nuclear foci (Figure 4A and B). Within the ovary, *Armi* foci were much more prominent in the germarium and younger egg chambers and became progressively smaller from stage 5–6 onwards (not shown). At this stage, follicle cells switch from a mitotic cycle to an endo-replication cycle in a Notch-signalling-dependent manner (Deng *et al*, 2001). Interestingly, disruption of the Notch-signalling pathway by RNAi against Notch led to a strong increase in *Armi* levels in older (>stage 6), but not in younger egg chambers (Figure 4C; not shown), suggesting that Notch signalling dampens *Armi* expression in endo-replicating cells.

The distinct localization of *Armi* in follicle cells was reminiscent of processing bodies (P bodies), cytoplasmic foci involved in RNA degradation and storage (Eulalio *et al*, 2007a). Co-labelling of *Armi* and the P-body marker DCP1 indicated that *Armi* foci are in most cases directly adjacent to P bodies (Figure 4D). There appear to be many more P bodies than *Armi* foci, but typically the most prominent P bodies flank *Armi* foci. In germline cells, we observed that areas with *Armi* staining typically exhibit also strong DCP1 labelling (Supplementary Figure S3A). The significance of this correlation remains to be determined, but we note that knockdown of two critical P-body components (*Lsm1* and *Me31b*) with *tj*-GAL4 did not lead to defects in *gypsy* sensor silencing. Moreover, RNAi flip-out clones for *Lsm1* did not affect number or integrity of the *Armi* foci (not shown).

The subcellular localization of *Armi* resembles that of the *Yb* protein. *Yb* is encoded by the *fs(1)Yb* gene and is a putative RNA helicase that contains a predicted Tudor domain (King and Lin, 1999; Szakmary *et al*, 2009). *Yb* is expressed in somatic support cells of the male and female gonad and accumulates in cytoplasmic foci that have been termed *Yb* bodies (Szakmary *et al*, 2009). *Yb* mutants are female sterile and exhibit defects in the maintenance of germline and somatic stem cells (King and Lin, 1999; King *et al*, 2001). Immuno-fluorescence experiments indicated that *Armi* and *Yb* proteins co-localize and accumulate in *Yb* bodies of follicle cells (Figure 4E) and OSCs (Supplementary Figure S3B). In addition, a specific interaction between the two proteins was observed in co-IP experiments from OSC lysate using FLAG-tagged *Yb* (Figure 4F). To test for a




Figure 3 Armitage is an essential player in the somatic piRNA pathway. (A) *gypsy-lacZ* reporter activity in *armi[1]/armi[72.1]* ovaries compared with ovaries in which *armi* was knocked down using *tj-GAL4*-mediated RNAi. (B) Shown are fold changes in steady-state RNA levels ($n = 3$) for the *ZAM*, *Tabor*, *HeT-A* and *TART* retro-elements in ovaries with *tj-GAL4*-driven RNAi against *armi*, *zuc* and *aub* (control). (C) Western blot analysis showing two Armi isoforms expressed in ovaries and OSCs, one isoform in early embryos, persistent expression of Armi in published mutant alleles and the absence of Armi in the *armi[Δ1]* allele. (D) Shown are Armi stainings of an ovariole expressing Armi-GFP under its endogenous promoter (left) and of ovarioles *trans*-heterozygous for the published alleles *armi[1]* and *armi[72.1]* over the *armi[Δ1]* allele. (E) Armi staining (red) of the follicular epithelium in which *armi*-RNAi has been clonally activated. GFP signal (green) labels cells expressing the *armi* dsRNA hairpin. (F) Cartoon of the *armi* genomic locus showing the extent of the *armi[Δ1]* deletion, the two identified 5'RACE products (red), their respective transcription start sites and the insertion site of the P-element in the *armi[1]* allele. To the left, an Agarose gel with the two sequenced RACE products is shown.

functional involvement of *Yb* in the somatic piRNA pathway, we crossed two independent *fs(1)Yb* RNAi lines to our tester strain and observed strong de-repression of the *gypsy* reporter

(Figure 4G). Depletion of *Yb* also led to a de-silencing of the *ZAM* and *Tabor* retro-transposons, but not the germline-controlled *HeT-A* and *TART* elements (Figure 4H). These




Figure 4 Armi localizes to Yb bodies and Yb is an essential pathway component. **(A)** Shown are immuno-fluorescence stainings for Armi (red) and Piwi (green) in somatic follicle cells. DNA was labelled with DAPI (blue). **(B)** Immuno-fluorescence staining for Armi (red), Tubulin (green) and DNA (blue) in OSCs. **(C)** Immuno-fluorescence analysis of Armi (red) in the follicular epithelium of an egg chamber containing a Notch dsRNA-expressing clone (clone marked with GFP (green) and clone borders indicated by the dashed line). DAPI staining (blue) confirms that Notch knockdown impairs endo-replication. **(D)** Co-labelling of Armi (red) and DCP1-GFP (green) in somatic follicle cells. **(E)** Co-labelling of Armi (red) and Yb (green) in somatic follicle cells. **(F)** Co-immunoprecipitation of endogenous Armi with FLAG-tagged Yb. Shown are western blots against FLAG to detect Yb (top) and against endogenous Armi (bottom). Control cells did not express FLAG-Yb. **(G)** Shown are β-Gal stainings of ovarioles expressing the *gypsy* reporter and two different dsRNA hairpins against *fs(1)Yb* with *tj-GAL4*. **(H)** Shown are fold changes in steady-state RNA levels ($n = 3$) of ZAM, Tabor, HeT-A and TART retro-elements in ovaries in which *fs(1)yb* and *aub* (control) were knocked down through *tj-GAL4*-mediated RNAi. Values are normalized to a *tj-GAL4* / + control.

results identified Yb as an essential component of the somatic piRNA pathway and link the previously described Yb bodies to piRNA biology.

Loss of Armitage, Zucchini and Yb impairs nuclear localization of Piwi

To gain insight into the defects exerted onto the somatic piRNA pathway upon loss of Armi, Zuc and Yb, we performed a systematic analysis of Armi, Piwi and Yb localization in cells mutant for each factor (a lack of reagents prevented localization studies for Zuc).

RNAi-mediated knockdown of Yb in follicle cell clones led to a loss of nuclear Piwi accumulation (Figure 5A). Furthermore, no Armi foci were detectable when Yb was depleted by RNAi. Instead, Armi was uniformly distributed throughout the cytoplasm (Figure 5A). Identical results were obtained in FRT-mediated mitotic clones using the *Yb[72]* loss of function allele (Supplementary Figure S4).

RNAi-mediated knockdown of Armi also caused a loss of Piwi from the nucleus (Figure 5B). In addition, Yb bodies were reduced in size, but did not disappear entirely (Figure 5C). We, therefore, analysed Yb localization in *armi* null mutants and observed a loss of detectable Yb bodies in nearly all cells (Supplementary Figure S5). Thus, Armitage and Yb are each required for their reciprocal localization to Yb bodies and potentially for Yb-body formation *per se*.

The most interesting effects were seen upon Zuc knockdown. Here, Piwi was not only lost from the nucleus, but instead accumulated in peri-nuclear foci (Figure 5D). Co-labelling experiments showed that these correspond to Armi bodies (Figure 5D, bottom). In addition, the size of Armi foci increased several fold in Zuc mutant cells (Figure 5D). We also observed elevated intensity of Yb foci in these cells, but the effects were much weaker than those observed for Armi (not shown). Identical results were obtained in FRT-mediated mitotic clones using the *zuc[HM27]* loss of function allele (Supplementary Figure S6). The co-localization of Piwi and Armi in *zuc* mutant cells suggests a rapid transit of Piwi through Armi/Yb bodies in wild-type cells, as we never observed accumulation of endogenous Piwi protein in cytoplasmic foci. However, OSCs chemically transfected with GFP-tagged Piwi often exhibit co-localization of Piwi with Armi foci, potentially as a consequence of Piwi over-expression (Supplementary Figure S7).

To test whether Armi is required for Piwi's accumulation in Yb bodies, we simultaneously knocked down Armi and Zuc in follicle cell clones. In these cells, Piwi did not accumulate in cytoplasmic foci (Supplementary Figure S8). We note, however, that Yb-body size is severely reduced upon Armi knockdown, thus complicating this interpretation. No defects on Armi localization were seen in cells expressing an RNAi construct against Piwi or in cells homozygous for the *piwi[1]* loss of function allele (Figures 5E and 8D). This indicates that Armi does not depend on Piwi for its localization to Yb bodies and that Yb-body formation occurs in the absence of Piwi.

Armitage and Piwi interact physically in OSCs and ovaries

In an attempt to identify Piwi-interacting proteins, we immuno-precipitated endogenous Piwi from OSC lysate using polyclonal antibodies directed against the Piwi N-terminus. Silver staining identified two protein bands at

~140 kDa that were specific for the Piwi-IP compared with a control IP (Figure 6A). Mass spectrometric analysis identified Armi as the major protein in both bands (32 and 27% sequence coverage). This result is at odds with Armi being a cytoplasmic protein, whereas Piwi is enriched in the nucleus under wild-type conditions. However, our observation that Piwi localizes transiently with Armi to Yb bodies offers an explanation for the observed interaction. Western blot analysis indicated that this interaction is also robust in IPs from ovary lysate (Figure 6B). The major interacting Armi isoform in ovaries is the germline-specific larger isoform, whereas the smaller isoform predominates in OSCs. To test the specificity of the Armi Piwi interaction, we immuno-precipitated transfected FLAG-tagged Piwi from OSC lysate. Again, endogenous Armi protein was only detected in the FLAG-Piwi-IP, but not in the FLAG-IP from non-transfected cells (not shown). Finally, endogenous Piwi was co-immuno-precipitated with Armi using flies expressing C-terminally GFP-tagged Armi under its own promoter (not shown). In this experiment, we also noted that Armi forms oligomeric complexes *in vivo*, as endogenous Armi protein was efficiently co-immuno-precipitated with GFP-tagged Armi (Figure 6C).

An immuno-precipitation of Piwi in the presence of RNase A and T1 indicates that the interaction between Armi and Piwi depends at least in part on RNA (Figure 6D). Although we cannot exclude incomplete RNA digestion, we suggest that the interaction between Armi and Piwi is direct, but enhanced by RNA. Such a direct interaction is supported by a co-immuno-precipitation of mouse PIWI proteins and the mouse Armi orthologue MOV10L1 in 293 cells that contain no active piRNA pathway (Zheng *et al*, 2010).

Armitage, Zucchini and Yb are required for primary piRNA biogenesis

Loss of Armi in germline cells leads to a failure of nuclear Piwi accumulation accompanied by a loss of germline-specific Piwi-bound piRNAs (Malone *et al*, 2009). A correlation between nuclear localization and intact small RNA biogenesis has also been observed for mouse MIWI2 and *Tetrahymena* Twi1 (Aravin *et al*, 2008; Noto *et al*, 2010; Zheng *et al*, 2010). We sequenced Piwi-bound piRNAs from ovaries in which we depleted Piwi, Armi, Zuc or Yb in all ovarian somatic support cells by RNAi using the *tj-GAL4* driver. RNAi against *aub* was used as a control. An immuno-fluorescence analysis showed the efficiency of the *tj-GAL4*-driven knockdown in follicle cells as they lacked Piwi-nuclear accumulation (Figure 7A). We prepared ovary lysates and immuno-precipitated Piwi-piRNA complexes. From these, small RNAs were isolated and used for the construction of cDNA libraries, which were sequenced on the Illumina G2 platform. After removal of small RNAs mapping to abundant cellular non-coding transcripts (rRNAs, tRNAs, snoRNAs), the composition of all libraries was highly similar with ~84% mapping to annotated repeats (Supplementary Table S2). As Piwi function within germline cells was unaffected, we used the germline-specific piRNA cluster at position 42AB to normalize the individual libraries. Three other peri-centromeric piRNA clusters (*cluster 20A*, *cluster 38C*, *cluster 80E*) with predominant or exclusive piRNA processing in the germline (Lau *et al*, 2009; Malone *et al*, 2009) showed similar piRNA levels across all libraries (Figure 7B). In contrast, piRNAs




Figure 5 Comprehensive analysis of Armi, Piwi and Yb localization upon their reciprocal knockdown. (A–E) Shown are surface views of the follicular epithelium of egg chambers containing clones expressing dsRNA hairpins against the indicated genes stained for Piwi (blue), Armi (red) and Yb (red). Clones are marked by the presence of GFP (green) and clone borders are indicated (dashed line). In all rows, the merged RGB image is shown alongside with the individual channels in black and white. In (D), the lower row shows higher magnification views.

uniquely mapping to the *flamenco* locus, a piRNA cluster that is exclusively or predominantly processed into piRNAs in somatic support cells, were 6- to 10-fold reduced in *piwi*, *armi*, *zuc* and *fs(1)Yb* knockdown ovaries. The *tj*-3'UTR is another source for

abundant piRNAs in somatic support cells (Robine *et al*, 2009; Saito *et al*, 2009). Levels of *tj*-derived piRNAs also decreased 6- to 10-fold in all libraries in comparison with the control (Figure 7B). Figure 7C depicts the normalized piRNA profiles




Figure 6 Piwi and Armitage interact physically. **(A)** SDS-PAGE of a Piwi immunoprecipitation (IP) alongside a control IP from OSC lysate stained with silver. A molecular weight marker is shown to the left and Piwi and Armi bands are indicated. **(B)** Western blot showing co-IP of endogenous Armi with Piwi from OSCs and ovaries. Pre-immune serum was used for the control IP. **(C)** Western blot showing co-IP of endogenous Armi with GFP-tagged Armi from ovary lysate. **(D)** Western blot showing co-IP of endogenous Armi with Piwi in the presence or absence of RNase.

(sense and antisense) originating from *cluster 42AB*, the *flamenco* cluster and the *tj* locus.

We further analysed piRNA levels mapping to transposons, which have been classified as germline dominant (*HeT-A*, *F-element*, *TART*, *GATE*) or soma dominant (*gypsy5*, *ZAM*, *Tabor*, *412*, *Idefix*) (Li *et al*, 2009; Malone *et al*, 2009). Using the same normalization from before (*cluster 42AB*), piRNA levels for the germline-dominant elements were essentially unchanged in all libraries (Supplementary Figure S9A). In contrast, piRNAs mapping to the soma-dominant elements were strongly reduced in number and those remaining had a higher proportion of sense-derived piRNAs, which in wild-type ovaries make up <5% of all *ZAM*- and *Tabor*-specific piRNAs (Supplementary Figure S9A,B). We speculate that the de-silencing of the *ZAM* and *Tabor* elements leads to disproportionate piRNA processing directly from their active sense transcripts.

Our combined results show that primary piRNA biogenesis or loading into Piwi is dependent on Armi, Yb and Zuc in ovarian somatic support cells.

Similarities between somatic and germline Piwi pathways

It has been suggested that Piwi biology in soma and germline of the *Drosophila* ovary differs (Malone *et al*, 2009). This was based, however, on *armi* alleles, which based on our data are germline specific. It has also been reported that *zuc* null mutants exhibit no defects in Piwi-nuclear accumulation (Malone *et al*, 2009). This prompted us to re-investigate the similarities of the somatic and the germline Piwi pathways using null mutants for *armi*, *piwi*, *fs(1)Yb* and *zuc*.

Other than previously reported, *zuc* mutant ovaries exhibited a severe loss of Piwi from the nucleus in somatic and germline cells of the ovary (Figure 8A and B). While Piwi protein accumulated in Yb bodies in follicle cells, it appeared to concentrate in clouds at the nuclear periphery of germline cells (asterisks in Figure 8B). These clouds were also prominently stained for Armi (Figure 8B, right). To test whether this could represent disintegrating nuage, we stained *zuc* mutant ovarioles for Aub and AGO3, two proteins localizing to nuage. No obvious defects in nuage pattern were detect-




Figure 7 piRNA biogenesis defects in somatic cells lacking Armi, Piwi, Zuc and Yb. **(A)** Shown are ovarioles stained for Piwi (green) and Traffic jam (magenta) from flies expressing the indicated dsRNAs with *tj*-GAL4. Note the lack of nuclear Piwi in follicle cells that are marked by Tj (*aub* knockdown serves as negative control). **(B)** Shown are relative levels of genome-unique Piwi-piRNAs mapping to the indicated piRNA clusters in ovaries of the indicated genotypes. Libraries were normalized through the genome-unique mappers to 42AB, a germline-specific piRNA cluster. The *tj*-GAL4 > *aub*-RNAi sample serves as control. The respective raw read numbers from the *tj* > *aub*-RNAi library are indicated above the black bars. **(C)** Detailed piRNA profiles (only genome-unique piRNAs are shown) across the piRNA clusters 42AB and *flamenco* as well as the major genic piRNA precursor *tj*. The colour code is as in **(B)** and the annotated *tj* gene is indicated in blue. The y axis indicates the number of sequences per 200 nt window normalized to 1 million sequenced reads and graphs within a vertical row are shown at the same scale.




Figure 8 Similarities of Piwi biology in soma and germline of the ovary. (A–E) Shown are optical sections through mid stage egg chambers of indicated genotypes (left) stained for DNA (blue), Piwi (green) and Armi (red). Peri-nuclear accumulation of Piwi in germline cells of *zuc* mutant ovaries is marked by asterisks in (B).

able based on Aub localization, but we observed frequent accumulation of AGO3 in cytoplasmic foci (Supplementary Figure S10). Loss of *Zuc* thus leads to a significant co-localization of Piwi and Armi in soma (Yb bodies) and germline (peri-nuclear clouds, distinct from nuage). As expected, loss of Armi de-localized Piwi from the nucleus in germline and soma (Figure 8C). No changes in Armi localization either to Yb bodies in somatic cells or peri-nuclear clouds in germline cells were seen in ovaries lacking Piwi (Figure 8D). These results suggest that Piwi biology is similar in somatic and germline cells of the ovary.

The only protein that acts differentially in the two cell types is Yb. Ovaries mutant for Yb lost nuclear Piwi accumulation in somatic but not in germline cells (Figure 8E).

Similarly, although somatic Armi localization to Yb bodies was lost, the pattern of Armi in germline cells was wild type (Figure 8E). This is consistent with Yb being specifically expressed in somatic cells of the ovary only (Szakmary *et al*, 2009), and raises a question to as whether a redundant protein might serve Yb's function in germline cells (see Discussion).

Our immuno-fluorescence analysis indicated that Piwi levels were reduced in cells mutant for *armi*, *zuc* or *fs(1)Yb*. To test this, we performed western blot analysis using ovary extracts obtained from null mutants for each respective gene (Supplementary Figure S11). This indicated indeed strongly reduced Piwi levels in *armi* and *zuc* mutants, whereas *fs(1)Yb* mutants were less affected, consistent with

it being only important for Piwi biology in the somatic support cells. No changes in Armi levels were observed in *piwi*, *zuc* or *fs(1)Yb* mutants (not shown). We conclude that failure to load Piwi with piRNAs leads to Piwi destabilization and failure in nuclear accumulation.

Discussion

In this study, we present genetic, cell biological and molecular data that shed light on primary piRNA biogenesis in *Drosophila*.

We describe a robust *in vivo* assay that allows the identification of genes with a critical involvement in the somatic piRNA pathway. It takes advantage of the recently constructed genome-wide RNAi library at the VDRC and, therefore, will allow the conduction of a genome-wide RNAi screen. The somatic support cells of the *Drosophila* ovary are the only described cells containing a piRNA pathway without ping-pong cycle. Elucidation of this pathway will, therefore, not only uncover the basic concepts behind the piRNA biogenesis and silencing machineries. It will also simplify the genetic and functional characterization of identified candidates, aided by the availability of the OSC culture system (Niki *et al*, 2006). Similar to any screen system, our assay has drawbacks such as availability of an RNAi line, potential off-target effects or inefficient knockdown. However, an RNAi screen has the major advantage that knockdown of genes essential for other vital functions in the cell might still allow the development of ovaries suitable for analysis, whereas EMS generated null alleles would often prevent this.

We used this assay to assign essential functions to the piRNA pathway proteins Armi and Zuc within the somatic pathway and identified Yb as a novel piRNA pathway gene. Detailed cell biological analyses in conjunction with protein interaction studies and piRNA sequencing efforts place all three factors upstream of the active Piwi-piRNA complex, either in piRNA biogenesis or in piRNA loading into Piwi. All three proteins are required for piRNA accumulation and for the nuclear accumulation of Piwi. The cytoplasmic localization of Armi, Yb and Zuc suggests that piRNA biogenesis/loading occurs in the cytoplasm. This is supported by the observation that N-terminally truncated Piwi that cannot translocate into the nucleus is still loaded with piRNAs (Saito *et al*, 2009). In addition, piRNAs derived from cellular mRNAs typically map to the 3'UTR (Robine *et al*, 2009), indicating that ribosomal association with those mRNAs did precede their funnelling into piRNA biogenesis.

Our data indicate that unloaded Piwi does not translocate into the nucleus. This is highly reminiscent to the situation in mouse and *Tetrahymena*. The mouse PIWI family protein MIWI2 as well as the *Tetrahymena* Twi1 protein accumulate in the cytoplasm under conditions that do not allow piRNA biogenesis or loading into these factors (Aravin *et al*, 2008; Noto *et al*, 2010; Zheng *et al*, 2010). Furthermore, unloaded Piwi appears to be unstable *in vivo*. This effect is most apparent in ovaries lacking Armi and Zuc, two factors that are essential for Piwi-piRNA biogenesis/loading in soma and germline.

Our results further indicate that aspects of piRNA biogenesis and/or piRNA loading take place in discrete peri-nuclear foci that have been previously termed Yb bodies (Szakmary

et al, 2009). The two RNA helicases Armi and Yb localize both to Yb bodies and interact physically. Both proteins are required for their reciprocal localization to cytoplasmic foci and potentially for Yb-body formation *per se*. A strong argument for the importance of Yb bodies in the piRNA pathway stems from the observation that *fs(1)Yb[1]* mutant ovaries also show Piwi loss from the nucleus (not shown). In this allele, a conserved Arginine residue in the Tudor domain is mutated. This still allows Yb protein expression, but blocks Yb-body formation (Szakmary *et al*, 2009). Ultra-structurally, Yb bodies were described as electron dense cytoplasmic spheres that are directly adjacent to an RNA containing electron dense structure (Szakmary *et al*, 2009). Our localization studies with DCP1 indicate that these previously identified neighbouring structures are likely a subset of cellular P bodies. The functional significance of this is unclear, as we do not detect any defects in *gypsy* silencing upon knockdown of two central P-body components LSM1 and Me31b, which have been reported to lead to a dispersal of P bodies in *Drosophila* S2 cells (Eulalio *et al*, 2007b). Nevertheless, the physical neighbourhood of these two bodies is intriguing and is reminiscent of the neighbouring and/or coinciding localization of a set of piRNA pathway proteins and P-body components in mouse germ cells and *Drosophila* nurse cells (Aravin *et al*, 2009; Lim *et al*, 2009).

Of the three identified proteins, Yb is the only factor that seems specific for the Piwi pathway in somatic support cells. This is consistent with genetic analyses showing that germline-specific mutants for *fs(1)Yb* are fertile (King *et al*, 2001). The function of Yb in the somatic support cells is required for maintenance of somatic and indirectly also for germline stem cells (King *et al*, 2001). In this, *fs(1)Yb* mutants strongly resemble *piwi* mutants and indicate once again that the Piwi-dependent piRNA pathway in the somatic support cells is essential for germline stem cell niche maintenance. The specificity of Yb for the somatic cells raises the question, whether there is a related protein with equivalent functions in germline cells. Protein BLAST analysis indeed identified two proteins (CG31755 and CG11133) with similarity to Yb over most of their sequence (Szakmary *et al*, 2009). Similar to Yb, CG31755 and CG11133 also contain a recognizable Tudor domain downstream of their helicase domain. Both proteins are selectively expressed in ovaries and testes (FlyAtlas). These observations indicate that CG31755 and CG11133 could serve Yb's function within the germline. We note that knockdown of these proteins individually does not impact the somatic piRNA pathway (not shown). CG31755 and CG11133 share considerable similarity to mouse TDRD12, a gene with testis and oocyte-specific expression.

The function of Armi as an essential factor in primary piRNA biogenesis is conserved in mouse (Frost *et al*, 2010; Zheng *et al*, 2010). In the absence of the germline-specific orthologue MOV10L1, spermatogenesis is blocked at a similar stage as in MILI mutants and MILI and MIWI2, which is thought to function downstream of MILI in the mouse ping-pong cycle are lacking bound piRNAs. In *Drosophila*, Armi is *per se* not required for the ping-pong cycle (Malone *et al*, 2009), indicating that the function of this RNA helicase is restricted to primary piRNA processing or Piwi loading. Two Armi isoforms are expressed in ovaries with germline cells expressing only the larger one. Although this might suggest functional specialization, we note that the two isoforms show

an identical distribution in a glycerol gradient, interact both with Piwi and show identical subcellular localization in follicle cells based on GFP fusions (data not shown).

Similar to *Armi*, *Zuc* is well conserved in vertebrates. The mouse orthologue *PLD6* is expressed specifically in testes (<http://www.biogps.gnf.org>) consistent with a function in the piRNA pathway. *Zuc* contains a single phospho-lipase D domain, which is predicted to confer endo-nucleolytic activity (Zhao *et al*, 1997). The three important residues for catalysis are conserved in *Zuc* and a point mutation in the active site (*zuc[SG]*; Pane *et al*, 2007) causes a similar delocalization of Piwi from the nucleus as the null mutant (not shown). Our data indicate that *Zuc* function (potentially endo-nucleolytic generation of piRNA 5' or 3' ends) is required to release Piwi and *Armi* from Yb bodies. We further show that *Zuc* is essential for Piwi-piRNA biogenesis. This casts doubts on the rather weak effects of *zuc* mutations on *flamenco* piRNA biogenesis reported previously (Malone *et al*, 2009). Genotyping of *trans*-heterozygous *zuc* mutants (*zuc[HM27]/zuc[Def]*) from stocks balanced over *CyO* is complicated by a dominant wing phenotype of the *zuc[Def]* allele interfering with confident selection against the *CyO* balancer. We suspect that this led to a mix of homozygous and heterozygous ovaries analysed in that study. It is unclear, however, how the defects in Piwi localization went unnoticed in such a situation.

The phenotype of *armi* and *zuc* mutant ovaries indicates that Piwi biology is similar in germline and somatic support cells. We note that the analysis of piRNAs bound by Piwi and AGO3 also showed a participation of Piwi in the germline-specific ping-pong cycle (Brennecke *et al*, 2007; Li *et al*, 2009). However, in *armi* germline mutants, the ping-pong cycle is still active, yet Piwi is not loaded effectively (Malone *et al*, 2009). This suggests that primary piRNA biogenesis through *Zuc/Armi* is the major pathway feeding into Piwi in germline cells.

In summary, we identified the RNA helicases *Armi* and *Yb* and the predicted nuclease *Zuc* as essential components of primary piRNA biogenesis. We further link peri-nuclear Yb bodies to piRNA biogenesis and show that Piwi biology in germline and soma follows a common logic. The described genetic assay system and the availability of the cultured OSC line will be important resources to work towards a mechanistic understanding of piRNA biogenesis and the silencing mechanism, two of the most mysterious open questions in the small RNA field.

Materials and methods

Drosophila stocks

All experiments were performed at 25°C. The used fly strains were *tj-GAL4* (DGRC stock 104055); *flam[restrictive]*; ; *gypsy-lacZ* (Sarot *et al*, 2004); attP2-UAST-Luciferase (Markstein *et al*, 2008); RNAi lines were from the Vienna *Drosophila* RNAi Centre: *armi* (16205GD; 103589KK), *zuc* (48764GD; 110430KK), *piwi* (22235GD; 101658KK), *fs(1)Yb* (25435GD; 110056KK), *tejas* (103264KK); *Notch* (100002KK); no RNAi lines are available at the VDRC for *krimp* and *AGO3*.
YFP-Dcp1 (Lin *et al*, 2006);
armi[1]/TM3 and *armi[72.1]/TM3* (Cook *et al*, 2004);
zuc[HM27]/CyO and *Df(2l)PRL/CyO* (Pane *et al*, 2007);
piwi[1]/CyO and *piwi[2]/CyO* (Lin and Spradling, 1997);
fs(1)Yb[72]/FM6 and *fs(1)Yb[1]/FM6* (Swan *et al*, 2001).

armi[Δ1] is an FRT-mediated deletion based on DGRC stocks e001160 and d07385; it deletes the entire *armi*, *CycJ* and *CG14971* genes;

Armi-GFP flies carry a genomic rescue construct with an EGFP cassette inserted at the C-terminus by bacterial recombineering; act-GAL4 flip-out clones were generated using *y,w,hsflp¹²²*; *act<CD2>GAL4,UAS-GFP*; newly enclosed females were heat shocked 1 h at 37°C and dissected 4–5 days later; FRT-based mitotic clones were based on *hsflp¹²²,FRT19A,ubi-GFP* and *hsflp¹²²;FRT40A* armadillo-lacZ; clones were induced by heat-shocking freshly enclosed females on two consecutive days for 1 h at 37°C; flies were dissected 5 days later.

Luciferase assay

Three female flies, carcasses or pairs of ovaries of the genotype [*tj-GAL4/+*; attP2-UAST-Luciferase/*+*] and three females of the genotype [attP2-UAST-Luciferase] were homogenized in 100 μl of Glo Lysis Buffer (Promega). A total of 20 μl lysate were mixed 1:1 with Steady-Glo Assay Reagent (Promega). Luminescence was measured with a Synergy Plate Reader after 10 min with a sensitivity of 160 and an integration time of 1 s (results were the average of three experiments).

β-Gal stainings

Ovaries from 2- to 3-day-old flies were dissected in PBS, kept on ice (maximum 30 min), fixed in 0.5% glutaraldehyde/PBS at room temperature (RT; 20 min) and rinsed with PBS. The chromogen reaction was run in staining solution (10 mM PBS, 1 mM MgCl₂, 150 mM NaCl, 3 mM potassium ferricyanide, 3 mM potassium ferrocyanide, 0.1% Triton, 0.1% X-Gal) for 2 h at RT.

Immuno-cytochemistry

Ovaries were dissected in PBS on ice, fixed with 4% paraformaldehyde/PBS with 0.2% Triton X-100 at RT for 20 min and rinsed three times with PBT (PBS with 0.2% Triton X-100). After blocking in BBT (PBT with 0.1% BSA) for 2 h at RT, the ovaries were incubated with primary antibodies in BBT overnight at 4°C. After three PBT washes, secondary antibodies (1:500; Molecular Probes) were incubated 3 h at RT.

OSCs were plated on Concanavalin-A-coated coverslips, fixed with 4% paraformaldehyde in PBS (RT, 20 min), permeabilized with 0.1% Triton X-100 in PBS (10 min) and blocked with 1% BSA in PBS.

Primary antibodies used were α-Tj (1:500, rat; Li *et al*, 2003), α-Piwi (1:1000, rabbit), α-Piwi (1:1000, mouse; Saito *et al*, 2006), α-*Armi* (1:1000, rabbit; Cook *et al*, 2004), α-β-Gal (1:50, mouse 40-1a from Developmental Studies Hybridoma Bank), α-Yb (1:1000, rabbit; Szakmary *et al*, 2009), α-gypsy-Env (1:1000, rabbit).

5' RACE

Total RNA from OSCs was used for the 5' RACE determination of *armi* mRNAs (Invitrogen). Primers used were TCCCAGGAATGCT CAAGG (reverse transcription) and CACATCCGACGTGTAGATCCA AG (PCR).

Cell culture

OSCs were cultured as described (Niki *et al*, 2006). OSCs were transfected with the Cell Line Nucleofector kit V (Amaxa Biosystems), selecting the programme T-029 (K Saito, personal communication).

Piwi-IP MS analysis

OSCs were harvested by trypsinization and lysed in two volumes of lysis buffer (10 mM Hepes pH7, 150 mM NaCl, 5 mM MgCl₂ 10% glycerol, 1% Triton X-100, complete protease inhibitors (Roche), 1 mM DTT, 1 mM EDTA, 0.1 mM PMSF), followed by manual disruption using a Wheaton dounce homogenizer and a 10 min centrifugation step at 4 degrees. The pellet was re-extracted twice and supernatants were pooled. A total of 3 mg of total protein were incubated (2 h, 4°C) with peptide purified Piwi antibodies (cross-linked to Dynabeads ProteinG (Invitrogen) using a standard dimethyl pimelimidate cross-linking protocol). Pre-immune serum from the same rabbit was used as control. Beads were washed five times with low detergent wash buffer (10 mM Hepes pH7, 150 mM NaCl, 5 mM MgCl₂ 10% glycerol, 0.1% Triton X-100, 1 mM DTT, 1 mM EDTA, 0.1 mM PMSF), five times with wash buffer (Hepes pH7, 150 mM NaCl, 5 mM MgCl₂, 1 mM DTT, 1 mM EDTA) and once

with 150 mM NaCl. The beads were boiled with SDS sample buffer and analysed by one-dimensional gel electrophoresis and silver staining.

Indicated bands were excised from the gel, processed by a standard in-gel tryptic digest protocol and further analysed by Tandem Mass Spectrometry. Data analysis was with Mascot 2.2.04.

Co-immuno-precipitation

act>3xFLAG-Yb and act>3xFLAG-Piwi constructs were based on the *Drosophila* Gateway collection (<http://www.ciwemb.edu/labs/murphy/Gateway%20vectors.html>) and were transfected in OSCs. Non-transfected OSCs were used as negative control. Per sample, a 10 cm plate of OSCs was harvested 4 days after transfection and lysed in 200 µl lysis buffer (10 mM Tris-HCl pH7.5, 150 mM NaCl, 0.5 mM EDTA, 0.5% Nonidet P40). Lysates were cleared for 10 min at 16 000 g at 4°C and the supernatant incubated for 4 h at 4°C with 7 µg of α-FLAG antibody (SIGMA M2) cross-linked to 30 µl of Invitrogen ProteinG Dynabeads. Beads were washed six times for 10 min with washing buffer (10 mM Tris-HCl pH7.5, 150 mM NaCl, 0.5 mM EDTA) and eluted by boiling in 2 × western blot sample buffer.

Small RNA cloning and analysis

Ovaries were dissected in PBS (on ice). Lysate from ~100 µl ovaries was obtained by extracting them three times in lysis buffer (10 mM Hepes pH7, 150 mM NaCl, 5 mM MgCl₂, 10% glycerol, 1% Triton X-100, complete protease inhibitors (Roche), 1 mM DTT, 1 mM EDTA, 0.1 mM PMSF). Piwi was immuno-precipitated using polyclonal rabbit antiserum and small RNAs were extracted with phenol chloroform. Small RNA libraries were generated as previously described (Brennecke *et al*, 2007) and sequenced using the Illumina G2 platform. Only sequences matching the *Drosophila* genome (release 5; excluding Uextra) 100% were retained. Libraries were normalized through their unique mappers to the 42AB piRNA cluster to allow for cross-comparisons.

Transposon QPCR analysis

First strand cDNA was obtained by using random primers on Trizol extracted total ovarian RNA from 2- to 3-day-old flies. Quantitative PCR was performed using BioRad IQ SYBR Green Super Mix. Steady-state RNA levels were calculated from the threshold cycle for amplification using the 2^{-ΔΔC_T} method (Livak and Schmittgen, 2001). Fold enrichments are in comparison with a *tj-GAL4*/+ control sample. Throughout rp49 was used for the normalization

and comparable results were obtained by using actin5C for normalization. Average levels and standard deviations were calculated from three biological replicates according to Livak and Schmittgen (2001). Control experiments with template dilution indicated that the efficiencies of target amplification and control genes (rp49 and actin5C) were comparable.

Primers for QPCR analysis were

rp49_for: CCGCTTCAAGGGACAGTATCTG
rp49_rev: ATCTCGCGCAGTAAACGC
actin5C_for: AAGTTGCTGCTCTGGTTGTCTG
actin5C_rev: GCCACACGCAGCTCATTGTAG
ZAM_for: CTACGAAATGGCAAGATTAATTCCACTTCC
ZAM_rev: CCGTTTCCTTTATGTCGCGAGTAGCT
Tabor_for: ACGTTGTTTCACGACATTAGCCG
Tabor_rev: GGGTTGGTTCGGATCTGACG
HeT-A_for: CGCGCGGAACCCATCTTCAGA
HeT-A_rev: CGCCGCAGTCGTTTGGTGAGT
TART_for: TTTTCCGGATCCAAGTGAAC
TART_rev: TCTGTCGTCGGAAGTTGTT

Supplementary data

Supplementary data are available at *The EMBO Journal* Online (<http://www.embojournal.org>).

Acknowledgements

We thank members of the Brennecke laboratory for helpful discussions and A Pelisson for valuable comments on the paper. We are grateful to Andreas Sommer for help with deep sequencing. We thank M Siomi, W Theurkauf, H Lin, D Godt, A Pelisson, M Lin, M Milan, T Schüpbach, M Markstein, the Developmental Studies Hybridoma Bank and the VDRC for flies and antibodies. We thank A Pelisson and A Bucheton for generously sharing their unpublished gypsy Env antibody and M Siomi for OSCs. Small RNA libraries are deposited at GEO (accession no. GSE23560, data sets GSM577958 to GSM577962).

Conflict of interest

The authors declare that they have no conflict of interest.

References

- Anne J, Ollo R, Ephrussi A, Mechler BM (2007) Arginine methyltransferase Capsulein is essential for methylation of spliceosomal Sm proteins and germ cell formation in *Drosophila*. *Development* **134**: 137–146
- Aravin AA, Lagos-Quintana M, Yalcin A, Zavolan M, Marks D, Snyder B, Gaasterland T, Meyer J, Tuschl T (2003) The small RNA profile during *Drosophila melanogaster* development. *Dev Cell* **5**: 337–350
- Aravin AA, Naumova NM, Tulin AV, Vagin VV, Rozovsky YM, Gvozdev VA (2001) Double-stranded RNA-mediated silencing of genomic tandem repeats and transposable elements in the *D. melanogaster* germline. *Curr Biol* **11**: 1017–1027
- Aravin AA, Sachidanandam R, Bourc'his D, Schaefer C, Pezic D, Toth KF, Bestor T, Hannon GJ (2008) A piRNA pathway primed by individual transposons is linked to *de novo* DNA methylation in mice. *Mol Cell* **31**: 785–799
- Aravin AA, van der Heijden GW, Castaneda J, Vagin VV, Hannon GJ, Bortvin A (2009) Cytoplasmic compartmentalization of the fetal piRNA pathway in mice. *PLoS Genet* **5**: e1000764
- Bergman CM, Quesneville H, Anxolabehere D, Ashburner M (2006) Recurrent insertion and duplication generate networks of transposable element sequences in the *Drosophila melanogaster* genome. *Genome Biol* **7**: R112
- Boswell RE, Mahowald AP (1985) tudor, a gene required for assembly of the germ plasm in *Drosophila melanogaster*. *Cell* **43**: 97–104
- Braslet E, Taddei AR, Arnaud F, Faye B, Fausto AM, Mazzini M, Giorgi F, Vaury C (2006) Viral particles of the endogenous retrovirus ZAM from *Drosophila melanogaster* use a pre-existing endosome/exosome pathway for transfer to the oocyte. *Retrovirology* **3**: 25
- Brennecke J, Aravin AA, Stark A, Dus M, Kellis M, Sachidanandam R, Hannon GJ (2007) Discrete small RNA-generating loci as master regulators of transposon activity in *Drosophila*. *Cell* **128**: 1089–1103
- Bucheton A, Paro R, Sang HM, Pelisson A, Finnegan DJ (1984) The molecular basis of I-R hybrid dysgenesis in *Drosophila melanogaster*: identification, cloning, and properties of the I factor. *Cell* **38**: 153–163
- Carmell MA, Girard A, van de Kant H, de Rooij D, Bestor T, Hannon GJ (2007) Miwi2 is essential for spermatogenesis and repression of transposons in the mouse male germline. *Dev Cell* **12**: 503–514
- Chalvet F, Teysset L, Terzian C, Prud'homme N, Santamaria P, Bucheton A, Pelisson A (1999) Proviral amplification of the Gypsy endogenous retrovirus of *Drosophila melanogaster* involves env-independent invasion of the female germline. *EMBO J* **18**: 2659–2669
- Chen Y, Pane A, Schupbach T (2007) Cutoff and aubergine mutations result in retrotransposon upregulation and checkpoint activation in *Drosophila*. *Curr Biol* **17**: 637–642
- Clegg NJ, Frost DM, Larkin MK, Subrahmanyam L, Bryant Z, Ruohola-Baker H (1997) maelstrom is required for an early step in the establishment of *Drosophila* oocyte polarity: posterior localization of grk mRNA. *Development* **124**: 4661–4671
- Cook HA, Koppetsch BS, Wu J, Theurkauf WE (2004) The *Drosophila* SDE3 homolog armitage is required for oskar mRNA silencing and embryonic axis specification. *Cell* **116**: 817–829

- Cox DN, Chao A, Baker J, Chang L, Qiao D, Lin H (1998) A novel class of evolutionarily conserved genes defined by piwi are essential for stem cell self-renewal. *Genes Dev* **12**: 3715–3727
- Cox DN, Chao A, Lin H (2000) piwi encodes a nucleoplasmic factor whose activity modulates the number and division rate of germline stem cells. *Development* **127**: 503–514
- Deng WM, Althausen C, Ruohola-Baker H (2001) Notch-Delta signaling induces a transition from mitotic cell cycle to endocycle in *Drosophila* follicle cells. *Development* **128**: 4737–4746
- Desset S, Meignin C, Dastugue B, Vaury C (2003) COM, a heterochromatic locus governing the control of independent endogenous retroviruses from *Drosophila melanogaster*. *Genetics* **164**: 501–509
- Dietzl G, Chen D, Schnorrer F, Su KC, Barinova Y, Fellner M, Gasser B, Kinsey K, Oettel S, Scheiblaue S, Couto A, Marra V, Keleman K, Dickson BJ (2007) A genome-wide transgenic RNAi library for conditional gene inactivation in *Drosophila*. *Nature* **448**: 151–156
- Eulalio A, Behm-Ansmant I, Izaurralde E (2007a) P bodies: at the crossroads of post-transcriptional pathways. *Nat Rev Mol Cell Biol* **8**: 9–22
- Eulalio A, Behm-Ansmant I, Schweizer D, Izaurralde E (2007b) P-body formation is a consequence, not the cause, of RNA-mediated gene silencing. *Mol Cell Biol* **27**: 3970–3981
- Frost RJ, Hamra FK, Richardson JA, Qi X, Bassel-Duby R, Olson EN (2010) MOV10L1 is necessary for protection of spermatocytes against retrotransposons by Piwi-interacting RNAs. *Proc Natl Acad Sci USA* **107**: 11847–11852
- Gillespie DE, Berg CA (1995) Homeless is required for RNA localization in *Drosophila* oogenesis and encodes a new member of the DE-H family of RNA-dependent ATPases. *Genes Dev* **9**: 2495–2508
- Girard A, Hannon GJ (2008) Conserved themes in small-RNA-mediated transposon control. *Trends Cell Biol* **18**: 136–148
- Gunawardane LS, Saito K, Nishida KM, Miyoshi K, Kawamura Y, Nagami T, Siomi H, Siomi MC (2007) A slicer-mediated mechanism for repeat-associated siRNA 5' end formation in *Drosophila*. *Science* **315**: 1587–1590
- Harris AN, Macdonald PM (2001) Aubergine encodes a *Drosophila* polar granule component required for pole cell formation and related to eIF2C. *Development* **128**: 2823–2832
- Horwich MD, Li C, Matranga C, Vagin V, Farley G, Wang P, Zamore PD (2007) The *Drosophila* RNA methyltransferase, DmHen1, modifies germline piRNAs and single-stranded siRNAs in RISC. *Curr Biol* **17**: 1265–1272
- Jensen S, Gassama MP, Dramard X, Heidmann T (2002) Regulation of I-transposon activity in *Drosophila*: evidence for cosuppression of nonhomologous transgenes and possible role of ancestral I-related pericentromeric elements. *Genetics* **162**: 1197–1209
- King FJ, Lin H (1999) Somatic signaling mediated by fs(1)Yb is essential for germline stem cell maintenance during *Drosophila* oogenesis. *Development* **126**: 1833–1844
- King FJ, Szakmary A, Cox DN, Lin H (2001) Yb modulates the divisions of both germline and somatic stem cells through piwi- and hh-mediated mechanisms in the *Drosophila* ovary. *Mol Cell* **7**: 497–508
- Kirino Y, Kim N, de Planell-Saguer M, Khandros E, Chiorean S, Klein PS, Rigoutsos I, Jongens TA, Mourelatos Z (2009) Arginine methylation of Piwi proteins catalysed by dPRMT5 is required for Ago3 and Aub stability. *Nat Cell Biol* **11**: 652–658
- Klattenhoff C, Theurkauf W (2008) Biogenesis and germline functions of piRNAs. *Development* **135**: 3–9
- Klattenhoff C, Xi H, Li C, Lee S, Xu J, Khurana JS, Zhang F, Schultz N, Koppetsch BS, Nowosielska A, Seitz H, Zamore PD, Weng Z, Theurkauf WE (2009) The *Drosophila* HP1 homolog Rhino is required for transposon silencing and piRNA production by dual-strand clusters. *Cell* **138**: 1137–1149
- Klenov MS, Lavrov SA, Stolyarenko AD, Ryazansky SS, Aravin AA, Tuschl T, Gvozdev VA (2007) Repeat-associated siRNAs cause chromatin silencing of retrotransposons in the *Drosophila melanogaster* germline. *Nucleic Acids Res* **35**: 5430–5438
- Kuramochi-Miyagawa S, Watanabe T, Gotoh K, Totoki Y, Toyoda A, Ikawa M, Asada N, Kojima K, Yamaguchi Y, Ijiri TW, Hata K, Li E, Matsuda Y, Kimura T, Okabe M, Sakaki Y, Sasaki H, Nakano T (2008) DNA methylation of retrotransposon genes is regulated by Piwi family members MILI and MIWI2 in murine fetal testes. *Genes Dev* **22**: 908–917
- Lau NC, Robine N, Martin R, Chung WJ, Niki Y, Berezikov E, Lai EC (2009) Abundant primary piRNAs, endo-siRNAs, and microRNAs in a *Drosophila* ovary cell line. *Genome Res* **19**: 1776–1785
- Leblanc P, Desset S, Giorgi F, Taddei AR, Fausto AM, Mazzini M, Dastugue B, Vaury C (2000) Life cycle of an endogenous retrovirus, ZAM, in *Drosophila melanogaster*. *J Virol* **74**: 10658–10669
- Li C, Vagin VV, Lee S, Xu J, Ma S, Xi H, Seitz H, Horwich MD, Syrzycka M, Honda BM, Kittler EL, Zapp ML, Klattenhoff C, Schultz N, Theurkauf WE, Weng Z, Zamore PD (2009) Collapse of germline piRNAs in the absence of Argonaute3 reveals somatic piRNAs in flies. *Cell* **137**: 509–521
- Li MA, Alls JD, Avancini RM, Koo K, Godt D (2003) The large Maf factor Traffic Jam controls gonad morphogenesis in *Drosophila*. *Nat Cell Biol* **5**: 994–1000
- Lim AK, Kai T (2007) Unique germ-line organelle, nuage, functions to repress selfish genetic elements in *Drosophila melanogaster*. *Proc Natl Acad Sci USA* **104**: 6714–6719
- Lim AK, Tao L, Kai T (2009) piRNAs mediate posttranscriptional retroelement silencing and localization to pi-bodies in the *Drosophila* germline. *J Cell Biol* **186**: 333–342
- Lin H, Spradling AC (1997) A novel group of pumilio mutations affects the asymmetric division of germline stem cells in the *Drosophila* ovary. *Development* **124**: 2463–2476
- Lin MD, Fan SJ, Hsu WS, Chou TB (2006) *Drosophila* decapping protein 1, dDcp1, is a component of the oskar mRNP complex and directs its posterior localization in the oocyte. *Dev Cell* **10**: 601–613
- Livak KJ, Schmittgen TD (2001) Analysis of relative gene expression data using real-time quantitative PCR and the 2^{(-Delta Delta C(T))} method. *Methods* **25**: 402–408
- Malone CD, Brennecke J, Dus M, Stark A, McCombie WR, Sachidanandam R, Hannon GJ (2009) Specialized piRNA pathways act in germline and somatic tissues of the *Drosophila* ovary. *Cell* **137**: 522–535
- Malone CD, Hannon GJ (2009) Small RNAs as guardians of the genome. *Cell* **136**: 656–668
- Markstein M, Pitsouli C, Villalta C, Celniker SE, Perrimon N (2008) Exploiting position effects and the gypsy retrovirus insulator to engineer precisely expressed transgenes. *Nat Genet* **40**: 476–483
- Niki Y, Yamaguchi T, Mahowald AP (2006) Establishment of stable cell lines of *Drosophila* germ-line stem cells. *Proc Natl Acad Sci USA* **103**: 16325–16330
- Nishida KM, Okada TN, Kawamura T, Mituyama T, Kawamura Y, Inagaki S, Huang H, Chen D, Kodama T, Siomi H, Siomi MC (2009) Functional involvement of Tudor and dPRMT5 in the piRNA processing pathway in *Drosophila* germlines. *EMBO J* **28**: 3820–3831
- Noto T, Kurth HM, Kataoka K, Aronica L, DeSouza LV, Siu KW, Pearlman RE, Gorovsky MA, Mochizuki K (2010) The Tetrahymena argonaute-binding protein Giw1p directs a mature argonaute-siRNA complex to the nucleus. *Cell* **140**: 692–703
- Pane A, Wehr K, Schupbach T (2007) zucchini and squash encode two putative nucleases required for rasiRNA production in the *Drosophila* germline. *Dev Cell* **12**: 851–862
- Patil VS, Kai T (2010) Repression of retroelements in *Drosophila* germline via piRNA pathway by the Tudor Domain Protein Tejas. *Curr Biol* **20**: 724–730
- Pelisson A, Sarot E, Payen-Groschene G, Bucheton A (2007) A novel rasiRNA-mediated silencing pathway downregulates sense gypsy transcripts in the somatic cells of the *Drosophila* ovary. *J Virol* **81**: 1951–1960
- Pelisson A, Song SU, Prud'homme N, Smith PA, Bucheton A, Corces VG (1994) Gypsy transposition correlates with the production of a retroviral envelope-like protein under the tissue-specific control of the *Drosophila* flamenco gene. *EMBO J* **13**: 4401–4411
- Prud'homme N, Gans M, Masson M, Terzian C, Bucheton A (1995) Flamenco, a gene controlling the gypsy retrovirus of *Drosophila melanogaster*. *Genetics* **139**: 697–711
- Robine N, Lau NC, Balla S, Jin Z, Okamura K, Kuramochi-Miyagawa S, Blower MD, Lai EC (2009) A broadly conserved pathway generates 3'UTR-directed primary piRNAs. *Curr Biol* **19**: 2066–2076
- Ronsseay S, Josse T, Boivin A, Anxolabehere D (2003) Telomeric transgenes and trans-silencing in *Drosophila*. *Genetica* **117**: 327–335
- Rubin GM, Kidwell MG, Bingham PM (1982) The molecular basis of P-M hybrid dysgenesis: the nature of induced mutations. *Cell* **29**: 987–994

- Saito K, Inagaki S, Mituyama T, Kawamura Y, Ono Y, Sakota E, Kotani H, Asai K, Siomi H, Siomi MC (2009) A regulatory circuit for piwi by the large Maf gene traffic jam in *Drosophila*. *Nature* **461**: 1296–1299
- Saito K, Nishida KM, Mori T, Kawamura Y, Miyoshi K, Nagami T, Siomi H, Siomi MC (2006) Specific association of Piwi with rasiRNAs derived from retrotransposon and heterochromatic regions in the *Drosophila* genome. *Genes Dev* **20**: 2214–2222
- Saito K, Sakaguchi Y, Suzuki T, Siomi H, Siomi MC (2007) Pimet, the *Drosophila* homolog of HEN1, mediates 2'-O-methylation of Piwi-interacting RNAs at their 3' ends. *Genes Dev* **21**: 1603–1608
- Sarot E, Payen-Groschene G, Bucheton A, Pelisson A (2004) Evidence for a piwi-dependent RNA silencing of the gypsy endogenous retrovirus by the *Drosophila melanogaster* flamenco gene. *Genetics* **166**: 1313–1321
- Schupbach T, Wieschaus E (1991) Female sterile mutations on the second chromosome of *Drosophila melanogaster*. II. Mutations blocking oogenesis or altering egg morphology. *Genetics* **129**: 1119–1136
- Slotkin RK, Martienssen R (2007) Transposable elements and the epigenetic regulation of the genome. *Nat Rev Genet* **8**: 272–285
- Swan A, Hijal S, Hilfiker A, Suter B (2001) Identification of new X-chromosomal genes required for *Drosophila* oogenesis and novel roles for fs(1)Yb, brainiac and dunce. *Genome Res* **11**: 67–77
- Szakmary A, Reedy M, Qi H, Lin H (2009) The Yb protein defines a novel organelle and regulates male germline stem cell self-renewal in *Drosophila melanogaster*. *J Cell Biol* **185**: 613–627
- Tanentzapf G, Devenport D, Godt D, Brown NH (2007) Integrin-dependent anchoring of a stem-cell niche. *Nat Cell Biol* **9**: 1413–1418
- Vagin VV, Klenov MS, Kalmykova AI, Stolyarenko AD, Kotelnikov RN, Gvozdev VA (2004) The RNA interference proteins and vasa locus are involved in the silencing of retrotransposons in the female germline of *Drosophila melanogaster*. *RNA Biol* **1**: 54–58
- Vagin VV, Sigova A, Li C, Seitz H, Gvozdev V, Zamore PD (2006) A distinct small RNA pathway silences selfish genetic elements in the germline. *Science* **313**: 320–324
- Vagin VV, Wohlschlegel J, Qu J, Jonsson Z, Huang X, Chuma S, Girard A, Sachidanandam R, Hannon GJ, Aravin AA (2009) Proteomic analysis of murine Piwi proteins reveals a role for arginine methylation in specifying interaction with Tudor family members. *Genes Dev* **23**: 1749–1762
- Zhao Y, Stuckey JA, Lohse DL, Dixon JE (1997) Expression, characterization, and crystallization of a member of the novel phospholipase D family of phosphodiesterases. *Protein Sci* **6**: 2655–2658
- Zheng K, Xioli J, Reuter M, Eckardt S, Leu NA, McLaughlin KJ, Stark A, Sachidanandam R, Pillai RS, Wang PJ (2010) Mouse MOV10L1 associates with Piwi proteins and is an essential component of the Piwi-interacting RNA (piRNA) pathway. *Proc Natl Acad Sci USA* **107**: 11841–11846
- Zhu X, Stein D (2004) RNAi-mediated inhibition of gene function in the follicle cell layer of the *Drosophila* ovary. *Genesis* **40**: 101–108