

catena-Poly[[diaquamanganese(II)]- μ_3 -pyridine-2,3-dicarboxylato- $\kappa^4N,O^2;O^3;O^{3'}$] dihydrate]

Zhong-Xiang Du* and Jun-Xia Li

Department of Chemistry and Chemical Engineering, Luoyang Normal University, Luoyang, Henan 471022, People's Republic of China
Correspondence e-mail: dzx101012@126.com


Received 4 September 2008; accepted 13 September 2008

Key indicators: single-crystal X-ray study; $T = 291$ K; mean $\sigma(C-C) = 0.005$ Å; R factor = 0.035; wR factor = 0.095; data-to-parameter ratio = 12.5.

In the title coordination polymer, $\{[Mn(C_7H_3NO_4)(H_2O)_2] \cdot 2H_2O\}_n$, the Mn^{II} ion is coordinated in a distorted octahedral environment by the O atoms of two water molecules, one N and one O atoms of the chelating pyridine-2,3-dicarboxylate (PDC) dianion, and two axial bridging carboxylate O atoms from two adjacent PDC ligands. The fully deprotonated PDC anion acts as a μ_3 -bridging ligand, establishing a chain structure along the a axis. These polymeric chains are connected into a three-dimensional framework via several intermolecular O—H···O hydrogen bonds.

Related literature

For related literature, see: Aghabozorg *et al.* (2007); Baruah *et al.* (2007); Drew *et al.* (1971); Ghoer & Youssef (1993); Kang *et al.* (2006); Li *et al.* (2006); Manteghi *et al.* (2007); Patrick *et al.* (2003); Sun *et al.* (2006); Takusagawa & Koetzle (1978); Turner & Batten (2007); Zhang & You (2003); Zhang *et al.* (2003).



Experimental

Crystal data

[Mn(C ₇ H ₃ NO ₄)(H ₂ O) ₂]·2H ₂ O	$V = 1034.9$ (2) Å ³
$M_r = 292.11$	$Z = 4$
Monoclinic, $P2_1/c$	Mo $K\alpha$ radiation
$a = 6.5719$ (8) Å	$\mu = 1.31$ mm ⁻¹
$b = 7.6703$ (9) Å	$T = 291$ (2) K
$c = 20.566$ (3) Å	$0.38 \times 0.15 \times 0.11$ mm
$\beta = 93.3540$ (10)°	

Data collection

Bruker APEXII CCD area-detector diffractometer	6428 measured reflections
Absorption correction: multi-scan (SADABS; Sheldrick, 1996)	1921 independent reflections
$T_{\min} = 0.639$, $T_{\max} = 0.865$	1774 reflections with $I > 2\sigma(I)$
	$R_{\text{int}} = 0.015$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.035$	3 restraints
$wR(F^2) = 0.095$	H-atom parameters constrained
$S = 1.07$	$\Delta\rho_{\max} = 0.49$ e Å ⁻³
1921 reflections	$\Delta\rho_{\min} = -0.74$ e Å ⁻³
154 parameters	

Table 1
Selected geometric parameters (Å, °).

Mn1—O3	2.098 (3)	Mn1—O5	2.160 (3)
Mn1—O4	2.139 (3)	Mn1—O6 ⁱⁱ	2.242 (3)
Mn1—O8 ⁱ	2.144 (2)	Mn1—N1 ⁱ	2.263 (3)
O3—Mn1—O4	96.07 (11)	O8 ⁱ —Mn1—O5	95.89 (9)
O3—Mn1—O8 ⁱ	168.80 (10)	O3—Mn1—O6 ⁱⁱ	84.63 (10)
O3—Mn1—O5	87.98 (10)	O5—Mn1—O6 ⁱⁱ	164.11 (10)
O4—Mn1—O5	87.54 (10)	O4—Mn1—N1 ⁱ	166.71 (11)

Symmetry codes: (i) $-x + 1, -y + 2, -z + 1$; (ii) $x + 1, y, z$.

Table 2
Hydrogen-bond geometry (Å, °).

D—H···A	D—H	H···A	D···A	D—H···A
O1—H1W···O8	0.84	1.95	2.793 (4)	177
O1—H2W···O2 ⁱⁱⁱ	0.84	2.09	2.845 (4)	149
O2—H4W···O6 ^{iv}	0.84	2.07	2.884 (4)	165
O2—H4W···O4 ^v	0.84	2.56	3.043 (4)	118
O2—H3W···O7 ^{vi}	0.85	1.94	2.788 (4)	180
O3—H5W···O7 ^{iv}	0.84	1.85	2.691 (4)	175
O3—H6W···O1 ^{iv}	0.85	1.92	2.730 (4)	159
O4—H8W···O1 ⁱ	0.85	2.61	3.457 (4)	180
O4—H7W···O1 ^{vii}	0.84	1.86	2.691 (4)	168
O4—H8W···O2 ^{vii}	0.85	2.37	2.788 (4)	111

Symmetry codes: (i) $-x + 1, -y + 2, -z + 1$; (iii) $-x + 1, y + \frac{1}{2}, -z + \frac{3}{2}$; (iv) $-x + 1, -y + 1, -z + 1$; (v) $-x + 2, -y + 1, -z + 1$; (vi) $x, -y + \frac{3}{2}, z + \frac{3}{2}$; (vii) $x, -y + \frac{3}{2}, z - \frac{1}{2}$.

Data collection: APEX2 (Bruker, 2004); cell refinement: APEX2; data reduction: SAINT (Bruker, 2004); program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: SHELXTL (Sheldrick, 2008); software used to prepare material for publication: SHELXTL.

This work was supported by the National Natural Science Foundation of China (grant No. 20471026) and the Natural

metal-organic compounds

Science Foundation of Henan Province (grant No. 0311021200).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: SG2261).

References

- Aghabozorg, H., Daneshvar, S., Motyeian, E., Ghadermazi, M. & Attar Gharamaleki, J. (2007). *Acta Cryst.* **E63**, m2468–m2469.
- Baruah, A. M., Karmakar, A. & Baruah, J. B. (2007). *Polyhedron*, **26**, 4518–4524.
- Bruker (2004). *APEX2* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Drew, M. B., Matthews, R. W. & Walton, R. A. (1971). *J. Chem. Soc. A*, pp. 2959–2962.
- Ghoer, M. S. & Youssef, A. A. (1993). *Polyhedron*, **12**, 1871–1878.
- Kang, Y., Zhang, J., Li, Z. J., Cheng, J. K. & Yao, Y. G. (2006). *Inorg. Chim. Acta*, **359**, 2201–2209.
- Li, M., Xiang, J. F., Yuan, L. J., Wu, S. M., Chen, S. P. & Sun, J. T. (2006). *Cryst. Growth Des.* **6**, 2036–2040.
- Manteghi, F., Ghadermazi, M. & Aghabozorg, H. (2007). *Acta Cryst.* **E63**, o2809.
- Patrick, B. O., Stevens, C. L., Storr, A. & Thompson, R. C. (2003). *Polyhedron*, **22**, 3025–3035.
- Sheldrick, G. M. (1996). *SADABS*. University of Göttingen, Germany.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Sun, L. P., Niu, S. Y., Jin, J., Yang, G. D. & Ye, L. (2006). *Inorg. Chem. Commun.* **9**, 679–682.
- Takusagawa, F. & Koetzle, T. F. (1978). *Acta Cryst.* **B34**, 1149–1154.
- Turner, D. R. & Batten, S. R. (2007). *Acta Cryst.* **E63**, m452–m454.
- Zhang, X.-M., Fang, R.-Q., Wu, H.-S. & Ng, S. W. (2003). *Acta Cryst.* **E59**, m1143–m1145.
- Zhang, H.-T. & You, X.-Z. (2003). *Acta Cryst.* **C59**, m313–m314.

supplementary materials

Acta Cryst. (2008). E64, m1295-m1296 [doi:10.1107/S1600536808029413]

catena-Poly[[[diaquamanganese(II)]- μ_3 -pyridine-2,3-dicarboxylato- $\kappa^4N,O^2;O^3;O^3'$] dihydrate]

Z.-X. Du and J.-X. Li

Comment

Pyridine-2,3-dicarboxylic acid (H₂PDC), being a potential multidentate bridging ligand, has aroused considerable interests in recent decades and a number of metal complexes have been reported. In these complexes, pyridine-2,3-dicarboxylic acid is partly or fully deprotonated and shows diverse coordination modes, such as non-coordinate (Takusagawa & Koetzle, 1978; Manteghi *et al.*, 2007), monodentate (Drew *et al.*, 1971; Ghoer & Youssef, 1993; Patrick *et al.*, 2003; Baruah *et al.*, 2007), μ_2 -bridging (Zhang *et al.*, 2003; Aghabozorg *et al.*, 2007; Sun *et al.*, 2006; Turner & Batten, 2007; Kang *et al.*, 2006), μ_3 -bridging (Zhang & You, 2003; Li *et al.*, 2006). Here we describe another new compound in which the PDC is μ_3 -bridging, (I),(Fig. 1).

Complex (I) is composed of {[Mn(C₇H₃NO₄)(H₂O)₂]_n.2H₂O} units, in which the Mn^{II} ion is six-coordinated in a distorted octahedral geometry (Table 1) formed by two coordinated water molecules, one N and one O atoms of a PDC dianion and two different carboxylate O atoms in the axial position from another two adjacent PDC ligands. The deprotonated PDC are μ_3 -bridging ligands and they join the neighbouring Mn^{II} ions to form this one-dimensional linear chain structure along *a* axis. This kind of μ_3 -bridging mode is different from that have been published (Zhang & You, 2003; Li *et al.*, 2006) as in this paper one bridging carboxylate O atom only links one metal ions, while in the latter one bridging carboxylate O atom simultaneously links two metal ions,respectively.

The carboxylate O atoms of PDC dianion and coordinate and non coordinated water molecules are all involved in rich O—H···O intermolecular hydrogen bonds (Table 2) and they connect polymeric chains into a three-dimensional framework (Fig. 2).

Experimental

The ligand H₂PDC (1 mmol, 0.17 g) and NaOH (2 mmol, 0.08 g) were dissolved in water and methanol mixed solvent (20 ml, *v/v* 1:1). To this solution, Mn(CH₃COO)₂.4H₂O (1 mmol, 0.25 g) was added and the resulting mixture was stirred and refluxed at 343 K for 5 h, then cooled to room temperature. After filtration and evaporation in air for a week, pink block-shaped crystals were obtained in a yield of 37%. Analysis, found (%): C, 28.70; H, 3.80; N, 4.71. C₇H₁₁Mn N O₈ requires (%): C,28.75; H,3.76; N,4.79. (CCDC number 668395)

Refinement

H atoms bonded to C atoms were positioned geometrically with C—H distance of 0.93 Å, and treated as riding atoms, with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}$. H atoms bonded to O atoms were located in a difference Fourier map and refined isotropically.

supplementary materials

Figures




Fig. 1. The structure of the building unit of the one-dimensional of (I), with the atom-numbering scheme. Displacement ellipsoids are drawn at the 30% probability level. Uncoordinate water molecules and H atoms on C atoms have been omitted. [Symmetry codes: (A) $2 - x, 2 - y, 1 - z$; (B) $1 - x, 2 - y, 1 - z$; (C) $-1 + x, y, z$.]




Fig. 2. The crystal packing of (I), showing hydrogen bonds as dashed lines. For the sake of clarity, H atoms on C atoms have been omitted.

catena-Poly[[[diaquamanganese(II)]- μ_3 -pyridine-2,3-dicarboxylato- $\kappa^4N,O^2:O^3:O^3'$] dihydrate]

Crystal data

[Mn(C ₇ H ₃ NO ₄)(H ₂ O) ₂] [·] 2H ₂ O	$F_{000} = 596$
$M_r = 292.11$	$D_x = 1.875 \text{ Mg m}^{-3}$
Monoclinic, $P2_1/c$	Mo $K\alpha$ radiation
Hall symbol: -P 2ybc	$\lambda = 0.71073 \text{ \AA}$
$a = 6.5719 (8) \text{ \AA}$	Cell parameters from 4039 reflections
$b = 7.6703 (9) \text{ \AA}$	$\theta = 2.8\text{--}28.1^\circ$
$c = 20.566 (3) \text{ \AA}$	$\mu = 1.31 \text{ mm}^{-1}$
$\beta = 93.3540 (10)^\circ$	$T = 291 (2) \text{ K}$
$V = 1034.9 (2) \text{ \AA}^3$	Block, pink
$Z = 4$	$0.38 \times 0.15 \times 0.11 \text{ mm}$

Data collection

Bruker APEXII CCD area-detector diffractometer	1921 independent reflections
Radiation source: fine-focus sealed tube	1774 reflections with $I > 2\sigma(I)$
Monochromator: graphite	$R_{\text{int}} = 0.015$
$T = 291(2) \text{ K}$	$\theta_{\max} = 25.5^\circ$
φ and ω scans	$\theta_{\min} = 2.8^\circ$
Absorption correction: multi-scan (SADABS; Sheldrick, 1996)	$h = -7 \rightarrow 7$

$T_{\min} = 0.640$, $T_{\max} = 0.865$
6428 measured reflections

$k = -9 \rightarrow 9$
 $l = -23 \rightarrow 24$

Refinement

Refinement on F^2	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.035$	H-atom parameters constrained
$wR(F^2) = 0.095$	$w = 1/[\sigma^2(F_o^2) + (0.042P)^2 + 2.0761P]$ where $P = (F_o^2 + 2F_c^2)/3$
$S = 1.07$	$(\Delta/\sigma)_{\max} = 0.001$
1921 reflections	$\Delta\rho_{\max} = 0.49 \text{ e \AA}^{-3}$
154 parameters	$\Delta\rho_{\min} = -0.74 \text{ e \AA}^{-3}$
3 restraints	Extinction correction: none
Primary atom site location: structure-invariant direct methods	

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
Mn1	0.69176 (6)	0.76827 (6)	0.38821 (2)	0.01743 (17)
O1	0.3592 (5)	0.7546 (3)	0.70493 (14)	0.0449 (7)
H1W	0.3455	0.8191	0.6719	0.067*
H2W	0.2695	0.7814	0.7307	0.067*
O2	0.9617 (5)	0.4500 (4)	0.74324 (14)	0.0543 (8)
H3W	0.8800	0.5269	0.7561	0.081*
H4W	0.9878	0.3937	0.7099	0.081*
O3	0.7081 (4)	0.4951 (3)	0.38849 (13)	0.0437 (7)
H5W	0.7244	0.4234	0.4193	0.066*
H6W	0.6950	0.4383	0.3532	0.066*
O4	0.6927 (4)	0.7979 (4)	0.28481 (13)	0.0420 (6)
H7W	0.5997	0.7749	0.2562	0.063*
H8W	0.6803	0.9080	0.2873	0.063*
O5	0.3643 (4)	0.7440 (3)	0.37627 (14)	0.0374 (6)
O6	0.0271 (4)	0.7556 (3)	0.37392 (13)	0.0362 (6)

supplementary materials

O7	0.2349 (5)	0.7489 (3)	0.51820 (13)	0.0393 (6)
O8	0.3057 (4)	0.9571 (3)	0.59238 (11)	0.0313 (5)
N1	0.2620 (4)	1.2081 (4)	0.50222 (14)	0.0272 (6)
C1	0.2629 (5)	0.9023 (4)	0.53482 (15)	0.0251 (7)
C2	0.2440 (4)	1.0408 (4)	0.48204 (15)	0.0234 (6)
C3	0.2117 (5)	1.0010 (4)	0.41594 (16)	0.0257 (7)
C4	0.1928 (4)	1.1277 (4)	0.37076 (14)	0.0212 (6)
H4	0.1690	1.1020	0.3268	0.025*
C5	0.2096 (6)	1.2911 (5)	0.39172 (18)	0.0374 (9)
H5	0.1971	1.3810	0.3614	0.045*
C6	0.2452 (6)	1.3332 (5)	0.45735 (17)	0.0331 (8)
H6	0.2574	1.4495	0.4699	0.040*
C7	0.2003 (5)	0.8171 (5)	0.38797 (15)	0.0255 (7)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Mn1	0.0193 (3)	0.0159 (3)	0.0170 (3)	-0.00007 (16)	0.00015 (17)	-0.00174 (16)
O1	0.0536 (18)	0.0450 (16)	0.0354 (15)	0.0076 (13)	-0.0033 (13)	0.0040 (12)
O2	0.0584 (19)	0.0623 (19)	0.0426 (16)	0.0277 (16)	0.0062 (13)	-0.0098 (14)
O3	0.0653 (19)	0.0273 (13)	0.0376 (15)	-0.0005 (12)	-0.0048 (13)	0.0011 (11)
O4	0.0416 (15)	0.0498 (16)	0.0339 (14)	-0.0068 (13)	-0.0048 (11)	0.0014 (12)
O5	0.0277 (13)	0.0371 (14)	0.0475 (16)	0.0024 (10)	0.0035 (11)	-0.0113 (11)
O6	0.0259 (13)	0.0410 (15)	0.0413 (15)	-0.0027 (10)	-0.0015 (11)	-0.0103 (11)
O7	0.0592 (18)	0.0225 (13)	0.0353 (14)	-0.0028 (11)	-0.0030 (13)	0.0002 (10)
O8	0.0405 (14)	0.0286 (12)	0.0242 (12)	0.0002 (10)	-0.0024 (10)	0.0017 (10)
N1	0.0272 (14)	0.0247 (13)	0.0297 (15)	0.0008 (11)	0.0024 (11)	-0.0001 (11)
C1	0.0223 (16)	0.0250 (16)	0.0279 (16)	0.0009 (12)	0.0010 (12)	-0.0004 (13)
C2	0.0182 (15)	0.0249 (16)	0.0273 (16)	0.0004 (12)	0.0016 (12)	0.0000 (13)
C3	0.0183 (15)	0.0292 (17)	0.0298 (17)	-0.0003 (12)	0.0016 (12)	-0.0001 (13)
C4	0.0262 (16)	0.0241 (15)	0.0133 (13)	0.0011 (12)	-0.0004 (11)	0.0015 (11)
C5	0.046 (2)	0.0328 (19)	0.033 (2)	0.0036 (16)	0.0031 (16)	0.0099 (15)
C6	0.043 (2)	0.0233 (16)	0.0325 (18)	-0.0003 (14)	0.0019 (15)	0.0017 (14)
C7	0.0232 (17)	0.0311 (17)	0.0220 (15)	0.0000 (13)	0.0002 (12)	-0.0007 (13)

Geometric parameters (\AA , $^\circ$)

Mn1—O3	2.098 (3)	O6—Mn1 ⁱⁱⁱ	2.243 (2)
Mn1—O4	2.139 (3)	O7—C1	1.236 (4)
Mn1—O8 ⁱ	2.144 (2)	O8—C1	1.272 (4)
Mn1—O5	2.160 (3)	O8—Mn1 ⁱ	2.144 (2)
Mn1—O6 ⁱⁱ	2.242 (3)	N1—C6	1.332 (4)
Mn1—N1 ^j	2.263 (3)	N1—C2	1.352 (4)
O1—H1W	0.8416	N1—Mn1 ⁱ	2.263 (3)
O1—H2W	0.8409	C1—C2	1.519 (4)
O2—H3W	0.8502	C2—C3	1.397 (4)
O2—H4W	0.8369	C3—C4	1.346 (4)

O3—H5W	0.8410	C3—C7	1.523 (5)
O3—H6W	0.8474	C4—C5	1.327 (5)
O4—H7W	0.8414	C4—H4	0.9300
O4—H8W	0.8497	C5—C6	1.394 (5)
O5—C7	1.250 (4)	C5—H5	0.9300
O6—C7	1.250 (4)	C6—H6	0.9300
O3—Mn1—O4	96.07 (11)	C1—O8—Mn1 ⁱ	119.8 (2)
O3—Mn1—O8 ⁱ	168.80 (10)	C6—N1—C2	118.0 (3)
O4—Mn1—O8 ⁱ	94.60 (10)	C6—N1—Mn1 ⁱ	129.3 (2)
O3—Mn1—O5	87.98 (10)	C2—N1—Mn1 ⁱ	112.7 (2)
O4—Mn1—O5	87.54 (10)	O7—C1—O8	126.4 (3)
O8 ⁱ —Mn1—O5	95.89 (9)	O7—C1—C2	117.6 (3)
O3—Mn1—O6 ⁱⁱ	84.63 (10)	O8—C1—C2	116.0 (3)
O4—Mn1—O6 ⁱⁱ	79.30 (10)	N1—C2—C3	120.8 (3)
O8 ⁱ —Mn1—O6 ⁱⁱ	94.02 (9)	N1—C2—C1	116.3 (3)
O5—Mn1—O6 ⁱⁱ	164.11 (10)	C3—C2—C1	122.9 (3)
O3—Mn1—N1 ⁱ	94.21 (10)	C4—C3—C2	121.1 (3)
O4—Mn1—N1 ⁱ	166.71 (11)	C4—C3—C7	114.1 (3)
O8 ⁱ —Mn1—N1 ⁱ	74.75 (9)	C2—C3—C7	124.8 (3)
O5—Mn1—N1 ⁱ	101.24 (10)	C5—C4—C3	117.1 (3)
O6 ⁱⁱ —Mn1—N1 ⁱ	93.33 (10)	C5—C4—H4	121.5
H1W—O1—H2W	108.6	C3—C4—H4	121.5
H3W—O2—H4W	140.9	C4—C5—C6	122.6 (3)
Mn1—O3—H5W	131.3	C4—C5—H5	118.7
Mn1—O3—H6W	120.6	C6—C5—H5	118.7
H5W—O3—H6W	108.1	N1—C6—C5	120.4 (3)
Mn1—O4—H7W	128.9	N1—C6—H6	119.8
Mn1—O4—H8W	92.3	C5—C6—H6	119.8
H7W—O4—H8W	100.4	O6—C7—O5	124.7 (3)
C7—O5—Mn1	143.5 (2)	O6—C7—C3	117.4 (3)
C7—O6—Mn1 ⁱⁱⁱ	147.3 (2)	O5—C7—C3	117.6 (3)
O3—Mn1—O5—C7	151.0 (4)	N1—C2—C3—C7	176.7 (3)
O4—Mn1—O5—C7	−112.9 (4)	C1—C2—C3—C7	−2.7 (5)
O8 ⁱ —Mn1—O5—C7	−18.5 (4)	C2—C3—C4—C5	1.1 (5)
O6 ⁱⁱ —Mn1—O5—C7	−146.8 (4)	C7—C3—C4—C5	−177.4 (3)
N1 ⁱ —Mn1—O5—C7	57.1 (4)	C3—C4—C5—C6	0.0 (5)
Mn1 ⁱ —O8—C1—O7	−171.8 (3)	C2—N1—C6—C5	0.2 (5)
Mn1 ⁱ —O8—C1—C2	7.9 (4)	Mn1 ⁱ —N1—C6—C5	−177.7 (3)
C6—N1—C2—C3	0.9 (5)	C4—C5—C6—N1	−0.7 (6)
Mn1 ⁱ —N1—C2—C3	179.1 (2)	Mn1 ⁱⁱⁱ —O6—C7—O5	165.6 (3)
C6—N1—C2—C1	−179.7 (3)	Mn1 ⁱⁱⁱ —O6—C7—C3	−20.0 (6)
Mn1 ⁱ —N1—C2—C1	−1.4 (3)	Mn1—O5—C7—O6	−177.6 (3)
O7—C1—C2—N1	175.7 (3)	Mn1—O5—C7—C3	8.0 (6)
O8—C1—C2—N1	−4.0 (4)	C4—C3—C7—O6	−81.4 (4)

supplementary materials

O7—C1—C2—C3	−4.8 (5)	C2—C3—C7—O6	100.2 (4)
O8—C1—C2—C3	175.5 (3)	C4—C3—C7—O5	93.4 (4)
N1—C2—C3—C4	−1.6 (5)	C2—C3—C7—O5	−85.0 (4)
C1—C2—C3—C4	179.0 (3)		


Symmetry codes: (i) $-x+1, -y+2, -z+1$; (ii) $x+1, y, z$; (iii) $x-1, y, z$.

Hydrogen-bond geometry (\AA , °)

$D\cdots H$	$H\cdots A$	$D\cdots A$	$D\cdots H\cdots A$
O1—H1W···O8	0.84	1.95	2.793 (4)
O1—H2W···O2 ^{iv}	0.84	2.09	2.845 (4)
O2—H4W···O6 ^v	0.84	2.07	2.884 (4)
O2—H4W···O4 ^{vi}	0.84	2.56	3.043 (4)
O2—H3W···O4 ^{vii}	0.85	1.94	2.788 (4)
O3—H5W···O7 ^v	0.84	1.85	2.691 (4)
O3—H6W···O1 ^v	0.85	1.92	2.730 (4)
O4—H8W···O1 ⁱ	0.85	2.61	3.457 (4)
O4—H7W···O1 ^{viii}	0.84	1.86	2.691 (4)
O4—H8W···O2 ^{viii}	0.85	2.37	2.788 (4)

Symmetry codes: (iv) $-x+1, y+1/2, -z+3/2$; (v) $-x+1, -y+1, -z+1$; (vi) $-x+2, -y+1, -z+1$; (vii) $x, -y+3/2, z+1/2$; (i) $-x+1, -y+2, -z+1$; (viii) $x, -y+3/2, z-1/2$.

Fig. 1



supplementary materials

Fig. 2

