

2-(4-Bromophenoxy)-3-isopropyl-5,6,7,8-tetrahydro-1-benzothieno[2,3-d]pyrimidin-4(3H)-one

Hong-Mei Wang, Li-Li Chen, Ting Hu and Xiao-Hua Zeng*

Department of Medicinal Chemistry, Yunyang Medical College, Shiyang 442000, People's Republic of China

Correspondence e-mail: zengken@126.com


Received 9 November 2008; accepted 11 November 2008

 Key indicators: single-crystal X-ray study; $T = 298$ K; mean $\sigma(\text{C}-\text{C}) = 0.014$ Å; R factor = 0.067; wR factor = 0.204; data-to-parameter ratio = 14.2.

In the title compound, $\text{C}_{19}\text{H}_{19}\text{BrN}_2\text{O}_2\text{S}$, the central thienopyrimidine ring system is essentially planar, with a maximum displacement of 0.068 (3) Å. The attached cyclohexene ring adopts a half-chair conformation. The molecular conformation and crystal packing are stabilized by three intramolecular $\text{C}-\text{H}\cdots\text{O}$ hydrogen bonds and two $\text{C}-\text{H}\cdots\pi$ interactions.

Related literature

For background to the use of pyrimidine derivatives as drugs, see: Ding *et al.* (2004). For a description of the Cambridge Structural Database, see: Allen (2002). For a related structure, see: Zeng *et al.* (2006).



Experimental

Crystal data

$\text{C}_{19}\text{H}_{19}\text{BrN}_2\text{O}_2\text{S}$
 $M_r = 418.32$
 Monoclinic, $P2_1$
 $a = 13.3957$ (7) Å

$b = 5.7366$ (3) Å
 $c = 13.3956$ (7) Å
 $\beta = 115.5410$ (10)°
 $V = 928.81$ (8) Å³

$Z = 2$
 Mo $K\alpha$ radiation
 $\mu = 2.34$ mm⁻¹

$T = 298$ (2) K
 $0.20 \times 0.10 \times 0.10$ mm

Data collection

Bruker SMART CCD area-detector diffractometer
 Absorption correction: multi-scan (*SADABS*; Sheldrick, 1996)
 $T_{\min} = 0.652$, $T_{\max} = 0.800$
 5798 measured reflections
 3228 independent reflections
 2346 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.106$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.067$
 $wR(F^2) = 0.204$
 $S = 1.07$
 3228 reflections
 228 parameters
 1 restraint

H-atom parameters constrained
 $\Delta\rho_{\max} = 7.69$ e Å⁻³
 $\Delta\rho_{\min} = -2.63$ e Å⁻³
 Absolute structure: Flack (1983),
 1424 Freidel pairs
 Flack parameter: 0.00 (8)

Table 1
 Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$\text{C11}-\text{H11}\cdots\text{O1}$	0.98	2.20	2.726 (10)	112
$\text{C12}-\text{H12B}\cdots\text{O2}$	0.96	2.43	2.915 (13)	111
$\text{C13}-\text{H13A}\cdots\text{O2}$	0.96	2.38	2.951 (10)	117
$\text{C12}-\text{H12A}\cdots\text{Cg1}^i$	0.96	2.92	3.854 (11)	165
$\text{C12}-\text{H12A}\cdots\text{Cg2}^i$	0.96	2.71	3.434 (11)	133

Symmetry code: (i) $x, y - 1, z$. Cg1 and Cg2 are the centroids of the thiophene (S1/C5-C8) and pyrimidine (N1/N2/C7-C10) rings, respectively.

Data collection: *SMART* (Bruker, 2001); cell refinement: *SAINT* (Bruker, 2001); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *PLATON* (Spek, 2003); software used to prepare material for publication: *SHELXTL* (Sheldrick, 2008).

We gratefully acknowledge financial support of this work by the Research Foundation for Students and Teachers of Yunyang Medical College (grant Nos. 2007QDJ15, 2007ZQB19, 2007ZQB20).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: AT2674).

References

- Allen, F. H. (2002). *Acta Cryst.* **B58**, 380–388.
 Bruker (2001). *SMART* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
 Ding, M. W., Xu, S. Z. & Zhao, J. F. (2004). *J. Org. Chem.* **69**, 8366–8371.
 Flack, H. D. (1983). *Acta Cryst.* **A39**, 876–881.
 Sheldrick, G. M. (1996). *SADABS*. University of Göttingen, Germany.
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
 Spek, A. L. (2003). *J. Appl. Cryst.* **36**, 7–13.
 Zeng, X.-H., Ding, M.-W. & He, H.-W. (2006). *Acta Cryst.* **E62**, o731–o732.

supplementary materials

Acta Cryst. (2008). E64, o2404 [doi:10.1107/S160053680803732X]

2-(4-Bromophenoxy)-3-isopropyl-5,6,7,8-tetrahydro-1-benzothieno[2,3-*d*]pyrimidin-4(3*H*)-one

H.-M. Wang, L.-L. Chen, T. Hu and X.-H. Zeng

Comment

Pyrimidine derivatives are attracting the increasing attention of synthetic community because of the important role played by such systems in many natural products, antibiotics and drugs (Ding *et al.*, 2004). In recent years, we have been engaged in the preparation of the derivatives of heterocycles *via* aza-Wittig reaction. The title compound, (I), was synthesized and structurally characterized in this context.

The molecular structure indicates that the thieno[2,3-*d*]pyrimidine moiety is a conjugated system (Fig. 1). All ring atoms in thieno[2,3-*d*]pyrimidine are essentially coplanar (Zeng *et al.*, 2006). The bond lengths and angles are within experimental error, in the ranges of values in previously reported structures in the Cambridge Structural Database (Version 5.26; Allen, 2002).

The cyclohexene ring adopts a half-chair conformation. The crystal packing is stabilized by three intramolecular C—H \cdots O hydrogen bonds and two C—H \cdots π interactions (Table 1). There exist no intermolecular hydrogen bonding interactions and no π - π stackings.

Experimental

To a solution of iminophosphorane (1.45 g, 3 mmol) in anhydrous dichloromethane (15 ml) was added iso-propyl isocyanate (3 mmol) under dry nitrogen at room temperature. After the reaction mixture was left unstirred for 48 h at room temperature, the solvent was removed off under reduced pressure and ether/petroleum ether (1:2 *v/v*, 20 ml) was added to precipitate triphenylphosphine oxide. After filtration, the solvent was removed, and the residue was dissolved in CH₃CN (15 ml). After adding 4-Br-PhOH (3.1 mmol) and excess K₂CO₃ to the solution of carbodiimide, The mixture was stirred for 24 h at room temperature, the solution was condensed and the residue was recrystallized by EtOH to give the title compound, (I), in yield of 80% (m.p. 478 K). Elemental analysis calculated for C₁₉H₁₉BrN₂O₂S: C 54.42, H 4.57, N 6.68. Found: C 54.56, H 4.42, N 6.53. Crystals suitable for single crystal X-ray diffraction were obtained by vapor diffusion of hexane and dichloromethane (1:3 *v/v*) at room temperature.

Refinement

H atoms were placed at calculated positions and treated as riding atoms, with C—H = 0.93–0.98 Å, and $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$ for CH or $1.5U_{\text{eq}}(\text{C})$ for CH₃.

Figures




Fig. 1. View of the molecule of (I) showing the atom-labeling scheme. Displacement ellipsoids are drawn at 50% probability level. H-atoms are represented by circles of arbitrary size.

2-(4-Bromo-phenoxy)-3-isopropyl-5,6,7,8-tetrahydro -benzothieno[2,3-d]pyrimidin-4(3H)-one

Crystal data

$C_{19}H_{19}BrN_2O_2S$	$F_{000} = 428$
$M_r = 419.33$	$D_x = 1.499 \text{ Mg m}^{-3}$
Monoclinic, $P2_1$	Melting point: 478K K
Hall symbol: P 2yb	Mo $K\alpha$ radiation
$a = 13.3957 (7) \text{ \AA}$	$\lambda = 0.71073 \text{ \AA}$
$b = 5.7366 (3) \text{ \AA}$	Cell parameters from 2048 reflections
$c = 13.3956 (7) \text{ \AA}$	$\theta = 2.9\text{--}24.5^\circ$
$\beta = 115.5410 (10)^\circ$	$\mu = 2.34 \text{ mm}^{-1}$
$V = 928.81 (8) \text{ \AA}^3$	$T = 298 (2) \text{ K}$
$Z = 2$	Block, colorless
	$0.20 \times 0.10 \times 0.10 \text{ mm}$

Data collection

Bruker SMART CCD area-detector diffractometer	3228 independent reflections
Radiation source: fine-focus sealed tube	2346 reflections with $I > 2\sigma(I)$
Monochromator: graphite	$R_{\text{int}} = 0.106$
$T = 298(2) \text{ K}$	$\theta_{\text{max}} = 25.0^\circ$
φ and ω scans	$\theta_{\text{min}} = 1.7^\circ$
Absorption correction: multi-scan (SADABS; Sheldrick, 1996)	$h = -13 \rightarrow 15$
$T_{\text{min}} = 0.652$, $T_{\text{max}} = 0.800$	$k = -6 \rightarrow 6$
5798 measured reflections	$l = -15 \rightarrow 15$

Refinement

Refinement on F^2	Hydrogen site location: inferred from neighbouring sites
Least-squares matrix: full	H-atom parameters constrained
$R[F^2 > 2\sigma(F^2)] = 0.067$	$w = 1/[\sigma^2(F_o^2) + (0.1151P)^2]$
$wR(F^2) = 0.204$	where $P = (F_o^2 + 2F_c^2)/3$
$S = 1.07$	$(\Delta/\sigma)_{\text{max}} = 0.001$
	$\Delta\rho_{\text{max}} = 0.67 \text{ e \AA}^{-3}$

3228 reflections	$\Delta\rho_{\min} = -0.62 \text{ e } \text{\AA}^{-3}$
228 parameters	Extinction correction: none
1 restraint	Absolute structure: Flack (1983), 1424 Friedel pairs
Primary atom site location: structure-invariant direct methods	Flack parameter: 0.00 (8)
Secondary atom site location: difference Fourier map	

Special details

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R-factor wR and goodness of fit S are based on F^2 , conventional R-factors R are based on F, with F set to zero for negative F^2 . The threshold expression of $F^2 > 2\sigma(F^2)$ is used only for calculating R-factors(gt) etc. and is not relevant to the choice of reflections for refinement. R-factors based on F^2 are statistically about twice as large as those based on F, and R- factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
Br1	0.84533 (6)	0.7555 (2)	0.71718 (7)	0.0762 (4)
C1	0.0942 (6)	1.2447 (18)	0.8813 (6)	0.0534 (18)
H1A	0.0444	1.3157	0.8118	0.064*
H1B	0.0650	1.0921	0.8853	0.064*
C2	0.0967 (9)	1.390 (3)	0.9738 (10)	0.103 (5)
H2A	0.0231	1.4514	0.9536	0.124*
H2B	0.1152	1.2908	1.0380	0.124*
C3	0.1719 (9)	1.578 (2)	1.0038 (11)	0.093 (4)
H3A	0.1730	1.6464	1.0704	0.111*
H3B	0.1416	1.6942	0.9459	0.111*
C4	0.2917 (7)	1.537 (2)	1.0246 (7)	0.067 (3)
H4A	0.3234	1.6797	1.0119	0.081*
H4B	0.3351	1.4885	1.1006	0.081*
C5	0.2930 (6)	1.3513 (13)	0.9472 (6)	0.0460 (19)
C6	0.2071 (5)	1.2157 (15)	0.8824 (5)	0.046 (2)
C7	0.2363 (6)	1.0548 (14)	0.8163 (6)	0.0405 (16)
C8	0.3446 (6)	1.0816 (16)	0.8335 (6)	0.052 (2)
C9	0.1666 (6)	0.9073 (15)	0.7312 (6)	0.0457 (18)
C10	0.3302 (6)	0.8295 (14)	0.7015 (6)	0.047 (2)
C11	0.1472 (6)	0.6545 (14)	0.5719 (6)	0.0459 (18)
H11	0.0726	0.6617	0.5683	0.055*
C12	0.1754 (9)	0.3997 (19)	0.5775 (8)	0.079 (3)
H12A	0.1906	0.3406	0.6497	0.119*
H12B	0.2395	0.3796	0.5638	0.119*
H12C	0.1142	0.3161	0.5228	0.119*
C13	0.1386 (8)	0.767 (2)	0.4657 (6)	0.071 (2)
H13A	0.2104	0.7707	0.4661	0.107*

supplementary materials

H13B	0.1112	0.9233	0.4608	0.107*
H13C	0.0889	0.6784	0.4033	0.107*
C14	0.4794 (6)	0.7242 (18)	0.6612 (7)	0.055 (2)
C15	0.5512 (8)	0.553 (2)	0.7233 (8)	0.073 (3)
H15	0.5268	0.4320	0.7536	0.087*
C16	0.6596 (8)	0.563 (2)	0.7394 (9)	0.074 (3)
H16	0.7096	0.4498	0.7815	0.088*
C17	0.6938 (5)	0.741 (2)	0.6934 (6)	0.061 (2)
C18	0.6221 (7)	0.9049 (19)	0.6317 (8)	0.066 (2)
H18	0.6458	1.0245	0.6002	0.079*
C19	0.5134 (8)	0.895 (2)	0.6152 (9)	0.073 (3)
H19	0.4634	1.0073	0.5720	0.088*
N1	0.3955 (5)	0.9634 (13)	0.7793 (6)	0.0527 (17)
N2	0.2193 (4)	0.7931 (13)	0.6717 (4)	0.0415 (13)
O1	0.0683 (4)	0.8637 (12)	0.7061 (5)	0.0664 (19)
O2	0.3666 (4)	0.7053 (13)	0.6384 (5)	0.072 (2)
S1	0.41333 (15)	1.2916 (5)	0.93045 (17)	0.0604 (6)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Br1	0.0420 (4)	0.1275 (10)	0.0622 (5)	-0.0010 (6)	0.0253 (4)	-0.0105 (6)
C1	0.048 (4)	0.055 (5)	0.061 (4)	-0.001 (5)	0.027 (3)	0.001 (5)
C2	0.063 (6)	0.162 (13)	0.089 (8)	-0.006 (7)	0.037 (6)	-0.052 (8)
C3	0.079 (7)	0.105 (10)	0.106 (9)	-0.008 (7)	0.051 (7)	-0.036 (8)
C4	0.059 (5)	0.090 (7)	0.043 (4)	0.009 (5)	0.013 (4)	-0.009 (5)
C5	0.045 (4)	0.050 (5)	0.038 (4)	0.004 (3)	0.014 (3)	0.001 (3)
C6	0.039 (4)	0.057 (6)	0.036 (3)	0.007 (4)	0.011 (3)	0.004 (4)
C7	0.032 (3)	0.052 (4)	0.033 (3)	0.007 (3)	0.010 (3)	0.002 (3)
C8	0.033 (4)	0.070 (6)	0.043 (4)	-0.004 (4)	0.006 (3)	-0.002 (4)
C9	0.037 (4)	0.055 (5)	0.040 (4)	-0.004 (4)	0.012 (3)	0.006 (3)
C10	0.038 (4)	0.053 (5)	0.049 (4)	0.004 (3)	0.018 (3)	-0.006 (3)
C11	0.037 (4)	0.053 (5)	0.044 (4)	-0.009 (3)	0.013 (3)	-0.004 (3)
C12	0.100 (8)	0.058 (6)	0.061 (6)	-0.002 (6)	0.016 (6)	0.001 (5)
C13	0.083 (6)	0.069 (6)	0.044 (4)	0.001 (6)	0.011 (4)	0.007 (5)
C14	0.039 (4)	0.068 (6)	0.061 (4)	-0.009 (4)	0.024 (3)	-0.022 (5)
C15	0.054 (5)	0.088 (8)	0.076 (7)	-0.002 (5)	0.028 (5)	0.010 (6)
C16	0.050 (5)	0.093 (8)	0.076 (6)	0.004 (5)	0.027 (5)	0.019 (6)
C17	0.034 (3)	0.106 (7)	0.043 (4)	0.005 (6)	0.016 (3)	-0.019 (6)
C18	0.049 (5)	0.072 (6)	0.074 (6)	0.000 (5)	0.023 (5)	0.016 (5)
C19	0.048 (5)	0.077 (7)	0.084 (7)	0.011 (5)	0.019 (5)	0.009 (5)
N1	0.034 (3)	0.063 (5)	0.057 (4)	-0.007 (3)	0.016 (3)	-0.022 (4)
N2	0.031 (3)	0.049 (4)	0.043 (3)	-0.005 (3)	0.013 (2)	0.002 (3)
O1	0.033 (3)	0.103 (6)	0.062 (3)	-0.012 (3)	0.018 (2)	-0.019 (3)
O2	0.039 (3)	0.098 (6)	0.079 (4)	-0.011 (3)	0.025 (3)	-0.041 (4)
S1	0.0373 (9)	0.0736 (16)	0.0600 (11)	-0.0053 (11)	0.0114 (8)	-0.0194 (12)

Geometric parameters (Å, °)

Br1—C17	1.916 (7)	C10—O2	1.347 (9)
C1—C2	1.481 (14)	C10—N2	1.378 (8)
C1—C6	1.515 (9)	C11—N2	1.497 (9)
C1—H1A	0.9700	C11—C12	1.504 (14)
C1—H1B	0.9700	C11—C13	1.521 (12)
C2—C3	1.410 (18)	C11—H11	0.9800
C2—H2A	0.9700	C12—H12A	0.9600
C2—H2B	0.9700	C12—H12B	0.9600
C3—C4	1.523 (14)	C12—H12C	0.9600
C3—H3A	0.9700	C13—H13A	0.9600
C3—H3B	0.9700	C13—H13B	0.9600
C4—C5	1.492 (12)	C13—H13C	0.9600
C4—H4A	0.9700	C14—C19	1.336 (15)
C4—H4B	0.9700	C14—C15	1.377 (15)
C5—C6	1.350 (11)	C14—O2	1.412 (9)
C5—S1	1.754 (8)	C15—C16	1.374 (13)
C6—C7	1.445 (10)	C15—H15	0.9300
C7—C8	1.375 (10)	C16—C17	1.370 (15)
C7—C9	1.404 (11)	C16—H16	0.9300
C8—N1	1.370 (10)	C17—C18	1.342 (14)
C8—S1	1.719 (9)	C18—C19	1.377 (12)
C9—O1	1.236 (9)	C18—H18	0.9300
C9—N2	1.431 (10)	C19—H19	0.9300
C10—N1	1.287 (10)		
C2—C1—C6	113.0 (7)	N2—C11—C12	114.8 (7)
C2—C1—H1A	109.0	N2—C11—C13	111.6 (7)
C6—C1—H1A	109.0	C12—C11—C13	112.1 (8)
C2—C1—H1B	109.0	N2—C11—H11	105.8
C6—C1—H1B	109.0	C12—C11—H11	105.8
H1A—C1—H1B	107.8	C13—C11—H11	105.8
C3—C2—C1	115.1 (10)	C11—C12—H12A	109.5
C3—C2—H2A	108.5	C11—C12—H12B	109.5
C1—C2—H2A	108.5	H12A—C12—H12B	109.5
C3—C2—H2B	108.5	C11—C12—H12C	109.5
C1—C2—H2B	108.5	H12A—C12—H12C	109.5
H2A—C2—H2B	107.5	H12B—C12—H12C	109.5
C2—C3—C4	120.1 (11)	C11—C13—H13A	109.5
C2—C3—H3A	107.3	C11—C13—H13B	109.5
C4—C3—H3A	107.3	H13A—C13—H13B	109.5
C2—C3—H3B	107.3	C11—C13—H13C	109.5
C4—C3—H3B	107.3	H13A—C13—H13C	109.5
H3A—C3—H3B	106.9	H13B—C13—H13C	109.5
C5—C4—C3	108.0 (8)	C19—C14—C15	121.0 (8)
C5—C4—H4A	110.1	C19—C14—O2	120.1 (9)
C3—C4—H4A	110.1	C15—C14—O2	118.6 (9)
C5—C4—H4B	110.1	C16—C15—C14	118.6 (10)

supplementary materials

C3—C4—H4B	110.1	C16—C15—H15	120.7
H4A—C4—H4B	108.4	C14—C15—H15	120.7
C6—C5—C4	126.6 (7)	C17—C16—C15	119.9 (10)
C6—C5—S1	112.4 (5)	C17—C16—H16	120.1
C4—C5—S1	121.0 (6)	C15—C16—H16	120.1
C5—C6—C7	112.3 (6)	C18—C17—C16	120.6 (7)
C5—C6—C1	120.9 (7)	C18—C17—Br1	119.9 (8)
C7—C6—C1	126.7 (7)	C16—C17—Br1	119.5 (8)
C8—C7—C9	119.2 (7)	C17—C18—C19	119.7 (10)
C8—C7—C6	111.8 (7)	C17—C18—H18	120.1
C9—C7—C6	128.4 (6)	C19—C18—H18	120.1
N1—C8—C7	125.8 (7)	C14—C19—C18	120.2 (9)
N1—C8—S1	121.3 (5)	C14—C19—H19	119.9
C7—C8—S1	112.8 (6)	C18—C19—H19	119.9
O1—C9—C7	126.9 (7)	C10—N1—C8	113.9 (6)
O1—C9—N2	118.7 (7)	C10—N2—C9	119.9 (6)
C7—C9—N2	114.4 (6)	C10—N2—C11	122.7 (6)
N1—C10—O2	121.4 (6)	C9—N2—C11	117.2 (5)
N1—C10—N2	126.6 (7)	C10—O2—C14	117.8 (6)
O2—C10—N2	112.0 (6)	C8—S1—C5	90.6 (4)
C6—C1—C2—C3	36.3 (16)	Br1—C17—C18—C19	-179.7 (8)
C1—C2—C3—C4	-50.1 (18)	C15—C14—C19—C18	-1.6 (16)
C2—C3—C4—C5	34.6 (16)	O2—C14—C19—C18	-175.9 (9)
C3—C4—C5—C6	-9.8 (13)	C17—C18—C19—C14	0.5 (16)
C3—C4—C5—S1	169.1 (8)	O2—C10—N1—C8	178.4 (7)
C4—C5—C6—C7	179.3 (8)	N2—C10—N1—C8	-1.7 (13)
S1—C5—C6—C7	0.4 (8)	C7—C8—N1—C10	4.8 (13)
C4—C5—C6—C1	1.0 (12)	S1—C8—N1—C10	-173.3 (6)
S1—C5—C6—C1	-177.9 (6)	N1—C10—N2—C9	-1.2 (12)
C2—C1—C6—C5	-13.3 (13)	O2—C10—N2—C9	178.8 (7)
C2—C1—C6—C7	168.6 (10)	N1—C10—N2—C11	173.0 (8)
C5—C6—C7—C8	-1.3 (10)	O2—C10—N2—C11	-7.0 (10)
C1—C6—C7—C8	176.9 (7)	O1—C9—N2—C10	-176.8 (7)
C5—C6—C7—C9	-172.1 (8)	C7—C9—N2—C10	1.2 (10)
C1—C6—C7—C9	6.1 (13)	O1—C9—N2—C11	8.7 (10)
C9—C7—C8—N1	-4.8 (13)	C7—C9—N2—C11	-173.3 (7)
C6—C7—C8—N1	-176.6 (8)	C12—C11—N2—C10	66.0 (10)
C9—C7—C8—S1	173.4 (6)	C13—C11—N2—C10	-63.0 (10)
C6—C7—C8—S1	1.6 (9)	C12—C11—N2—C9	-119.6 (9)
C8—C7—C9—O1	179.4 (8)	C13—C11—N2—C9	111.4 (8)
C6—C7—C9—O1	-10.4 (14)	N1—C10—O2—C14	0.8 (13)
C8—C7—C9—N2	1.5 (11)	N2—C10—O2—C14	-179.2 (7)
C6—C7—C9—N2	171.7 (7)	C19—C14—O2—C10	-86.5 (11)
C19—C14—C15—C16	1.5 (15)	C15—C14—O2—C10	99.1 (10)
O2—C14—C15—C16	175.9 (9)	N1—C8—S1—C5	177.1 (8)
C14—C15—C16—C17	-0.4 (16)	C7—C8—S1—C5	-1.2 (7)
C15—C16—C17—C18	-0.6 (16)	C6—C5—S1—C8	0.4 (6)
C15—C16—C17—Br1	179.7 (8)	C4—C5—S1—C8	-178.6 (7)
C16—C17—C18—C19	0.6 (15)		

Hydrogen-bond geometry (Å, °)

<i>D—H...A</i>	<i>D—H</i>	<i>H...A</i>	<i>D...A</i>	<i>D—H...A</i>
C11—H11...O1	0.98	2.20	2.726 (10)	112
C12—H12B...O2	0.96	2.43	2.915 (13)	111
C13—H13A...O2	0.96	2.38	2.951 (10)	117
C12—H12B...Cg1	0.96	2.92	3.854 (11)	165
C12—H12B...Cg2	0.96	2.71	3.434 (11)	133

Fig. 1

