

N-(2,6-Dichlorophenyl)benzamide

B. Thimme Gowda,^{a*} Miroslav Tokarčík,^b Jozef Kožíšek,^b
B. P. Sowmya^a and Hartmut Fuess^c

^aDepartment of Chemistry, Mangalore University, Mangalagangotri-574 199, Mangalore, India, ^bFaculty of Chemical and Food Technology, Slovak Technical University, Radlinského 9, SK-812 37 Bratislava, Slovak Republic, and ^cInstitute of Materials Science, Darmstadt University of Technology, Petersenstrasse 23, D-64287, Darmstadt, Germany
Correspondence e-mail: gowdabt@yahoo.com


Received 15 January 2008; accepted 28 January 2008

Key indicators: single-crystal X-ray study; $T = 295$ K; mean $\sigma(\text{C}-\text{C}) = 0.003$ Å; R factor = 0.043; wR factor = 0.127; data-to-parameter ratio = 15.8.

The conformation of the N—H and C=O bonds in the structure of the title compound (N26DCPBA), $\text{C}_{13}\text{H}_9\text{Cl}_2\text{NO}$, are *anti* to each other, similar to that observed in *N*-phenylbenzamide (NPBA), *N*-(2-chlorophenyl)benzamide (N2CPBA), *N*-(2,3-dichlorophenyl)benzamide (N23DCPBA) and other benzanilides. The asymmetric unit of N26DCPBA contains two molecules. The bond parameters in N26DCPBA are similar to those in NPBA, N2CPBA, N23DCPBA and other benzanilides. The amide group, —NHCO—, makes a dihedral angle of 30.8 (1)° with the benzoyl ring in the first molecule and 35.1 (2)° in the second molecule of the asymmetric unit. The dihedral angle between the two benzene rings (benzoyl and aniline) is 56.8 (1)° in the first molecule and 59.1 (1)° in the second molecule. N—H...O hydrogen bonds give rise to infinite chains running along the *a* axis of the crystal structure.

Related literature

For related literature, see: Gowda *et al.* (2003, 2007*a,b,c*, 2008).



Experimental

Crystal data

$\text{C}_{13}\text{H}_9\text{Cl}_2\text{NO}$
 $M_r = 266.11$
Monoclinic, $P2_1/c$
 $a = 10.0431$ (2) Å
 $b = 13.7150$ (3) Å

$c = 18.4585$ (4) Å
 $\beta = 93.623$ (2)°
 $V = 2537.41$ (9) Å³
 $Z = 8$
Mo $K\alpha$ radiation

$\mu = 0.49$ mm⁻¹
 $T = 295$ (2) K

0.26 × 0.24 × 0.21 mm

Data collection

Oxford Diffraction Xcalibur System
diffractometer
Absorption correction: analytical
[(Oxford Diffraction, 2007);
analytical numeric absorption
correction using a multifaceted
crystal model (Clark & Reid,
1995)]
 $T_{\min} = 0.882$, $T_{\max} = 0.904$
4956 measured reflections
4956 independent reflections
3810 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.023$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.043$
 $wR(F^2) = 0.127$
 $S = 1.10$
4956 reflections
314 parameters
2 restraints

H atoms treated by a mixture of
independent and constrained
refinement
 $\Delta\rho_{\max} = 0.40$ e Å⁻³
 $\Delta\rho_{\min} = -0.35$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

<i>D</i> —H... <i>A</i>	<i>D</i> —H	H... <i>A</i>	<i>D</i> ... <i>A</i>	<i>D</i> —H... <i>A</i>
N1—H1N...O2	0.860 (19)	2.09 (2)	2.9035 (18)	158 (2)
N2—H2N...O1 ¹	0.843 (18)	2.060 (19)	2.8831 (18)	165.1 (19)

Symmetry code: (i) $x + 1, y, z$.

Data collection: *CrysAlis CCD* (Oxford Diffraction, 2007); cell refinement: *CrysAlis RED* (Oxford Diffraction, 2007); data reduction: *CrysAlis RED*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3* (Farrugia, 1997) and *DIAMOND* (Brandenburg, 2002); software used to prepare material for publication: *SHELXL97*, *PLATON* (Spek, 2003) and *WinGX* (Farrugia, 1999).

MT and JK thank the Grant Agency of the Slovak Republic (Grant No. VEGA 1/0817/08) and the Structural Funds, Interreg IIIA, for financial support in purchasing the diffractometer.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: OM2209).

References

- Brandenburg, K. (2002). *DIAMOND*. Crystal Impact GbR, Bonn, Germany.
Clark, R. C. & Reid, J. S. (1995). *Acta Cryst.* **A51**, 887–897.
Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
Farrugia, L. J. (1999). *J. Appl. Cryst.* **32**, 837–838.
Gowda, B. T., Jyothi, K., Paulus, H. & Fuess, H. (2003). *Z. Naturforsch. Teil A*, **58**, 225–230.
Gowda, B. T., Sowmya, B. P., Kožíšek, J., Tokarčík, M. & Fuess, H. (2007*a*). *Acta Cryst.* **E63**, o2906.
Gowda, B. T., Sowmya, B. P., Tokarčík, M., Kožíšek, J. & Fuess, H. (2007*b*). *Acta Cryst.* **E63**, o3326.
Gowda, B. T., Sowmya, B. P., Tokarčík, M., Kožíšek, J. & Fuess, H. (2007*c*). *Acta Cryst.* **E63**, o3365.
Gowda, B. T., Tokarčík, M., Kožíšek, J., Sowmya, B. P. & Fuess, H. (2008). *Acta Cryst.* **E64**, o462.
Oxford Diffraction (2007). *CrysAlis CCD* and *CrysAlis RED*. Oxford Diffraction Ltd, Abingdon, Oxfordshire, England.
Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
Spek, A. L. (2003). *J. Appl. Cryst.* **36**, 7–13.

supplementary materials

Acta Cryst. (2008). E64, o540 [doi:10.1107/S160053680800305X]

N-(2,6-Dichlorophenyl)benzamide

B. T. Gowda, M. Tokarcík, J. Kozísek, B. P. Sowmya and H. Fuess

Comment

In the present work, the structure of *N*-(2,6-dichlorophenyl)-benzamide (N26DCPBA) has been determined to explore the effect of substituents on the structure of *N*-aromatic amides (Gowda *et al.*, 2003,2007a,b,c,2008). The conformation of the N—H and C=O bonds in the structure of N26DCPBA (Fig.1) are anti to each other, similar to that observed in *N*-(phenyl)-benzamide (NPBA)(Gowda *et al.*, 2003), *N*-(2-chlorophenyl)-benzamide (N2CPBA), *N*-(2,3-dichlorophenyl)-benzamide (N23DCPBA) and other benzanilides(Gowda *et al.*, 2007a,b,c, 2008). The title compound crystallizes in the space group P21/c, with two molecules in the asymmetric unit. The bond parameters in N26DCPBA are similar to those in NPBA, N2CPBA, N23DCPBA and other benzanilides. The amide group —NHCO— has the dihedral angle of 30.8 (1)° with the benzoyl ring in the first molecule and 35.1 (2)° in the second molecule of the asymmetric unit. The dihedral angle between the two benzene rings (benzoyl and aniline) is 56.8 (1)° in the first molecule and 59.1 (1)° in the second molecule. One-dimensional chains of N26DCPBA along the base vector [1 0 0] formed by hydrogen bonds N1—H1N···O2(i) and N2—H2N···O1 (Table 1) as viewed down the *b* axis is shown in Fig.2. Symmetry code (i): $x + 1, y, z$.

Experimental

The title compound was prepared according to the literature method (Gowda *et al.*, 2003). The purity of the compound was checked by determining its melting point. It was characterized by recording its infrared and NMR spectra. Single crystals of the title compound were obtained from an ethanolic solution and used for X-ray diffraction studies at room temperature.

Refinement

H atoms bonded to C atoms were placed in geometrically calculated positions and subsequently treated as riding with C—H bond distance 0.93 Å. H(N) atoms were visible in the difference map. In the refinement the N—H distance was restrained to 0.86 (4) Å. The $U_{\text{iso}}(\text{H})$ values were set at 1.2 $U_{\text{eq}}(\text{C}, \text{N})$.

Figures




Fig. 1. Molecular structure of the title compound showing the atom labelling scheme. Numbers of the C atoms in the second molecule are increased by 20. Displacement ellipsoids are drawn at the 30% probability level. The hydrogen bond N2—H2N···O1 is shown as a dashed line.




Fig. 2. Crystal structure of the title compound viewed down the *b* axis. One-dimensional chains along the base vector [1 0 0] are formed by hydrogen bonds N1–H1N···O2 and N2–H2N···O1(i). Symmetry code (i): $x + 1, y, z$. H atoms not involved in hydrogen bonding are omitted.

N-(2,6-Dichlorophenyl)benzamide

Crystal data

$C_{13}H_9Cl_2NO$

$M_r = 266.11$

Monoclinic, $P2_1/c$

Hall symbol: $-P\ 2_1/c$

$a = 10.0431\ (2)\ \text{\AA}$

$b = 13.7150\ (3)\ \text{\AA}$

$c = 18.4585\ (4)\ \text{\AA}$

$\beta = 93.623\ (2)^\circ$

$V = 2537.41\ (9)\ \text{\AA}^3$

$Z = 8$

$F_{000} = 1088$

$D_x = 1.393\ \text{Mg m}^{-3}$

Mo $K\alpha$ radiation

$\lambda = 0.71073\ \text{\AA}$

Cell parameters from 24495 reflections

$\theta = 3.1\text{--}29.5^\circ$

$\mu = 0.49\ \text{mm}^{-1}$

$T = 295\ (2)\ \text{K}$

Block, colorless

$0.26 \times 0.24 \times 0.21\ \text{mm}$

Data collection

Oxford Diffraction Xcalibur System
diffractometer

Radiation source: Enhance (Mo) X-ray Source

Monochromator: graphite

Detector resolution: $10.434\ \text{pixels mm}^{-1}$

$T = 295(2)\ \text{K}$

φ scans, and ω scans with κ offsets

Absorption correction: analytical

[(Oxford Diffraction, 2007); analytical numeric absorption correction using a multifaceted crystal model (Clark & Reid, 1995)]

$T_{\min} = 0.883, T_{\max} = 0.904$

54956 measured reflections

4956 independent reflections

3810 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.023$

$\theta_{\max} = 26.0^\circ$

$\theta_{\min} = 5.1^\circ$

$h = -12 \rightarrow 12$

$k = -16 \rightarrow 16$

$l = -22 \rightarrow 22$

Refinement

Refinement on F^2

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.043$

$wR(F^2) = 0.127$

Secondary atom site location: difference Fourier map

Hydrogen site location: inferred from neighbouring sites

H atoms treated by a mixture of independent and constrained refinement

$w = 1/[\sigma^2(F_o^2) + (0.0646P)^2 + 0.4979P]$

$S = 1.10$	where $P = (F_o^2 + 2F_c^2)/3$
4956 reflections	$(\Delta/\sigma)_{\max} = 0.001$
314 parameters	$\Delta\rho_{\max} = 0.40 \text{ e } \text{\AA}^{-3}$
2 restraints	$\Delta\rho_{\min} = -0.35 \text{ e } \text{\AA}^{-3}$
Primary atom site location: structure-invariant direct methods	Extinction correction: SHELXL97 (Sheldrick, 2008), $F_c^* = kFc[1+0.001xFc^2\lambda^3/\sin(2\theta)]^{-1/4}$ Extinction coefficient: 0.0083 (11)

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > 2\sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
C1	0.29163 (15)	0.88930 (13)	0.13557 (10)	0.0468 (4)
C2	0.31340 (17)	0.99515 (13)	0.12131 (10)	0.0496 (4)
C3	0.2344 (2)	1.06295 (18)	0.1520 (2)	0.0957 (10)
H3	0.1718	1.0433	0.184	0.115*
C4	0.2478 (3)	1.1613 (2)	0.1353 (2)	0.1205 (13)
H4	0.1942	1.2072	0.1564	0.145*
C5	0.3392 (3)	1.19053 (19)	0.08819 (19)	0.0957 (10)
H5	0.3447	1.2559	0.0752	0.115*
C6	0.4206 (4)	1.1252 (2)	0.06070 (14)	0.0979 (9)
H6	0.4856	1.1456	0.0304	0.118*
C7	0.4090 (3)	1.02747 (17)	0.07699 (13)	0.0798 (7)
H7	0.4667	0.9828	0.0577	0.096*
C8	0.39430 (16)	0.73180 (13)	0.15565 (12)	0.0555 (5)
C9	0.41165 (19)	0.70599 (15)	0.22837 (14)	0.0661 (6)
C10	0.4153 (2)	0.60990 (18)	0.25095 (17)	0.0805 (7)
H10	0.4284	0.5945	0.3	0.097*
C11	0.3993 (2)	0.53776 (18)	0.19996 (19)	0.0853 (8)
H11	0.4011	0.4729	0.2146	0.102*
C12	0.3805 (2)	0.55993 (17)	0.12743 (18)	0.0822 (8)
H12	0.3702	0.5102	0.0933	0.099*
C13	0.3771 (2)	0.65690 (16)	0.10524 (14)	0.0664 (6)
N1	0.39953 (14)	0.83121 (11)	0.13436 (10)	0.0535 (4)
H1N	0.4772 (19)	0.8565 (16)	0.1310 (11)	0.064*
O1	0.18181 (11)	0.85646 (10)	0.14687 (9)	0.0635 (4)

supplementary materials

C11	0.42425 (7)	0.79839 (5)	0.29312 (4)	0.0924 (2)
C12	0.35242 (9)	0.68484 (5)	0.01367 (4)	0.1030 (3)
C21	0.79317 (15)	0.84885 (12)	0.13788 (10)	0.0457 (4)
C22	0.81608 (17)	0.75379 (13)	0.10084 (11)	0.0505 (4)
C23	0.7373 (2)	0.67477 (16)	0.11555 (17)	0.0803 (8)
H23	0.6724	0.6804	0.1491	0.096*
C24	0.7548 (3)	0.58684 (18)	0.0802 (2)	0.1034 (11)
H24	0.7023	0.5334	0.0906	0.124*
C25	0.8487 (3)	0.57822 (19)	0.03020 (18)	0.0922 (9)
H25	0.8588	0.5193	0.0062	0.111*
C26	0.9277 (3)	0.65546 (18)	0.01526 (13)	0.0761 (7)
H26	0.9919	0.6491	-0.0186	0.091*
C27	0.9124 (2)	0.74372 (15)	0.05076 (11)	0.0588 (5)
H27	0.9669	0.7963	0.0409	0.071*
C28	0.89884 (14)	0.99789 (11)	0.18305 (9)	0.0412 (4)
C29	0.92285 (17)	1.01514 (13)	0.25670 (10)	0.0501 (4)
C30	0.9290 (2)	1.10850 (16)	0.28488 (12)	0.0622 (5)
H30	0.9468	1.1184	0.3344	0.075*
C31	0.9085 (2)	1.18598 (15)	0.23909 (13)	0.0670 (6)
H31	0.9121	1.249	0.2578	0.08*
C32	0.8826 (2)	1.17230 (14)	0.16579 (13)	0.0653 (5)
H32	0.8683	1.2255	0.1351	0.078*
C33	0.87794 (18)	1.07878 (13)	0.13833 (10)	0.0511 (4)
N2	0.90279 (14)	0.90255 (10)	0.15368 (8)	0.0450 (4)
H2N	0.9794 (18)	0.8817 (14)	0.1457 (10)	0.054*
O2	0.68208 (11)	0.87638 (10)	0.15285 (9)	0.0677 (4)
C13	0.94534 (8)	0.91674 (5)	0.31518 (3)	0.0865 (2)
C14	0.84784 (8)	1.06116 (5)	0.04585 (3)	0.0869 (2)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C1	0.0336 (8)	0.0493 (10)	0.0569 (10)	-0.0039 (7)	-0.0009 (7)	0.0105 (8)
C2	0.0427 (9)	0.0468 (10)	0.0576 (10)	-0.0041 (8)	-0.0106 (8)	0.0080 (8)
C3	0.0452 (11)	0.0592 (14)	0.185 (3)	0.0027 (10)	0.0251 (15)	0.0048 (16)
C4	0.0598 (15)	0.0540 (15)	0.247 (4)	0.0129 (12)	0.004 (2)	-0.001 (2)
C5	0.0892 (19)	0.0524 (14)	0.139 (3)	-0.0148 (13)	-0.0468 (18)	0.0265 (16)
C6	0.163 (3)	0.0603 (15)	0.0716 (15)	-0.0325 (18)	0.0185 (17)	0.0136 (12)
C7	0.125 (2)	0.0535 (12)	0.0637 (13)	-0.0217 (13)	0.0321 (13)	0.0015 (10)
C8	0.0290 (8)	0.0467 (10)	0.0905 (15)	-0.0013 (7)	0.0021 (8)	0.0144 (10)
C9	0.0424 (10)	0.0583 (12)	0.0961 (16)	-0.0013 (9)	-0.0076 (10)	0.0177 (11)
C10	0.0614 (13)	0.0643 (15)	0.114 (2)	0.0050 (11)	-0.0110 (13)	0.0311 (14)
C11	0.0612 (13)	0.0495 (13)	0.145 (3)	0.0099 (10)	0.0081 (14)	0.0293 (15)
C12	0.0664 (14)	0.0485 (12)	0.133 (2)	-0.0020 (10)	0.0192 (15)	-0.0005 (14)
C13	0.0478 (10)	0.0548 (12)	0.0976 (16)	-0.0053 (9)	0.0116 (10)	0.0053 (11)
N1	0.0312 (7)	0.0452 (8)	0.0843 (11)	-0.0067 (6)	0.0043 (7)	0.0129 (8)
O1	0.0320 (6)	0.0573 (8)	0.1014 (11)	-0.0022 (5)	0.0056 (6)	0.0213 (7)
Cl1	0.1026 (5)	0.0785 (4)	0.0927 (5)	-0.0146 (3)	-0.0221 (4)	0.0098 (3)

C12	0.1321 (6)	0.0836 (5)	0.0940 (5)	-0.0293 (4)	0.0131 (4)	-0.0028 (4)
C21	0.0333 (8)	0.0438 (9)	0.0596 (10)	0.0012 (7)	0.0003 (7)	-0.0075 (8)
C22	0.0382 (9)	0.0435 (9)	0.0679 (12)	0.0031 (7)	-0.0111 (8)	-0.0117 (8)
C23	0.0445 (11)	0.0524 (12)	0.145 (2)	-0.0056 (9)	0.0107 (12)	-0.0206 (13)
C24	0.0651 (15)	0.0515 (14)	0.192 (3)	-0.0098 (11)	-0.0021 (18)	-0.0332 (17)
C25	0.0757 (16)	0.0631 (15)	0.134 (2)	0.0173 (13)	-0.0220 (16)	-0.0463 (15)
C26	0.0871 (16)	0.0718 (15)	0.0681 (14)	0.0202 (13)	-0.0060 (12)	-0.0257 (11)
C27	0.0664 (12)	0.0531 (11)	0.0563 (11)	0.0077 (9)	-0.0004 (9)	-0.0102 (9)
C28	0.0288 (7)	0.0404 (9)	0.0547 (10)	-0.0010 (6)	0.0049 (7)	-0.0089 (7)
C29	0.0464 (9)	0.0506 (10)	0.0536 (10)	0.0008 (8)	0.0050 (8)	-0.0060 (8)
C30	0.0624 (12)	0.0631 (13)	0.0613 (12)	-0.0083 (10)	0.0060 (10)	-0.0219 (10)
C31	0.0693 (13)	0.0448 (11)	0.0887 (16)	-0.0085 (9)	0.0194 (11)	-0.0236 (11)
C32	0.0701 (13)	0.0409 (10)	0.0862 (16)	-0.0001 (9)	0.0148 (11)	0.0002 (10)
C33	0.0468 (9)	0.0513 (10)	0.0554 (10)	0.0013 (8)	0.0054 (8)	-0.0051 (8)
N2	0.0290 (6)	0.0427 (8)	0.0634 (9)	0.0026 (6)	0.0029 (6)	-0.0147 (6)
O2	0.0316 (6)	0.0583 (8)	0.1137 (12)	-0.0017 (6)	0.0088 (7)	-0.0256 (8)
Cl3	0.1228 (5)	0.0719 (4)	0.0638 (4)	0.0113 (3)	-0.0019 (3)	0.0093 (3)
Cl4	0.1163 (5)	0.0878 (4)	0.0553 (3)	0.0093 (4)	-0.0057 (3)	0.0019 (3)

Geometric parameters (Å, °)

C1—O1	1.2214 (19)	C21—O2	1.2258 (19)
C1—N1	1.346 (2)	C21—N2	1.341 (2)
C1—C2	1.494 (2)	C21—C22	1.497 (2)
C2—C3	1.368 (3)	C22—C23	1.379 (3)
C2—C7	1.373 (3)	C22—C27	1.386 (3)
C3—C4	1.391 (4)	C23—C24	1.388 (3)
C3—H3	0.93	C23—H23	0.93
C4—C5	1.364 (5)	C24—C25	1.366 (4)
C4—H4	0.93	C24—H24	0.93
C5—C6	1.335 (4)	C25—C26	1.362 (4)
C5—H5	0.93	C25—H25	0.93
C6—C7	1.380 (3)	C26—C27	1.390 (3)
C6—H6	0.93	C26—H26	0.93
C7—H7	0.93	C27—H27	0.93
C8—C9	1.389 (3)	C28—C29	1.386 (2)
C8—C13	1.389 (3)	C28—C33	1.391 (3)
C8—N1	1.421 (2)	C28—N2	1.417 (2)
C9—C10	1.382 (3)	C29—C30	1.382 (3)
C9—Cl1	1.741 (3)	C29—Cl3	1.734 (2)
C10—C11	1.368 (4)	C30—C31	1.365 (3)
C10—H10	0.93	C30—H30	0.93
C11—C12	1.374 (4)	C31—C32	1.374 (3)
C11—H11	0.93	C31—H31	0.93
C12—C13	1.391 (3)	C32—C33	1.379 (3)
C12—H12	0.93	C32—H32	0.93
C13—Cl2	1.736 (3)	C33—Cl4	1.732 (2)
N1—H1N	0.860 (19)	N2—H2N	0.843 (18)
O1—C1—N1	121.41 (16)	O2—C21—N2	121.88 (16)

supplementary materials

O1—C1—C2	122.15 (16)	O2—C21—C22	122.66 (15)
N1—C1—C2	116.43 (14)	N2—C21—C22	115.46 (14)
C3—C2—C7	118.2 (2)	C23—C22—C27	119.09 (18)
C3—C2—C1	119.51 (18)	C23—C22—C21	119.21 (18)
C7—C2—C1	122.29 (18)	C27—C22—C21	121.68 (16)
C2—C3—C4	120.1 (3)	C22—C23—C24	119.9 (2)
C2—C3—H3	120	C22—C23—H23	120
C4—C3—H3	120	C24—C23—H23	120
C5—C4—C3	120.3 (3)	C25—C24—C23	120.4 (2)
C5—C4—H4	119.9	C25—C24—H24	119.8
C3—C4—H4	119.9	C23—C24—H24	119.8
C6—C5—C4	119.9 (2)	C26—C25—C24	120.4 (2)
C6—C5—H5	120.1	C26—C25—H25	119.8
C4—C5—H5	120.1	C24—C25—H25	119.8
C5—C6—C7	120.5 (3)	C25—C26—C27	119.9 (2)
C5—C6—H6	119.8	C25—C26—H26	120.1
C7—C6—H6	119.8	C27—C26—H26	120.1
C2—C7—C6	121.0 (3)	C22—C27—C26	120.3 (2)
C2—C7—H7	119.5	C22—C27—H27	119.9
C6—C7—H7	119.5	C26—C27—H27	119.9
C9—C8—C13	117.49 (18)	C29—C28—C33	117.15 (15)
C9—C8—N1	120.5 (2)	C29—C28—N2	121.70 (16)
C13—C8—N1	122.0 (2)	C33—C28—N2	121.06 (16)
C10—C9—C8	122.3 (2)	C30—C29—C28	121.88 (18)
C10—C9—C11	119.2 (2)	C30—C29—C13	119.07 (15)
C8—C9—C11	118.51 (16)	C28—C29—C13	119.05 (13)
C11—C10—C9	118.8 (3)	C31—C30—C29	119.13 (19)
C11—C10—H10	120.6	C31—C30—H30	120.4
C9—C10—H10	120.6	C29—C30—H30	120.4
C10—C11—C12	120.9 (2)	C30—C31—C32	120.99 (18)
C10—C11—H11	119.6	C30—C31—H31	119.5
C12—C11—H11	119.6	C32—C31—H31	119.5
C11—C12—C13	119.8 (3)	C31—C32—C33	119.3 (2)
C11—C12—H12	120.1	C31—C32—H32	120.4
C13—C12—H12	120.1	C33—C32—H32	120.4
C8—C13—C12	120.7 (2)	C32—C33—C28	121.57 (18)
C8—C13—C12	119.53 (16)	C32—C33—C14	119.47 (16)
C12—C13—C12	119.8 (2)	C28—C33—C14	118.95 (14)
C1—N1—C8	121.29 (14)	C21—N2—C28	123.19 (14)
C1—N1—H1N	119.9 (14)	C21—N2—H2N	121.5 (13)
C8—N1—H1N	117.2 (15)	C28—N2—H2N	115.3 (13)
O1—C1—C2—C3	30.4 (3)	O2—C21—C22—C23	-34.2 (3)
N1—C1—C2—C3	-150.5 (2)	N2—C21—C22—C23	145.9 (2)
O1—C1—C2—C7	-148.3 (2)	O2—C21—C22—C27	144.2 (2)
N1—C1—C2—C7	30.8 (3)	N2—C21—C22—C27	-35.7 (3)
C7—C2—C3—C4	3.0 (4)	C27—C22—C23—C24	-0.2 (4)
C1—C2—C3—C4	-175.7 (3)	C21—C22—C23—C24	178.3 (2)
C2—C3—C4—C5	0.2 (5)	C22—C23—C24—C25	-0.8 (4)
C3—C4—C5—C6	-3.3 (5)	C23—C24—C25—C26	1.0 (5)

C4—C5—C6—C7	3.0 (5)	C24—C25—C26—C27	-0.4 (4)
C3—C2—C7—C6	-3.3 (4)	C23—C22—C27—C26	0.9 (3)
C1—C2—C7—C6	175.4 (2)	C21—C22—C27—C26	-177.54 (18)
C5—C6—C7—C2	0.4 (4)	C25—C26—C27—C22	-0.6 (3)
C13—C8—C9—C10	1.5 (3)	C33—C28—C29—C30	-1.4 (3)
N1—C8—C9—C10	-176.03 (18)	N2—C28—C29—C30	175.06 (16)
C13—C8—C9—C11	-176.52 (14)	C33—C28—C29—C13	178.37 (13)
N1—C8—C9—C11	5.9 (2)	N2—C28—C29—C13	-5.1 (2)
C8—C9—C10—C11	-1.0 (3)	C28—C29—C30—C31	1.2 (3)
C11—C9—C10—C11	177.08 (18)	C13—C29—C30—C31	-178.63 (16)
C9—C10—C11—C12	0.3 (4)	C29—C30—C31—C32	-0.3 (3)
C10—C11—C12—C13	-0.3 (4)	C30—C31—C32—C33	-0.3 (3)
C9—C8—C13—C12	-1.5 (3)	C31—C32—C33—C28	0.0 (3)
N1—C8—C13—C12	176.04 (18)	C31—C32—C33—C14	-179.03 (16)
C9—C8—C13—C12	179.02 (14)	C29—C28—C33—C32	0.8 (3)
N1—C8—C13—C12	-3.4 (2)	N2—C28—C33—C32	-175.69 (17)
C11—C12—C13—C8	0.9 (3)	C29—C28—C33—C14	179.88 (12)
C11—C12—C13—C12	-179.60 (17)	N2—C28—C33—C14	3.4 (2)
O1—C1—N1—C8	-8.5 (3)	O2—C21—N2—C28	-5.4 (3)
C2—C1—N1—C8	172.35 (18)	C22—C21—N2—C28	174.49 (16)
C9—C8—N1—C1	-84.0 (2)	C29—C28—N2—C21	98.8 (2)
C13—C8—N1—C1	98.5 (2)	C33—C28—N2—C21	-84.8 (2)

Hydrogen-bond geometry (\AA , $^\circ$)

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
N1—H1N \cdots O2	0.860 (19)	2.09 (2)	2.9035 (18)	158 (2)
N2—H2N \cdots O1 ⁱ	0.843 (18)	2.060 (19)	2.8831 (18)	165.1 (19)

Symmetry codes: (i) $x+1, y, z$.

Fig. 1




Fig. 2

