

Poly[[bis[μ_2 -8-ethyl-5-oxo-2-(piperazin-1-yl)-5,8-dihydropyrido[2,3-*d*]pyrimidine-6-carboxylato]manganese(II)] dihydrate]

Jing Huang,^a Wei-Ping Hu^b and Zhe An^{a*}

^aSchool of Pharmaceutical Science, Harbin Medical University, Harbin 150086, People's Republic of China, and ^bSecond Hospital, Harbin Medical University, Harbin 150086, People's Republic of China
Correspondence e-mail: anzhe6409@sina.com


Received 18 February 2008; accepted 8 March 2008

Key indicators: single-crystal X-ray study; $T = 295\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.004\text{ \AA}$; H-atom completeness 89%; disorder in solvent or counterion; R factor = 0.061; wR factor = 0.192; data-to-parameter ratio = 17.3.

In the title compound, $\{[\text{Mn}(\text{C}_{14}\text{H}_{16}\text{N}_5\text{O}_3)_2]\cdot 2\text{H}_2\text{O}\}_n$, the Mn^{II} atom (site symmetry $\bar{1}$) exhibits a distorted *trans*- MnN_2O_4 octahedral geometry defined by two monodentate *N*-bonded and two bidentate *O,O'*-bonded 8-ethyl-5-oxo-2-(piperazin-1-yl)-5,8-dihydropyrido[2,3-*d*]pyrimidine-6-carboxylate anions. An N—H···O hydrogen bond is present in the crystal structure. The extended two-dimensional structure is a square grid and the disordered uncoordinated water molecules occupy cavities within the grid.

Related literature

For background, see: Mizuki *et al.* (1996).



Experimental

Crystal data

$[\text{Mn}(\text{C}_{14}\text{H}_{16}\text{N}_5\text{O}_3)_2]\cdot 2\text{H}_2\text{O}$	$V = 1636.8 (1)\text{ \AA}^3$
$M_r = 695.58$	$Z = 2$
Monoclinic, $P2_1/c$	$\text{Mo } K\alpha$ radiation
$a = 6.0422 (2)\text{ \AA}$	$\mu = 0.47\text{ mm}^{-1}$
$b = 21.5673 (8)\text{ \AA}$	$T = 295 (2)\text{ K}$
$c = 12.7395 (5)\text{ \AA}$	$0.34 \times 0.26 \times 0.18\text{ mm}$
$\beta = 99.617 (1)^\circ$	

Data collection

Bruker SMART CCD diffractometer	10030 measured reflections
Absorption correction: multi-scan (<i>SADABS</i> ; Bruker, 1998)	3938 independent reflections
$T_{\min} = 0.861$, $T_{\max} = 0.910$	3465 reflections with $I > 2\sigma(I)$
	$R_{\text{int}} = 0.019$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.060$	H atoms treated by a mixture of independent and constrained refinement
$wR(F^2) = 0.191$	$\Delta\rho_{\text{max}} = 0.97\text{ e \AA}^{-3}$
$S = 1.10$	$\Delta\rho_{\text{min}} = -0.48\text{ e \AA}^{-3}$
3938 reflections	
227 parameters	
1 restraint	

Table 1
Selected bond lengths (Å).

Mn1—O1	$2.106 (2)$	Mn1—N5^{i}	$2.372 (2)$
Mn1—O3	$2.1667 (16)$		

Symmetry code: (i) $x + 1$, $-y + \frac{3}{2}$, $z + \frac{1}{2}$.

Table 2
Hydrogen-bond geometry (Å, °).

$D—H \cdots A$	$D—H$	$H \cdots A$	$D \cdots A$	$D—H \cdots A$
$\text{N5—H5N} \cdots \text{O2}^{\text{ii}}$	$0.893 (10)$	$2.268 (12)$	$3.149 (3)$	$169 (3)$

Symmetry code: (ii) $-x + 1$, $y + \frac{1}{2}$, $-z + \frac{1}{2}$.

Data collection: *SMART* (Bruker, 1998); cell refinement: *SAINT-Plus* (Bruker, 1998); data reduction: *SAINT-Plus*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL*.

The authors thank the Innovation Science Foundation of Harbin Medical University for financial support (grant No. 060041).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HB2703).

References

- Bruker (1998). *SMART*, *SAINT-Plus* and *SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Mizuki, Y., Fujiwara, I. & Yamaguchi, T. (1996). *J. Antimicrob. Chemother. Suppl. A*, **37**, 41–45.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.

supplementary materials

Acta Cryst. (2008). E64, m547 [doi:10.1107/S1600536808006533]

Poly[[bis(μ_2 -8-ethyl-5-oxo-2-(piperazin-1-yl)-5,8-dihdropyrido[2,3-*d*]pyrimidine-6-carboxylato]manganese(II)] dihydrate]

J. Huang, W.-P. Hu and Z. An

Comment

Pipemidic acid (Hppa, C₁₄H₁₇N₅O₃, 8-Ethyl-5,8-dihydro-5-oxo-2-(1-piperazinyl)-pyrido(2,3-*d*)-pyrimidine-6-carboxylic acid) is member of a class of quinolones used to treat infections (Mizuki *et al.*, 1996). The metal complexes of the ppa anion have not been reported; the title manganese(II) complex, (I), is reported here (Fig. 1).

The Mn^{II} atom in (I) with site symmetry $\bar{1}$ is coordinated by four oxygen atoms and two N atoms from four ppa ligands (two monodentate-N and two O,O-bidentate) (Table 1) to form a square grid propagating in (Fig. 2). An N—H···O hydrogen bond (Table 2) helps to stabilize this arrangement.

The disordered, uncoordinated, water molecules occupy cavities within the grid. In the present study, their attached H atoms could not be located.

Experimental

A mixture of Mn(CH₃COO)₂·4H₂O (0.061 g, 0.25 mmol), Hppa (0.15 g, 0.5 mmol), sodium hydroxide (0.04 g, 1 mmol) and water (12 ml) was stirred for 30 min in air. The mixture was then transferred to a 23 ml Teflon-lined hydrothermal bomb. The bomb was kept at 433 K for 72 h under autogenous pressure. Upon cooling, colourless prisms of (I) were obtained from the reaction mixture.

Refinement

The carbon-bound H atoms were positioned geometrically (C—H = 0.93–0.97 Å) and refined as riding with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$. The N-bound H atom was located in a difference map and its position was freely refined with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{N})$.

The water H atoms could not be placed due to disorder.

Figures




Fig. 1. The asymmetric unit of (I) showing the showing 50% displacement ellipsoids (water molecule O atoms have been omitted for clarity).

supplementary materials




Fig. 2. A view of part of a two-dimensional polymeric sheet in (I) showing the square-grid connectivity (H atoms and water molecule O atoms omitted for clarity).

Poly[[bis[μ_2 -8-ethyl-5-oxo-2-(piperazin-1-yl)-5,8-dihydropyrido[2,3-*d*]pyrimidine-6-carboxylato]manganese(II)] dihydrate]

Crystal data

[Mn(C ₁₄ H ₁₆ N ₅ O ₃) ₂] ₂ H ₂ O	$F_{000} = 718.0$
$M_r = 695.58$	$D_x = 1.399 \text{ Mg m}^{-3}$
Monoclinic, $P2_1/c$	Mo $K\alpha$ radiation
Hall symbol: -P 2ybc	$\lambda = 0.71073 \text{ \AA}$
$a = 6.0422 (2) \text{ \AA}$	Cell parameters from 4747 reflections
$b = 21.5673 (8) \text{ \AA}$	$\theta = 2.5\text{--}28.3^\circ$
$c = 12.7395 (5) \text{ \AA}$	$\mu = 0.47 \text{ mm}^{-1}$
$\beta = 99.617 (1)^\circ$	$T = 295 (2) \text{ K}$
$V = 1636.8 (1) \text{ \AA}^3$	Prism, colorless
$Z = 2$	$0.34 \times 0.26 \times 0.18 \text{ mm}$

Data collection

Bruker SMART CCD diffractometer	3938 independent reflections
Radiation source: fine-focus sealed tube	3465 reflections with $I > 2\sigma(I)$
Monochromator: graphite	$R_{\text{int}} = 0.019$
$T = 295(2) \text{ K}$	$\theta_{\text{max}} = 28.3^\circ$
ω scans	$\theta_{\text{min}} = 2.5^\circ$
Absorption correction: multi-scan (SADABS; Sheldrick, 1996)	$h = -6 \rightarrow 8$
$T_{\text{min}} = 0.861$, $T_{\text{max}} = 0.910$	$k = -27 \rightarrow 28$
10030 measured reflections	$l = -15 \rightarrow 16$

Refinement

Refinement on F^2	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.061$	H atoms treated by a mixture of independent and constrained refinement
$wR(F^2) = 0.191$	$w = 1/[\sigma^2(F_o^2) + (0.1066P)^2 + 1.0983P]$ where $P = (F_o^2 + 2F_c^2)/3$
$S = 1.11$	$(\Delta/\sigma)_{\text{max}} < 0.001$
3938 reflections	$\Delta\rho_{\text{max}} = 0.98 \text{ e \AA}^{-3}$

227 parameters $\Delta\rho_{\min} = -0.48 \text{ e } \text{\AA}^{-3}$
 1 restraint Extinction correction: none
 Primary atom site location: structure-invariant direct
 methods

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > 2\text{sigma}(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$	Occ. (<1)
Mn1	0.5000	0.5000	0.5000	0.02831 (19)	
O1W	0.388 (4)	0.5235 (7)	0.9613 (10)	0.244 (9)	0.50
O1	0.7068 (3)	0.49893 (7)	0.38228 (16)	0.0341 (4)	
O2W	-0.081 (2)	0.4426 (8)	0.9246 (7)	0.194 (6)	0.50
O2	0.8721 (5)	0.51863 (13)	0.2451 (2)	0.0717 (8)	
O3	0.3574 (3)	0.58106 (8)	0.41354 (14)	0.0378 (4)	
N1	0.5078 (5)	0.67158 (12)	0.1504 (2)	0.0533 (7)	
N2	-0.0007 (4)	0.73723 (11)	0.2985 (2)	0.0464 (6)	
N3	0.2395 (4)	0.74681 (10)	0.16596 (18)	0.0432 (5)	
N4	-0.0102 (4)	0.82368 (10)	0.19027 (19)	0.0384 (5)	
N5	-0.2339 (3)	0.93726 (9)	0.11005 (17)	0.0322 (4)	
H5N	-0.140 (4)	0.9649 (12)	0.146 (2)	0.048*	
C1	0.7282 (4)	0.52991 (11)	0.3018 (2)	0.0358 (5)	
C2	0.5782 (4)	0.58537 (11)	0.2731 (2)	0.0354 (5)	
C3	0.4044 (4)	0.60559 (10)	0.33118 (19)	0.0313 (5)	
C4	0.2852 (4)	0.66087 (11)	0.28824 (19)	0.0336 (5)	
C5	0.1042 (5)	0.68542 (12)	0.3304 (2)	0.0425 (6)	
H5A	0.0546	0.6635	0.3848	0.051*	
C6	0.0805 (4)	0.76813 (12)	0.2189 (2)	0.0360 (5)	
C7	0.3383 (5)	0.69362 (12)	0.2010 (2)	0.0395 (6)	
C8	0.6170 (5)	0.61896 (14)	0.1875 (2)	0.0496 (7)	
H8A	0.7282	0.6046	0.1512	0.059*	
C9	0.5717 (8)	0.7043 (2)	0.0566 (3)	0.0725 (12)	
H9A	0.5531	0.7486	0.0645	0.087*	
H9B	0.7285	0.6963	0.0537	0.087*	
C10	0.4373 (10)	0.6840 (4)	-0.0400 (5)	0.116 (2)	
H10A	0.4583	0.6403	-0.0488	0.174*	
H10B	0.4811	0.7059	-0.0989	0.174*	
H10C	0.2822	0.6923	-0.0375	0.174*	

supplementary materials

C11	-0.1446 (6)	0.85820 (14)	0.2560 (2)	0.0499 (7)
H11A	-0.0477	0.8852	0.3045	0.060*
H11B	-0.2170	0.8295	0.2980	0.060*
C12	-0.3226 (5)	0.89702 (13)	0.1855 (2)	0.0433 (6)
H12A	-0.4326	0.8693	0.1460	0.052*
H12B	-0.3996	0.9225	0.2308	0.052*
C13	-0.1018 (4)	0.89946 (12)	0.0466 (2)	0.0372 (5)
H13A	-0.0347	0.9264	-0.0003	0.045*
H13B	-0.2015	0.8710	0.0025	0.045*
C14	0.0812 (4)	0.86281 (12)	0.1146 (2)	0.0382 (6)
H14A	0.1573	0.8371	0.0693	0.046*
H14B	0.1906	0.8911	0.1530	0.046*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Mn1	0.0350 (3)	0.0187 (3)	0.0304 (3)	-0.00041 (16)	0.0030 (2)	0.00255 (16)
O1W	0.47 (3)	0.155 (12)	0.111 (9)	0.000 (15)	0.058 (14)	-0.057 (9)
O1	0.0407 (10)	0.0237 (9)	0.0383 (10)	0.0041 (6)	0.0073 (8)	0.0040 (6)
O2W	0.203 (11)	0.307 (17)	0.088 (6)	-0.110 (12)	0.076 (7)	-0.028 (8)
O2	0.0854 (18)	0.0730 (16)	0.0674 (16)	0.0491 (15)	0.0441 (14)	0.0338 (13)
O3	0.0448 (10)	0.0297 (9)	0.0403 (9)	0.0085 (7)	0.0110 (7)	0.0122 (7)
N1	0.0730 (17)	0.0433 (13)	0.0500 (14)	0.0245 (12)	0.0294 (12)	0.0196 (11)
N2	0.0498 (13)	0.0386 (12)	0.0550 (14)	0.0152 (10)	0.0211 (11)	0.0213 (11)
N3	0.0549 (13)	0.0339 (11)	0.0434 (12)	0.0165 (10)	0.0157 (10)	0.0150 (9)
N4	0.0418 (11)	0.0301 (10)	0.0458 (12)	0.0107 (9)	0.0146 (9)	0.0133 (9)
N5	0.0344 (10)	0.0231 (9)	0.0382 (10)	0.0029 (7)	0.0034 (8)	0.0012 (8)
C1	0.0410 (13)	0.0311 (12)	0.0355 (12)	0.0080 (10)	0.0071 (10)	0.0029 (9)
C2	0.0429 (13)	0.0281 (11)	0.0360 (12)	0.0084 (9)	0.0092 (10)	0.0040 (9)
C3	0.0367 (11)	0.0227 (10)	0.0337 (11)	0.0032 (9)	0.0033 (9)	0.0042 (8)
C4	0.0396 (12)	0.0265 (11)	0.0352 (12)	0.0060 (9)	0.0073 (9)	0.0067 (9)
C5	0.0491 (15)	0.0339 (13)	0.0475 (15)	0.0103 (11)	0.0167 (12)	0.0168 (11)
C6	0.0377 (12)	0.0307 (12)	0.0398 (13)	0.0066 (9)	0.0066 (10)	0.0090 (10)
C7	0.0506 (14)	0.0313 (12)	0.0386 (13)	0.0109 (11)	0.0132 (11)	0.0082 (10)
C8	0.0609 (17)	0.0438 (15)	0.0485 (16)	0.0220 (13)	0.0219 (13)	0.0120 (12)
C9	0.091 (3)	0.071 (2)	0.065 (2)	0.035 (2)	0.041 (2)	0.0293 (19)
C10	0.111 (4)	0.154 (6)	0.085 (4)	0.032 (4)	0.022 (3)	0.021 (4)
C11	0.0604 (17)	0.0465 (15)	0.0478 (15)	0.0251 (14)	0.0234 (13)	0.0176 (13)
C12	0.0427 (13)	0.0368 (13)	0.0535 (16)	0.0129 (11)	0.0169 (12)	0.0133 (12)
C13	0.0394 (13)	0.0308 (12)	0.0414 (13)	0.0087 (10)	0.0071 (10)	0.0078 (10)
C14	0.0365 (12)	0.0304 (12)	0.0497 (14)	0.0075 (9)	0.0128 (11)	0.0136 (10)

Geometric parameters (\AA , $^\circ$)

Mn1—O1	2.106 (2)	C2—C8	1.361 (4)
Mn1—O1 ⁱ	2.106 (2)	C2—C3	1.450 (3)
Mn1—O3	2.1667 (16)	C3—C4	1.452 (3)
Mn1—O3 ^j	2.1667 (16)	C4—C7	1.399 (3)

Mn1—N5 ⁱⁱ	2.372 (2)	C4—C5	1.400 (4)
Mn1—N5 ⁱⁱⁱ	2.3723 (19)	C5—H5A	0.9300
O1—C1	1.248 (3)	C8—H8A	0.9300
O2—C1	1.244 (3)	C9—C10	1.425 (8)
O3—C3	1.249 (3)	C9—H9A	0.9700
N1—C8	1.357 (3)	C9—H9B	0.9700
N1—C7	1.382 (4)	C10—H10A	0.9600
N1—C9	1.493 (4)	C10—H10B	0.9600
N2—C5	1.315 (3)	C10—H10C	0.9600
N2—C6	1.371 (4)	C11—C12	1.531 (4)
N3—C7	1.335 (3)	C11—H11A	0.9700
N3—C6	1.344 (3)	C11—H11B	0.9700
N4—C6	1.343 (3)	C12—H12A	0.9700
N4—C14	1.457 (3)	C12—H12B	0.9700
N4—C11	1.463 (3)	C13—C14	1.510 (3)
N5—C12	1.461 (3)	C13—H13A	0.9700
N5—C13	1.474 (3)	C13—H13B	0.9700
N5—Mn1 ^{iv}	2.3723 (19)	C14—H14A	0.9700
N5—H5N	0.90 (3)	C14—H14B	0.9700
C1—C2	1.508 (3)		
O1—Mn1—O1 ⁱ	180.0	N4—C6—N3	117.5 (2)
O1—Mn1—O3	83.09 (6)	N4—C6—N2	117.0 (2)
O1 ⁱ —Mn1—O3	96.91 (6)	N3—C6—N2	125.4 (2)
O1—Mn1—O3 ⁱ	96.91 (6)	N3—C7—N1	117.7 (2)
O1 ⁱ —Mn1—O3 ⁱ	83.09 (6)	N3—C7—C4	123.4 (2)
O3—Mn1—O3 ⁱ	180.0	N1—C7—C4	118.9 (2)
O1—Mn1—N5 ⁱⁱ	90.17 (7)	N1—C8—C2	125.8 (3)
O1 ⁱ —Mn1—N5 ⁱⁱ	89.83 (7)	N1—C8—H8A	117.1
O3—Mn1—N5 ⁱⁱ	90.74 (7)	C2—C8—H8A	117.1
O3 ⁱ —Mn1—N5 ⁱⁱ	89.26 (7)	C10—C9—N1	111.2 (5)
O1—Mn1—N5 ⁱⁱⁱ	89.83 (7)	C10—C9—H9A	109.4
O1 ⁱ —Mn1—N5 ⁱⁱⁱ	90.17 (7)	N1—C9—H9A	109.4
O3—Mn1—N5 ⁱⁱⁱ	89.26 (7)	C10—C9—H9B	109.4
O3 ⁱ —Mn1—N5 ⁱⁱⁱ	90.73 (7)	N1—C9—H9B	109.4
N5 ⁱⁱ —Mn1—N5 ⁱⁱⁱ	180.0	H9A—C9—H9B	108.0
C1—O1—Mn1	137.22 (16)	C9—C10—H10A	109.5
C3—O3—Mn1	129.93 (16)	C9—C10—H10B	109.5
C8—N1—C7	118.7 (2)	H10A—C10—H10B	109.5
C8—N1—C9	119.8 (3)	C9—C10—H10C	109.5
C7—N1—C9	121.4 (2)	H10A—C10—H10C	109.5
C5—N2—C6	115.3 (2)	H10B—C10—H10C	109.5
C7—N3—C6	116.3 (2)	N4—C11—C12	110.2 (2)
C6—N4—C14	120.9 (2)	N4—C11—H11A	109.6
C6—N4—C11	122.6 (2)	C12—C11—H11A	109.6
C14—N4—C11	113.1 (2)	N4—C11—H11B	109.6

supplementary materials

C12—N5—C13	108.90 (19)	C12—C11—H11B	109.6
C12—N5—Mn1 ^{iv}	116.15 (15)	H11A—C11—H11B	108.1
C13—N5—Mn1 ^{iv}	111.54 (14)	N5—C12—C11	114.3 (2)
C12—N5—H5N	110 (2)	N5—C12—H12A	108.7
C13—N5—H5N	107 (2)	C11—C12—H12A	108.7
Mn1 ^{iv} —N5—H5N	103 (2)	N5—C12—H12B	108.7
O2—C1—O1	123.4 (2)	C11—C12—H12B	108.7
O2—C1—C2	117.6 (2)	H12A—C12—H12B	107.6
O1—C1—C2	118.9 (2)	N5—C13—C14	112.8 (2)
C8—C2—C3	119.0 (2)	N5—C13—H13A	109.0
C8—C2—C1	116.1 (2)	C14—C13—H13A	109.0
C3—C2—C1	124.9 (2)	N5—C13—H13B	109.0
O3—C3—C2	126.0 (2)	C14—C13—H13B	109.0
O3—C3—C4	119.8 (2)	H13A—C13—H13B	107.8
C2—C3—C4	114.3 (2)	N4—C14—C13	111.1 (2)
C7—C4—C5	114.3 (2)	N4—C14—H14A	109.4
C7—C4—C3	123.3 (2)	C13—C14—H14A	109.4
C5—C4—C3	122.4 (2)	N4—C14—H14B	109.4
N2—C5—C4	124.8 (2)	C13—C14—H14B	109.4
N2—C5—H5A	117.6	H14A—C14—H14B	108.0
C4—C5—H5A	117.6		
O3—Mn1—O1—C1	-1.4 (3)	C7—N3—C6—N4	175.3 (3)
O3 ⁱ —Mn1—O1—C1	178.6 (3)	C7—N3—C6—N2	-6.2 (4)
N5 ⁱⁱ —Mn1—O1—C1	-92.1 (3)	C5—N2—C6—N4	-174.7 (3)
N5 ⁱⁱⁱ —Mn1—O1—C1	87.9 (3)	C5—N2—C6—N3	6.8 (4)
O1—Mn1—O3—C3	1.4 (2)	C6—N3—C7—N1	-177.9 (3)
O1 ⁱ —Mn1—O3—C3	-178.6 (2)	C6—N3—C7—C4	-0.5 (4)
N5 ⁱⁱ —Mn1—O3—C3	91.5 (2)	C8—N1—C7—N3	177.3 (3)
N5 ⁱⁱⁱ —Mn1—O3—C3	-88.5 (2)	C9—N1—C7—N3	-2.9 (5)
Mn1—O1—C1—O2	179.3 (2)	C8—N1—C7—C4	-0.1 (5)
Mn1—O1—C1—C2	1.5 (4)	C9—N1—C7—C4	179.7 (3)
O2—C1—C2—C8	-1.0 (4)	C5—C4—C7—N3	5.6 (4)
O1—C1—C2—C8	177.0 (3)	C3—C4—C7—N3	-175.2 (3)
O2—C1—C2—C3	-179.1 (3)	C5—C4—C7—N1	-177.1 (3)
O1—C1—C2—C3	-1.2 (4)	C3—C4—C7—N1	2.1 (4)
Mn1—O3—C3—C2	-1.8 (4)	C7—N1—C8—C2	-1.3 (5)
Mn1—O3—C3—C4	-179.39 (16)	C9—N1—C8—C2	178.8 (4)
C8—C2—C3—O3	-176.6 (3)	C3—C2—C8—N1	0.8 (5)
C1—C2—C3—O3	1.5 (4)	C1—C2—C8—N1	-177.5 (3)
C8—C2—C3—C4	1.1 (4)	C8—N1—C9—C10	92.7 (5)
C1—C2—C3—C4	179.2 (2)	C7—N1—C9—C10	-87.1 (5)
O3—C3—C4—C7	175.3 (2)	C6—N4—C11—C12	-148.9 (3)
C2—C3—C4—C7	-2.5 (4)	C14—N4—C11—C12	51.9 (3)
O3—C3—C4—C5	-5.6 (4)	C13—N5—C12—C11	54.0 (3)
C2—C3—C4—C5	176.6 (3)	Mn1 ^{iv} —N5—C12—C11	-179.1 (2)
C6—N2—C5—C4	-0.7 (5)	N4—C11—C12—N5	-52.9 (4)
C7—C4—C5—N2	-5.0 (4)	C12—N5—C13—C14	-55.0 (3)

C3—C4—C5—N2	175.8 (3)	Mn1 ^{iv} —N5—C13—C14	175.47 (16)
C14—N4—C6—N3	−8.0 (4)	C6—N4—C14—C13	146.2 (3)
C11—N4—C6—N3	−165.6 (3)	C11—N4—C14—C13	−54.2 (3)
C14—N4—C6—N2	173.4 (3)	N5—C13—C14—N4	55.9 (3)
C11—N4—C6—N2	15.8 (4)		

Symmetry codes: (i) $-x+1, -y+1, -z+1$; (ii) $x+1, -y+3/2, z+1/2$; (iii) $-x, y-1/2, -z+1/2$; (iv) $-x, y+1/2, -z+1/2$.

Hydrogen-bond geometry (\AA , °)

$D\text{—H}\cdots A$	$D\text{—H}$	$H\cdots A$	$D\cdots A$	$D\text{—H}\cdots A$
N5—H5N \cdots O2 ^v	0.893 (10)	2.268 (12)	3.149 (3)	169 (3)

Symmetry codes: (v) $-x+1, y+1/2, -z+1/2$.

supplementary materials

Fig. 1




Fig. 2

