

N-(Fluoren-9-ylmethoxycarbonyl)-L-leucine

Kazuhiko Yamada,^{a*} Daisuke Hashizume^b and Tadashi Shimizu^a

^aNational Institute for Materials Science, 3-13 Sakura, Tsukuba 305-0003, Japan, and

^bAdvanced Technology Support Division, RIKEN, 2-1 Hirosawa, Wako, Saitama 351-0198, Japan

Correspondence e-mail: yamada.kazuhiko@nims.go.jp

Received 18 April 2008; accepted 13 May 2008


Key indicators: single-crystal X-ray study; $T = 150\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.003\text{ \AA}$;

R factor = 0.040; wR factor = 0.111; data-to-parameter ratio = 9.8.

The title compound [systematic name: fluoren-9-yl *N*-(1-carboxy-3-methylbutyl)carbamate], $\text{C}_{21}\text{H}_{23}\text{NO}_4$, exhibits torsion angles that vary from the typical values found in other Fmoc-protected amino acids, *viz.* the orientations of the fluorene and carboxyl groups [$\text{C}-\text{O}-\text{C}-\text{C} = 93.8(2)$ and $\text{N}-\text{C}-\text{C}=\text{O} = -23.6(2)^\circ$]. The crystal structure exhibits two intermolecular hydrogen bonds ($\text{O}-\text{H}\cdots\text{O}$ and $\text{N}-\text{H}\cdots\text{O}$) that link the molecules into two-dimensional sheets parallel to the *ab* plane.

Related literature

For related literature on the structures of *N*- α -Fmoc-protected amino acids, see: Valle *et al.* (1984); Yamada *et al.* (2008).



Experimental

Crystal data

$\text{C}_{21}\text{H}_{23}\text{NO}_4$	$V = 1911.51(7)\text{ \AA}^3$
$M_r = 353.40$	$Z = 4$
Orthorhombic, $P2_12_12_1$	Mo $K\alpha$ radiation
$a = 5.4953(1)\text{ \AA}$	$\mu = 0.09\text{ mm}^{-1}$
$b = 14.2700(3)\text{ \AA}$	$T = 150\text{ K}$
$c = 24.3759(6)\text{ \AA}$	$0.40 \times 0.08 \times 0.06\text{ mm}$

Data collection

Rigaku AFC-8 diffractometer with Saturn70 CCD detector	3207 independent reflections
Absorption correction: none	2906 reflections with $I > 2\sigma(I)$
40257 measured reflections	$R_{\text{int}} = 0.055$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.039$	327 parameters
$wR(F^2) = 0.111$	All H-atom parameters refined
$S = 1.09$	$\Delta\rho_{\text{max}} = 0.27\text{ e \AA}^{-3}$
3207 reflections	$\Delta\rho_{\text{min}} = -0.26\text{ e \AA}^{-3}$

Table 1

Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{O}2-\text{H}2\text{H}\cdots\text{O}3^{\text{i}}$	0.85 (3)	1.82 (3)	2.6558 (17)	167 (3)
$\text{N}1-\text{H}1\text{N}\cdots\text{O}1^{\text{ii}}$	0.87 (3)	2.24 (3)	3.0751 (18)	161 (2)
$\text{C}8-\text{H}8\text{A}\cdots\text{O}1^{\text{iii}}$	0.90 (2)	2.51 (2)	3.392 (2)	166 (2)

Symmetry codes: (i) $-x, y + \frac{1}{2}, -z + \frac{1}{2}$; (ii) $x + 1, y, z$; (iii) $-x, y - \frac{1}{2}, -z + \frac{1}{2}$.

Data collection: *CrystalClear* (Rigaku/MSC, 2005); cell refinement: *HKL-2000* (Otwinowski & Minor, 1997); data reduction: *HKL-2000*; program(s) used to solve structure: *STR2004* (Burla *et al.*, 2005); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3 for Windows* (Farrugia, 1997); software used to prepare material for publication: *SHELXL97*.

KY thanks the Ministry of Education, Science, Sports, Culture, and Technology (MEXT) of Japan for funding this work [Young Scientists (B), 20750022].

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: FL2198).

References

- Burla, M. C., Caliandro, R., Camalli, M., Carrozzini, B., Cascarano, G. L., De Caro, L., Giacovazzo, C., Polidori, G. & Spagna, R. (2005). *J. Appl. Cryst.* **38**, 381–388.
- Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
- Otwinowski, Z. & Minor, W. (1997). *Methods in Enzymology*, Vol. 276, *Macromolecular Crystallography*, Part A, edited by C. W. Carter Jr & R. M. Sweet, pp. 307–326. New York: Academic Press.
- Rigaku/MSC (2005). *CrystalClear*. Rigaku/MSC Inc., The Woodlands, Texas, USA.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Valle, G., Bonora, G. M. & Toniolo, C. (1984). *Can. J. Chem.* **62**, 2661–2666.
- Yamada, K., Hashizume, D., Shimizu, T., Ohiki, S. & Yokoyama, S. (2008). *J. Mol. Struct.* In the press.

supplementary materials

Acta Cryst. (2008). E64, o1112 [doi:10.1107/S1600536808014372]

N-(Fluoren-9-ylmethoxycarbonyl)-L-leucine

K. Yamada, D. Hashizume and T. Shimizu

Comment

The fluoren-9-ylmethoxycarbonyl (Fmoc) group is currently one of the most frequently used protecting groups for peptide synthesis since rapid cleavages can be readily achieved under mild basic conditions with racemization-free results. Almost all the Fmoc-protected twenty-*L*-amino acids are commercially available. The crystal structures of *N*- α -Fmoc-protected-*L*-alanine monohydrate (II, Valle *et al.*, 1984) and *L*-serine (III, Yamada *et al.*, 2008) have been reported so far. In the present study, we have carried out the crystal structure analysis of *N*- α -Fmoc-*L*-leucine, (I).

The bond distances and bond angles of (I, Fig. 1) are consistent with the typical values of Fmoc-protected amino acids found in the other crystal structures. Some torsion angles, however, are found to be quite different. The torsion angle of O2—C6—C1—N1, for example, is -23.6 (2) $^{\circ}$, which is in disagreement with the previous observations in the Fmoc-protected amino acids in which the corresponding angles are 150.6 $^{\circ}$ and 175.8 $^{\circ}$ for (II) (Valle *et al.*, 1984) and (III) (Yamada *et al.*, 2008), respectively. Another example is that the torsion angle of C6—C1—N1—C7 in (I) is found to be -134.51 (15) $^{\circ}$, which is in reasonable agreement with that of (II), -151.6 $^{\circ}$, but is inconsistent with that found in (II), -65.6 $^{\circ}$. Each angle between the fluorene ring and the NC(δ b O)O plane is found to be different among the three Fmoc-protected amino acids. The torsion angles C7—O4—C8—C9 and O4—C8—C9—C10 for the title compound, for instance, are 93.78 (16) $^{\circ}$ and 60.54 (17) $^{\circ}$, respectively. On the other hand, the corresponding angles are -179.7 $^{\circ}$ and -172.1 $^{\circ}$, and 121.9 $^{\circ}$ and -68.2 $^{\circ}$ for (II) and (III), respectively.

Crystals of (I) contain two intermolecular hydrogen bonds (Table 1), which are formed between the carboxyl (O2—H2H) and amide oxygens (O3), and between the amide (N1—H1N) and the carbonyl (O1). The molecules are linked by O2—H2H \cdots O3 hydrogen bonds to form a chain structure along the *b* axis. The linkage is supported by an additional C—H \cdots O interaction (C8—H8A \cdots O1). The chains are joined together by the N1—H1N \cdots O1 hydrogen bonds to form a sheet structure parallel to the *ab* plane. The Fmoc and *i*-butyl moieties are packed between the sheets (Fig. 2).

Experimental

A powdered sample (I) was obtained from Wako Pure Chemical Industries, Ltd. (Osaka, Japan) and was used for crystallization without further purifications. Colourless needle like crystals of (I) were slowly grown from a saturated dichloromethane solution.

Refinement

All H atoms were found in difference maps and refined with isotropic thermal parameters.

supplementary materials

Figures




Fig. 1. A view of the molecular structure of (I), showing the atom labeling scheme. Displacement ellipsoids are drawn at the 50% probability level.




Fig. 2. A packing diagram of (I) viewed. The hydrogen atoms were omitted for clarity, except for those forming the hydrogen bonds. Broken lines indicate the hydrogen bonds.

fluoren-9-yl N-(1-carboxy-3-methylbutyl)carbamate

Crystal data

C ₂₁ H ₂₃ NO ₄	$F_{000} = 752$
$M_r = 353.40$	$D_x = 1.228 \text{ Mg m}^{-3}$
Orthorhombic, $P2_12_12_1$	Mo $K\alpha$ radiation
Hall symbol: P 2ac 2ab	$\lambda = 0.71073 \text{ \AA}$
$a = 5.49530 (10) \text{ \AA}$	Cell parameters from 40402 reflections
$b = 14.2700 (3) \text{ \AA}$	$\theta = 2.2\text{--}30.0^\circ$
$c = 24.3759 (6) \text{ \AA}$	$\mu = 0.09 \text{ mm}^{-1}$
$V = 1911.51 (7) \text{ \AA}^3$	$T = 150 \text{ K}$
$Z = 4$	Needle, colourless
	$0.40 \times 0.08 \times 0.06 \text{ mm}$

Data collection

Rigaku AFC-8	3207 independent reflections
diffractometer with Saturn70 CCD detector	2906 reflections with $I > 2\sigma(I)$
Radiation source: fine-focus rotating anode	$R_{\text{int}} = 0.055$
Monochromator: confocal	$\theta_{\text{max}} = 30.0^\circ$
Detector resolution: 28.5714 pixels mm^{-1}	$\theta_{\text{min}} = 2.2^\circ$
$T = 150 \text{ K}$	$h = -7 \rightarrow 7$
ω scans	$k = -20 \rightarrow 20$
Absorption correction: none	$l = -34 \rightarrow 34$
40257 measured reflections	

Refinement

Refinement on F^2	Secondary atom site location: difference Fourier map
---------------------	--

Least-squares matrix: full	Hydrogen site location: difference Fourier map
$R[F^2 > 2\sigma(F^2)] = 0.039$	All H-atom parameters refined
$wR(F^2) = 0.111$	$w = 1/[\sigma^2(F_o^2) + (0.0687P)^2 + 0.1717P]$
$S = 1.09$	where $P = (F_o^2 + 2F_c^2)/3$
3207 reflections	$(\Delta/\sigma)_{\max} < 0.001$
327 parameters	$\Delta\rho_{\max} = 0.27 \text{ e } \text{\AA}^{-3}$
Primary atom site location: structure-invariant direct methods	$\Delta\rho_{\min} = -0.25 \text{ e } \text{\AA}^{-3}$
	Extinction correction: none

Special details

Experimental. All Friedel pairs were merged, and all f's of containing atoms were set to zero.

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R-factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
O1	-0.3755 (2)	0.42665 (9)	0.27853 (7)	0.0421 (3)
O2	0.0123 (2)	0.47101 (9)	0.26982 (7)	0.0455 (4)
H2H	-0.047 (5)	0.526 (2)	0.2667 (11)	0.055 (7)*
O3	0.1016 (2)	0.15166 (8)	0.23545 (6)	0.0360 (3)
O4	0.4680 (2)	0.22208 (8)	0.22403 (5)	0.0306 (2)
N1	0.1792 (2)	0.29706 (9)	0.26957 (6)	0.0284 (3)
H1N	0.280 (5)	0.3438 (18)	0.2738 (9)	0.044 (6)*
C1	-0.0651 (3)	0.31083 (10)	0.29096 (7)	0.0287 (3)
H1	-0.171 (4)	0.2659 (15)	0.2718 (9)	0.035 (5)*
C2	-0.0783 (4)	0.29621 (13)	0.35339 (8)	0.0406 (4)
H2A	0.047 (5)	0.3372 (19)	0.3712 (10)	0.048 (7)*
H2B	-0.225 (6)	0.322 (2)	0.3692 (13)	0.065 (8)*
C3	-0.0497 (4)	0.19389 (13)	0.37135 (8)	0.0413 (4)
H3	0.102 (5)	0.1685 (18)	0.3516 (10)	0.048 (7)*
C4	-0.2695 (5)	0.13470 (17)	0.35502 (14)	0.0601 (7)
H4A	-0.312 (6)	0.1369 (19)	0.3131 (11)	0.060 (8)*
H4B	-0.387 (8)	0.163 (3)	0.3755 (15)	0.088 (11)*
H4C	-0.262 (6)	0.065 (2)	0.3677 (13)	0.072 (9)*
C5	0.0014 (9)	0.18903 (19)	0.43291 (10)	0.0705 (9)
H5A	0.163 (8)	0.227 (3)	0.4437 (15)	0.093 (12)*
H5B	0.017 (6)	0.124 (2)	0.4463 (11)	0.060 (8)*

supplementary materials

H5C	-0.127 (7)	0.223 (2)	0.4509 (13)	0.072 (9)*
C6	-0.1606 (3)	0.40869 (10)	0.27831 (7)	0.0299 (3)
C7	0.2373 (3)	0.21843 (10)	0.24241 (6)	0.0261 (3)
C8	0.5484 (3)	0.14752 (11)	0.18786 (6)	0.0289 (3)
H8A	0.476 (5)	0.0932 (16)	0.1976 (9)	0.034 (5)*
H8B	0.721 (5)	0.1412 (15)	0.1960 (9)	0.034 (5)*
C9	0.5153 (3)	0.17588 (11)	0.12778 (7)	0.0291 (3)
H9	0.337 (5)	0.1858 (16)	0.1193 (9)	0.043 (6)*
C10	0.6613 (3)	0.26233 (11)	0.11325 (7)	0.0308 (3)
C11	0.6335 (4)	0.35445 (12)	0.13131 (8)	0.0385 (4)
H11	0.504 (5)	0.3707 (16)	0.1573 (9)	0.035 (5)*
C12	0.8001 (4)	0.42145 (13)	0.11327 (9)	0.0456 (4)
H12	0.783 (5)	0.4852 (19)	0.1253 (10)	0.054 (7)*
C13	0.9889 (4)	0.39782 (15)	0.07795 (8)	0.0459 (5)
H13	1.117 (5)	0.4495 (19)	0.0642 (11)	0.057 (7)*
C14	1.0166 (4)	0.30626 (15)	0.05954 (8)	0.0408 (4)
H14	1.148 (5)	0.2884 (16)	0.0338 (10)	0.048 (6)*
C15	0.8514 (3)	0.23871 (12)	0.07728 (7)	0.0323 (3)
C16	0.8333 (3)	0.13832 (12)	0.06424 (6)	0.0329 (3)
C17	0.9738 (4)	0.08170 (16)	0.03029 (8)	0.0437 (4)
H17	1.114 (5)	0.1107 (17)	0.0111 (10)	0.048 (7)*
C18	0.9091 (5)	-0.01199 (16)	0.02442 (8)	0.0505 (5)
H18	1.003 (6)	-0.0545 (19)	0.0001 (11)	0.060 (7)*
C19	0.7076 (5)	-0.04829 (14)	0.05165 (8)	0.0473 (5)
H19	0.671 (5)	-0.1136 (19)	0.0461 (10)	0.053 (7)*
C20	0.5670 (4)	0.00808 (12)	0.08599 (8)	0.0392 (4)
H20	0.423 (5)	-0.0159 (18)	0.1060 (11)	0.053 (7)*
C21	0.6305 (3)	0.10141 (11)	0.09189 (6)	0.0317 (3)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
O1	0.0236 (6)	0.0258 (5)	0.0769 (9)	0.0009 (4)	0.0034 (6)	-0.0057 (6)
O2	0.0258 (6)	0.0197 (5)	0.0910 (11)	-0.0004 (4)	-0.0005 (7)	0.0087 (6)
O3	0.0236 (5)	0.0220 (5)	0.0623 (7)	-0.0025 (4)	0.0012 (5)	-0.0085 (5)
O4	0.0245 (5)	0.0259 (5)	0.0415 (6)	-0.0027 (4)	0.0051 (5)	-0.0058 (4)
N1	0.0232 (6)	0.0190 (5)	0.0430 (7)	-0.0026 (5)	0.0012 (5)	-0.0035 (5)
C1	0.0246 (7)	0.0194 (6)	0.0423 (8)	-0.0004 (5)	0.0036 (6)	-0.0008 (5)
C2	0.0521 (11)	0.0276 (7)	0.0422 (8)	0.0008 (8)	0.0104 (8)	0.0005 (6)
C3	0.0447 (10)	0.0325 (8)	0.0466 (9)	0.0005 (8)	0.0056 (8)	0.0076 (7)
C4	0.0468 (13)	0.0424 (11)	0.0911 (19)	-0.0093 (10)	0.0005 (13)	0.0225 (12)
C5	0.115 (3)	0.0485 (12)	0.0480 (11)	0.0009 (17)	0.0066 (16)	0.0115 (10)
C6	0.0243 (7)	0.0214 (6)	0.0440 (8)	-0.0003 (5)	0.0012 (6)	-0.0026 (5)
C7	0.0229 (6)	0.0204 (6)	0.0351 (7)	0.0002 (5)	-0.0014 (5)	0.0006 (5)
C8	0.0270 (7)	0.0235 (6)	0.0361 (7)	0.0019 (6)	0.0027 (6)	-0.0013 (5)
C9	0.0249 (7)	0.0254 (7)	0.0370 (7)	-0.0001 (6)	-0.0018 (6)	-0.0007 (5)
C10	0.0282 (8)	0.0287 (7)	0.0353 (7)	-0.0019 (6)	-0.0020 (6)	0.0030 (6)
C11	0.0407 (10)	0.0293 (8)	0.0455 (9)	-0.0012 (7)	0.0029 (8)	0.0011 (6)

C12	0.0529 (12)	0.0301 (8)	0.0537 (10)	-0.0085 (9)	0.0003 (10)	0.0034 (7)
C13	0.0463 (11)	0.0412 (10)	0.0500 (10)	-0.0140 (9)	-0.0004 (9)	0.0089 (8)
C14	0.0346 (9)	0.0490 (10)	0.0389 (8)	-0.0064 (8)	0.0021 (7)	0.0082 (7)
C15	0.0305 (8)	0.0338 (7)	0.0326 (7)	-0.0004 (6)	-0.0026 (6)	0.0037 (6)
C16	0.0325 (8)	0.0364 (8)	0.0296 (6)	0.0047 (7)	-0.0040 (6)	0.0006 (6)
C17	0.0427 (10)	0.0523 (11)	0.0360 (8)	0.0134 (9)	-0.0003 (8)	-0.0033 (7)
C18	0.0625 (14)	0.0489 (11)	0.0401 (9)	0.0206 (11)	-0.0069 (9)	-0.0108 (8)
C19	0.0671 (14)	0.0341 (9)	0.0406 (8)	0.0092 (9)	-0.0122 (10)	-0.0079 (7)
C20	0.0483 (11)	0.0299 (8)	0.0395 (8)	-0.0001 (8)	-0.0082 (8)	-0.0026 (6)
C21	0.0330 (8)	0.0298 (7)	0.0324 (7)	0.0042 (6)	-0.0060 (6)	-0.0021 (6)

Geometric parameters (\AA , $^\circ$)

O1—C6	1.208 (2)	C8—H8B	0.97 (2)
O2—C6	1.318 (2)	C9—C10	1.514 (2)
O2—H2H	0.85 (3)	C9—C21	1.515 (2)
O3—C7	1.2217 (18)	C9—H9	1.01 (3)
O4—C7	1.3460 (19)	C10—C11	1.395 (2)
O4—C8	1.4506 (18)	C10—C15	1.405 (2)
N1—C7	1.3413 (18)	C11—C12	1.395 (3)
N1—C1	1.453 (2)	C11—H11	0.98 (2)
N1—H1N	0.87 (3)	C12—C13	1.390 (3)
C1—C6	1.523 (2)	C12—H12	0.96 (3)
C1—C2	1.538 (2)	C13—C14	1.390 (3)
C1—H1	0.98 (2)	C13—H13	1.07 (3)
C2—C3	1.532 (3)	C14—C15	1.393 (3)
C2—H2A	1.00 (3)	C14—H14	0.99 (3)
C2—H2B	0.96 (3)	C15—C16	1.471 (2)
C3—C4	1.527 (3)	C16—C17	1.391 (3)
C3—C5	1.528 (3)	C16—C21	1.405 (3)
C3—H3	1.03 (3)	C17—C18	1.391 (3)
C4—H4A	1.05 (3)	C17—H17	0.99 (3)
C4—H4B	0.91 (4)	C18—C19	1.391 (4)
C4—H4C	1.04 (3)	C18—H18	0.99 (3)
C5—H5A	1.08 (4)	C19—C20	1.395 (3)
C5—H5B	0.98 (3)	C19—H19	0.96 (3)
C5—H5C	0.96 (4)	C20—C21	1.384 (2)
C8—C9	1.530 (2)	C20—H20	0.99 (3)
C8—H8A	0.90 (2)		
C6—O2—H2H	111.1 (19)	C9—C8—H8B	109.5 (13)
C7—O4—C8	117.44 (12)	H8A—C8—H8B	107 (2)
C7—N1—C1	120.66 (13)	C10—C9—C21	102.41 (13)
C7—N1—H1N	123.1 (16)	C10—C9—C8	112.11 (13)
C1—N1—H1N	116.1 (16)	C21—C9—C8	108.51 (13)
N1—C1—C6	111.69 (13)	C10—C9—H9	110.6 (13)
N1—C1—C2	112.36 (15)	C21—C9—H9	112.6 (13)
C6—C1—C2	107.96 (13)	C8—C9—H9	110.4 (13)
N1—C1—H1	106.7 (13)	C11—C10—C15	120.30 (16)
C6—C1—H1	107.3 (13)	C11—C10—C9	129.51 (16)

supplementary materials

C2—C1—H1	110.7 (13)	C15—C10—C9	110.16 (14)
C3—C2—C1	114.03 (14)	C10—C11—C12	118.34 (19)
C3—C2—H2A	111.3 (15)	C10—C11—H11	120.5 (13)
C1—C2—H2A	108.5 (14)	C12—C11—H11	121.1 (13)
C3—C2—H2B	109.2 (18)	C13—C12—C11	121.27 (19)
C1—C2—H2B	112.6 (18)	C13—C12—H12	119.4 (17)
H2A—C2—H2B	100 (2)	C11—C12—H12	119.3 (17)
C4—C3—C5	112.1 (2)	C14—C13—C12	120.65 (18)
C4—C3—C2	111.80 (19)	C14—C13—H13	118.2 (15)
C5—C3—C2	110.04 (18)	C12—C13—H13	121.1 (14)
C4—C3—H3	108.9 (14)	C13—C14—C15	118.62 (19)
C5—C3—H3	107.1 (14)	C13—C14—H14	121.8 (14)
C2—C3—H3	106.6 (15)	C15—C14—H14	119.6 (14)
C3—C4—H4A	114.4 (17)	C14—C15—C10	120.82 (17)
C3—C4—H4B	100 (2)	C14—C15—C16	130.61 (18)
H4A—C4—H4B	111 (3)	C10—C15—C16	108.56 (15)
C3—C4—H4C	115 (2)	C17—C16—C21	120.53 (18)
H4A—C4—H4C	109 (2)	C17—C16—C15	131.08 (19)
H4B—C4—H4C	107 (3)	C21—C16—C15	108.36 (15)
C3—C5—H5A	112 (2)	C16—C17—C18	118.6 (2)
C3—C5—H5B	112.7 (16)	C16—C17—H17	118.0 (14)
H5A—C5—H5B	109 (3)	C18—C17—H17	123.4 (14)
C3—C5—H5C	107 (2)	C17—C18—C19	120.8 (2)
H5A—C5—H5C	104 (3)	C17—C18—H18	121.1 (16)
H5B—C5—H5C	112 (3)	C19—C18—H18	118.0 (16)
O1—C6—O2	124.21 (15)	C18—C19—C20	120.84 (19)
O1—C6—C1	122.02 (15)	C18—C19—H19	117.2 (17)
O2—C6—C1	113.70 (14)	C20—C19—H19	121.9 (17)
O3—C7—N1	125.15 (15)	C21—C20—C19	118.5 (2)
O3—C7—O4	123.98 (14)	C21—C20—H20	118.9 (15)
N1—C7—O4	110.87 (13)	C19—C20—H20	122.6 (15)
O4—C8—C9	110.58 (13)	C20—C21—C16	120.73 (16)
O4—C8—H8A	109.7 (14)	C20—C21—C9	129.00 (17)
C9—C8—H8A	115.2 (14)	C16—C21—C9	110.22 (14)
O4—C8—H8B	104.0 (13)		
C7—N1—C1—C6	-134.51 (15)	C13—C14—C15—C10	-0.3 (3)
C7—N1—C1—C2	103.99 (16)	C13—C14—C15—C16	178.99 (18)
N1—C1—C2—C3	-70.9 (2)	C11—C10—C15—C14	0.7 (3)
C6—C1—C2—C3	165.48 (17)	C9—C10—C15—C14	-177.72 (16)
C1—C2—C3—C4	-69.5 (3)	C11—C10—C15—C16	-178.72 (16)
C1—C2—C3—C5	165.2 (2)	C9—C10—C15—C16	2.82 (18)
N1—C1—C6—O1	159.29 (17)	C14—C15—C16—C17	-0.5 (3)
C2—C1—C6—O1	-76.7 (2)	C10—C15—C16—C17	178.86 (18)
N1—C1—C6—O2	-23.6 (2)	C14—C15—C16—C21	-178.76 (18)
C2—C1—C6—O2	100.39 (19)	C10—C15—C16—C21	0.63 (18)
C1—N1—C7—O3	-3.6 (2)	C21—C16—C17—C18	-0.1 (3)
C1—N1—C7—O4	177.08 (13)	C15—C16—C17—C18	-178.11 (18)
C8—O4—C7—O3	8.3 (2)	C16—C17—C18—C19	0.3 (3)
C8—O4—C7—N1	-172.31 (13)	C17—C18—C19—C20	-0.7 (3)

C7—O4—C8—C9	93.78 (16)	C18—C19—C20—C21	0.8 (3)
O4—C8—C9—C10	60.54 (17)	C19—C20—C21—C16	-0.5 (3)
O4—C8—C9—C21	172.91 (13)	C19—C20—C21—C9	-177.57 (17)
C21—C9—C10—C11	176.89 (18)	C17—C16—C21—C20	0.2 (2)
C8—C9—C10—C11	-67.0 (2)	C15—C16—C21—C20	178.61 (16)
C21—C9—C10—C15	-4.84 (17)	C17—C16—C21—C9	177.72 (15)
C8—C9—C10—C15	111.28 (16)	C15—C16—C21—C9	-3.83 (18)
C15—C10—C11—C12	-0.7 (3)	C10—C9—C21—C20	-177.46 (17)
C9—C10—C11—C12	177.40 (17)	C8—C9—C21—C20	63.8 (2)
C10—C11—C12—C13	0.3 (3)	C10—C9—C21—C16	5.24 (17)
C11—C12—C13—C14	0.1 (3)	C8—C9—C21—C16	-113.45 (15)
C12—C13—C14—C15	-0.1 (3)		

Hydrogen-bond geometry (Å, °)

<i>D</i> —H··· <i>A</i>	<i>D</i> —H	H··· <i>A</i>	<i>D</i> ··· <i>A</i>	<i>D</i> —H··· <i>A</i>
O2—H2H···O3 ⁱ	0.85 (3)	1.82 (3)	2.6558 (17)	167 (3)
N1—H1N···O1 ⁱⁱ	0.87 (3)	2.24 (3)	3.0751 (18)	161 (2)
C8—H8A···O1 ⁱⁱⁱ	0.90 (2)	2.51 (2)	3.392 (2)	166 (2)

Symmetry codes: (i) $-x, y+1/2, -z+1/2$; (ii) $x+1, y, z$; (iii) $-x, y-1/2, -z+1/2$.

supplementary materials

Fig. 1




Fig. 2

