



Published in final edited form as:

Ther Drug Monit. 1986 ; 8(3): 380–381.

Diurnal Variation in Cyclosporine Kinetics

Raman Venkataraman^{*}, Shuin Yang[†], Gilbert J. Burckart^{*}, Richard J. Ptachcinski^{*}, David H. Van Thiel[‡], and Thomas E. Starzl[†]

^{*} Clinical Pharmacokinetics Laboratory, School of Pharmacy, University of Pittsburgh, Pittsburgh, Pennsylvania, U.S.A

[‡] Department of Gastroenterology, School of Medicine, University of Pittsburgh, Pittsburgh, Pennsylvania, U.S.A

[†] Department of Surgery, School of Medicine, University of Pittsburgh, Pittsburgh, Pennsylvania, U.S.A

Cyclosporine (CyA) is now considered a primary immunosuppressant used to prevent organ rejection in liver, kidney, and heart transplant patients (1–3). It has been recommended that the blood level of CyA be monitored in these patients to minimize the possibility of rejection due to low CyA blood concentrations or potential toxicity due to high CyA concentrations (4,5). This is based on the observation of marked variations in the trough CyA blood levels between and within patients. Such variations in blood levels are the result of pronounced differences in the pharmacokinetic parameters, such as clearance and bioavailability (6–8). We report on one factor, diurnal variation, that may contribute to the observed variability in the pharmacokinetics of CyA in transplant patients.

CyA pharmacokinetics were studied during two or three different dosing intervals (8 h) in two patients who underwent orthotopic liver transplantation at the University of Pittsburgh. The same maintenance dose of CyA (patient 1, 140 mg; patient 2, 150 mg) was administered over 1 h as an intravenous infusion in the morning and at night. Hourly blood samples were drawn and analyzed for cyclosporine following modifications of a high pressure liquid chromatographic assay (9). Kinetic parameters were calculated according to standard techniques (10).

The biochemical profiles on different study periods in a given patient were similar.

Table 1 summarizes the pharmacokinetic parameters obtained. The 8-h trough levels (C_{\min}) obtained following drug administration at night were lower than the trough levels obtained following drug administration during the day. The apparent blood CyA clearance was higher following drug administration at night as compared with the clearance obtained following daytime drug administration in both subjects. Patient 2 was studied on a third occasion (day 5 postoperatively), during the day, and was found to clear the drug in a pattern similar to that observed previously. Other pharmacokinetic parameters, such as the elimination rate constant and volume of distribution, could not be calculated because of the short dosing interval employed in treating these patients.

In both the subjects studied, the clearance of CyA was higher during the night as compared with the clearance during the day. This phenomenon cannot be attributed to any variation in the analytical methodology, as our analytical procedure is highly reproducible. The coefficient

of variation at 600 ng/ml is 3.4% (n = 6). The inpatient variability in CyA kinetics cannot account for the observed differences in CyA kinetics, as demonstrated by the data obtained from Patient 2. During the study period, the patients did not receive any drugs known to induce or inhibit drug-metabolizing enzymes. The presence of circadian rhythms in various biological systems has long been recognized. Recently, diurnal variations in the pharmacokinetics of theophylline (11), valproic acid (12), prednisolone (13), and ethanol (14) have been reported. The actual mechanism(s) responsible for the observed diurnal variation in CyA kinetics is not clear at this point. It may be related either to (a) circadian variations in the activities of one or more hepatic drug-metabolizing enzymes, as CyA is metabolized to a significant extent by the liver in humans (15) or to (b) possible diurnal changes in plasma lipoprotein profiles, as CyA is primarily transported by lipoproteins. Regardless of the mechanism involved in the diurnal variation observed in the kinetics of CyA, it is essential to standardize the time of blood sampling for accurate CyA blood level monitoring.

Acknowledgments

We wish to acknowledge the financial support provided by Sandoz, Inc., for conducting studies in liver transplant patients.

References

1. Starzl TE, Klintmalm GBG, Porter KA, Iwatsuki S, Schroter GPJ. Liver transplantation with use of cyclosporin A and prednisone. *N Engl J Med* 1981;305:266–9. [PubMed: 7017414]
2. Starzl TE, Weil R, Iwatsuki S, et al. The use of cyclosporine A and prednisone in cadaver kidney transplantation. *Surg GynecolObstet* 1980;151:17–26.
3. Griffith BP, Hardesty RL, Thompson ME, Dummer JS, Bahnson HT. Cardiac transplantation with cyclosporine: the Pittsburgh experience. *Heart Transplant* 1983;2:251–6.
4. Burckart GJ, Venkataramanan R, Ptachcinski RJ. Usefulness of cyclosporine monitoring questioned [letter]. *Clin Pharm* 1984;3:243.
5. Keown PA, Stiller CR, Ulan RA, et al. Immunological and pharmacological monitoring in the clinical use of cyclosporin A. *Lancet* 1981;1:686–9. [PubMed: 6110913]
6. Kahan BD, Ried M, Newburger J, et al. Pharmacokinetics of cyclosporine in human renal transplantation. *Transplant Proc* 1983;15:446–53.
7. Burckart G, Starzl T, Williams L, et al. Cyclosporine monitoring and pharmacokinetics in pediatric liver transplant patients. *Transplant Proc* 1985;17:1172–5. [PubMed: 20640180]
8. Ptachcinski RJ, Venkataramanan R, Rosenthal JT, Burckart GJ, Taylor RJ, Hakala TR. Cyclosporine pharmacokinetics in renal transplant recipients. *Clin Pharmacol Ther* 1985;38:296–300. [PubMed: 3896612]
9. Sawchuk RJ, Cartier LL. Liquid chromatographic determination of cyclosporin A in blood and plasma. *Clin Chem* 1981;27:1368–71. [PubMed: 7273396]
10. Gibaldi, M.; Perrier, D. *Pharmacokinetics*. New York: Marcel Dekker; 1983.
11. Decourt S, Fodor F, Flouvat B, Pradalier A, Dry J. Pharmacokinetics of theophylline in night-workers. *Br J Clin Pharm* 1982;13:567–9.
12. Lockard JS, Levy RH, DuCharme LL, Congdon WC, Patel IH. Diurnal variation of valproic acid plasma level and day-night reversal in monkey. *Epilepsia* 1977;18:183–9. [PubMed: 406137]
13. McAllister WAC, Mitchell DM, Collins FV. Prednisolone pharmacokinetics compared between night and day in asthmatic and normal subjects. *Br J Clin Pharm* 1981;11:303–4.
14. Sturtevant FM. Chronopharmacokinetics of ethanol. *Chronobiologia* 1976;3:237. [PubMed: 797559]
15. Wood AJ, Maurer G, Niederberger W, Beveridge T. Cyclosporine: pharmacokinetics, metabolism, and drug interactions. *Transplant Proc* 1983;15:2409–10.

TABLE 1

Pharmacokinetics of cyclosporine in liver transplant patients

Subject	Day post-transplant	Time of study	Dose (mg)	C _{min} (ng/ml)	CyA clearance	
					(ml/min)	(ml/min/kg)
1	21	2 p.m.-10 p.m.	140	519	385	5.13
	18	10 p.m.-6 a.m.	140	301	536	7.14
2	3	10 a.m.-6 p.m.	150	391	337	4.32
	3	8 p.m.-4 a.m.	150	328	479	6.20
	5	9 a.m.-5 p.m.	150	453	369	4.82

CyA, cyclosporine.