



Published in final edited form as:

Neurosci Lett. 2010 October 4; 482(3): 225–229. doi:10.1016/j.neulet.2010.07.044.

In Vitro Interaction of Tubulin with the Photoreceptor cGMP Phosphodiesterase γ -Subunit

Uyen B. Chu[†], Jikui Song[§], Timur A. Mavlyutov[†], and Lian-Wang Guo^{†,*}

[†] Department of Pharmacology, University of Wisconsin School of Medicine and Public Health, Madison, Wisconsin 53706, USA

[§] Center for Eukaryotic Structural Genomics, Department of Biochemistry, University of Wisconsin, Madison, WI 53706, USA

Abstract

The α and β tubulins compose the microtubule cytoskeleton which is involved in many cellular processes such as vesicular transport. The photoreceptor cells in the retina are neurons specialized for phototransduction. Here we report a novel interaction between tubulin and the photoreceptor cGMP phosphodiesterase (PDE6) gamma subunit (PDE γ). The specificity and molecular details of the PDE γ :tubulin interaction were analyzed through the experiments of pull down, microtubule co-sedimentation, and NMR spectroscopy. The tubulin-interacting site was identified to be in the PDE γ C-terminal I67-G85 region, and the interaction interface appeared to be distinct from those with the other PDE γ targets in phototransduction. We also observed that PDE γ interacted with tubulin in a GTP-dependent manner. Our findings offer implications for non-phototransduction role(s) of PDE γ in the photoreceptor neurons.

Keywords

Photoreceptor cells; tubulin/microtubules; PDE6; PDE γ

Microtubules, a major component of the cytoskeleton, participate in numerous cellular processes ranging from cell division, organelle positioning, intracellular transport, to neuronal differentiation [5]. They usually consist of laterally associated protofilaments, each made up of α/β -tubulin dimers that are able to self-assemble in a GTP-dependent way. There are a large number of tubulin/microtubule-interacting proteins that do not necessarily share sequence homology or structural similarity. While many of them regulate microtubule stability/dynamics, some undertake intracellular protein transport such as kinesins and dyneins. Noticeably, many of the microtubule-associated proteins contain intrinsically disordered domains, for example, MAP2 and tau [4], doublecortin and RP1 [12], TPPP/p25 [14], α -synuclein [21], and stathmin [19]. In this study, we have identified another intrinsically disordered tubulin/microtubule-interacting protein, the γ -subunit (PDE γ) of the photoreceptor cGMP phosphodiesterase (PDE6) [8].

*Address correspondence to: Lian-Wang Guo, Department of Pharmacology, University of Wisconsin School of Medicine and Public Health, 1300 University Ave. Madison, WI 53706, Tel.: 608-263-3980, Fax: 608-262-1257, lianwangguo@wisc.edu.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

The rod photoreceptor PDE6 is composed of two similar catalytic subunits (PDE $\alpha\beta$) and two identical inhibitory PDE γ subunits. PDE γ is a small protein of 87 amino acids containing two distinct functional domains: the positively charged central domain (residues G19-G49) and a negatively charged but relatively hydrophobic C-terminal domain (T62-I87) (for review see [8]). The canonical function of PDE γ in phototransduction has been well defined. PDE γ keeps PDE6 inactive in the dark but regulates the turn-on as well as turn-off of visual signaling upon photoresponse, *via* interactions with PDE $\alpha\beta$, the transducin α subunit (Gat) and the regulator of G protein signaling (RGS9-1). Crystal structures revealed distinct PDE γ C-terminal interactions with the chimeric PDE5/6 catalytic domain [2], Gat and the RGS9-1 catalytic core [16]. An NMR study indicated that some structural elements for interacting with these targets were preconfigured in the free PDE γ molecules in solution although PDE γ was overall disordered [18].

Consistent with the feature of intrinsically disordered proteins that they can interact with distinct partners to achieve various functions, increasing evidence implicates non-phototransduction PDE γ targets [8]. A recent study reported that the N-terminal proline rich region of PDE γ interacts with PACSIN, suggesting a possible role of PDE γ in endocytosis in the photoreceptor cell [10]. We were therefore motivated to find additional PDE γ -interacting proteins in the retinal photoreceptors.

Methods

Bovine and mouse retinal homogenates were prepared according to the method described previously [6]. Biotinylated PDE γ (Btn-PDE γ) was generated by covalently linking maleimide-PEO₂-biotin (Pierce) to the single cysteine placed at the PDE γ N-terminus. The peptide Btn-Bz30 containing a PDE γ N-terminal polycationic sequence (BtnG₁₉-Q₃₂) was custom-synthesized at the University of Wisconsin Peptide Synthesis Facility. Btn-CytC (cytochrome C) and Btn-BSA were prepared by first reacting maleimide-PEO₂-biotin with pure proteins and then removing the unreacted biotin molecules with a G-50 spin column (Amersham).

Pull down from retinal homogenates was performed using the streptavidin beads (Pierce) pre-bound with Btn-PDE γ . The beads were first incubated with retinal homogenates for 1 h at 4°C in buffer A (20 mM Tris-HCl, pH 7.4, 150 mM NaCl, 1mM EDTA, 10 mM MgSO₄, 50 μ M AlCl₃, 30 mM NaF, and 50 μ M GDP), and then washed 3 \times with the same buffer. Proteins on the beads were eluted using the SDS/DTT sample buffer and resolved on a 15% SDS gel. In the control, equal amounts of retina homogenates were incubated with the streptavidin beads with no Btn-PDE γ bound.

Western blotting was performed with antibody dilutions as follows: Anti-PDE α (Affinity Bioreagents), 1,000 fold; anti-tubulin- α (Sigma-Aldrich), 10,000 fold; anti-Gat (Affinity Bioreagents), 5,000 fold.

For pulling down microtubules, purified bovine brain tubulin (Cytoskeleton, purity >99%) was used. Tubulin polymerization was performed according to the manufacturer's instruction, and Taxol was added to stabilize microtubules. For each pull down reaction, 5 μ g of microtubules was added to 1 μ l of streptavidin beads with 2 μ g Btn-PDE γ pre-bound. Following incubation of the reactions for 30 min at room temperature in the BST01 buffer (80 mM PIPES, pH 7.0, 1 mM EGTA, 1 mM MgCl₂) supplemented with AlF₄⁻-GDP, the beads were washed 3 \times with 50 μ l of buffer each time.

Microtubule co-sedimentation assays were performed according to the published method [22] with minor modifications. In each reaction, 20 μ g of microtubules was incubated with 2 μ g of PDE γ for 30 min at room temperature in Buffer A. Microtubules were pelleted by

centrifugation at 20,000 $\times g$ for 30 min at room temperature, and then washed twice by repeating centrifugation and removal of the supernatants. The protein pellets were resolved on a SDS gel.

NMR analysis of the tubulin-PDE γ interactions was performed based on the published methods [12,18]. ^{15}N -labeled PDE γ was prepared as previously described [18]. The NMR experiments were performed at 20°C using a 600-MHz Varian Inova spectrometer equipped with ^1H , ^{15}N , and ^{13}C triple-resonance cryogenic probe at the National Magnetic Resonance Facility at Madison, WI (NMRFAM). [^1H , ^{15}N]-HSQC (heteronuclear single quantum correlation) spectra were first collected for 30 μM ^{15}N -labeled free PDE γ in the buffer containing 10 mM PIPES, 1 mM EGTA, 1 mM Mg^{2+} , 90% $\text{H}_2\text{O}/10\% \text{D}_2\text{O}$, pH 6.2. The data collection was repeated under the same experimental conditions following addition of 10 μM tubulin dimer and 1 mM GTP or GDP.

Results

We have observed a novel PDE γ :tubulin interaction. This finding resulted from the effort to identify additional PDE γ targets in the photoreceptor cells through pull down experiments using Btn-PDE γ (Figure 1A, B, C, D, E).

The major pull down from bovine retinal homogenates appeared as a prominent band of ~55 kDa (A, lane 3; C, lane 2). N-terminal micro-sequencing of this band resulted in two sequences: MRECISIH and MREIVHIQ. Blast search indicated that these sequences match the bovine tubulin α and β subunits, respectively. Apparently, this 55 kDa band contained both α and β -tubulin because these two subunits form a constitutive heterodimer [5].

The specificity of the observed tubulin pull down is manifested by the following facts: 1) There was no prominent tubulin pull down in the control, in which no Btn-PDE γ was bound on the beads (A, lane 2; B, lane 4; C, lane 1). 2) Tubulin was not pulled down by Btn-30Bz (lane 3 in C), indicating that tubulin was pulled down by a specific PDE γ sequence other than the N-terminal polycationic G19-Q32 region. In contrast, arrestin in photoreceptor cells is reported to bind microtubules with its positively charged surface [9]. 3) Tubulin was not pulled down by Btn-CytC either (lane 4 in C), which like PDE γ is also a positively charged protein with an isoelectric point at pH 9.25. This result together with the lack of pull down by Btn-30Bz argues against the possibility that tubulin was pulled down merely due to a nonspecific charge effect. 4) While tubulin is abundant in the retina, actin, another abundant cytoskeleton protein (42 kDa), was not efficiently pulled down (Lane 3 in A). 5) PDE $\alpha\beta$ and Gat, however, were both pulled down, as confirmed by Western blotting (Figure 1E). This result served as a positive control because PDE $\alpha\beta$ and Gat are well-documented PDE γ -interacting proteins [8]. It is unlikely that PDE γ pulled down tubulin indirectly by binding to Gat because Gat does not interact with tubulin although Gas, Gai and Gaq do interact [3]. 6) In addition, Btn-PDE γ also readily pulled down tubulin from mouse retinal homogenates (Figure 1D, lane 4), rat brain homogenates and CHO-k1 cell lysates (Figure 1B), further supporting a specific PDE γ -tubulin interaction.

We then further confirmed the specific PDE γ :tubulin interaction using pure proteins. As seen in Figure 2A, in comparison to the background control (lane 4), Btn-PDE γ readily pulled down microtubules which were prepared with the pure bovine brain tubulin (lane 2). Btn-BSA which is irrelevant to PDE γ , however, pulled down microtubules only to the background level (lane 6). Btn-CytC did not pull down microtubules efficiently either (lane 5). This is in agreement with the data obtained with retinal homogenates (Figure 1C, lane 4), further eliminating a possibility of nonspecific charge effect in the observed PDE γ :tubulin interaction. Remarkably, when added as competitors, the C-terminal PDE γ peptide

PDE γ_{62-87} but not the N-terminal peptide PDE γ_{1-61} , significantly reduced microtubule pull down (Figure 2B and lane 3 in A). This is also in agreement with the observation that the PDE γ central peptide Btn-30Bz was not able to pull down tubulin from retinal homogenates (Figure 1C, lane 3), suggesting that the PDE γ C-terminal domain was responsible for the observed PDE γ -tubulin interaction. Thus the PDE γ domain-dependence of this interaction shows an additional layer of specificity.

One useful feature of tubulin is that the polymerized form (microtubules) can be readily pelleted down by centrifugation, and appears as visible glassy pellet at the bottom of the micro-centrifuge tube (Cytoskeleton Inc.). Thus the normally formed microtubule matrix is distinguishable from the opaque appearance of possible protein precipitations. Taking advantage of this property, we have further characterized the specificity of the PDE γ -tubulin interaction by co-sedimentation assays. Consistent with the Btn-PDE γ pull down results (Figure 2A), PDE γ co-sedimented with microtubules (Figure 2C, lanes 2 and 3); CytC however, did not (lanes 4 and 5). PDE γ itself as a highly soluble protein was not pelleted in the absence of microtubules (lane 1).

These pull down and co-sedimentation experiments together indicated that tubulin interacted with PDE γ specifically, probably with its C-terminal domain.

Since tubulin is a GTP/GDP binding protein, we examined a possible GTP dependence of the PDE γ :tubulin interaction by pull down assays and NMR spectroscopy. As shown in Figure 2D, while Btn-PDE γ pulled down tubulin 1.6 fold over the background in the presence of GDP, the pull down was 3.2 fold over the background in the presence of GTP, indicating a GTP dependence of the PDE γ :tubulin interaction.

This feature was also revealed by NMR spectroscopy, a method that can provide molecular details in PDE γ interactions [18]. Identification of residues that exhibit a large change in NMR chemical shift or peak intensity upon complex formation offers an informative approach for mapping the protein-protein interaction interface [12,18]. We have therefore performed the NMR titrations on full-length ^{15}N -labeled PDE γ with pure tubulin (Figure 3). The ^1H , ^{15}N -HSQC spectra were collected for free PDE γ (black), PDE γ in the presence of tubulin and GTP (blue), as well as PDE γ in the presence of tubulin and GDP (red). The NMR peaks were assigned based on the published results [18]. The residues of PDE γ in contact with tubulin were identified by analysis of signal broadening.

Inspection of the NMR spectra indicated that, whereas most of the NMR peaks retained nearly identical intensities upon addition of tubulin in the presence of GTP, some PDE γ residues experienced significant signal loss (Figure 3A). When mapped to the sequence of PDE γ , these residues were located to the PDE γ C-terminal I67-G85 region. This is in accord with the observation that the PDE γ C-terminal peptide PDE γ_{62-87} but not the N-terminal peptide PDE γ_{1-61} eliminated the tubulin pull down (Figure 2B). Residues that displayed most prominent signal broadening included I67, E71, A72, F73, N74, A82, and Q83. Interestingly, these residues are not critically important for either the G α GTPase stimulation [17] or PDE $\alpha\beta$ inhibition [7]. By contrast, residues I86 and I87, which are key for the inhibitory PDE γ interaction with the PDE6 catalytic domain [2, 7], exhibited minimum signal broadening. Meanwhile, the indole group of W70, which is critical for binding of PDE γ with G α [16, 17], was also unaffected by the presence of tubulin. Thus this NMR study suggests that the PDE γ :tubulin interaction is distinguished from the established PDE γ C-terminal interactions with its visual signaling targets [2, 7, 16] (Figure 3D).

Furthermore, the GDP-bound tubulin appeared to impose less line broadening effect than the GTP-bound tubulin (Figure 3A, B,C), confirming a GTP-dependence of the PDE γ :tubulin interaction observed through pull down assays (Figure 2D).

Discussion

In this study we have identified a novel PDE γ interaction with tubulin/microtubules. This interaction was determined to be specific for the PDE γ C-terminus, and was enhanced in the presence of GTP. The tubulin-interacting interface, however, was distinct from the PDE γ interaction with either transducin, the PDE6 catalytic domain, or the RGS9-1 catalytic core. The uniqueness of the PDE γ :tubulin interaction may provide useful clues for possible non-phototransduction PDE γ functions in the photoreceptor cells.

An important feature of the photoreceptor cell is that all the phototransduction proteins are synthesized in the inner segment, and must be transported through the narrow connecting cilium into the outer segment, the default destination, to perform their functions [1]. Microtubules constitute the “railway” for this inter-segmental transport, with the minus ends anchored in the basal body [15] and the plus ends extended either into the inner segment or through the connecting cilium into the outer segment (Figure S1). The mechanism underlying transport inside the inner segment, from the ER (endoplasmic reticulum) to the basal body, is largely unknown. Recent studies using mouse models led to a hypothesis that PDE6 and other membrane proteins transport along microtubules on cargo vesicles, which are moved from the ER region to the basal body most likely by the minus end-directed dynein II [11].

Our results showed that the PDE γ :tubulin interaction is GTP-dependent (Figure 2D and Figure 3) and is differential from the PDE $\alpha\beta$ interaction (Figure 3). We thus speculate that through the interaction with GTP-tubulin, PDE γ may help load the PDE6-carrying vesicles to microtubules and thus facilitate PDE6 transport from ER to the basal body (see the diagram in Figure S1). Since fatty acids are attached to the PDE6 α and β C-termini by a post-translational process on the ER [11], the PDE6 $\alpha\beta\gamma\gamma$ heterotetramer may be assembled and then tethered to the membrane of carrier vesicles in the ER region. The PDE6-laden vesicles can then be “docked” through the PDE γ :tubulin interaction to nearby microtubule plus ends where GTP-bound tubulins are accessible [5]. Thus temporary immobilization of the PDE6 vesicle to microtubules should facilitate its efficient recognition by dyneins, probably also with the assistance of the dynactin complex. In this case, PDE γ could reasonably bind tubulin and PDE $\alpha\beta$ simultaneously, because it is the PDE γ polycationic region rather than the tubulin-interacting C-terminal domain (Figure 2B and Figure 3) that provides most of the binding strength with PDE $\alpha\beta$ [13]. In addition, the key PDE6-inhibiting residues did not interact with tubulin (Figure 3), suggesting that tubulin may not compete with the PDE6 catalytic domain for interaction with the PDE γ C-terminus. Once GTP-tubulin is converted into GDP-tubulin by its GTPase activity [5], PDE γ dissociates from microtubules and the PDE6-carrying vesicles can now be moved along microtubules toward the connecting cilium by the motor proteins.

Microtubule dynamics is a highly regulated event that plays a critical role in neurite extension. The microtubule plus end is stabilized by the loading of GTP-tubulin subunits [5]. Therefore, in a scenario alternative to a possible PDE γ role in PDE6 trafficking, PDE γ which recognizes GTP-tubulin (Figures 2D and 3) may modulate microtubule dynamics and thus regulate the morphogenesis of the photoreceptor synaptic terminal.

In either case, the novel interaction between the PDE γ C-terminus and tubulin that we have observed may provide important implications for understanding the degenerative retinal diseases caused by the lack of PDE γ [20]. Further investigations are warranted to unravel possible cellular functions of the PDE γ :tubulin interaction in the photoreceptor neurons. Distribution of the PDE6 α and β subunits in the retinal outer segment of the PDE γ -knockout mouse should be determined. In the absence of PDE γ , PDE6 may not be able to

transport normally from the inner segment to the outer segment, which may partially explain the abnormal development of the rod outer segment prior to a rapid retinal degeneration in the PDE γ -knockout mouse [20]. In addition, the morphology of the rod photoreceptor synaptic terminal in the PDE γ -knockout mouse should also be examined. A lack of possible regulation of microtubule dynamics by PDE γ could lead to abnormal synapse formation.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

References

1. Baker SA, Haeri M, Yoo P, Gospe SM 3rd, Skiba NP, Knox BE, Arshavsky VY. The outer segment serves as a default destination for the trafficking of membrane proteins in photoreceptors. *J Cell Biol.* 2008; 183:485–98. [PubMed: 18981232]
2. Barren B, Gakhar L, Muradov H, Boyd KK, Ramaswamy S, Artemyev NO. Structural basis of phosphodiesterase 6 inhibition by the C-terminal region of the gamma-subunit. *Embo J.* 2009; 28:3613–22. [PubMed: 19798052]
3. Chen NF, Yu JZ, Skiba NP, Hamm HE, Rasenick MM. A specific domain of Gialpha required for the transactivation of Gialpha by tubulin is implicated in the organization of cellular microtubules. *J Biol Chem.* 2003; 278:15285–90. [PubMed: 12582171]
4. Dehmelt L, Halpain S. The MAP2/Tau family of microtubule-associated proteins. *Genome Biol.* 2005; 6:204. [PubMed: 15642108]
5. Etienne-Manneville S. From signaling pathways to microtubule dynamics: the key players. *Curr Opin Cell Biol.* 2009
6. Ferreira PA. Characterization of RanBP2-associated molecular components in neuroretina. *Methods Enzymol.* 2000; 315:455–68. [PubMed: 10736720]
7. Granovsky AE, Artemyev NO. A conformational switch in the inhibitory gamma-subunit of PDE6 upon enzyme activation by transducin. *Biochemistry.* 2001; 40:13209–15. [PubMed: 11683629]
8. Guo LW, Ruoho AE. The retinal cGMP phosphodiesterase gamma-subunit - a chameleon. *Curr Protein Pept Sci.* 2008; 9:611–25. [PubMed: 19075750]
9. Hanson SM, Francis DJ, Vishnivetskiy SA, Klug CS, Gurevich VV. Visual arrestin binding to microtubules involves a distinct conformational change. *J Biol Chem.* 2006; 281:9765–72. [PubMed: 16461350]
10. Houdart F, Girard-Nau N, Morin F, Voisin P, Vannier B. The regulatory subunit of PDE6 interacts with PACSIN in photoreceptors. *Mol Vis.* 2005; 11:1061–70. [PubMed: 16357825]
11. Karan S, Zhang H, Li S, Frederick JM, Baehr W. A model for transport of membrane-associated phototransduction polypeptides in rod and cone photoreceptor inner segments. *Vision Res.* 2008; 48:442–52. [PubMed: 17949773]
12. Kim MH, Cierpicki T, Derewenda U, Krowarsch D, Feng Y, Devedjiev Y, Dauter Z, Walsh CA, Otlewski J, Bushweller JH, Derewenda ZS. The DCX-domain tandems of doublecortin and doublecortin-like kinase. *Nat Struct Biol.* 2003; 10:324–33. [PubMed: 12692530]
13. Mou H, Cote RH. The catalytic and GAF domains of the rod cGMP phosphodiesterase (PDE6) heterodimer are regulated by distinct regions of its inhibitory gamma subunit. *J Biol Chem.* 2001; 276:27527–34. [PubMed: 11375400]
14. Ovadi J, Orosz F. An unstructured protein with destructive potential: TPPP/p25 in neurodegeneration. *Bioessays.* 2009; 31:676–86. [PubMed: 19382230]
15. Rosenbaum JL, Witman GB. Intraflagellar transport. *Nat Rev Mol Cell Biol.* 2002; 3:813–25. [PubMed: 12415299]
16. Slep KC, Kercher MA, He W, Cowan CW, Wensel TG, Sigler PB. Structural determinants for regulation of phosphodiesterase by a G protein at 2.0 Å. *Nature.* 2001; 409:1071–7. [PubMed: 11234020]

17. Slepak VZ, Artemyev NO, Zhu Y, Dumke CL, Sabacan L, Sondek J, Hamm HE, Bownds MD, Arshavsky VY. An effector site that stimulates G-protein GTPase in photoreceptors. *J Biol Chem.* 1995; 270:14319–24. [PubMed: 7782290]
18. Song J, Guo LW, Muradov H, Artemyev NO, Ruoho AE, Markley JL. Intrinsically disordered gamma-subunit of cGMP phosphodiesterase encodes functionally relevant transient secondary and tertiary structure. *Proc Natl Acad Sci U S A.* 2008; 105:1505–10. [PubMed: 18230733]
19. Steinmetz MO. Structure and thermodynamics of the tubulin-stathmin interaction. *J Struct Biol.* 2007; 158:137–47. [PubMed: 17029844]
20. Tsang SH, Gouras P, Yamashita CK, Kjeldbye H, Fisher J, Farber DB, Goff SP. Retinal degeneration in mice lacking the gamma subunit of the rod cGMP phosphodiesterase. *Science.* 1996; 272:1026–9. [PubMed: 8638127]
21. Wersinger C, Sidhu A. Disruption of the interaction of alpha-synuclein with microtubules enhances cell surface recruitment of the dopamine transporter. *Biochemistry.* 2005; 44:13612–24. [PubMed: 16216085]
22. Xuan C, Qiao W, Gao J, Liu M, Zhang X, Cao Y, Chen Q, Geng Y, Zhou J. Regulation of microtubule assembly and stability by the transactivator of transcription protein of Jemrana disease virus. *J Biol Chem.* 2007; 282:28800–6. [PubMed: 17656359]




Figure 1. Specific pull down of tubulin by PDE γ from bovine retinal homogenates
 Shown are Coomassie-stained SDS gels (A, B, C) and Western blots (D, E), each is a representative of 2–4 similar experiments. In each reaction pull down from 100 μ g retinal homogenates was performed using 2 μ g (or otherwise stated) of Btn-PDE γ immobilized on 1 μ l streptavidin beads.
A. The tubulin pull down by Btn-PDE γ from retinal homogenates appears as a prominent band of ~55 kDa (lane 3). Lane 2 is the control with no Btn-PDE γ . MW, molecular weight.
B. Tubulin was pulled down by Btn-PDE γ from bovine retinal homogenates (lane 1), rat brain homogenates (lane 2), as well as CHO-k1 cell lysates (lane 3). The gel was silver-stained using the Pierce SilverSNAP Stain kit II.
C. Btn-PDE γ on the streptavidin beads pulled down tubulin specifically from retinal homogenates (lane 2), as compared to the beads with no Btn-PDE γ (lane 1), the beads bound with Btn-30Bz (lane 3), and the beads bound with Btn-CytC (lane 4).
D. Btn-PDE γ specifically pulled down tubulin from both

bovine (see lanes 1 and 2) and mouse (see lanes 3 and 4) retinal homogenates, as revealed by immunoblotting. The lower panel shows unbound tubulin in the supernatants. **E.** Immunoblotting of PDE α , tubulin- α (tub α), and transducin- α (Gat) in the pull down from bovine retinal homogenates. For each reaction 6 μ g of Btn-PDE γ and 2 μ l of streptavidin beads were used. Lane 1 is the Btn-PDE γ minus control; lane 2 is pull down on the Btn-PDE γ beads; lane 3 is the homogenates used for each reaction; lane 4 is the unbound protein in the supernatant. Semi-quantitation of tubulin pull down relative to input (100%, lane 3): 6.7%, 34.4%, and 58.2% in lanes 1, 2, and 4, respectively.




Figure 2. Specific PDE γ :tubulin interaction revealed by assays using pure proteins
 Streptavidin beads pull down assays (A, B, D) or microtubule co-sedimentation assay (C) were performed using pure tubulin proteins.
A. Btn-PDE γ pulled down microtubules specifically (lane 2) as compared with various controls including the beads with no Btn-PDE γ bound (lane 4), the beads with Btn-CytC bound (lane 5), and the beads with Btn-BSA bound (lane 6). In lane 3, the PDE γ C-terminal peptide PDE γ 62-87 in a 20 fold molar excess over Btn-PDE γ was added. **B.** Microtubules pull down by Btn-PDE γ on the streptavidin beads (lane 1) was reduced in the presence of PDE γ 62-87 (lane 2) but not PDE γ 1-61 (lane 3) in a 20 fold molar excess. Pull down was quantitated as percent of the Coomassie staining in the absence of PDE γ peptides (lane 1). Error bar is a standard deviation from 3 separated experiments. Student's t-tests indicate that the difference between bars 2 and 1 is insignificant ($p > 0.5$); whereas the difference between bars 3 and 1 is very significant ($p < 0.1$). **C.** Co-sedimented proteins are shown in lanes 2–5; the supernatants of the corresponding conditions are shown in lanes 7–10. Each condition is duplicates except lane 1, which is the control of PDE γ pellet from the reaction with no microtubules. **D.** GTP dependence of tubulin pull down by Btn-PDE γ -beads. Tubulin (9 μ g) was incubated with beads alone (lane 1) or beads prebound with 1 μ g Btn-PDE γ for 1 hr at room temperature in the buffer containing 20 mM HEPES, pH 7.4, 120 mM NaCl, and 5 mM MgCl₂, in the presence of 2 mM GDP (lane 2) or 2 mM GTP (lane 3). Pull down was

quantitated as fold of the Coomassie staining in the control (lane 1). Error bar is a standard deviation of 2 separate experiments. The difference between the GDP condition and the GTP condition is significance ($p= 0.207$).




Figure 3. NMR HSQC spectra indicating the PDE γ :tubulin interaction

The [¹H,¹⁵N]-HSQC spectra were collected on 30 μ M ¹⁵N-labeled PDE γ , in the absence (black) or presence of 10 μ M tubulin dimer in the buffer containing 10 mM PIPES, 1 mM EGTA, 1 mM Mg²⁺, 90% H₂O/10%D₂O, pH 6.2, including either 1 mM GTP (blue) or 1 mM GDP (red).

A. Overlay of the HSQC spectra of free ¹⁵N-PDE γ (black) and ¹⁵N-PDE γ in the presence of tubulin-GTP (blue). The cross peaks corresponding to the indole group of W70 is denoted as W70 ϵ . **B.** Overlay of the HSQC spectra of free ¹⁵N-PDE γ (black) and ¹⁵N-PDE γ in the presence of tubulin-GDP (red). **C.** Overlay of the HSQC spectra of ¹⁵N-PDE γ +tubulin-GTP (blue) and ¹⁵N-PDE γ +tubulin-GDP (red). **D.** A diagram of PDE γ interactions with different partners. The PDE γ residues that showed most significant interactions with tubulin-GTP are highlighted in orange. The polycationic region is shown in blue. I86 and I87 are labeled red.

The G α t-interacting (green) and RGS9-interacting (cyan) residues are shown based on the crystal structure including the C-terminal half of the PDE γ molecule [16].