

Liposomal Amphotericin B for Treatment of Cutaneous Leishmaniasis

Glenn Wortmann,* Michael Zapor, Roseanne Ressler, Susan Fraser, Josh Hartzell, Joseph Pierson,
Amy Weintrob, and Alan Magill

Infectious Diseases Service, Walter Reed Army Medical Center, Washington, District of Columbia; Uniformed Services University of the Health Sciences, Bethesda, Maryland; Infectious Diseases Clinical Research Program, Bethesda, Maryland; Guthrie Army Health Clinic, Fort Drum, New York; Division of Experimental Therapeutics, Walter Reed Army Institute of Research, Silver Spring, Maryland

Abstract. Treatment options for cutaneous leishmaniasis in the United States are problematic because the available products are either investigational, toxic, and/or of questionable effectiveness. A retrospective review of patients receiving liposomal amphotericin B through the Walter Reed Army Medical Center for the treatment of cutaneous leishmaniasis during 2007–2009 was conducted. Twenty patients who acquired disease in five countries and with five different strains of *Leishmania* were treated, of whom 19 received a full course of treatment. Sixteen (84%) of 19 experienced a cure with the initial treatment regimen. Three patients did not fully heal after an initial treatment course, but were cured with additional dosing. Acute infusion-related reactions occurred in 25% and mild renal toxicity occurred in 45% of patients. Although the optimum dosing regimen is undefined and the cost and toxicity may limit widespread use, liposomal amphotericin B is a viable treatment alternative for cutaneous leishmaniasis.

INTRODUCTION

Leishmaniasis refers to a broad spectrum of disease caused by protozoan parasites belonging to the genus *Leishmania*. Cutaneous leishmaniasis (CL) usually occurs in the Old World and is caused by *Leishmania major* and *L. tropica*. In the New World, it is caused by *L. (Viannia) braziliensis*, *L. (V.) panamensis*, and *L. mexicana*. Infected patients typically have a non-healing ulcer on exposed skin but the diversity of clinical manifestations includes all dermatologic syndromes.¹

The treatment of CL in the United States is problematic because there is no U.S. Food and Drug Administration (FDA)-approved drug for this indication. Sodium stibogluconate (Pentostam; Glaxo-Smith-Kline, Brentford, United Kingdom) is the therapy suggested by many experts and is available to clinicians under an Investigational New Drug (IND) protocol from the Centers for Disease Control and Prevention (CDC). For U.S. Department of Defense beneficiaries, Pentostam is available under an IND protocol at the Walter Reed Army Medical Center (WRAMC) in Washington, DC, and more than 500 patients have been treated at WRAMC since 2000. The recommended treatment of moderate-to-severe CL requires daily intravenous infusions (20 doses) and is associated with increased levels of pancreatic enzymes in > 90% of patients, increased levels of liver enzymes in > 50%, and significant arthralgias and myalgias in > 50%.^{2,3} Rarer side effects include development of herpes zoster infection during or shortly after treatment and electrocardiographic changes.^{4,5} A safety release from the manufacturer in 2006 warned of particulate matter in the vials (from an interaction with the stopper), and required that the medication be strained through a filter before administration.⁶

Because Pentostam is not an approved drug in the United States, the administration of Pentostam is only possible under an approved protocol administered under Institutional Review Board oversight and an IND application with the FDA with the attendant documentation for sponsor regulatory oversight. Pentostam can only be given with individual written informed consent, and it is often a time-consuming challenge to explain

why Pentostam is the drug of choice for CL and yet not FDA approved for this indication. Because of the rarity of CL in the United States, few clinicians and nurses are experienced with the administration and side effect profile of Pentostam and they may seek expert consultation. For busy clinicians, the combination of the additional documentation burden, written informed consent, an extensive side effect profile that requires close monitoring and inexperience with the drug present significant limitations to the use of Pentostam.

Amphotericin B deoxycholate has been used as a second-line treatment for mucosal leishmaniasis and CL (especially with pentavalent antimony treatment failures) in the New World since the early 1960s. The systemic and renal toxicity, cost, and difficulty of intravenous administration in leishmaniasis-endemic areas prevented more widespread use. The introduction of lipid-associated amphotericin B products with less renal toxicity has enabled more widespread use.

In 1997, the FDA approved liposomal amphotericin B (AmBisome; Astellus Pharam US Inc., Deerfield, IL) for treatment of visceral leishmaniasis in otherwise immunocompetent adults at the dose of 3 mg/kg/day for 7 doses given on days 1–5, 14, and 21 (total dose = 21 mg/kg).⁷ In a search for a more tolerable therapy for CL, some clinicians have reported success with the use of AmBisome (Table 1).^{8–18} Most of these reports include only one or small numbers of patients, include persons with immunosuppressive conditions, persons who had shown initial treatment failures with pentavalent antimony, or report efficacy against just one species of *Leishmania* or from just one geographic region. We report our experience with the use of AmBisome as drug therapy for the treatment of CL in 20 non-immune, immunocompetent returning travelers.

METHODS

A retrospective chart review was conducted on all patients receiving AmBisome for laboratory-confirmed CL who were either directly managed or managed in consultation with the WRAMC Infectious Disease Service during 2007–2009. Documentation of health care within the Department of Defense is accomplished through a centralized electronic medical record system termed the Armed Forces Health Longitudinal Technology Application system. Patients receiving health care by any Department of Defense health care

*Address correspondence to Glenn Wortmann, Infectious Diseases Clinic, Walter Reed Army Medical Center, 6900 Georgia Avenue, NW, Washington, DC 20307-5001. E-mail: glenn.wortmann@us.army.mil

TABLE 1
Published studies using liposomal amphotericin B (AmBisome) for the treatment of cutaneous leishmaniasis*

Study	Treatment regimen	Total dose of AmBisome (mg/kg)	No. patients evaluated/age/sex	Clinical manifestation	Disease-endemic area resident or returned traveler (non-immune)	Immunocompromised	Prior anti-leishmanial drug treatment	Country in which acquired	Species	Clinical cure	Duration of follow-up (months)
Amato and others ⁸	50 mg every 36 h for a total of 300 mg, followed by daily dose of 25 mg	600 mg (weight NR)	1/50 years/M	Single ulcer at base of finger	Endemic resident	History of renal transplant and receiving dialysis; diabetes	Oral azithromycin	Brazil	NR	1/1	9
Brown and others ⁹	3.125 mg/kg/d or 200 mg/d for 7 days, followed by 200 mg biweekly for 3 weeks	41	1/19 years/M	6-cm ulcer on left knee and 2 nodules on right hand	Non-immune returned traveler	No	No	Belize	<i>L. (Viannia) braziliensis</i>	1/1	12
Gündüz and others ¹⁰	3 mg/kg/d biweekly for 6 weeks	36	1/60 years/F	3-year history of extensive facial plaque (leishmaniasis recidivans)	Endemic area resident	No	Pentavalent antimony and ketoconazole	Turkey	NR (likely <i>L. tropica</i>)	0/1, inadequate clinical response	NR
Mirzabeigi and others ¹¹	5 mg/kg/d for 2 months	300	1/50 years/F	Multiple painful nodules on both lower extremities 2 months after transplant surgery (reactivation)	Resident of Bolivia, immigrated to USA	History of renal transplantation and receiving immunosuppressive drugs	No	Bolivia	NR	1/1	12
Paradisi and others ¹²	3 mg/kg/d on days 1-5, 14, and 21	21	1/56 years/M	Three large leg ulcers	Endemic area resident	Type II diabetes; lesions previously treated with topical steroids	Topical paromomycin and itraconazole	Italy	<i>L. infantum</i>	1/1	9
Rapp and others ¹³	3 mg/kg/d on days 1-5 and 10	18	1	4-cm crusted lesion on left elbow for 3 years	Non-immune returned traveler	No	Pentavalent antimony and ketoconazole	Djibouti	NR	1/1	18
Rongioletti and others ¹⁴	3 mg/kg/d for 5 days	15	1/82 years/M	Lip swelling for 2 years	Endemic area resident	No	Pentavalent antimony	Italy	<i>L. infantum</i>	1/1	12
Rosal and others ¹⁵	5 mg/kg/d for 10 days	50	1/4 months/M and 1/9 years/M	3 cm hyperkeratotic nodule cheek; ulcers on jaw and right hand	Endemic area resident; non-immune returned traveler	No	Intralesional pentavalent antimony plus topical imiquimod and amphotericin B; itraconazole	Spain, Bolivia, or Peru	1 NR, 1 <i>L. (Viannia) braziliensis</i>	2/2	24, NR

Continued

TABLE 1
Continued

Study	Treatment regimen	Total dose of AmBisome (mg/kg)	No. patients evaluated/age/sex	Clinical manifestation	Disease-endemic area resident or returned traveler (non-immune)	Immunocompromised	Prior anti-leishmanial drug treatment	Country in which acquired	Species	Clinical cure	Duration of follow-up (months)
Solomon and others ⁶	3 mg/kg/d on days 1–5 and 10	18	7, 5/M and 2/F age range = 21–24 years	Upper extremity ulcers (4) and facial ulcers (3) with lymph node enlargement	Non-immune returned travelers	No	5 pentavalent antimony treatment failures, 2 initial treatment with AmBisome	Bolivia	<i>L. (Viannia) braziliensis</i>	7/7	3–17
Torre-Cisneros and others ¹⁷	1.5 mg/kg/d for 2 weeks, followed by 4 weekly doses of amphotericin B	31.5 mg AmBisome; 2.5 grams amphotericin B	1/40 years/F	0.5-cm erythematous scaling lesion in right ear	Endemic area resident	No	Pentavalent antimony	Spain	NR	1/1	NR
Perez-Ayala and others ¹⁸	3 mg/kg/d on days 1–5, 14, and 21	21	1/27 years/M and 1/39 years/F	Two ulcers left thigh; one ulcer lower extremity	Immigrant; non-immune returned traveler	HIV+, No	No	Burkina Faso, French Guiana	NR, <i>L. braziliensis</i>	0/2 side effects; inadequate clinical response	NR

* NR = not reported; HIV = human immunodeficiency virus.

provider have their encounter documented in the electronic medical record. The study was reviewed and approved by the WRAMC Institutional Review Board.

Laboratory diagnosis was made by lesion sampling (aspirate or scraping) and microscopic visualization of amastigotes, isolation of promastigotes in culture, or detection of *Leishmania* rRNA by real-time polymerase chain reaction amplification performed in a College of American Pathologists/Clinical Laboratory Improvement Amendments–certified laboratory at the Walter Reed Army Institute of Research (Silver Spring, MD).¹⁹ Identification of *Leishmania* species was performed by using cellulose acetate electrophoresis for isoenzyme analysis and comparison with World Health Organization reference strains.^{20,21}

Treatment with intravenous AmBisome was prescribed at a dose of 3mg/kg/day, with up to 10 doses given within a 21-day period. All patients received premedication with oral acetaminophen (650 mg) and diphenhydramine (50 mg one hour before administration of AmBisome) and intravenous infusion of one liter of physiologic saline. AmBisome was initially infused at a rate of 200 mL/hour. However, because of development of acute infusion-related reactions in two patients, this rate was changed to a dose-escalating infusion (20 mL/hour for 15 minutes, and then 40 mL/hour for 30 minutes, then doubled every 30 minutes until a maximum rate of 200 mL/hour was reached). The infusion duration varied depending on the dose of AmBisome, but lasted approximately two hours.

Efficacy of drug activity was assessed by a review of lesion photographs and/or clinical notes documented in Armed Forces Health Longitudinal Technology Application. Patients who experienced complete healing of their skin lesion(s) after an initial course of AmBisome were scored as cured.

Toxicity of therapy was assessed by a review of the medical record for documentation of adverse events. Nephrotoxicity was assessed by the Common Terminology Criteria (CTC), in which grade 1 indicates values greater than the upper limit of normal (ULN) to 1.5 times the ULN, grade 2 is 1.5–3.0 times the ULN, and grade 3 is > 3.0–6.0 times the ULN.²²

RESULTS

Twenty patients with confirmed CL received treatment with AmBisome. The epidemiologic characteristics and outcome of the cohort are shown in Table 2. Two patients had concurrent nasal mucosa and CL (determined by endoscopic nasal biopsies of intranasal lesions) and AmBisome was suggested as the recommended therapy. Three patients had received prior treatment with Pentostam at a dose of 20 mg/kg/day for their disease. One patient had received 20 doses without cure of his lesion, and two patients had adverse events (hepatitis and pancreatitis), which resulted in cessation of therapy after 9 and 4 doses, respectively. The remaining 15 patients chose treatment with AmBisome after a discussion of the risks/benefits and treatment alternatives.

Ten patients acquired their infection in the Old World (5 in Iraq and 5 in Afghanistan). Isoenzyme analysis of five samples identified *L. major* (3) and *L. tropica* (2). Ten infections were acquired in the New World (1 in Peru, 1 in French Guiana, 2 in Honduras, and 6 in Columbia). Isoenzyme analysis of samples from these 10 patients identified *L. (V.) braziliensis* (3), *L. (V.) guyanensis* (3), and *L. (L.) V. panamensis* (4).

TABLE 2
Cohort description and outcome of patients with cutaneous leishmaniasis treated with amphotericin B (AmBisome)

Characteristic	Result
Mean age, years (range)	29 (19–46)
Sex (% male)	95
Median no. lesions (range)	1 (1–11)
Median maximum lesion size, cm (range)	2 (1–5)
Countries of acquisition (number of cases)	Iraq (5), Afghanistan (5), Peru (1), French Guiana (1), Honduras (2), Columbia (6)
Strains of <i>Leishmania</i> recovered (number)	<i>L. (Viannia) braziliensis</i> (3), <i>L. (V.) guyanensis</i> (3), <i>L. (V.) panamensis</i> (4), <i>L. tropica</i> (2), <i>L. major</i> (3)
Median total dose of AmBisome for initial treatment course, mg and mg/kg (range)	1,748 (530–2,670), 21 (10–30)
Patients receiving a full treatment course	19
Patients with a cure after initial treatment regimen	16/19 (84%)
Median cumulative dose of AmBisome prescribed for 3 patients not healing with an initial treatment course, mg and mg/kg (range)	3,146 (3,000–3,430), 36 (33–42)

The initial treatment regimen for AmBisome was 3 mg/kg/day for a median number of 7 doses (range = 2–10 doses) and a median total dose of 1,748 mg (range = 530–2670 mg). By milligrams/kilogram, the median total treatment dose was 21 mg/kg (range = 6–30 mg/kg). Thirteen patients (65%) had documentation of at least one adverse reaction to AmBisome. Five (25%) patients experienced infusion-related reactions, with development of chest pain, dyspnea, flank pain, flushing, and/or urticaria. For one patient, AmBisome was stopped after two doses because of these side effects, and the remaining patients tolerated further infusions. Some evidence of renal toxicity developed in 9 (45%) patients; 8 had a grade 1 toxicity value and 1 had a grade 2 toxicity value. Creatinine levels were either normalized (2 patients) or were returning to normal (4 patients) at time of discharge from medical care. Three patients did not return for follow-up laboratory testing.

Excluding one patient who received only two doses of AmBisome secondary to an acute infusion-related reaction and who then received treatment with Pentostam, 19 patients were included in the efficacy analysis. Median length of follow-up was 4 months (range = 1–27 months). Sixteen (84%) patients experienced a cure of their skin lesion after receiving an initial treatment course of AmBisome. Unfortunately, lack of standardized follow-up time points (caused by re-deployment of patients to combat zones or delay in responding to the queries of the investigators) prevented determination of a time to cure after administration of AmBisome. An illustrative example of a patient with *L. (V.) braziliensis* who experienced a cure after receiving seven doses of AmBisome is shown in Figure 1.

Three patients (16%) required additional doses of AmBisome for cure of their lesions. The median total dose of AmBisome given during the initial treatment course was 1,750 mg (range = 1,260–2,670 mg) for those patients who were cured with one course of AmBisome and 1,745 mg (range = 1,250–2,002 mg) for those who required a second course. Of those patients who required a second course, one patient with *L. (V.) braziliensis* acquired in Honduras initially experienced healing of his single skin lesion (4 × 2.5 cm) after 5 doses totaling 1,250 mg (doses 6 and 7 were not given because of development of gastroenteritis), but subsequent re-ulceration at the same site developed approximately 6 weeks after his treatment course. The patient was re-treated with a seven-dose regimen of AmBisome (re-treatment dose = 1,750 mg [21 mg/kg], with a cumulative dose of 3,000 mg [36 mg/kg]) with subse-

quent clinical cure. Two patients with *L. (V.) panamensis* acquired in Columbia were treated with seven doses of AmBisome each (total doses = 1,745 mg and 2,002 mg [21 mg/kg]) with transient improvement in their solitary lesions (1 × 1 cm and 2 × 2 cm). Both lesions failed to completely heal, but were successfully cured with seven and four dose re-treatment courses of AmBisome (cumulative doses = 3,430 mg [42 mg/kg] and 3,146 mg [33 mg/kg], respectively) approximately 10 weeks and 7 months, respectively, after the initial treatment course. One patient refused more than four additional doses of AmBisome because of infusion-related reactions, but nonetheless experienced healing of his lesion.

DISCUSSION

Our results demonstrate that AmBisome is an efficacious treatment for CL. Although encompassing a small number of patients, this study represents the largest published cohort of patients treated with AmBisome for CL to date. In addition, our study encompasses greater diversity than prior reports and describes disease acquired in five countries with five different strains of *Leishmania*. Sixteen (84%) of 19 patients were cured with an initial course of AmBisome. The 16% failure rate and the fact that the three failures were successfully treated with a second course of AmBisome suggests that the optimum dosing regimen for treatment of CL with AmBisome is still undefined.

In this study, administration of AmBisome was associated with mild-to-moderate toxicity; thus, this drug should be administered with caution. Acute infusion-related reactions associated with liposomal amphotericin B have been described, and a recent review reported a frequency of 20%, which is similar to the rate observed in this study.²³ Nephrotoxicity is a recognized complication of therapy with AmBisome, and 45% of patients in our study experienced some evidence of renal dysfunction as measured by Common Terminology Criteria. Fortunately, renal dysfunction among our cohort was mild and transient. However, careful monitoring of renal function during therapy is prudent.

A direct comparison between AmBisome and the gold standard treatment of Pentostam was not conducted as part of this study, but previous studies have reported similar cure rates with Pentostam (depending on the patient population and geographic locale). Toxicity was observed with the administration of AmBisome, but Pentostam has its own host of recognized



FIGURE 1. Patient with cutaneous leishmaniasis caused by *Leishmania (Viannia) braziliensis* baseline and three weeks and five months after treatment with seven doses of AmBisome. This figure appears in color at www.ajtmh.org.

side-effects. A prospective, randomized controlled study comparing AmBisome and Pentostam would be required to more clearly establish the strength and quality of the evidence base to support a recommendation to use AmBisome in non-immune returned travelers. However, the low numbers of patients in the United States receiving CL each year would necessitate a long-term, multi-site, and multi-national effort

that would be costly and unlikely to be undertaken. Given our results, one could argue that such a study is not needed because AmBisome, although not the panacea for CL, appears to be efficacious for CL caused by several different infecting strains.

Unfortunately, the expense (pharmacy cost of \$821 per 218-mg dose or approximately \$6,500 for the median dose in our cohort of 1,748 mg) and potential toxicity of AmBisome argue that an improved therapy for CL is still needed, with perhaps an inexpensive, well-tolerated oral regimen as the goal. Because AmBisome is a costly off-label prescription in the United States, some insurance and health care plans may be reluctant to reimburse for the cost of treatment. However, alternatives are limited. Pentostam is provided free of charge to requesting civilian physicians by the Centers for Disease Control and Prevention and to Department of Defense beneficiaries by the U.S. Army, but this drug carries the cost of regulatory oversight and has defined toxicities, and there are increasing reports of therapeutic failures with pentavalent antimonial drugs.^{24,25} In addition, although Pentostam can be administered in the outpatient setting, some providers may not have access to an infusion clinic, necessitating hospitalization for approximately three weeks. This cost and that of laboratory toxicity testing add to the expense of Pentostam. Oral agents (such as miltefosine and azole antifungal medications) have variable efficacy, may not be readily available, or are not FDA approved for a CL indication. Local destructive therapies (such as cryotherapy and the Thermomed[®] device) (Thermosurgery Technologies, Inc., Phoenix, AZ) are, in the authors' opinion, unsuitable for large lesions. Other lipid-associated amphotericin B products have extremely limited data for use in CL in humans, and animal data have demonstrated variable efficacy among the different products.²⁶ There is also a single case report of a treatment failure for CL with amphotericin B lipid complex.²⁷ At this time, there is insufficient information to recommend any other lipid-associated amphotericin B formulation for treatment of CL.

The limitations of our data reflect the demographic characteristic of our adult military population (19 of 20 male patients). Thus, we have data for only one female patient and no children. In addition, the U.S. military population is otherwise healthy, well nourished, and immunocompetent. The efficacy of all drug treatments for immunocompromised patients is likely to be lower.

Faced with a multitude of less-than-optimal choices, we suggest the following strategy for the treatment of CL in otherwise healthy non-immune returning travelers. For patients with small lesions (< 1 cm) caused by strains of *Leishmania* that often self-heal (such as *L. mexicana* and *L. major*), the best therapy may be no therapy because the risks of treatment may outweigh the benefits.¹ Local treatments such as cryotherapy, the Thermomed[®] device, and intra-lesional Pentostam (as practiced at some locations outside the United States) may be appropriate for limited disease. Therapy with Pentostam or AmBisome could be offered to patients with cosmetically concerning lesions (e.g., on the face), or with strains of *Leishmania* that self-heal slowly or that can metastasize (such as *L. tropica* and *L. (V). braziliensis*). Our experience suggests that AmBisome be considered as an option for initial therapy in patients requiring systemic therapy for CL. The optimal regimen (daily dose and schedule) for each infecting strain of

Leishmania has not been determined, but the current FDA-approved regimen for the treatment of visceral leishmaniasis in immunocompetent patients of 3 mg/kg/day on days 1–5, 14, and 21 (21 mg/kg) is a reasonable starting point for most patients. The day 14 and 21 dose may not be logistically feasible for some patients, and administering the total dose of 21 mg/kg over a shorter period can be considered.

Received March 23, 2010. Accepted for publication July 2, 2010.

Acknowledgments: We thank Kelly Hummer, Cyrilla Smalls, and Angela Tyler for their outstanding nursing support.

Disclaimer: The views expressed in this article are those of the authors and do not reflect the official policy of the Department of Army, the Department of Defense, or the US Government.

Authors' addresses: Glenn Wortmann, Michael Zupor, Roseanne Ressler, Susan Fraser, Josh Hartzell, and Amy Weintrob, Infectious Diseases Clinic, Washington, DC, E-mails: glenn.wortmann@us.army.mil, michael.zapor@us.army.mil, rose.ressner@us.army.mil, susan.fraser@us.army.mil, joshua.hartzell@us.army.mil, and amy.weintrob@us.army.mil. Joseph Pierson, Guthrie Army Health Clinic, Fort Drum, NY, E-mail: joe.c.pierson@afghan.swa.army.mil. Alan Magill, Division of Experimental Therapeutics, Walter Reed Army Institute of Research, Silver Spring, MD, E-mail: alan.magill@us.army.mil.

REFERENCES

- Magill A, 2010. *Leishmania* species: visceral (kala-azar), cutaneous, and mucosal leishmaniasis. Mandell GL, Bennett JE, Dolin R, eds. *Principles and Practices of Infectious Diseases*. Seventh edition. Volume 2. Philadelphia, PA: Churchill Livingstone Elsevier, 3463–3480.
- Aronson NE, Wortmann GW, Johnson SC, Jackson JE, Gasser RA, Magill AJ, Endy TP, Coyne PE, Grogl M, Benson PM, Beard JS, Tally JD, Gambel JM, Kreutzer RD, Oster CN, 1998. Safety and efficacy of intravenous sodium stibogluconate in the treatment of leishmaniasis: recent U.S. military experience. *Clin Infect Dis* 27: 1457–1464.
- Wortmann G, Miller RS, Oster C, Jackson J, Aronson N, 2002. A randomized, double-blind study of the efficacy of a 10- or 20-day course of sodium stibogluconate for treatment of cutaneous leishmaniasis in United States military personnel. *Clin Infect Dis* 35: 261–267.
- Wortmann GW, Aronson NE, Byrd JC, Grever MR, Oster CN, 1998. Herpes zoster and lymphopenia associated with sodium stibogluconate therapy for cutaneous leishmaniasis. *Clin Infect Dis* 27: 509–512.
- Lawn SD, Armstrong M, Chilton D, Whitty CJ, 2006. Electrocardiographic and biochemical adverse effects of sodium stibogluconate during treatment of cutaneous and mucosal leishmaniasis among returned travelers. *Trans R Soc Trop Med Hyg* 100: 264–269.
- Drug Alert MHRA, 2006. Available at: <https://www.cas.dh.gov.uk/ViewandAcknowledgment/ViewAlert.aspx?AlertID=100876>. Accessed March 15, 2010.
- Meyerhoff A, 1999. U.S. Food and Drug Administration approval of AmBisome (liposomal amphotericin B) for treatment of visceral leishmaniasis. *Clin Infect Dis* 28: 42–48.
- Amato VS, Rabello A, Rotondo-Silva A, Kono A, Maldonado T, Alves I, Floeter-Winter L, Neto V, Shikanai-Yasuda M, 2004. Successful treatment of cutaneous leishmaniasis with lipid formulations of amphotericin B in two immunocompromised patients. *Acta Trop* 92: 127–132.
- Brown M, Noursadeghi M, Boyle J, Davidson RN, 2005. Successful liposomal amphotericin B treatment of *Leishmania braziliensis* cutaneous leishmaniasis. *Br J Dermatol* 153: 203–205.
- Gündüz K, Afsar S, Ayhan S, Kandilolu A, Turel A, Filiz E, Ok U, 2000. Recidivans cutaneous leishmaniasis unresponsive to liposomal amphotericin B (AmBisome). *J Eur Acad Dermatol Venereol* 14: 11–13.
- Mirzabeigi M, Farooq U, Baraniak S, Dowdy L, Ciancio G, Vincek V, 2006. Reactivation of dormant cutaneous *Leishmania* infection in a kidney transplant patient. *J Cutan Pathol* 33: 701–704.
- Paradisi A, Capizzi R, Zampetti A, Proietti I, De Simone C, Feliciani C, Amerio P, 2005. Atypical multifocal cutaneous leishmaniasis in an immunocompetent patient treated by liposomal amphotericin B. *J Infect* 51: e261–e264.
- Rapp C, Imbert P, Darie H, Simon F, Gros P, Debord T, Roue R, 2003. Liposomal amphotericin B treatment of cutaneous leishmaniasis contracted in Djibouti and resistant to meglumine antimoniate. *Bull Soc Pathol Exot* 96: 209–211.
- Rongioletti F, Cannata GE, Parodi A, 2009. Leishmaniasis due to *L. infantum* presenting as macrocheilitis and responding to liposomal amphotericin B. *Eur J Dermatol* 19: 281–282.
- Del Rosal T, Artigao FB, Miguel MJ, de Lucas R, Del Castillo F, 2010. Successful treatment of childhood cutaneous leishmaniasis with liposomal amphotericin B: report of two cases. *J Trop Pediatr* 56: 122–124.
- Solomon M, Baum S, Barzilay A, Pavlotsky F, Trau H, Schwartz E, 2007. Liposomal amphotericin B in comparison to sodium stibogluconate for cutaneous infection due to *Leishmania braziliensis*. *J Am Acad Dermatol* 56: 612–616.
- Torre-Cisneros J, Prada JL, Villanueva JL, Valverde F, Sanchez-Guijo P, 1994. Successful treatment of antimony-resistant cutaneous leishmaniasis with liposomal amphotericin B. *Clin Infect Dis* 18: 1024–1025.
- Pérez-Ayala A, Norman F, Pérez-Molina JA, Herrero JM, Monge B, López-Vélez R, 2009. Imported leishmaniasis: a heterogeneous group of diseases. *J Travel Med* 16: 395–401.
- Wortmann G, Sweeney C, Houg HS, Aronson N, Stiteler J, Jackson J, Ockenhouse C, 2001. Rapid diagnosis of leishmaniasis by fluorogenic polymerase chain reaction. *Am J Trop Med Hyg* 65: 583–587.
- Kreutzer RD, Christensen HA, 1980. Characterization of *Leishmania* spp. by isozyme electrophoresis. *Am J Trop Med Hyg* 29: 199–208.
- Kreutzer RD, Souraty N, Semko ME, 1987. Biochemical identities and differences among *Leishmania* species and subspecies. *Am J Trop Med Hyg* 36: 22–32.
- National Cancer Institute, 2006. *Common Terminology Criteria for Adverse Events (CTCAE)*. Bethesda, MD. Available at: https://webapps.ctep.nci.nih.gov/webobjs/ctc/webhelp/Common_Terminology_Criteria_for_Adverse_Events_CTCAE_v3.htm. Accessed April 22, 2010.
- Roden MM, Nelson LD, Knudsen TA, Jarosinski PF, Starling JM, Shiflett SE, Calis K, DeChristoforo R, Donowitz GR, Buell D, Walsh TJ, 2003. Triad of acute infusion-related reactions associated with liposomal amphotericin B: analysis of clinical and epidemiological characteristics. *Clin Infect Dis* 36: 1213–1220.
- Llanos-Cuentas A, Tulliano G, Auaujo-Castillo R, Miranda-Verastegui C, Santamaria-Castrellon G, Ramirez L, Lazo M, de Doncker S, Boelaert M, Robays J, Dujarkin J, Arevalo J, Chappuis F, 2008. Clinical and parasite species risk factors for pentavalent antimonial treatment failure in cutaneous leishmaniasis in Peru. *Clin Infect Dis* 46: 223–231.
- Hadighi R, Boucher P, Khamesipour A, Meamar AR, Roy G, Ouellette M, Mohebbi M, 2007. Glucantime-resistant *Leishmania tropica* isolated from Iranian patients with cutaneous leishmaniasis are sensitive to alternative antileishmaniasis drugs. *Parasitol Res* 101: 1319–1322.
- Yardley C, Croft SL, 2000. A comparison of the activities of three amphotericin B lipid formulations against experimental visceral and cutaneous leishmaniasis. *Int J Antimicrob Agents* 13: 243–248.
- Wortmann GW, Fraser SL, Aronson NE, Davis C, Miller RS, Jackson JD, Oster CN, 1998. Failure of amphotericin B lipid complex in the treatment of cutaneous leishmaniasis. *Clin Infect Dis* 26: 1006–1007.