

# Cyclin-dependent kinase 5 activity is required for T cell activation and induction of experimental autoimmune encephalomyelitis

Tej K. Pareek,<sup>1</sup> Eric Lam,<sup>1</sup> Xiaojing Zheng,<sup>2</sup> David Askew,<sup>1</sup> Ashok B. Kulkarni,<sup>3</sup> Mark R. Chance,<sup>2</sup> Alex Y. Huang,<sup>1</sup> Kenneth R. Cooke,<sup>1</sup> and John J. Letterio<sup>1</sup>

<sup>1</sup>Division of Pediatric Hematology/Oncology, Department of Pediatrics, University Hospitals Case Medical Center and The Case Comprehensive Cancer Center, and <sup>2</sup>Case Center for Proteomics and Bioinformatics, Case Western Reserve University, Cleveland, OH 44106

<sup>3</sup>National Institute of Dental and Craniofacial Research, National Institutes of Health, Bethesda, MD 20892

**Cyclin-dependent kinase 5 (Cdk5) is a ubiquitously expressed serine/threonine kinase. However, a requirement for Cdk5 has been demonstrated only in postmitotic neurons where there is abundant expression of its activating partners p35 and/or p39. Although hyperactivation of the Cdk5–p35 complex has been found in a variety of inflammatory neurodegenerative disorders, the potential contribution of nonneuronal Cdk5–p35 activity has not been explored in this context. We describe a previously unknown function of the Cdk5–p35 complex in T cells that is required for induction of experimental autoimmune encephalomyelitis (EAE). T cell receptor (TCR) stimulation leads to a rapid induction of Cdk5–p35 expression that is required for T lymphocyte activation. Chimeric mice lacking Cdk5 gene expression in hematopoietic tissues (Cdk5<sup>-/-c</sup>) are resistant to induction of EAE, and adoptive transfer of either Cdk5<sup>-/-c</sup> or p35<sup>-/-</sup> encephalitogenic lymphocytes fails to transfer disease. Moreover, our data reveal a novel mechanism involving Cdk5-mediated phosphorylation of the actin modulator coronin 1a on threonine 418. Cdk5-deficient lymphocytes lack this posttranslational modification of coronin 1a and exhibit defective TCR-induced actin polarization and reduced migration toward CCL-19. These data define a distinct role for Cdk5 in lymphocyte biology and suggest that inhibition of this kinase may be beneficial in the treatment of T cell-mediated inflammatory disorders.**

## CORRESPONDENCE


John J. Letterio:  
John.Letterio@uhhospitals.org

Abbreviations used: ANOVA, analysis of variance; Cdk5, cyclin-dependent kinase 5; CNS, central nervous system; EAE, experimental autoimmune encephalomyelitis; IS, immune synapse; MOG, myelin oligodendrocyte glycoprotein; MPM-2, mitotic protein monoclonal 2; MS, multiple sclerosis; NF-H, neurofilament H; PTX, pertussis toxin.

Multiple sclerosis (MS) is a major cause of neurological disability in young adults and the most common chronic demyelinating disorder of the central nervous system (CNS; Noseworthy et al., 2000). Our understanding of the cellular and molecular mechanisms mediating MS has been advanced by studies in the murine MS preclinical model experimental autoimmune encephalomyelitis (EAE). Development and progression of this and other autoimmune disorders completely depends on the dynamic nature of immune cells, particularly their ability to migrate and to rapidly form an immune synapse (IS) with antigen-presenting cells. Upon stimulation, lymphocytes undergo marked actin-dependent changes in shape that are required for productive cellular interactions and movement during an immune response. The posttranslational modification of proteins triggered by TCR signaling is a fundamental requirement for successful IS

formation and includes the convergence of several signaling molecules at the plasma membrane. Conformational changes induced by protein phosphorylation affect function by modifying binding motifs essential for recruiting proteins into signaling networks or by placing enzymes within proximity to substrates (Pawson and Scott, 1997). Both tyrosine (Thome and Acuto, 1995) and serine/threonine kinases (Matthews and Cantrell, 2006) have been reported to be key modulators during lymphocyte activation, and several novel small molecules designed to inhibit these kinases are currently under investigation in clinical trials involving patients with inflammatory and autoimmune disorders (Cohen, 2002).

© 2010 Pareek et al. This article is distributed under the terms of an Attribution-Noncommercial-Share Alike-No Mirror Sites license for the first six months after the publication date (see <http://www.rupress.org/terms>). After six months it is available under a Creative Commons License (Attribution-Noncommercial-Share Alike 3.0 Unported license, as described at <http://creativecommons.org/licenses/by-nc-sa/3.0/>).



**Figure 1. Increased Cdk5 activity in the CNS of mice with EAE.** Mice were sacrificed 4 wk after EAE induction. (A) Western blotting shows increased p35 and p25 in lysates of diseased lumbar spinal cord and brain stem. Data represent three independent experiments. (B) Quantitative PCR for Cdk5 and p35 transcripts shows an increase in mononuclear cells harvested from lumbar spinal cord and brain stem of affected mice. Data represent 10 pairs of mice (control) and 6 pairs of mice (EAE) of two independent experiments. (C and D) Cdk5 kinase activity assays were measured as  $\gamma$ -<sup>32</sup>P incorporation in the Cdk5-specific substrate neurofilament H (NF-H) peptide. Tissue lysates prepared from brain stem and lumbar spinal cord show a correlation between Cdk5 expression and kinase activity (C). Data represent three independent experiments. Specificity of Cdk5 kinase activity was confirmed using 10  $\mu$ M roscovitine (C) and Cdk5<sup>-/-</sup> embryo (D; E16.5) brain lysates as controls. All data are the mean  $\pm$  SEM. \*\*,  $P < 0.01$ , Student's *t* test.

Cyclin-dependent kinase 5 (Cdk5), a ubiquitously expressed proline-directed serine/threonine kinase, is mainly active in postmitotic neurons as a result of abundant expression of its obligate activating partners p35 and/or p39 in these cells. Cdk5 has been considered a neuron-specific kinase and narrowly viewed as an essential regulator of neuronal function (Dhavan and Tsai, 2001). This perception has been supported by gene KO studies in which germ line deletion of the genes encoding either Cdk5 or p35 lead to inverted cortical neuronal layering during brain development (Ohshima et al., 1996; Chae et al., 1997). Normal activity of Cdk5 is required for proper neuronal migration, synapse formation, and neuronal survival. However, aberrant or hyperactivation of Cdk5 is associated with severe neurodegenerative disorders including Alzheimer's disease (Shelton and Johnson, 2004).

Recently, Cdk5-p35 has been linked with disease induction in nonneuronal lineages (Rosales and Lee, 2006), with examples which include malignant transformation in cancer (Strock et al., 2006; Lin et al., 2007; Upadhyay et al., 2008), induction of inflammatory pain (Pareek et al., 2006), and other inflammation-mediated disorders (Kitazawa et al., 2005). Neurons and immune cells share functional similarities, such as the ability to migrate and form a functional synapse with neighboring cells. Understanding that a common molecular mechanism may underlie the link between inflammation and distinct conditions such as neurodegeneration (Wyss-Coray and Mucke, 2002) and cancer (Coussens and Werb, 2002), we hypothesized that the enhanced Cdk5 activity observed in these conditions may actually reflect an essential role for Cdk5 in the immune cells that traffic to sites of disease.

Cdk5-p35 activity has been reported in human leukemic cell lines and is thought to play a role in monocytic differentiation (Chen and Studzinski, 2001; Studzinski and Harrison, 2003). However, a role for the Cdk5-p35 complex in the activation and function of normal nontransformed immune

cells has not been established. In this paper, we provide the first demonstration that the Cdk5-p35 complex is essential for T cell activation and for the induction of EAE. We have generated immune chimeric mice (Cdk5<sup>-/-C</sup>) by reconstituting WT mice with hematopoietic progenitors from Cdk5-deficient mice embryos (Cdk5<sup>-/-</sup>) after lethal irradiation. Characterization of the Cdk5<sup>-/-C</sup> and p35<sup>-/-</sup> mice shows that Cdk5 activity is dispensable for immune cell development and lineage differentiation. Using phosphoproteomics, we show that Cdk5 phosphorylates coronin 1a, a leukocyte-specific protein and actin modulator (Ferrari et al., 1999), at the threonine 418 residue within a critical actin binding domain. Furthermore, Cdk5-deficient lymphocytes lack this phosphorylation and are hyporesponsive to mitogenic signals, including TCR ligation, and also fail to migrate after CCL19 chemokine stimulation. Most importantly, we show the relevance of these observations by demonstrating the reduced susceptibility of Cdk5<sup>-/-C</sup> mice to EAE. Encephalitogenic lymphocytes derived from either the Cdk5<sup>-/-C</sup> or p35<sup>-/-</sup> mice fail to transfer disease to normal mice. These results establish Cdk5 as an essential regulator of lymphocyte activation and as a potential therapeutic target for autoimmune disorders, including MS.

## RESULTS

### Increased Cdk5-p35 expression in CNS mononuclear infiltrates in EAE


Deregulated Cdk5 activity triggers progressive neurodegeneration and neurofibrillary tangle formation in mice (Cruz and Tsai, 2004). Tau pathology associated with axonal loss during EAE is linked to up-regulation of Cdk5 activity (Schneider et al., 2004). Furthermore, proteomic analysis of chronic active plaques of MS patients has shown significantly increased expression of Cdk5 (Han et al., 2008). In agreement with these results, we also observed increased expression of p35 protein and of its cleaved product p25 in the lumbar spinal cord and brain stem of mice with EAE (Fig. 1 A). To determine the contribution of inflammatory cells to this process,

we isolated mononuclear cell infiltrates (including T cells, B cells, and macrophages) from these tissues and examined the level of Cdk5 and p35 transcript in these cells. Quantitative RT-PCR demonstrated an increase in both Cdk5 (10-fold) and p35 (fivefold) transcripts (Fig. 1 B). The observed increase in expression of Cdk5 and p35 correlated with a more than twofold elevation in Cdk5 activity in total tissue lysates collected from brain stem and lumbar spinal cord (Fig. 1 C). Specificity of Cdk5 kinase activity in this assay was confirmed using Cdk5 WT and KO embryo (embryonic day [E] 16.5) brain lysates (Fig. 1 D). These results suggest a potentially important role for Cdk5 in the function of immune cells that may be necessary for induction of EAE.

### Induction of Cdk5 activity is a requirement for lymphocyte activation

We next determined the presence and importance of this kinase during lymphocyte activation. Although both Cdk5 and p35 were found in very low abundance in normal T cells, their mRNA and protein expression increased within the first 24 h after CD3/CD28 stimulation and remained elevated until 48 h after stimulation (Fig. 2 A). Although mRNA expression of Cdk5–p35 was consistently high for up to 72 h, the levels of the Cdk5–p35 protein started to decline after 48 h. This late reduction in the levels of Cdk5–p35 protein likely reflects an increase in their degradation, a process which is known to be calcium dependent. We also observed a two- to threefold increase in Cdk5 kinase activity within 24–72 h of CD3/CD28-mediated T cell activation (Fig. 2 B). Similarly, a threefold increase in Cdk5 activity was observed in response to IL-7, a nonredundant cytokine required for T cell survival and development which is also implicated in MS (Lundmark et al., 2007; Fig. 2 C). This effect of IL-7 is enhanced in the presence of CD3/CD28 stimulation. Lastly, protein lysates prepared from total splenocytes (Fig. 2 D) or from CD4<sup>+</sup> T cells (Fig. 2 E) collected from mice with EAE showed a three- to fourfold increase in Cdk5 activity. The restimulation of these lymphocytes *in vitro* with myelin oligodendrocyte glycoprotein (MOG) antigen further increased the activity of Cdk5 when assayed at 72 h. The specificity of Cdk5 kinase activity in these assays was confirmed using Cdk5 WT and KO embryo (E16.5) brain lysates (Fig. 2 F).

To analyze, if the overall increase in Cdk5 activity observed in these experiments is a result of the true increase in specific kinase activity of Cdk5 or merely a result of the increased expression of Cdk5 and p35 protein, we repeated these experiments in the presence or absence of a pharmacologic inhibitor of Cdk5 activity (roscovitine), which is known to inhibit Cdk5 activity (by competitive ATP binding) without altering its protein expression. The total splenocytes or purified T cells were stimulated under different conditions in the presence or absence of roscovitine. T cells stimulated with CD3/CD28, PMA/ionomycin, or allogeneic dendritic cells showed a significant decline in their proliferation when treated with 10  $\mu$ M roscovitine (Fig. S1 A). Results obtained from an MTT assay ruled out the possibility of roscovitine



**Figure 2. Cdk5 activity is up-regulated in lymphocytes after TCR or antigen-specific stimulation.** Total T cells were isolated from lymph node and spleen and then stimulated with CD3/CD28. (A) PCR and Western blot analyses showed increased expression of both Cdk5 and p35 upon stimulation. Data represent three independent experiments of six pairs of mice. (B–F) Cdk5 kinase activity was measured as  $\gamma$ -<sup>32</sup>P incorporation in the Cdk5-specific substrate NF-H peptide. (B) Cdk5 kinase activity in T cells was increased after stimulation with or without CD3/CD28. Data represent two independent experiments of six pairs of mice. (C) Addition of IL-7 alone or with CD3/CD28 further enhanced Cdk5 activity. Data represent eight pairs of mice. (D and E) Total splenocytes (D) or CD4<sup>+</sup> T cells (E) collected after EAE induction and cultured with or without MOG (35–55) showed increased Cdk5 activity. Data are derived from three independent experiments with six to eight pairs of mice. (F) Specificity of Cdk5 kinase activity was confirmed by using Cdk5<sup>-/-</sup> embryo (E16.5) brain lysates as a control. All data are the mean  $\pm$  SEM. \*\*,  $P < 0.01$ , one-way analysis of variance (ANOVA) followed by Student's *t* test.

having any adverse effect on lymphocyte viability (Fig. S1 B), and there was no adverse effect of roscovitine on the expression of Cdk5 and p35 protein levels (not depicted). It should be noted that in each kinase assay the activity of Cdk5 is normalized on a per-milligram protein basis. Although it may also be informative to analyze the specific kinase activity on a per-mole basis, we observed a clear increase in Cdk5 kinase activity upon TCR stimulation.


Although roscovitine is viewed as a selective inhibitor of Cdk5 in the dose range used for these experiments (Mapelli et al., 2005), nonselectivity of this compound cannot be ruled out, as this is the case for many small molecule inhibitors of kinase activity. Therefore, we next tested the impact of Cdk5 and p35 gene deletion on T cell activation. Germ line deletion of the Cdk5 gene is associated with embryonic lethality in mice (Ohshima et al., 1996); thus, we generated Cdk5-null immune chimeric mice (Cdk5<sup>-/-</sup>C; Fig. S2). We observed a 60–70% reduction in the proliferative response (as measured

by  $^3\text{H}$ -thymidine uptake) of  $\text{Cdk5}^{-/-C}$  (Fig. 3 A) or  $\text{p35}^{-/-}$  T cells (collected from  $\text{p35}^{-/-}$  mice; Fig. 3 B) to CD3/CD28 stimulation, suggesting an essential requirement for Cdk5 activity in T cell activation. To determine the potential relevance of this observation in the EAE model, we collected lymphocytes 4 wk after MOG (35–55)/CFA immunization and subsequently cultured them either in the presence or absence of 33  $\mu\text{g}/\text{ml}$  MOG and/or 10  $\mu\text{M}$  roscovitine. These MOG-primed lymphocytes harvested from immunized WT mice exhibited a robust proliferative response when restimulated in vitro with MOG antigen and this effect was completely suppressed by roscovitine treatment (Fig. S1 C). More importantly, encephalitogenic lymphocytes harvested from either the MOG-immunized  $\text{Cdk5}^{-/-C}$  mice (Fig. 3 C) or the MOG-immunized  $\text{p35}^{-/-}$  mice (Fig. 3 D) did not respond when reexposed to MOG in vitro. Interestingly, we observed a gene dose effect for p35 in these experiments. Proliferation after restimulation with MOG in vitro was reduced by 50% in  $\text{p35}^{+/-}$  lymphocytes and by 70% in  $\text{p35}^{-/-}$  lymphocytes relative to  $\text{p35}^{+/+}$  controls. This represents the first demonstration of true haploinsufficiency for p35.

#### Disruption of Cdk5 or p35 gene expression ameliorates clinical signs of EAE

We next analyzed the susceptibility of  $\text{Cdk5}^{-/-C}$  and  $\text{p35}^{-/-}$  mice to EAE. During the first 4 wk after immunization, the survival of  $\text{Cdk5}^{-/-C}$  mice (75%) was significantly greater than that of either the  $\text{Cdk5}^{+/+C}$  mice or C57Bl6 $^{+/+C}$  mice (radiation control) in whom survival was <25% as a result of the severity of disease (Fig. 4 A). There was also a significant delay in the development and a reduction in the clinical severity of EAE in  $\text{Cdk5}^{-/-C}$  mice (Fig. 4 B). Histological examination of brain and spinal cord of  $\text{Cdk5}^{-/-C}$  mice also revealed significantly less inflammation, demyelination, and axonal loss (Fig. 4 C). Immunohistochemical analysis of the immune infiltrates in the CNS demonstrated a significant reduction in the numbers of T cells, macrophages, and neutrophils in  $\text{Cdk5}^{-/-C}$  mice (Fig. S3). In contrast, there was no significant difference in either disease progression or disease severity among genotypes when EAE was induced in  $\text{p35}^{+/+}$ ,  $\text{p35}^{+/-}$ , and  $\text{p35}^{-/-}$  mice (not depicted), despite the objective differences in the proliferative response of lymphocytes reexposed to MOG in vitro (Fig. 3). It is important to note that there are significant defects in neuronal development in  $\text{p35}^{-/-}$  mice, and these mice have a shorter life span (Chae et al., 1997). Thus, interpretation of the results of MOG immunization in the  $\text{p35}^{-/-}$  mice may be complicated either by the ability of p39 to compensate for disruption of p35 or by the underlying CNS disease that enhances susceptibility to EAE.

To obviate the impact of these variables that exist in the  $\text{p35}^{-/-}$  model, as well as the potential influence of the radiation required for establishing the Cdk5 chimeric mice, we next used an adoptive transfer approach, assessing the ability of encephalitogenic lymphocytes from these models to transfer disease to naive animals. Lymphocytes harvested from immunized  $\text{Cdk5}^{-/-C}$  mice completely failed to transfer disease (Fig. 4 D). More importantly, disease progression was significantly delayed




**Figure 3. Disruption of Cdk5 activity abrogates antigen receptor-mediated lymphocyte activation.** Lymphocyte activation and proliferation were measured by  $^3\text{H}$ -thymidine incorporation. (A and B)  $\text{Cdk5}^{+/+C}$  and  $\text{Cdk5}^{-/-C}$  (A) and  $\text{p35}^{+/+}$  and  $\text{p35}^{-/-}$  (B) T cells were isolated and activated with CD3/CD28. (C and D) Lymphocytes were collected from  $\text{Cdk5}^{+/+C}$  and  $\text{Cdk5}^{-/-C}$  (C) and  $\text{p35}^{+/+}$ ,  $\text{p35}^{+/-}$ , and  $\text{p35}^{-/-}$  (D) mice 4 wk after EAE induction and stimulated in vitro with MOG. Each data set is derived from a minimum of four independent experiments with four to six pairs of mice. All data are mean  $\pm$  SEM. \*\*,  $P < 0.01$ ; \*\*\*,  $P < 0.001$ , one-way ANOVA followed by Student's *t* test.

in recipients of  $\text{p35}^{-/-}$  encephalitogenic lymphocytes (Fig. 4 E). This observation is supported by histopathology, which shows reduced inflammatory infiltrates within the spinal cord of recipients of the  $\text{p35}^{-/-}$  encephalitogenic lymphocytes (Fig. 4 F). It is noteworthy that encephalitogenic lymphocytes require restimulation with IL-12 before adoptive transfer (Kroenke et al., 2008). Therefore it may be interesting to explore whether a Th17-favoring culture condition might restore the encephalitogenic potential to the  $\text{Cdk5}^{-/-}$  T cells.

#### Cdk5 directly phosphorylates coronin 1a at threonine 418

To identify potential biological substrates of the Cdk5 kinase in activated immune cells during EAE, we collected lymphocytes from  $\text{Cdk5}^{+/+C}$  and  $\text{Cdk5}^{-/-C}$  mice 4 wk after MOG immunization. Deletion of Cdk5 in these cells was confirmed by performing Cdk5-specific RT-PCR (not depicted) and Western blotting (Fig. 5 A). The mitotic protein monoclonal 2 (MPM-2)-specific antibody, which is known to recognize phosphorylated proline-directed serine/threonine residues in proteins (Yaffe et al., 1997), was used to analyze the impact of Cdk5 deletion on proline-directed serine/threonine phosphorylation in lymphocytes. We observed multiple bands with low abundance on our Western blot in  $\text{Cdk5}^{-/-C}$  lymphocytes compared with  $\text{Cdk5}^{+/+C}$  lymphocytes (Fig. 5 B). Analysis of these bands with liquid chromatography–tandem mass spectrometry and Mascot database search identified a 51-kD band as coronin 1a (a leukocyte-specific protein) with sequence coverage of 63% (Fig. S4 A) and a Mascot search score of 3557. Our liquid chromatography–tandem mass spectrometry findings clearly confirmed phosphorylation of coronin 1A






**Figure 4. Abrogation of Cdk5 activity in lymphocytes ameliorates clinical signs and associates with reduced pathology in EAE.** (A and B)  $Cdk5^{+/+C}$ ,  $Cdk5^{-/-C}$ , and  $C57Bl6^{+/+C}$  mice were immunized with MOG to induce EAE and routinely observed for survival (Kaplan-Meier survival curve; A) and for development of clinical signs of EAE (B). Data represent three independent experiments with a total of 26 pairs of mice. (C) 4 wk after induction of EAE, mice were sacrificed and spinal cord was processed and stained with hematoxylin and eosin to assess inflammation, with Luxol fast blue to analyze demyelination, and with Bielschowsky stain to measure axonal loss (each indicated by arrows). Bars, 200  $\mu$ m. Data represent three independent experiments. (D–F)  $C57Bl6$  WT mice received encephalitogenic lymphocytes from either  $Cdk5^{+/+C}$  or  $Cdk5^{-/-C}$  mice (D) or from  $p35^{+/+}$ ,  $p35^{+/-}$ , and  $p35^{-/-}$  mice (E) and were observed closely for the development of clinical signs of EAE. Data represent two independent experiments with nine pairs of mice. Error bars are mean  $\pm$  SEM. (F) Lumbar spinal cord from mice receiving either  $p35^{+/+}$  or  $p35^{-/-}$  encephalitogenic immune cells was removed and stained with hematoxylin and eosin to evaluate for inflammation (arrows). Bars, 200  $\mu$ m. Data represent two independent experiments.

phosphorylation of threonine 418 was observed in  $Cdk5^{-/-C}$  lymphocytes and was confirmed by selected ion chromatography (Fig. 5 D). However, Western blot analyses for total coronin 1a protein showed identical expression in  $Cdk5^{+/+C}$  and  $Cdk5^{-/-C}$  lymphocytes (Fig. 5 C), demonstrating that this posttranslational modification does not affect the abundance of coronin 1a.

To determine whether coronin 1a is a direct substrate for Cdk5, coronin 1a was immunoprecipitated from lysates of  $Cdk5^{+/+C}$  and  $Cdk5^{-/-C}$  encephalitogenic lymphocytes.

Total immunoprecipitates were then assayed by Western blotting with both Cdk5 and coronin 1a antibodies (Fig. 5 E). The results suggest a direct association of Cdk5 and coronin 1a in lymphocytes. Next, to identify if Cdk5 can directly phosphorylate coronin 1a, immunoprecipitates of coronin 1a obtained from normal lymphocytes were mixed with active Cdk5 complex obtained from  $Cdk5^{+/+}$  embryo brain lysates and then subjected to *in vitro* kinase assays in the

and pinpointed the phosphorylation site at threonine 418. A total of two phosphopeptides containing this phosphorylation site were detected and their sequences are  $^{415}RRATPEPSGTPSSDTVSR$ <sup>432</sup> and  $^{416}RATPEPSGTPSSDTVSR$ <sup>432</sup>. The tandem mass spectrometry spectrum of peptide 416–432 is shown in Fig. S4 B. We further analyzed the status of threonine 418 phosphorylation in both  $Cdk5^{+/+C}$  and  $Cdk5^{-/-C}$  encephalitogenic lymphocytes. A threefold reduction in



**Figure 5. Cdk5 physically interacts with and phosphorylates coronin 1a.** (A–C) Protein lysates were prepared from lymphocytes collected 10 d after EAE induction in Cdk5<sup>+/+</sup>C and Cdk5<sup>-/-</sup>C mice and Western blots were probed with anti-Cdk5 (A), –MPM-2 (B), and –coronin 1a (C) antibodies. Data represents three independent experiments with six to eight pairs of mice. (D) Coronin 1a–threonine 418 phosphorylation of Cdk5<sup>+/+</sup>C and Cdk5<sup>-/-</sup>C encephalitogenic lymphocytes as confirmed by selected ion chromatography. Data represent six pairs of mice of two independent experiments. (E) Coronin 1a protein was immunoprecipitated from Cdk5<sup>+/+</sup>C and Cdk5<sup>-/-</sup>C encephalitogenic lymphocytes and Western blotting was performed using antibodies against Cdk5 (top) and coronin 1a (bottom). Data represent three independent experiments. (F) Coronin 1a immunoprecipitated from normal lymphocytes was used as a substrate for Cdk5 in a kinase assay. The top depicts the autoradiograph of phosphorylated coronin 1a by Cdk5 and the bottom is the Coomassie blue stained gel showing total coronin 1a used in the kinase reaction. Data represent three independent experiments. (G) Coronin 1a peptide harboring amino acid sequence of WT protein from 416–432 (418<sup>T</sup> and 424<sup>T</sup>) or mutant peptide where threonine was replaced with alanine or aspartate. Data represent five independent experiments. (H) Cdk5-mediated coronin 1a WT or mutant peptide phosphorylation was measured through  $\gamma$ -[<sup>32</sup>P] incorporation. Cdk5 used in this kinase assay was collected from mouse embryo brain and Cdk5 KO brain served as control. Data represent three independent experiments. All data are the mean  $\pm$  SEM. \*\*, P < 0.01; \*, P < 0.05, one way ANOVA followed by Student's *t* test.


presence of  $\gamma$ -[<sup>32</sup>P]ATP. Autoradiographs of the kinase assay products clearly show that Cdk5 directly phosphorylates coronin 1a (Fig. 5 F). To confirm the relative importance of threonine residues identified by mass spectrometry analysis, we synthesized 17-mer coronin 1a peptides (416–432) and replaced threonine 418 and/or threonine 424 with alanine (Fig. 5 G). These peptides were then used in an *in vitro* kinase assay as substrates for Cdk5. All three mutations were associated with a significant reduction in Cdk5-mediated phosphorylation of the coronin 1a peptide compared with WT, with threonine 418 being most important (Fig. 5 G).

**Disruption of Cdk5 activity impairs actin polarization and migration of lymphocytes to specific chemokine signals**

Genetic deletion of coronin 1a in mice established the absolute requirement of this protein in T cell antigen receptor function (Föger et al., 2006). Coronin 1a has been shown to bind directly to F-actin *in vitro* and to colocalize with F-actin structures *in vivo* (de Hostos et al., 1991). To determine the impact of Cdk5-mediated coronin 1a phosphorylation on F-actin and coronin 1a association, we collected lymphocytes from C57BL6<sup>+/+</sup>C, Cdk5<sup>+/+</sup>C, and Cdk5<sup>-/-</sup>C mice after EAE induction. As expected, both normal (Fig. 6 A) and control (Fig. 6 B) lymphocytes showed homogeneous colocalization of coronin 1a and F-actin with slight polarization in control cells but no discernible difference among lymphocytes isolated

from the C57BL6<sup>+/+</sup>C, Cdk5<sup>+/+</sup>C, and Cdk5<sup>-/-</sup>C mice. Interestingly, Cdk5<sup>+/+</sup>C encephalitogenic lymphocytes exhibited a distinct polarization of F-actin and coronin 1a, with most of the coronin 1a clustered on the pole directly opposite of F-actin clustering, and this phenomenon was completely lost in Cdk5<sup>-/-</sup>C encephalitogenic lymphocytes (Fig. 6, C and D). In contrast, Arp2/3 polarization was not compromised in Cdk5<sup>-/-</sup>C encephalitogenic lymphocytes (Fig. S5). These results support an essential role for Cdk5-mediated coronin 1a phosphorylation in actin dynamics, revealing an important mechanism through which Cdk5 may control both T cell activation and migration.

Several chemokines and their receptors have been shown to play a major role in the recruitment of lymphocytes to the CNS during EAE (Proudfoot et al., 2008). Moreover coronin 1a-deficient lymphocytes fail to migrate toward CCL19 (Föger et al., 2006). Therefore, we next analyzed the impact of either pharmacologic inhibition or genetic deletion of Cdk5 on lymphocyte migration toward the chemokine CCL19. In transmigration assays, there was a 55% reduction in the migration of Jurkat cells toward CCL19 (MIP3- $\beta$ ) and ~10-fold decrease in migration toward of SDF1 $\alpha$  and SDF1 $\beta$  (CXCL12) in the presence of 10  $\mu$ M roscovitine (Fig. S6).



**Figure 6. Disruption of Cdk5 activity diminishes actin polarization and migration toward CCL19.** (A–C) Cdk5<sup>+/+C</sup>, Cdk5<sup>-/-C</sup>, and C57BL6<sup>+/+C</sup> mice were immunized with PBS (normal; A), CFA + pertussis toxin (PTX; control; B), or MOG(35–55) + CFA + PTX (EAE; C) and, 4 wk later, lymphocytes were collected and stained for F-actin using phalloidin (red) and coronin 1a (green) antibody. The white arrows indicate polarization of F actin and the yellow arrows indicate polarization of coronin 1a on the opposite end. (D) These polarized cells were counted from six different regions by a person blinded to sample identity. The data represent two independent experiments of six pairs of mice. (E) The number of cells harvested from the lower chamber containing 100 ng CCL19 in transmigration assays, where the upper chamber contains primary lymphocytes collected from either Cdk5<sup>+/+C</sup> or Cdk5<sup>-/-C</sup> mice. Bars, 100  $\mu$ m. Data represent three independent experiments of five pairs of mice, mean  $\pm$  SEM. \*\*\*,  $P < 0.001$ , one way ANOVA followed by Student's  $t$  test.



Finally, migration of  $Cdk5^{-/-C}$  lymphocytes toward CCL19 was significantly reduced when compared with migration of  $Cdk5^{+/+C}$  control lymphocytes (Fig. 6 E). Collectively, these results suggest that Cdk5-mediated coronin 1a phosphorylation is required for proper lymphocyte activation and migration.

## DISCUSSION

More than a century after the first description of the clinical and pathological characteristics of MS (Pearce, 2005), we have gained immense knowledge regarding the potential molecular and cellular mechanisms mediating disease pathogenesis. Aberrant activation of immune cells is a major hallmark of this autoimmune disease. Thus, therapeutic efforts in MS patients have been largely directed toward suppression of the activated immune system. The initiation of the immune cell response requires dynamic processing of the actin cytoskeleton and involves the recruitment of different proteins within the cell to form the IS, enabling the cell to migrate toward a specific chemokine signal. Several protein kinases have been implicated in this process, but how these proteins collaborate to enable lymphocyte activation and promote the genesis of an autoimmune response has not been fully elucidated.

In the current study, we explore the role of Cdk5 in immune cells. Our current knowledge suggests that kinase activity of ubiquitously expressed Cdk5 is mainly restricted to postmitotic neurons as a result of the predominant expression of its activating partner proteins p35 and p39 in these cells. However, the abundance of Cdk5 expression in nonneuronal cells argues against a functional irrelevance of Cdk5 in these lineages and, rather, points to a potentially important role for this kinase in their normal cell physiology. It is likely that Cdk5 is recruited or activated by cells in response to external stimuli that either influence their state of activation or trigger a differentiated function such as the secretion of cytokines or cell migration. In this context, the absence of Cdk5 kinase activity in nonneuronal cells may not impair normal development or distribution of cells but, rather, affect cellular events that are required for host responses to harmful stimuli or to an external challenge.

The relevance of Cdk5 kinase activity in immune cells has not been extensively studied. However, there are several common signaling partners among neurons and immune cells that are participants in the control of gene transcription and cytoskeletal architecture, and the Cdk5-mediated phosphorylation of these proteins is known to modulate their function in neurons. The dependency of these proteins on Cdk5 function in immune cell signaling has not been explored. For example, Cdk5-mediated phosphorylation induces STAT3 transcriptional activity (Fu et al., 2004) and suppresses MEF2-mediated transcription (Gong et al., 2003) in neurons, and both of these transcription factors are critical regulators of T cell (Lu et al., 2008) and B cell (Wilker et al., 2008) activation. Moreover, Cdk5 is known to modulate actin dynamics in neurons through phosphorylation of proteins involved in maintaining cytoskeletal architecture and promoting neuronal migration such as the ERM protein ezrin (Yang and Hinds,

2003), WAVE1 (Kim et al., 2006), WAVE2 (Miyamoto et al., 2008), FAK (Xie et al., 2003), and also Rac and Pak (Nikolic et al., 1998; Rashid et al., 2001). These proteins are similarly known to regulate reorganization of the actin cytoskeleton in immune cells during TCR signaling, lymphocyte differentiation, and migration and also to orchestrate effector function (Burkhardt et al., 2008). Posttranslational modification of these proteins by their phosphorylation is known to regulate cytoskeletal dynamics in immune cells (Kurosaki and Hikida, 2009; Prince et al., 2009; Salmond et al., 2009). However, the functional relevance of Cdk5 in modulating cytoskeletal proteins during lymphocyte activation has not been explored.

The results presented herein suggest that Cdk5 is a non-redundant kinase in immune cells whose activity is required for lymphocyte activation. When total lymphocytes or purified T cells are activated *in vitro* by antigenic stimulation or by TCR ligation, they demonstrate a significant increase in Cdk5 activity. It is important to note that the up-regulation of Cdk5 activity observed in lymphocytes of MOG/CFA-immunized mice is not seen after immunization with CFA alone. The most likely explanation for this difference is that CFA alone induces only a transient activation of lymphocytes that wanes within 3 wk, unlike the sustained adaptive lymphocyte response to MOG. Regardless, it may be informative to see whether there is a short-term induction of Cdk5 activity (either local or systemic) after exposure to CFA alone and, if so, to explore whether this plays a role in the establishment of disease in this model.

Furthermore, pharmacologic suppression of Cdk5 activity or targeted disruption of Cdk5–p35 gene expression impairs the lymphocyte response to TCR ligation or antigenic stimulation. Hematopoietic organs of  $Cdk5^{-/-}$  embryos develop normally, and there are no obvious phenotypic differences in either the number or ratio of immune cell populations in either  $Cdk5^{-/-}$  or  $p35^{-/-}$  mice compared with their WT littermates (Fig. S2, D and F–H). Similarly, immunophenotyping of the  $Cdk5^{+/+C}$  and  $Cdk5^{-/-C}$  mice shows no differences in gross anatomical structure, cellularity, or lineage distribution in hematopoietic organs (bone marrow, spleen, thymus, and lymph nodes; Fig. S2 E). However, when these mice are challenged through induction of EAE, the spleen and lymph nodes of  $Cdk5^{-/-C}$  mice are significantly smaller with the substantial reduction in tissue weight, cellularity, and disorganized tissue architecture compared with the  $Cdk5^{+/+C}$  control mice (unpublished data). Thus, although  $Cdk5^{-/-C}$  naive lymphocytes develop normally, their coordinated response to antigenic stimuli is hampered as demonstrated by the delayed development and reduced severity of EAE.

Interestingly,  $Cdk5^{-/-C}$  encephalitogenic splenocytes, collected either during the active phase (10 d after EAE induction) or chronic phase (21 d after EAE induction) of EAE, show significant reduction in the production of IL-2, IL-6, IL-17, TNF, and GM-CSF, when compared with  $Cdk5^{+/+C}$  cells, after restimulation with MOG antigen *in vitro*. We did observe a significant drop in IFN- $\gamma$  cytokine production from these cells during active phase but the difference was



not significant in chronic phase (Fig. S7). The observed difference in cytokine production may reflect a reduction in the number of MOG-specific T cells, impaired TCR stimulation, or a skewing in the differentiation of Cdk5<sup>-/-C</sup> lymphocytes. Regardless, the observed reduction in Th1 and Th17 cytokines production provides a plausible explanation for the reduced severity of disease in these mice.

The phosphoproteomic studies described herein show that Cdk5-dependent modulation of immune cell function is partially mediated through the posttranslational modification of coronin 1a. Coronin is a conserved actin binding protein that promotes different cellular processes that rely on rapid remodeling of the actin cytoskeleton (Clemen et al., 2008). Deletion of the coronin 1a gene in mice is associated with alterations in cellular steady-state F-actin formation in lymphocytes (Föger et al., 2006) and reductions in lymphocyte survival, migration, and Ca<sup>2+</sup> release from intracellular stores (Mueller et al., 2008). Coronin 1a gene deletions are associated with severe combined immunodeficiency in humans (Shiow et al., 2008, 2009), and nonsense mutation of the coronin 1a gene (*Lmb3* locus) protects against the induction of systemic lupus in mice (Haraldsson et al., 2008). The crystal structures of a C-terminal truncated form of coronin 1a (residues 1–402), along with a C-terminal fragment (residues 430–461), have been solved (Appleton et al., 2006); however, the structure information with respect to residues 403–429, which contains the Cdk5-targeted threonine 418 phosphorylation site, has not been resolved. It is noteworthy that this domain contains a linker region (residues 356–429) of this protein, which is considered essential for interaction of the coronin 1a homotrimer complex with cytoskeleton (Gatfield et al., 2005). In this paper, we report for the first time that Cdk5 directly phosphorylates coronin 1a on threonine 418. It is quite possible that phosphorylation of threonine 418 is required for the association of a positively charged stretch of linker region with F-actin. Interestingly, Pho85, a functional homologue of Cdk5 in budding yeast (Huang et al., 1999), has been shown to phosphorylate yeast coronin 1 on multiple sites (Gandhi and Goode, 2008). In the current study, we demonstrate that coronin 1a is directly associated with and is phosphorylated by Cdk5 and that this phosphorylation is required for F-actin nucleation but not for Arp2/3 polarization in activated lymphocytes during EAE.

Finally, specific chemokine receptors have been implicated in MS and their roles have been demonstrated in the EAE model (Karpus and Ransohoff, 1998). In addition to regulating lymphocyte homing to secondary lymphoid tissue, the chemokine CCL19 participates in the pathophysiology of EAE by controlling T cell migration into the CNS (Alt et al., 2002). Similar to coronin 1a KO lymphocytes (Föger et al., 2006), either the pharmacologic suppression or genetic deletion of Cdk5 in lymphocytes impairs migration toward CCL19. Whether these results are a direct and specific consequence of altered Cdk5-mediated phosphorylation of coronin 1a remains to be proven, and thus future studies will explore the functional consequences of Cdk5-mediated phosphorylation of coronin 1a on threonine 418.

In summary, Cdk5 now joins a growing list of proteins that were first discovered in the nervous system, such as Dscam and semaphorins, that have since been found to have an important function in immune cells (Boulanger, 2009). This study provides novel insight into the regulation of lymphocyte activation and function through Cdk5-mediated coronin 1a phosphorylation. This discovery of an important extraneuronal role for Cdk5 as a biochemical intermediate in T cell signaling expands the repertoire of nonneuronal activities for a molecule that was previously assigned to a predominantly lineage-restricted function in the CNS. Our results have implications for several neurodegenerative and mental health disorders, wherein immune-mediated activation of Cdk5 may contribute to the predisposition and pathogenesis of diseases including MS. These data establish a new paradigm that links activation of Cdk5 in immune cells to the pathogenesis of disorders associated with inflammation and provide a strong rationale for the development and clinical evaluation of novel inhibitors of Cdk5 in the context of immune-mediated diseases such as MS.

## MATERIALS AND METHODS

**Animals.** 6–8-wk-old C57BL/6 or B6D21 mice were purchased from The Jackson Laboratory and used for collection of lymphocytes or allogeneic APC collection and EAE induction. B6-Ly5.2Cr mice (referred to as CD45.1<sup>+</sup>) were purchased from the National Cancer Institute (Charles River Laboratories) and used for generation of chimeric mice. Cdk5<sup>+/-</sup> mice were used to generate Cdk5<sup>+/+</sup> and Cdk5<sup>-/-</sup> embryos, which were further used for collection of brain or hematopoietic cells. 3-mo-old p35<sup>+/+</sup>, p35<sup>+/-</sup>, or p35<sup>-/-</sup> mice were used for lymphocyte collection and for induction of EAE. All animals were housed in microisolator cages and maintained in climate- and light-controlled rooms (22 ± 0.5°C, 12/12-h dark/light cycle) with free access to food and water. Studies were performed in compliance with procedures approved by the Case Western Reserve University School of Medicine's Institutional Animal Care and Use Committee.

**Generation of Cdk5-null immune chimeric mice.** Hematopoietic cells were collected from liver and spleen of E16.5 Cdk5<sup>+/+</sup> and Cdk5<sup>-/-</sup> littermate embryos, all offspring of male and female Cdk5<sup>+/-</sup> breeder mice. Approximately 10 million cells per mouse were then injected into 12-wk-old lethally irradiated (1,400 rad; Gammacel 137Cs; J.L. Shepard & Associates) CD45.1 female mice by tail vein injection. 8 wk later, peripheral blood was collected from all recipients and tested for CD45.1 versus CD45.2 expression by flow cytometry and PCR analysis for Cdk5 using the following primers: 5'-TGAGGGTGTGCCAAGTTCAGC-3' and 5'-GGCATTGAGTTGG-GCAGCAGAC-3'. These Cdk5 WT and Cdk5-null immune chimeric mice were further recognized as Cdk5<sup>+/+C</sup> and Cdk5<sup>-/-C</sup>, respectively. C57BL/6 chimeras were generated by injecting lethally irradiated CD45.1 mice with 8 × 10<sup>6</sup> bone marrow cells isolated from 10–12-wk-old C57BL/6 female mice and recognized as C57BL6<sup>+/+C</sup>. Upon establishment of chimeric mice, these lines were then maintained by transferring 8 × 10<sup>6</sup> bone marrow cells from each subsequent generation into lethally irradiated CD45.1 female mice. Cdk5 chimeric mice were maintained for a maximum of three generations.

**EAE induction.** EAE was induced in mice by MOG (35–55) immunization and disease was adoptively transferred to unaffected WT animals as previously described (Mi et al., 2007). In brief, Cdk5<sup>+/+C</sup>, Cdk5<sup>-/-C</sup>, and C57BL6<sup>+/+C</sup> or p35<sup>+/+</sup>, p35<sup>+/-</sup>, and p35<sup>-/-</sup> mice were immunized with subcutaneous injections on both flanks with 200 µl of emulsion containing either CFA consisting of 100 µl of incomplete Freund's adjuvant with 8 mg/ml *Mycobacterium tuberculosis* and 100 µl PBS (DIFCO Laboratories) or CFA with 200 µg MOG (35–55). At the time of injection and 48 h later, 200 ng PTX (List Biological Laboratories) was injected intraperitoneally in 100 µl PBS and

mice were monitored daily for clinical signs of EAE. For adoptive transfer of EAE, *Cdk5<sup>+/+C</sup>* and *Cdk5<sup>-/-C</sup>* or *p35<sup>+/+</sup>*, *p35<sup>+/-</sup>*, and *p35<sup>-/-</sup>* mice were euthanized 10 d after MOG (35–55) immunization and cell suspensions were prepared from spleen and regional lymph nodes (axillary, branchial, and inguinal). About  $5 \times 10^6$  cells were cultured in the presence of 33  $\mu\text{g}/\text{ml}$  MOG (35–55) and 20 ng/ml of mouse IL-12. 3 d later  $\sim 2\text{--}3 \times 10^7$  of the cultured encephalitogenic cells were then injected into 3-mo-old female C57BL6 mice via intraperitoneal injection. 200 ng PTX was injected on the same day and again 48 h later. The following criteria were used to grade clinical scores: 0, no signs of disease; 1, limp tail; 2, moderate hind limb weakness; 3, severe hind limb weakness; 4, complete hind limb paralysis; 5, quadriplegia or premoribund state; 6, death.

**Peptide synthesis.** The following peptides were used in this study: MOG (35–55) peptide, MEVGWYRSPFSRVVHLYRNGK; NF-H peptide, VK-SPAKEKAKSPVK; coronin 1a WT, RATPEPSGTPSSDTVSR; coronin 1a mutant 1, RAAPEPSGTPSSDTVSR; coronin 1a mutant 2, RATPEPSGAPSSDTVSR; and coronin 1a mutant 3, RAAPEPSGAPSSDTVSR. All peptides were synthesized at 21st Century Biochemical using Fmoc chemistry and were HPLC purified to a minimum >90%, and the mass and sequence were verified by nanospray mass spectrometry and collision-induced dissociation tandem mass spectrometry.

**Antibodies.** The anti-Cdk5 antibody (C-8), anti-p35 antibody (C-19; Santa Cruz Biotechnology, Inc.), and MPM-2 antibody (Millipore) were each used at 1:200 dilution for Western blots. 5  $\mu\text{g}$  of the anti-Cdk5 antibody (C-8) was used for immunoprecipitation. Hamster monoclonal anti-coronin 1a antibody was provided by A.C. Chen (Genentech, San Francisco, CA) and used at a 1:1,000 dilution for Western blotting and a 1:200 dilution for immunoprecipitation. Rabbit polyclonal anti-coronin 1a serum was provided by J. Pieters (University of Basel, Basel, Switzerland) and used for Western blotting at a 1:5,000 dilution and for immunohistochemistry at a 1:500 dilution. Alexa Fluor 568 phalloidin (Invitrogen) was used to detect F-actin at a 1:25 dilution. Horseradish peroxidase-conjugated anti-hamster (BD) was used as a secondary antibody at 1:5,000 and horseradish peroxidase-conjugated anti-mouse and anti-rabbit antibodies (Jackson ImmunoResearch Laboratories, Inc.) were used at a 1:10,000 dilution for Western blots. The secondary antibodies anti-rabbit Alexa Fluor 488, anti-mouse Alexa Fluor 546, and anti-mouse Alexa Fluor 633 (Invitrogen) were used at a 1:200 dilution for immunofluorescence.

**Roscovitine.** All roscovitine ( $\text{C}_{19}\text{H}_{26}\text{N}_6\text{O}$ ; 6-Benzylamino-2-(R)-((1-ethyl)-2-hydroxyethylamino)-9-isopropylpurine; 2(R)-(1-Ethyl-2-hydroxyethylamino)-6-benzylamino-9-isopropylpurine) used in this study was purchased from Enzo Life Sciences, Inc. A 10-mM stock solution was prepared in DMSO and used at 10  $\mu\text{M}$  for in vitro kinase assays, cell proliferation, and cell migration assays and in the range of 5 to 80  $\mu\text{M}$  for cell viability assays.

**Western blot analyses.** Tissue or cell lysates were prepared in RIPA buffer (Thermo Fisher Scientific) containing a protease inhibitor cocktail tablet (Roche) as well as phosphatase inhibitor cocktail I and II (Sigma-Aldrich). Proteins were denatured by heating for 10 min at 95°C in sample buffer (2% SDS, 10% glycerol, 80 mM Tris, pH 6.8, and 1 mM DTT, and 50–100  $\mu\text{g}$  of proteins were electrophoresed in 4–20% bis-Tris/polyacrylamide gels (Invitrogen). Proteins were then transferred to 0.2- $\mu\text{m}$  nitrocellulose membranes (Invitrogen), which were then blocked for 2 h in blocking solution (TBS containing 10% nonfat dry milk and 0.05% Tween 20) and incubated overnight at 4°C with primary antibody diluted in blocking solution. On the next day, membranes were probed with horseradish peroxidase-conjugated secondary antibody for 1 h at room temperature, and immunoreactivity was detected by using enhanced chemiluminescence (Thermo Fisher Scientific).

**Immunoprecipitation assays.** Protein lysates (at the concentration of 1  $\mu\text{g}/\mu\text{l}$ ) were prepared from *Cdk5<sup>+/+C</sup>* and *Cdk5<sup>-/-C</sup>* lymphocytes collected from mice 4 wk after the induction of EAE. Lysates were precleared with

protein A-Agarose beads and then incubated overnight with anti-hamster coronin 1a antibody at 4°C. On the next day, lysates were incubated for an additional 4 h at 4°C with 50  $\mu\text{l}$  of a 50% protein A-Agarose bead slurry prepared in lysis buffer. After incubation, the protein A-Agarose beads were spun down and washed three times with lysis buffer and then subjected to Western blot analysis by using antibody against Cdk5 (C-8) and coronin 1a (anti-rabbit coronin 1a).

**Cdk5 kinase activity assay.** Cdk5 kinase activity assays were performed as described earlier (Pareek et al., 2007). In brief, tissue or cell lysates were prepared in RIPA buffer containing a protease inhibitor tablet (Roche) and the phosphatase inhibitor cocktails I and II (Sigma-Aldrich). Protein lysates (500  $\mu\text{g}$  of lumbar spinal cord, brain stem, or embryo brain lysates or 1 mg of spleen, lymphocyte, or T cell lysates) were dissolved in lysis buffer to achieve a 1- $\mu\text{g}/\mu\text{l}$  concentration and then precleared with normal rabbit IgG followed with 50–100  $\mu\text{l}$  of 50% protein A-Agarose slurry (Sigma-Aldrich) prepared in lysis buffer. These lysates were then incubated overnight at 4°C with 0.01  $\mu\text{g}/\mu\text{l}$  of anti-Cdk5 IgG. On the next day, lysates were subjected to a 3-h incubation with 50  $\mu\text{l}$  of a 50% protein A-Agarose slurry at 4°C. Immunoprecipitates were washed three times with lysis buffer followed with kinase buffer (20 mM Tris-HCl, pH 7.4, 10 mM  $\text{MgCl}_2$ , 1 mM EDTA, 10  $\mu\text{M}$  NaF, and 1  $\mu\text{M}$   $\text{Na}_2\text{VO}_3$ ) and resuspended in 10  $\mu\text{l}$  of 5 $\times$  kinase assay mixture (100 mM Tris-HCl, pH 7.4, 50 mM  $\text{MgCl}_2$ , 5 mM EDTA, 50  $\mu\text{M}$  NaF, 5  $\mu\text{M}$   $\text{Na}_2\text{VO}_3$ , and 5 mM DTT), 30  $\mu\text{l}$  of water, and 20  $\mu\text{M}$  of either the NF-H peptide or immunoprecipitated coronin 1a or coronin 1a peptides (WT or mutant 1, 2, or 3). Samples were kept at 30°C for 60 min after adding 5  $\mu\text{Ci}$   $\gamma$ -[ $^{32}\text{P}$ ]ATP (0.5 mM), and the reaction was stopped by adding 10% trichloroacetic acid in peptide kinase assay buffer and by adding sample buffer (2% SDS, 10% glycerol, 80 mM Tris, pH 6.8, and 1 mM DTT) and boiling for the 10 min for the coronin 1a kinase reaction. To detect coronin 1a phosphorylation, 20- $\mu\text{l}$  aliquots were electrophoresed on a 10% polyacrylamide gel, which was then stained with Coomassie blue, destained, dried, and exposed overnight to a phosphoscreen. The phosphoscreen was scanned on a Storm PhosphorImager (Molecular Dynamics). For peptide kinase assay, supernatants were collected and spotted in triplicate onto P81 phosphocellulose squares, air dried, and washed five times with 75 mM phosphoric acid and once in 95% ethanol. Phosphocellulose squares were then soaked in ultima gold liquid scintillation buffer (PerkinElmer), and the incorporated  $\gamma$ -[ $^{32}\text{P}$ ] was counted on a MicroBeta TriLux (PerkinElmer) as a measurement of Cdk5 activity.

**Lymphocyte and T cell isolation.** Mononuclear cells infiltrating mouse CNS tissues were collected by discontinuous percoll gradient as previously described (Cua et al., 2003). Total lymphocytes were collected from spleen and regional lymph nodes (axillary, branchial, and inguinal) by passing minced tissues through a 40- $\mu\text{m}$  cell strainer (BD) and thereafter incubating with ACK lysing buffer (Lonza) on ice for 5 min to deplete erythrocytes. For further purification, these cells were then subjected to MACS separation columns using a pan T isolation kit (Miltenyi Biotec), as per the manufacturer's protocol, to isolate total T cells.

**Thymidine assay of cell proliferation.** Total lymphocytes or T cells were stimulated with CD3/CD28, PMA-ionomycin, allogeneic dendritic cells, or MOG (35–55) for 72 h in a 96-well plate in RPMI-1640 culture media (Invitrogen) containing 50  $\mu\text{M}$  2-mercaptoethanol and 10% fetal bovine serum. During the last 16–18 h of culture, 1  $\mu\text{Ci}$   $^3\text{H}$ -thymidine was added to each well and cells were harvested on a Unifilter-96 Harvester (Perkin Elmer). Incorporation of  $^3\text{H}$ -thymidine was measured as counts per minute on a 1450 MicroBeta TriLux (Perkin Elmer).

**Histology.** Histology and immunocytochemical localization were performed as described previously (Pareek et al., 2006). In brief, 4–6 wk after EAE induction, mice were anesthetized and perfused with PBS followed with 4% paraformaldehyde in PBS (PFA-PBS). Spinal cords were collected and fixed in 4% PFA-PBS and embedded in paraffin to obtain 5- $\mu\text{m}$ -thick sections. The spinal cord was divided into four parts (cervical, thoracic, lumbar,

and sacral). At least five sections from each part and longitudinal serial sections of all regions were obtained for the entire representation of spinal cord. Hematoxylin/eosin stain was performed to assess routine histology and inflammation, and Luxol Fast blue counterstained with Periodic Acid Schiff reagent was used to analyze myelin content. Bielschowsky stain (silver stain) was performed to detect nerve fibers and analyze axonal loss. Images were captured with a digital slide microscope (DX-40; DMetrix, Inc.) and analyzed with eyepiece software.

**Immunocytochemical localization.** Lymphocytes were collected from spleen and regional lymph nodes and fixed with 4% PFA-PBS for 30 min. After three washes with PBS, cells were then permeabilized with 1% Triton X-100 (MP Biomedicals) and 10% normal goat serum in PBS for 10 min and further washed three times with PBS and blocked in blocking solution (10% vol/vol normal goat serum/PBS and 0.1% Tween 20). After blocking, cells were incubated overnight at 4°C with primary antibodies diluted in blocking solution. The next day, cells were washed three times with PBS and incubated for 1 h at room temperature with secondary conjugated goat anti-mouse or goat anti-rabbit IgGs. Cells were then mounted on glass slides using Vectashield mounting medium (Vector laboratories). Immunofluorescence images were acquired using a confocal microscope with a 63- $\mu$ m oil-immersion objective (LSM 510; Carl Zeiss, Inc.). Images were combined using LSM 510 imaging software (Carl Zeiss, Inc.) and managed in Photoshop (Adobe).

**Flow cytometry analyses.** Cells were isolated from bone marrow, spleen, and lymph node by passing tissue through a 40- $\mu$ m cell strainer. 200,000 cells were first incubated with FACS blocking buffer (DPBS with 0.1% BSA and 10% normal mouse serum) and stained with 1  $\mu$ g of CD3 (clone 145-2C11), CD4 (clone RM4-5), CD8 (clone 53-6.7), CD11b (clone 37.51), CD45R (clone RA3-6B2), CD45 (clone 30-F11), CD45.1 (clone A20), CD45.2 (clone 104), and CD34 (clone RAM-34) at room temperature and subjected to FACS analysis. All antibodies used for FACS analysis were purchased from either eBioscience or BD.

**Mass spectrometric analyses.** After the reduction and S-alkylation, sequencing grade modified trypsin (Promega) was used for the overnight digestion of cut gel slices at 37°C. Phosphopeptides were further enriched by MonoTip (GL Sciences, Inc.) according to the manufacturer's protocol. Trypsin-digested samples were next analyzed online using nanoflow high performance liquid chromatography–nanoelectrospray ionization on an LTQ–Orbitrap mass spectrometer (Thermo Fisher Scientific) coupled with Ultimate 3000 high performance liquid chromatography (Dionex). All data were acquired in positive ion mode. Tandem mass spectrometry spectra were searched against the mouse IPI protein database using the Mascot algorithm (Version 2.1; Matrix Science Ltd.). Search parameters included a static modification of 57.02146 D (carboxyamidomethylation) on Cys, dynamic modifications of 79.96633 D (phosphorylation) on Ser, Thr, and Tyr, and 15.99491 D (oxidation) on Met. All significant hits were manually inspected.

**Lymphocyte chemotaxis migration assay.** The ability of lymphocytes to move toward a chemoattractant signal was tested in a transmigration assay performed in a 6.5-mm transwell with a 5- $\mu$ m-pore polycarbonate membrane insert (Costar; Corning). In brief,  $\sim$ 10 million Jurkat cells, or Cdk5<sup>+/+C</sup> or Cdk5<sup>-/-C</sup> lymphocytes, were serum starved for 4 h and treated with or without 10  $\mu$ M roscovitine. These cells were cultured in the upper chamber of a transwell plate and the lower chamber was filled with 100 ng MIP-3  $\beta$ /CCL19 (PeproTech). After 15 h of culture, cells migrated in the lower chamber were counted by flow cytometry. Data from triplicate samples were plotted as the percent of input cells migrating in response to a chemokine.

**Statistical analyses.** Statistical evaluation was done with Prism (GraphPad Software, Inc.). Significant differences between experiments were assessed by univariate ANOVA (more than two groups) or unpaired Student's *t* test (two groups). ANOVA was followed by Student's *t* tests with a Bonferroni  $\alpha$  correction for multiple comparisons, where  $\alpha$  was set at 0.05.

**Online supplemental material.** Fig. S1 shows suppression of antigen receptor-mediated lymphocyte activation by roscovitine treatment. Fig. S2 shows the strategy for the generation of immune chimeric mice and presents the immunophenotyping data demonstrating no difference among Cdk5<sup>+/+</sup> and Cdk5<sup>-/-</sup>, p35<sup>+/+</sup> and p35<sup>-/-</sup>, and Cdk5<sup>+/+C</sup> and Cdk5<sup>-/-C</sup> at baseline. Fig. S3 shows representative sections of immunostaining performed to characterize immune cell infiltrates in spinal cord after EAE induction. Fig. S4 shows the mass spectrometry<sup>2</sup> spectrum and identification of coronin 1a with peptide mapping. Fig. S5 shows the results of immunostaining for Arp2/3 and F-actin in encephalitogenic lymphocytes revealing normal Arp2/3 polarization in lymphocytes from the Cdk5<sup>-/-C</sup> mice. Fig. S6 provides results of transmigration assays in which migration of Jurkat cells toward CCL19 is suppressed in the presence of roscovitine. Fig. S7 shows the results of assays of in vitro cytokine production by encephalitogenic splenocytes, during both the active and chronic phase of disease, after restimulation with MOG (35–55) ex vivo. Online supplemental material is available at <http://www.jem.org/cgi/content/full/jem.20100876/DC1>.

We thank Drs. Jean Pieters (University of Basel, Switzerland) and Andrew Chen (Genentech, USA) for providing coronin 1a antibody, Drs. Harish Pant (National Institute of Neurological Disorders and Stroke/National Institutes of Health), Veeranna (New York University) and Sashi Kesavapany (National University of Singapore) for valuable discussions, Michael Sramkoski (Case Comprehensive Cancer Center confocal core facility) for confocal microscopy, and Janet Robinson for experimental help.

We would like to acknowledge the support from the Jane and Lee Seidman Chair in Pediatric Cancer Innovation and The Case Research Institute.

The authors have no conflicting financial interests.

Submitted: 3 May 2010

Accepted: 20 September 2010

## REFERENCES

- Alt, C., M. Laschinger, and B. Engelhardt. 2002. Functional expression of the lymphoid chemokines CCL19 (ELC) and CCL 21 (SLC) at the blood-brain barrier suggests their involvement in G-protein-dependent lymphocyte recruitment into the central nervous system during experimental autoimmune encephalomyelitis. *Eur. J. Immunol.* 32:2133–2144. doi:10.1002/1521-4141(200208)32:8<2133::AID-IMMU2133>3.0.CO;2-W
- Appleton, B.A., P. Wu, and C. Wiesmann. 2006. The crystal structure of murine coronin-1: a regulator of actin cytoskeletal dynamics in lymphocytes. *Structure.* 14:87–96. doi:10.1016/j.str.2005.09.013
- Boulanger, L.M. 2009. Immune proteins in brain development and synaptic plasticity. *Neuron.* 64:93–109. doi:10.1016/j.neuron.2009.09.001
- Burkhardt, J.K., E. Carrizosa, and M.H. Shaffer. 2008. The actin cytoskeleton in T cell activation. *Annu. Rev. Immunol.* 26:233–259. doi:10.1146/annurev.immunol.26.021607.090347
- Chae, T., Y.T. Kwon, R. Bronson, P. Dikkes, E. Li, and L.H. Tsai. 1997. Mice lacking p35, a neuronal specific activator of Cdk5, display cortical lamination defects, seizures, and adult lethality. *Neuron.* 18:29–42. doi:10.1016/S0896-6273(01)80044-1
- Chen, F., and G.P. Studzinski. 2001. Expression of the neuronal cyclin-dependent kinase 5 activator p35Nck5a in human monocytic cells is associated with differentiation. *Blood.* 97:3763–3767. doi:10.1182/blood.V97.12.3763
- Clemen, C.S., V. Rybakina, and L. Eichinger. 2008. The coronin family of proteins. *Subcell. Biochem.* 48:1–5. doi:10.1007/978-0-387-09595-0\_1
- Cohen, P. 2002. Protein kinases—the major drug targets of the twenty-first century? *Nat. Rev. Drug Discov.* 1:309–315. doi:10.1038/nrd773
- Coussens, L.M., and Z. Werb. 2002. Inflammation and cancer. *Nature.* 420:860–867. doi:10.1038/nature01322
- Cruz, J.C., and L.H. Tsai. 2004. Cdk5 deregulation in the pathogenesis of Alzheimer's disease. *Trends Mol. Med.* 10:452–458. doi:10.1016/j.molmed.2004.07.001
- Cua, D.J., J. Sherlock, Y. Chen, C.A. Murphy, B. Joyce, B. Seymour, L. Lucian, W. To, S. Kwan, T. Churakova, et al. 2003. Interleukin-23 rather than interleukin-12 is the critical cytokine for autoimmune inflammation of the brain. *Nature.* 421:744–748. doi:10.1038/nature01355



- de Hostos, E.L., B. Bradtke, F. Lottspeich, R. Guggenheim, and G. Gerisch. 1991. Coronin, an actin binding protein of *Dictyostelium discoideum* localized to cell surface projections, has sequence similarities to G protein beta subunits. *EMBO J.* 10:4097–4104.
- Dhavan, R., and L.H. Tsai. 2001. A decade of CDK5. *Nat. Rev. Mol. Cell Biol.* 2:749–759. doi:10.1038/35096019
- Ferrari, G., H. Langen, M. Naito, and J. Pieters. 1999. A coat protein on phagosomes involved in the intracellular survival of mycobacteria. *Cell.* 97:435–447. doi:10.1016/S0092-8674(00)80754-0
- Föger, N., L. Rangell, D.M. Danilenko, and A.C. Chan. 2006. Requirement for coronin 1 in T lymphocyte trafficking and cellular homeostasis. *Science.* 313:839–842. doi:10.1126/science.1130563
- Fu, A.K., W.Y. Fu, A.K. Ng, W.W. Chien, Y.P. Ng, J.H. Wang, and N.Y. Ip. 2004. Cyclin-dependent kinase 5 phosphorylates signal transducer and activator of transcription 3 and regulates its transcriptional activity. *Proc. Natl. Acad. Sci. USA.* 101:6728–6733. doi:10.1073/pnas.0307606100
- Gandhi, M., and B.L. Goode. 2008. Coronin: the double-edged sword of actin dynamics. *Subcell. Biochem.* 48:72–87. doi:10.1007/978-0-387-09595-0\_7
- Gatfield, J., I. Albrecht, B. Zanolari, M.O. Steinmetz, and J. Pieters. 2005. Association of the leukocyte plasma membrane with the actin cytoskeleton through coiled coil-mediated trimeric coronin 1 molecules. *Mol. Biol. Cell.* 16:2786–2798. doi:10.1091/mbc.E05-01-0042
- Gong, X., X. Tang, M. Wiedmann, X. Wang, J. Peng, D. Zheng, L.A. Blair, J. Marshall, and Z. Mao. 2003. Cdk5-mediated inhibition of the protective effects of transcription factor MEF2 in neurotoxicity-induced apoptosis. *Neuron.* 38:33–46. doi:10.1016/S0896-6273(03)00191-0
- Han, M.H., S.I. Hwang, D.B. Roy, D.H. Lundgren, J.V. Price, S.S. Ousman, G.H. Fernald, B. Gerlitz, W.H. Robinson, S.E. Baranzini, et al. 2008. Proteomic analysis of active multiple sclerosis lesions reveals therapeutic targets. *Nature.* 451:1076–1081. doi:10.1038/nature06559
- Haraldsson, M.K., C.A. Louis-Dit-Sully, B.R. Lawson, G. Sternik, M.L. Santiago-Raber, N.R. Gascoigne, A.N. Theofilopoulos, and D.H. Kono. 2008. The lupus-related Lmb3 locus contains a disease-suppressing Coronin-1A gene mutation. *Immunity.* 28:40–51. doi:10.1016/j.immuni.2007.11.023
- Huang, D., G. Patrick, J. Moffat, L.H. Tsai, and B. Andrews. 1999. Mammalian Cdk5 is a functional homologue of the budding yeast Pho85 cyclin-dependent protein kinase. *Proc. Natl. Acad. Sci. USA.* 96:14445–14450. doi:10.1073/pnas.96.25.14445
- Karpus, W.J., and R.M. Ransohoff. 1998. Chemokine regulation of experimental autoimmune encephalomyelitis: temporal and spatial expression patterns govern disease pathogenesis. *J. Immunol.* 161:2667–2671.
- Kim, Y., J.Y. Sung, I. Ceglia, K.W. Lee, J.H. Ahn, J.M. Halford, A.M. Kim, S.P. Kwak, J.B. Park, S. Ho Ryu, et al. 2006. Phosphorylation of WAVE1 regulates actin polymerization and dendritic spine morphology. *Nature.* 442:814–817. doi:10.1038/nature04976
- Kitazawa, M., S. Oddo, T.R. Yamasaki, K.N. Green, and F.M. LaFerla. 2005. Lipopolysaccharide-induced inflammation exacerbates tau pathology by a cyclin-dependent kinase 5-mediated pathway in a transgenic model of Alzheimer's disease. *J. Neurosci.* 25:8843–8853. doi:10.1523/JNEUROSCI.2868-05.2005
- Kroenke, M.A., T.J. Carlson, A.V. Andjelkovic, and B.M. Segal. 2008. IL-12- and IL-23-modulated T cells induce distinct types of EAE based on histology, CNS chemokine profile, and response to cytokine inhibition. *J. Exp. Med.* 205:1535–1541. doi:10.1084/jem.20080159
- Kurosaki, T., and M. Hikida. 2009. Tyrosine kinases and their substrates in B lymphocytes. *Immunol. Rev.* 228:132–148. doi:10.1111/j.1600-065X.2008.00748.x
- Lin, H., M.C. Chen, C.Y. Chiu, Y.M. Song, and S.Y. Lin. 2007. Cdk5 regulates STAT3 activation and cell proliferation in medullary thyroid carcinoma cells. *J. Biol. Chem.* 282:2776–2784. doi:10.1074/jbc.M607234200
- Lu, S.X., O. Alpdogan, J. Lin, R. Balderas, R. Campos-Gonzalez, X. Wang, G.J. Gao, D. Suh, C. King, M. Chow, et al. 2008. STAT-3 and ERK 1/2 phosphorylation are critical for T-cell alloactivation and graft-versus-host disease. *Blood.* 112:5254–5258. doi:10.1182/blood-2008-03-147322
- Lundmark, F., K. Duvefelt, E. Jacobaeus, I. Kockum, E. Wallström, M. Khademi, A. Oturai, L.P. Ryder, J. Saarela, H.F. Harbo, et al. 2007. Variation in interleukin 7 receptor alpha chain (IL7R) influences risk of multiple sclerosis. *Nat. Genet.* 39:1108–1113. doi:10.1038/ng2106
- Mapelli, M., L. Massimiliano, C. Crovace, M.A. Seeliger, L.H. Tsai, L. Meijer, and A. Musacchio. 2005. Mechanism of CDK5/p25 binding by CDK inhibitors. *J. Med. Chem.* 48:671–679. doi:10.1021/jm049323m
- Matthews, S.A., and D.A. Cantrell. 2006. The role of serine/threonine kinases in T-cell activation. *Curr. Opin. Immunol.* 18:314–320. doi:10.1016/j.coi.2006.03.006
- Mi, S., B. Hu, K. Hahm, Y. Luo, E.S. Kam Hui, Q. Yuan, W.M. Wong, L. Wang, H. Su, T.H. Chu, et al. 2007. LINGO-1 antagonist promotes spinal cord remyelination and axonal integrity in MOG-induced experimental autoimmune encephalomyelitis. *Nat. Med.* 13:1228–1233. doi:10.1038/nm1664
- Miyamoto, Y., J. Yamauchi, and A. Tanoue. 2008. Cdk5 phosphorylation of WAVE2 regulates oligodendrocyte precursor cell migration through nonreceptor tyrosine kinase Fyn. *J. Neurosci.* 28:8326–8337. doi:10.1523/JNEUROSCI.1482-08.2008
- Mueller, P., J. Massner, R. Jayachandran, B. Combaluzier, I. Albrecht, J. Gatfield, C. Blum, R. Ceredig, H.R. Rodewald, A.G. Rolink, and J. Pieters. 2008. Regulation of T cell survival through coronin-1-mediated generation of inositol-1,4,5-trisphosphate and calcium mobilization after T cell receptor triggering. *Nat. Immunol.* 9:424–431. doi:10.1038/ni1570
- Nikolic, M., M.M. Chou, W. Lu, B.J. Mayer, and L.H. Tsai. 1998. The p35/Cdk5 kinase is a neuron-specific Rac effector that inhibits Pak1 activity. *Nature.* 395:194–198. doi:10.1038/26034
- Noseworthy, J.H., C. Lucchinetti, M. Rodriguez, and B.G. Weinshenker. 2000. Multiple sclerosis. *N. Engl. J. Med.* 343:938–952. doi:10.1056/NEJM200009283431307
- Ohshima, T., J.M. Ward, C.G. Huh, G. Longenecker, Veeranna, H.C. Pant, R.O. Brady, L.J. Martin, and A.B. Kulkarni. 1996. Targeted disruption of the cyclin-dependent kinase 5 gene results in abnormal corticogenesis, neuronal pathology and perinatal death. *Proc. Natl. Acad. Sci. USA.* 93:11173–11178. doi:10.1073/pnas.93.20.11173
- Pareek, T.K., J. Keller, S. Kesavapany, H.C. Pant, M.J. Iadarola, R.O. Brady, and A.B. Kulkarni. 2006. Cyclin-dependent kinase 5 activity regulates pain signaling. *Proc. Natl. Acad. Sci. USA.* 103:791–796. doi:10.1073/pnas.0510405103
- Pareek, T.K., J. Keller, S. Kesavapany, N. Agarwal, R. Kuner, H.C. Pant, M.J. Iadarola, R.O. Brady, and A.B. Kulkarni. 2007. Cyclin-dependent kinase 5 modulates nociceptive signaling through direct phosphorylation of transient receptor potential vanilloid 1. *Proc. Natl. Acad. Sci. USA.* 104:660–665. doi:10.1073/pnas.0609916104
- Pawson, T., and J.D. Scott. 1997. Signaling through scaffold, anchoring, and adaptor proteins. *Science.* 278:2075–2080. doi:10.1126/science.278.5346.2075
- Pearce, J.M. 2005. Historical descriptions of multiple sclerosis. *Eur. Neurol.* 54:49–53. doi:10.1159/000087387
- Prince, A.L., C.C. Yin, M.E. Enos, M. Felices, and L.J. Berg. 2009. The Tec kinases Itk and Rlk regulate conventional versus innate T-cell development. *Immunol. Rev.* 228:115–131. doi:10.1111/j.1600-065X.2008.00746.x
- Proudfoot, A.E., A.L. de Souza, and V. Muzio. 2008. The use of chemokine antagonists in EAE models. *J. Neuroimmunol.* 198:27–30. doi:10.1016/j.jneuroim.2008.04.007
- Rashid, T., M. Banerjee, and M. Nikolic. 2001. Phosphorylation of Pak1 by the p35/Cdk5 kinase affects neuronal morphology. *J. Biol. Chem.* 276:49043–49052. doi:10.1074/jbc.M105599200
- Rosales, J.L., and K.Y. Lee. 2006. Extraneuronal roles of cyclin-dependent kinase 5. *Bioessays.* 28:1023–1034. doi:10.1002/bies.20473
- Salmond, R.J., A. Filby, I. Qureshi, S. Caserta, and R. Zamojska. 2009. T-cell receptor proximal signaling via the Src-family kinases, Lck and Fyn, influences T-cell activation, differentiation, and tolerance. *Immunol. Rev.* 228:9–22. doi:10.1111/j.1600-065X.2008.00745.x
- Schneider, A., G.W. Araújo, K. Trajkovic, M.M. Herrmann, D. Merkler, E.M. Mandelkow, R. Weissert, and M. Simons. 2004. Hyperphosphorylation and aggregation of tau in experimental autoimmune encephalomyelitis. *J. Biol. Chem.* 279:55833–55839. doi:10.1074/jbc.M409954200

- Shelton, S.B., and G.V. Johnson. 2004. Cyclin-dependent kinase-5 in neurodegeneration. *J. Neurochem.* 88:1313–1326. doi:10.1111/j.1471-4159.2003.02328.x
- Shiow, L.R., D.W. Roadcap, K. Paris, S.R. Watson, I.L. Grigorova, T. Lebet, J. An, Y. Xu, C.N. Jenne, N. Föger, et al. 2008. The actin regulator coronin 1A is mutant in a thymic egress-deficient mouse strain and in a patient with severe combined immunodeficiency. *Nat. Immunol.* 9:1307–1315. doi:10.1038/ni.1662
- Shiow, L.R., K. Paris, M.C. Akana, J.G. Cyster, R.U. Sorensen, and J.M. Puck. 2009. Severe combined immunodeficiency (SCID) and attention deficit hyperactivity disorder (ADHD) associated with a Coronin-1A mutation and a chromosome 16p11.2 deletion. *Clin. Immunol.* 131:24–30. doi:10.1016/j.clim.2008.11.002
- Strock, C.J., J.I. Park, E.K. Nakakura, G.S. Bova, J.T. Isaacs, D.W. Ball, and B.D. Nelkin. 2006. Cyclin-dependent kinase 5 activity controls cell motility and metastatic potential of prostate cancer cells. *Cancer Res.* 66:7509–7515. doi:10.1158/0008-5472.CAN-05-3048
- Studzinski, G.P., and J.S. Harrison. 2003. The neuronal cyclin-dependent kinase 5 activator p35Nck5a and Cdk5 activity in monocytic cells. *Leuk. Lymphoma.* 44:235–240. doi:10.1080/1042819021000030009
- Thome, M., and O. Acuto. 1995. Molecular mechanism of T-cell activation: role of protein tyrosine kinases in antigen receptor-mediated signal transduction. *Res. Immunol.* 146:291–307. doi:10.1016/0923-2494(96)80263-X
- Upadhyay, A.K., A.K. Ajay, S. Singh, and M.K. Bhat. 2008. Cell cycle regulatory protein 5 (Cdk5) is a novel downstream target of ERK in carboplatin induced death of breast cancer cells. *Curr. Cancer Drug Targets.* 8:741–752. doi:10.2174/156800908786733405
- Wilker, P.R., M. Kohyama, M.M. Sandau, J.C. Albring, O. Nakagawa, J.J. Schwarz, and K.M. Murphy. 2008. Transcription factor Mef2c is required for B cell proliferation and survival after antigen receptor stimulation. *Nat. Immunol.* 9:603–612. doi:10.1038/ni.1609
- Wyss-Coray, T., and L. Mucke. 2002. Inflammation in neurodegenerative disease—a double-edged sword. *Neuron.* 35:419–432. doi:10.1016/S0896-6273(02)00794-8
- Xie, Z., K. Sanada, B.A. Samuels, H. Shih, and L.H. Tsai. 2003. Serine 732 phosphorylation of FAK by Cdk5 is important for microtubule organization, nuclear movement, and neuronal migration. *Cell.* 114:469–482. doi:10.1016/S0092-8674(03)00605-6
- Yaffe, M.B., M. Schutkowski, M. Shen, X.Z. Zhou, P.T. Stukenberg, J.U. Rahfeld, J. Xu, J. Kuang, M.W. Kirschner, G. Fischer, et al. 1997. Sequence-specific and phosphorylation-dependent proline isomerization: a potential mitotic regulatory mechanism. *Science.* 278:1957–1960. doi:10.1126/science.278.5345.1957
- Yang, H.S., and P.W. Hinds. 2003. Increased ezrin expression and activation by CDK5 coincident with acquisition of the senescent phenotype. *Mol. Cell.* 11:1163–1176. doi:10.1016/S1097-2765(03)00135-7