



Published in final edited form as:

Curr Opin Pediatr. 2010 August ; 22(4): 508–515. doi:10.1097/MOP.0b013e32833b7c23.

Hypercalcemia in Children and Adolescents

Steven A. Lietman, M.D.¹, Emily L. Germain-Lee, M.D.², and Michael A. Levine, M.D.³

¹Orthopaedic and Rheumatologic Institute, Cleveland Clinic

²Division of Pediatric Endocrinology, Johns Hopkins University School of Medicine, and Bone Center & Osteogenesis Imperfecta Program, Kennedy Krieger Institute

³Division of Endocrinology and Diabetes, The Children's Hospital of Philadelphia and Department of Pediatrics, University of Pennsylvania School of Medicine

Abstract

Purpose of the review—In this review we define hypercalcemia levels, common etiologies for hypercalcemia in children, and treatment in order to aid the practicing pediatrician.

Recent Findings—One rare cause of hypercalcemia in the child is *Familial hypocalciuric hypercalcemia* (FHH) [also termed familial benign hypercalcemia (FBH)]. Mutations that inactivate the *Ca²⁺-sensing receptor gene* FHH have been described as an autosomal dominant disorder, but recently milder mutations in the *CASR* have been shown to cause hypercalcemia when homozygous.

Summary—Normal serum levels of calcium are maintained through the interplay of parathyroid, renal, and skeletal factors. In this review, we have distinguished the neonate and infant from the older child and adolescent because the causes and clinical features of hypercalcemia can differ in these two age groups. However the initial approach to the medical treatment of severe or symptomatic hypercalcemia is to increase the urinary excretion of calcium in both groups. In most cases, hypercalcemia is due osteoclastic bone resorption, and agents that inhibit or destroy osteoclasts are therefore effective treatments. Parathyroid surgery, the conventional treatment for adults with symptomatic primary hyperparathyroidism, is recommended for all children with primary hyperparathyroidism.

Keywords

hypercalcemia; pediatrics; bisphosphonates; hyperparathyroidism

Introduction

Hypercalcemia is less common in children than in adults, but is more likely to be clinically significant in younger patients as routine biochemical screening tests are rarely performed in children. Normal serum levels of calcium are maintained through the interplay of parathyroid, renal, and skeletal factors. The principle calciotropic hormone is parathyroid hormone (PTH), which is synthesized and secreted from the parathyroid glands and at a rate inversely proportional to the circulating level of ionized calcium. PTH secretion is regulated through the interaction of extracellular calcium with specific calcium-sensing receptors (CaSR's) that are present on the surface of the parathyroid cell. In turn, PTH regulates mineral metabolism and skeletal homeostasis by activating specific membrane receptors on target cells in bone and kidney that bind PTH and the related hormone parathyroid hormone-related peptide (PTHrP)

Address correspondence to: Dr. Steven A. Lietman, A41 Crile Building, 9500 Euclid Avenue, Cleveland, Ohio 44195, (T)216-445-2742, lietmas@ccf.org.

with equal affinity. In the kidney, PTH stimulates conversion of 25(OH)D to calcitriol (1,25(OH)₂D), the active metabolite of vitamin D. Both PTH and calcitriol activate osteoclastic bone resorption and increase renal absorption of filtered calcium, while calcitriol can also increase active transport of calcium in the intestine. While absorption of too much calcium from the intestine can account for some cases of hypercalcemia, in most patients hypercalcemia is due to excessive osteoclastic activity. PTH and other factors that stimulate osteoclastic activity interact directly with receptors on osteoblasts, the bone forming cells, to increase their expression of RANKL, a ligand that binds to the receptor activator of NF-kappa B (RANK) on osteoclast precursors, and to decrease production of osteoprotegerin (OPG), a circulating decoy receptor for RANKL. RANKL stimulates activation, migration, differentiation, and fusion of hematopoietic cells of the osteoclast lineage to stimulate bone resorption (1,2).

Serum calcium concentrations are greater in children than in adults (Table 1 (3)), therefore the diagnosis of hypercalcemia in younger patients requires use of age-appropriate normal ranges. In this review, we have distinguished the neonate and infant from the older child and adolescent because the causes and clinical features of hypercalcemia can differ in these two age groups.

Diagnosis of hypercalcemia in neonates and infants

The differential diagnosis of hypercalcemia in neonates and infants up to age 2 years is listed in Table 2. (4,5) Clinical features of hypercalcemia may be nonspecific in neonates and infants, and hypercalcemia is often discovered when a chemistry panel is obtained to evaluate failure-to-thrive. Severe hypercalcemia can affect the nervous system and cause weakness, hypotonia with proximal myopathy, lethargy and stupor, and rarely seizures. Hypercalcemia can also induce polyuria and dehydration, as stimulation of the renal CaSRs in the collecting tubule decreases expression of aquaporin-2 at the apical plasma membrane (6,7) and leads to resistance to vasopressin and diabetes insipidus. In addition, there can be damage to the kidneys from nephrocalcinosis (4,5).

In the small infant, *enriched formulas that provide excess calcium* can quickly lead to hypercalcemia. Phosphate depletion can cause a defect in mineralization and hypercalcemia, and in the past often resulted from human milk feeding in preterm, very-low-birth weight infants (4,5). The introduction of breast milk fortifiers, which contain 30-40 mg/kg/day of phosphate, for these infants has all but eliminated this problem. By contrast, a far more common cause of phosphate depletion is from inappropriately supplemented parenteral nutrition (8,9). Hypophosphatemia suppresses circulating levels of fibroblast growth factor 23 (FGF23), a phosphate-regulating hormone or “phosphatonin,” with subsequent disinhibition of calcitriol production. Elevated serum levels of calcitriol stimulate intestinal absorption of calcium and activate osteoclastic bone resorption. Finally, *extracorporeal membrane oxygenation* can also cause hypercalcemia, but the mechanism is unclear (10).

Neonatal hyperparathyroidism

Neonatal hyperparathyroidism is uncommon, and is usually due to multi-gland hyperplasia rather than to parathyroid adenoma as in older patients (11). Severe bone deformities, as well as fractures, may be present at birth and reflect impaired mineralization and intense osteoclastic bone resorption. Respiratory difficulties can occur if the rib cage is affected. There may be hepatosplenomegaly and anemia. Failure to normalize the serum calcium level can have profound developmental implications (12).

In some infants neonatal hyperparathyroidism represents an adaptation to maternal hypocalcemia, most commonly from maternal hypoparathyroidism or vitamin D deficiency (13) but also from pseudohyperparathyroidism (14) and renal tubular acidosis (15). Serum

calcium levels are generally normal but hypercalcemia has been reported in 25% of these infants (16). Hyperparathyroidism typically resolves within a few weeks.

Familial hypocalciuric hypercalcemia (FHH) [also termed familial benign hypercalcemia (FBH)] is characterized by asymptomatic hypercalcemia and very low levels of urinary calcium. Mutations that inactivate the *Ca²⁺-sensing receptor gene (CASR)*, localized to chromosome 3q, are the most common cause of FHH, and are also associated with neonatal severe hyperparathyroidism (NSHPT), a life threatening form of primary hyperparathyroidism that is associated with very high levels of serum calcium and PTH. FHH and NSHPT can occur in the same family, and represent the presence of one or two defective *CASR* alleles, respectively (17-21). NSHPT can also occur in heterozygous offspring born to affected fathers but unaffected normocalcemic mothers, or in neonates with an apparent *de novo* heterozygous mutations in the *CASR* gene (22,23). In addition to hypercalcemia, infants with NSHPT have hypophosphatemia and osteopenia at birth (20). The bone disease is likely a reflection of both elevated PTH and decreased *CASR* activity in osteoblasts and osteoclasts (24).

Reduced expression of CaSRs decreases the sensitivity of the parathyroid cells to extracellular Ca^{2+} and leads to mild hyperplasia of all parathyroid glands. The CaSR is also expressed in the kidney, and decreased receptor activity in the distal nephron accounts for the relative hypocalciuria (i.e., the fractional excretion of calcium is less than 1%) that is the hallmark of the disorder.

Conventional treatment of NSHPT has been urgent subtotal parathyroidectomy, but intravenous bisphosphonates may be used to reduce the serum calcium level quickly and either delay or obviate surgery (23,25). Recent studies of calcimimetic compounds, which activate the parathyroid cell CaSR and inhibit PTH secretion, in adult patients (26) and a single adolescent (27) with primary hyperparathyroidism suggest that these agents may be useful medical alternatives or adjuncts to surgery for control of hypercalcemia. Type I calcimimetics are direct receptor agonists while type II calcimimetics are allosteric activators that interact with the membrane-spanning segments of the CaSR and enhance signal transduction, presumably by inducing conformational changes in the receptor. The presumed conformational change reduces the threshold for CaSR activation by the endogenous ligand, Ca^{2+} , thereby reducing PTH secretion in the absence of a change in the level of extracellular Ca^{2+} . Type II calcimimetics have been shown *in vitro* to enhance the potency of extracellular Ca^{2+} to activate mutant CaSRs (28,29) and have been used successfully in an older patient with FHH and symptomatic hypercalcemia (30), but efficacy in children with FHH or NSPHT has not been reported.

FHH is typically an autosomal dominant disorder, but some milder mutations in the *CASR* only cause hypercalcemia when homozygous (31). In nearly all cases FHH (and NSHPT) is due to mutation of the *CASR* gene, but the disorder has been also linked to the long (32,33) and short (34) arms of chromosome 19, indicating genetic heterogeneity for this disorder.

Non-parathyroid causes of hypercalcemia

Hypercalcemia may be a manifestation of *vitamin D intoxication* in newborns whose mothers ingested *excessive amounts of vitamin D* and/or its derivatives during pregnancy (14). Due to the tight regulation of renal 25(OH)D-1 alpha hydroxylase in the kidney, serum levels of 25 (OH)D but not calcitriol are elevated in these infants. By contrast, serum levels of calcitriol are markedly elevated in hypercalcemic infants with *subcutaneous fat necrosis*, an unusual disorder that occurs in some neonates after a complicated delivery. Hypercalcemia can occur days or weeks after birth, and results from production of excess calcitriol by macrophages within the granulomatous reaction to the necrotic fat. Calcium released from necrotic fat tissue and increased prostaglandin E activity further aggravates hypercalcemia (35-37). Failure-to-

thrive is the most common clinical sign associated with subcutaneous fat necrosis, which is associated with a significant 15% mortality (37).

Williams syndrome is associated with hypercalcemia in approximately 15% of cases. Hypercalcemia typically occurs during infancy and resolves between 2 and 4 years of age. There are cases, however, of older children and adults who have persistent hypercalcemia. Circulating levels of calcitriol are elevated in some but not all patients with Williams syndrome, and PTH levels are low (38). A common finding is increased sensitivity to vitamin D, which appears to be due to decreased ability of ligand-bound Vitamin D receptor (VDR) to mediate transrepression of the 25(OH)D₃ 1 α -hydroxylase gene, *CYP27B1*. A potential explanation for this defect is haploinsufficiency of the Williams syndrome transcription factor (WSTF) that is located within the 7q11.23 deleted region. The VDR interacts with a multifunctional, ATP-dependent chromatin remodeling complex termed WINAC in a ligand-independent manner through WSTF, and loss of WSTF prevents calcitriol-bound VDR-induced transrepression of *CYP27B1* (39,40).

Inborn errors of metabolism that cause hypercalcemia

Hypophosphatasia, a metabolic bone disease characterized by bone and teeth hypomineralization due to defective function of tissue-nonspecific alkaline phosphatase (TNSALP), is also associated with hypercalcemia. Hypophosphatasia is caused by various mutations in the *ALPL* gene at 1p34–36 (41–44), and is classified into six clinical forms depending on the age at diagnosis and the severity of the symptoms: perinatal lethal; infantile; childhood; adult; odontohypophosphatasia; and perinatal benign. Infantile hypophosphatasia presents before age six months and can cause severe hypercalcemia. Deficiency of alkaline phosphatase impairs skeletal mineralization and calcium uptake, leading to hypercalcemia with hypercalciuria and nephrocalcinosis. Affected infants have respiratory complications due to rachitic deformities of the chest. Despite the presence of an open fontanelle, premature craniosynostosis is a common finding that may result in increased intracranial pressure. Serum levels of total and bone-specific alkaline phosphatase are low, and the diagnosis is supported by the presence of very high levels urinary phosphoethanolamine. Mutational analysis of the *ALPL* gene can provide molecular confirmation. In infants who survive, there is often spontaneous improvement in mineralization and remission of clinical problems, with the exception of craniosynostosis. Although there is no known treatment of hypophosphatasia, some patients have shown improvement after transplantation of bone marrow, bone or mesenchymal stem cells(45). A bone-targeted form of enzyme replacement therapy (46) is now undergoing clinical trials.

Blue diaper syndrome is an uncommon metabolic disorder that is due to a defect in tryptophan metabolism (47). The block in tryptophan metabolism leads to urinary excretion of excessive amounts of indole derivatives, including a derivative called “indican” that gives the urine-soaked diaper a blue tint. Affected infants have hypercalcemia, hypercalciuria and nephrocalcinosis, but the mechanism is unknown.

Hypercalcemia has been described in infants with *congenital lactase deficiency*. Hypercalcemia typically resolves after initiation of a lactose-free diet, but hypercalciuria and nephrocalcinosis may persist (48). The etiology of the hypercalcemia is unclear, but may be due to metabolic acidosis and/or an increase in intestinal calcium absorption secondary to increased gut lactose (48). A similar mechanism may explain hypercalcemia in infants with disaccharide intolerance (49).

Bartter syndrome is commonly associated with hypercalciuria (50), but hypercalcemia has been described in some infants with homozygous inactivation in the gene for either the

furosemide-sensitive NaK-2Cl-cotransporter NKCC2 (*SLC12A1*) or the inwardly rectifying potassium channel ROMK (*KCNJ1*) (51,52).

Hypercalcemia and/or hypercalciuria can also occur in children with the *IMAGe* syndrome, a poorly defined disorder that consists of Intrauterine growth retardation, Metaphyseal dysplasia, Adrenal hypoplasia congenital, and Genital defects (53). Neither the molecular defect, or the basis for hypercalcemia, has been identified.

Hypercalcemia in older children

A list of biochemical features that aid in the differential diagnosis of hypercalcemia in children is presented in Table 3, and an algorithm for evaluation of the child with hypercalcemia is presented in Figure 1 (54). In older children primary hyperparathyroidism, immobilization and malignancy are the principle causes of hypercalcemia.

Primary hyperparathyroidism in children and adolescents

Primary hyperparathyroidism is usually *sporadic* in the child or adolescent, and is nearly always (65%) due to a single parathyroid adenoma. The age range is from 3-19 years with a mean of 12.8 years and a 3:2 female:male incidence (55). Primary hyperparathyroidism is far less common in children and adolescents than in adults. Nearly all patients (79%) are symptomatic at presentation and end-organ damage (nephrocalcinosis, nephrolithiasis, acute pancreatitis, or bone involvement) is common (44%).

Primary hyperparathyroidism can also be an autosomal dominant *genetic* disorder that is typically associated with multi-gland hyperplasia (56). Primary hyperparathyroidism is most commonly (57%) the presenting manifestation of MEN type I (55), but can also be the initial feature of the hyperparathyroidism-jaw tumor syndrome, which is associated with parathyroid carcinoma (57,58). Primary hyperparathyroidism is less often a manifestation of MEN type II. Children with asymptomatic, mild hypercalcemia are likely to have FHH.

Non-parathyroid causes of hypercalcemia in older children and adolescents

Hypervitaminosis D can occur in children who ingest *excessive amounts of vitamin D* (or its metabolites). Although the upper limit of vitamin D tolerability for adults is 10,000 units per day, over time young children and infants may develop vitamin D intoxication when receiving only 2000-4000 units daily (14,54). Serum levels of 25(OH)D are elevated, but serum levels of calcitriol are usually normal; PTH levels are suppressed. Endogenous vitamin D intoxication can occur in patients with *granulomatous* disease and other inflammatory disorders. Infectious diseases such as cat scratch fever (59) as well as histoplasmosis, coccidiomycosis, leprosy and tuberculosis have all been associated with hypercalcemia in children. In these conditions activated T cells and macrophages express 25-(OH)D-1-alpha hydroxylase activity, which converts 25(OH)D to calcitriol (60).

Immobilization

Immobilization is a common cause of hypercalciuria in children and adolescents and can also cause hypercalcemia. When a rapidly growing child is immobilized or placed on bed rest there is a marked decrease in osteoblastic bone formation and a corresponding increase in osteoclastic bone resorption. This imbalance in bone remodeling leads to excessive mobilization of calcium (and phosphate) from the skeleton; the consequent net loss of bone mass is designated disuse osteoporosis (61).

Hypercalcemia of Malignancy

Hypercalcemia occurs in less than 1% of children with cancer, and has been reported with leukemia, lymphoma, myeloma, neuroblastoma, hepatocellular carcinoma, ovarian carcinoma, hepatoblastoma, rhabdomyosarcoma, brain cancer and dysgerminomas. Malignancy-associated hypercalcemia can be attributed to two general mechanisms: 1) osteolytic due to direct invasion of the skeleton by tumor cells and 2) humoral due to tumor production of circulating factors that activate osteoclastic bone resorption. The most commonly identified humoral factor that causes hypercalcemia of malignancy is PTHrP. PTHrP normally acts as a paracrine and intracrine factor, but some tumors can secrete sufficient PTHrP into the circulation to induce hypercalcemia via interaction with the type 1 PTH/PTHrP receptor. The specific tumors that characteristically produce humoral hypercalcemia of malignancy via secretion of PTHrP include squamous cell carcinoma of the lung, head and neck, renal cell carcinoma, breast and ovarian carcinoma, adult T cell leukemia, and dysgerminoma. Other tumor-produced factors that play a role in producing hypercalcemia include calcitriol, prostaglandins, interleukin 1 and interleukin 6, transforming growth factor β , and tumor necrosis factor.

Miscellaneous Causes

A variety of unusual causes of hypercalcemia must be considered in children who do not have any of the disorders discussed above (Table 4). Hypercalcemia can occur after *chronic ingestion of vitamin A*. The child develops anorexia, pruritus, irritability, bone pain and tender swellings of bone. Associated features include osteopenia due to increased osteoclastic bone resorption, hyperostosis of the shafts of the long bones, and osteophyte formation, particularly in the thoracic spine. Severe hypercalcemia has also been associated with the administration of the vitamin A analog all-trans-retinoic acid (ATRA) during therapy for acute promyelocytic leukemia.

Children with *Jansen metaphyseal chondrodysplasia* have hypercalcemia and bone lesions that are typical of primary hyperparathyroidism, but have suppressed PTH levels. This unusual disorder is due to mutations in the *PTHrP gene* encoding the type 1 PTH/PTHrP receptor that lead to ligand-independent activation of signaling. This causes increased bone resorption, metaphyseal defects, elevated serum levels of calcitriol, and growth delay.

Treatment

The initial approach to the medical treatment of severe or symptomatic hypercalcemia is to increase the urinary excretion of calcium. Infants are frequently dehydrated, and two thirds to full strength saline containing 30 mEq of potassium chloride per liter should be infused to correct dehydration and maximize glomerular filtration rate. Furosemide and other powerful loop diuretics are rarely necessary, and can induce excessive diuresis and dehydration; the consequent fall in the glomerular filtration rate can worsen hypercalcemia.

In most cases, hypercalcemia is due osteoclastic bone resorption, and agents that inhibit or destroy osteoclasts will be effective treatments. Calcitonin (2-4 U/kg q12h) given by subcutaneous injection is effective at first, but resistance to the hormone occurs quite rapidly. The nitrogen-containing bisphosphonates, including alendronate, ibandronate, pamidronate disodium, risedronate, and zoledronic acid, induce osteoclast apoptosis and are potent inhibitors of bone resorption. Zoledronic acid is a new-generation, heterocyclic nitrogen-containing bisphosphonate and the most potent inhibitor of bone resorption identified to date. Both pamidronate disodium and zoledronic acid can rapidly lower serum and urinary calcium levels in patients with hypercalcemia due to a variety of causes, and the effects can last for weeks. Patients must be monitored carefully, as these powerful agents can cause severe

hypocalcemia, hypophosphatemia, and hypomagnesemia. In addition, approximately 20% of patients will experience an untoward acute phase reaction after receiving an initial intravenous infusion. There is increasing concern about the development of osteonecrosis of the jaw in patients who are receiving chronic or prolonged bisphosphonate therapy. Adult patients with multiple myeloma and metastatic carcinoma to the skeleton who receive multiple, frequent doses of intravenous, nitrogen-containing bisphosphonates appear to be at greatest risk for osteonecrosis of the jaw, and these patients account for nearly all published cases. The mandible is more commonly affected than the maxilla (2:1 ratio), and most cases are preceded by a recent dental surgical procedure (62). Oversuppression of bone turnover is probably the primary mechanism for the development of this condition, although there may be contributing comorbid factors. Osteonecrosis of the jaw has not been reported in children who are receiving bisphosphonates.

Due to the low frequency of hypercalcemia in children, comprehensive clinical trials on the safety and efficacy of the bisphosphonates in children are lacking, although several small studies have reported promising results for these agents in the treatment of young patients with hypercalcemia (25,63-65). In particular, bisphosphonates can rapidly reverse the hypercalcemia and hypercalciuria of immobilization (66).

Because of concerns regarding potential late adverse effects of bisphosphonates on growth and development of the skeleton, calcitonin tends to be used more frequently in children because it has no long-term sequelae. Glucocorticoid steroids, which are effective treatments for hypercalcemia associated with vitamin D excess, are used with caution in children because they impair linear growth and bone mineralization. Children with Williams syndrome or idiopathic infantile hypercalcemia often have mildly elevated serum levels of calcitriol, and a low calcium formula in the infant or reduced calcium diet in the older child may be all that is needed to treat the hypercalcemia and/or hypercalciuria, particularly when long term treatment will be necessary. CalciloXD (Ross Laboratories, North Chicago, Ill.) is a low calcium infant formula without vitamin D that is commonly used. As the hypercalcemia improves, the CalciloXD can be gradually mixed with regular formula or breast milk. The infants and children on the low calcium diet need to be followed closely, however, for the possible development of hypocalcemia and rickets.

Parathyroid surgery, the conventional treatment for adults with symptomatic primary hyperparathyroidism, is recommended for all children with primary hyperparathyroidism. Indeed, the younger age of pediatric patients provides an even more compelling justification for recommending surgery for most patients. Of course, hypercalcemic children with FHH will rarely require any intervention unless they have NSPHT, and then parathyroidectomy may be necessary. The calcimimetic cinacalcet has been used in small studies of children and adolescents with hyperparathyroidism secondary to renal failure, and effectively lowers serum calcium and PTH, but does not alter bone turnover or increase bone mineral density. (67)

Summary

The etiology of hypercalcemia in children is age-dependent and includes a broad differential diagnosis (Table 3 and Figure 1). Although these conditions are not common, it is nevertheless important not to overlook them, as untreated hypercalcemia can have a profound impact on a child's growth and development.

Acknowledgments

This work has been supported in part by grants from the National Institutes of Health (DK-34281, DK-56178, and DE-18237).

Bibliography

1. Lean JM, Matsuo K, Fox SW, Fuller K, Gibson FM, Draycott G, et al. Osteoclast lineage commitment of bone marrow precursors through expression of membrane-bound TRANCE \dagger . *Bone* 2000;27(1):29–40. [PubMed: 10865206]
2. Fuller K, Wong B, Fox S, Choi Y, Chambers TJ. TRANCE \dagger is necessary and sufficient for osteoblast-mediated activation of bone resorption in osteoclasts. *JExpMed* 1998;188(5):997–1001. \dagger TRANCE (Tumor necrosis factor-related activation-induced cytokine) is another term for RANKL.
3. Portale, AA. Blood calcium, phosphorus, and magnesium. In: Favus, MJ., editor. *Primer on the Metabolic Bone Diseases and Disorders of Mineral Metabolism*. Fourth. Lippincott: Williams, & Wilkins; 1999. p. 116
4. Nishiyama S. Hypercalcemia in children: an overview. *Acta Paediatr Jpn* 1997 Aug;39(4):479–84. [PubMed: 9316297]
5. Rodd C, Goodyer P. Hypercalcemia of the newborn: etiology, evaluation, and management. *Pediatr Nephrol* 1999 Aug;13(6):542–7. [PubMed: 10452286]
6. Hasler U, Leroy V, Martin PY, Feraille E. Aquaporin-2 abundance in the renal collecting duct: new insights from cultured cell models. *Am J Physiol Renal Physiol* 2009 Jul;297(1):F10–8. [PubMed: 19244407]
7. Procino G, Mastrofrancesco L, Mira A, Tamma G, Carmosino M, Emma F, et al. Aquaporin 2 and apical calcium-sensing receptor: new players in polyuric disorders associated with hypercalciuria. *Semin Nephrol* 2008 May;28(3):297–305. [PubMed: 18519090]
8. Kimura S, Nose O, Harada T, Maki I, Kanaya S, Tajiri H, et al. Serum levels of vitamin D metabolites in children receiving total parenteral nutrition. *J Parenter Enteral Nutr* 1986 Mar-Apr;10(2):191–4.
9. Kimura S, Nose O, Seino Y, Harada T, Kanaya S, Yabuuchi H, et al. Effects of alternate and simultaneous administrations of calcium and phosphorus on calcium metabolism in children receiving total parenteral nutrition. *J Parenter Enteral Nutr* 1986 Sep-Oct;10(5):513–6.
10. Hak EB, Crill CM, Bugnitz MC, Mouser JF, Chesney RW. Increased parathyroid hormone and decreased calcitriol during neonatal extracorporeal membrane oxygenation. *Intensive Care Med* 2005 Feb;31(2):264–70. [PubMed: 15685444]
11. Damiani D, Aguiar CH, Bueno VS, Montenegro FL, Koch VH, Cocozza AM, et al. Primary hyperparathyroidism in children: patient report and review of the literature. *J Pediatr Endocrinol Metab* 1998 Jan-Feb;11(1):83–6. [PubMed: 9642634]
12. Cole D, Forsythe CR, Dooley JM, Grantmyre EB, Salisbury SR. Primary neonatal hyperparathyroidism: a devastating neurodevelopmental disorder if left untreated. *J Craniofac Genet Dev Biol* 1990;10(2):205–14. [PubMed: 2211966]
13. Demirel N, Aydin M, Zenciroglu A, Okumus N, Cetinkaya S, Yildiz YT, et al. Hyperparathyroidism secondary to maternal hypoparathyroidism and vitamin D deficiency: an uncommon cause of neonatal respiratory distress. *Ann Trop Paediatr* 2009 Jun;29(2):149–54. [PubMed: 19460269]
14. Glass EJ, Barr DG. Transient neonatal hyperparathyroidism secondary to maternal pseudohypoparathyroidism. *Arch Dis Child* 1981 Jul;56(7):565–8. [PubMed: 7271292]
15. Savani RC, Mimouni F, Tsang RC. Maternal and neonatal hyperparathyroidism as a consequence of maternal renal tubular acidosis. *Pediatrics* 1993 Mar;91(3):661–3. [PubMed: 8441579]
16. Loughhead JL, Mughal Z, Mimouni F, Tsang RC, Oestreich AE. Spectrum and natural history of congenital hyperparathyroidism secondary to maternal hypocalcemia. *Am J Perinatol* 1990 Oct;7(4):350–5. [PubMed: 2222627]
17. Cole DE, Janicic N, Salisbury SR, Hendy GN. Neonatal severe hyperparathyroidism, secondary hyperparathyroidism, and familial hypocalciuric hypercalcemia: multiple different phenotypes associated with an inactivating Alu insertion mutation of the calcium-sensing receptor gene. *Am J Med Genet* 1997 Aug 8;71(2):202–10. [PubMed: 9217223]
18. Chou YH, Pollak MR, Brandi ML, Toss G, Arnqvist H, Atkinson AB, et al. Mutations in the human Ca(2+)-sensing-receptor gene that cause familial hypocalciuric hypercalcemia. *Am J Hum Genet* 1995 May;56(5):1075–9. [PubMed: 7726161]
19. Pollak MR, Brown EM, Chou YH, Hebert SC, Marx SJ, Steinmann B, et al. Mutations in the human Ca(2+)-sensing receptor gene cause familial hypocalciuric hypercalcemia and neonatal severe

- hyperparathyroidism. *Cell* 1993 Dec 31;75(7):1297–303. [PubMed: 7916660] . **This paper is the landmark paper in which CASR mutations are described.
20. Powell BR, Blank E, Benda G, Buist NR. Neonatal hyperparathyroidism and skeletal demineralization in an infant with familial hypocalciuric hypercalcemia. *Pediatrics* 1993 Jan;91(1):144–5. [PubMed: 8416480]
 21. Pearce SH, Trump D, Wooding C, Besser GM, Chew SL, Grant DB, et al. Calcium-sensing receptor mutations in familial benign hypercalcemia and neonatal hyperparathyroidism. *J Clin Invest* 1995 Dec;96(6):2683–92. [PubMed: 8675635]
 22. Bai M, Pearce SH, Kifor O, Trivedi S, Stauffer UG, Thakker RV, et al. In vivo and in vitro characterization of neonatal hyperparathyroidism resulting from a de novo, heterozygous mutation in the Ca²⁺-sensing receptor gene: normal maternal calcium homeostasis as a cause of secondary hyperparathyroidism in familial benign hypocalciuric hypercalcemia. *J Clin Invest* 1997 Jan 1;99(1): 88–96. [PubMed: 9011580]
 23. Obermannova B, Banghova K, Sumnik Z, Dvorakova HM, Betka J, Fencel F, et al. Unusually severe phenotype of neonatal primary hyperparathyroidism due to a heterozygous inactivating mutation in the CASR gene. *Eur J Pediatr* 2009 May;168(5):569–73. [PubMed: 18751724] . *A severe form of neonatal primary hyperparathyroidism is described here.
 24. Chang W, Tu C, Chen TH, Bikle D, Shoback D. The extracellular calcium-sensing receptor (CaSR) is a critical modulator of skeletal development. *Sci Signal* 2008;1(35):1–13. [PubMed: 18364512] . *A review of the calcium-sensing receptor is given here.
 25. Waller S, Kurzwinski T, Spitz L, Thakker R, Cranston T, Pearce S, et al. Neonatal severe hyperparathyroidism: genotype/phenotype correlation and the use of pamidronate as rescue therapy. *Eur J Pediatr* 2004 Oct;163(10):589–94. [PubMed: 15241688]
 26. Peacock M, Bolognese MA, Borofsky M, Scumpia S, Sterling LR, Cheng S, et al. Cinacalcet treatment of primary hyperparathyroidism: biochemical and bone densitometric outcomes in a five-year study. *J Clin Endocrinol Metab* 2009 Dec;94(12):4860–7. [PubMed: 19837909]
 27. Henrich LM, Rogol AD, D'Amour P, Levine MA, Hanks JB, Bruns DE. Persistent hypercalcemia after parathyroidectomy in an adolescent and effect of treatment with cinacalcet HCl. *Clin Chem* 2006 Dec;52(12):2286–93. [PubMed: 17105782]
 28. Lu JY, Yang Y, Gnacadja G, Christopoulos A, Reagan JD. Effect of the calcimimetic R-568 [3-(2-chlorophenyl)-N-((1R)-1-(3-methoxyphenyl)ethyl)-1-propanamine] on correcting inactivating mutations in the human calcium-sensing receptor. *J Pharmacol Exp Ther* 2009 Dec;331(3):775–86. [PubMed: 19759318]
 29. Rus R, Haag C, Bumke-Vogt C, Bahr V, Mayr B, Mohlig M, et al. Novel inactivating mutations of the calcium-sensing receptor: the calcimimetic NPS R-568 improves signal transduction of mutant receptors. *J Clin Endocrinol Metab* 2008 Dec;93(12):4797–803. [PubMed: 18796518]
 30. Timmers HJ, Karperien M, Hamdy NA, de Boer H, Hermus AR. Normalization of serum calcium by cinacalcet in a patient with hypercalcaemia due to a de novo inactivating mutation of the calcium-sensing receptor. *J Intern Med* 2006 Aug;260(2):177–82. [PubMed: 16882283]
 31. Lietman SA, Tenenbaum-Rakover Y, Jap TS, Yi-Chi W, De-Ming Y, Ding C, et al. A novel loss-of-function mutation, Gln459Arg, of the calcium-sensing receptor gene associated with apparent autosomal recessive inheritance of familial hypocalciuric hypercalcemia. *J Clin Endocrinol Metab* 2009 Nov;94(11):4372–9. [PubMed: 19789209] . *This is the first paper describing autosomal recessive inheritance of familial hypocalciuric hypercalcemia
 32. Nesbit MA, Hannan FM, Graham U, Whyte MP, Morrison PJ, Hunter SJ, et al. Identification of a Second Kindred with Familial Hypocalciuric Hypercalcemia Type 3 (FHH3) Narrows Localization to a <3.5 Megabase Pair Region on Chromosome 19q13.3. *J Clin Endocrinol Metab*. Feb 4;
 33. Lloyd SE, Pannett AA, Dixon PH, Whyte MP, Thakker RV. Localization of familial benign hypercalcemia, Oklahoma variant (FBHOk), to chromosome 19q13. *Am J Hum Genet* 1999 Jan;64(1):189–95. [PubMed: 9915958]
 34. Heath H 3rd, Jackson CE, Otterud B, Leppert MF. Genetic linkage analysis in familial benign (hypocalciuric) hypercalcemia: evidence for locus heterogeneity. *Am J Hum Genet* 1993 Jul;53(1): 193–200. [PubMed: 8317484]

35. Burden AD, Krafchik BR. Subcutaneous fat necrosis of the newborn: a review of 11 cases. *Pediatr Dermatol* 1999 Sep-Oct;16(5):384–7. [PubMed: 10571840]
36. Finne PH, Sanderud J, Aksnes L, Bratlid D, Aarskog D. Hypercalcemia with increased and unregulated 1,25-dihydroxyvitamin D production in a neonate with subcutaneous fat necrosis. *J Pediatr* 1988 May;112(5):792–4. [PubMed: 3361394]
37. Hicks MJ, Levy ML, Alexander J, Flaitz CM. Subcutaneous fat necrosis of the newborn and hypercalcemia: case report and review of the literature. *Pediatr Dermatol* 1993 Sep;10(3):271–6. [PubMed: 8415307]
38. Cagle AP, Waguespack SG, Buckingham BA, Shankar RR, Dimeglio LA. Severe infantile hypercalcemia associated with Williams syndrome successfully treated with intravenously administered pamidronate. *Pediatrics* 2004 Oct;114(4):1091–5. [PubMed: 15466114]
39. Kato S, Fujiki R, Kim MS, Kitagawa H. Ligand-induced transrepressive function of VDR requires a chromatin remodeling complex, WINAC. *J Steroid Biochem Mol Biol* 2007 Mar;103(3-5):372–80. [PubMed: 17368181]
40. Kitagawa H, Fujiki R, Yoshimura K, Mezaki Y, Uematsu Y, Matsui D, et al. The chromatin-remodeling complex WINAC targets a nuclear receptor to promoters and is impaired in Williams syndrome. *Cell* 2003 Jun 27;113(7):905–17. [PubMed: 12837248]
41. Fedde KN, Blair L, Silverstein J, Coburn SP, Ryan LM, Weinstein RS, et al. Alkaline phosphatase knock-out mice recapitulate the metabolic and skeletal defects of infantile hypophosphatasia. *J Bone Miner Res* 1999 Dec;14(12):2015–26. [PubMed: 10620060]
42. Mochizuki H, Saito M, Michigami T, Ohashi H, Koda N, Yamaguchi S, et al. Severe hypercalcaemia and respiratory insufficiency associated with infantile hypophosphatasia caused by two novel mutations of the tissue-nonspecific alkaline phosphatase gene. *Eur J Pediatr* 2000 May;159(5):375–9. [PubMed: 10834525]
43. Whyte MP, Magill HL, Fallon MD, Herrod HG. Infantile hypophosphatasia: normalization of circulating bone alkaline phosphatase activity followed by skeletal remineralization. Evidence for an intact structural gene for tissue nonspecific alkaline phosphatase. *J Pediatr* 1986 Jan;108(1):82–8. [PubMed: 3944698]
44. Teree TM, Klein LR. Hypophosphatasia: clinical and metabolic studies. *J Pediatr* 1968 Jan;72(1):41–50. [PubMed: 4294575]
45. Tadokoro M, Kanai R, Taketani T, Uchio Y, Yamaguchi S, Ohgushi H. New bone formation by allogeneic mesenchymal stem cell transplantation in a patient with perinatal hypophosphatasia. *J Pediatr* 2009 Jun;154(6):924–30. [PubMed: 19446101]
46. Millan JL, Narisawa S, Lemire I, Loisel TP, Boileau G, Leonard P, et al. Enzyme replacement therapy for murine hypophosphatasia. *J Bone Miner Res* 2008 Jun;23(6):777–87. [PubMed: 18086009]
47. Drummond KN, Michael AF, Ulstrom RA, Good RA. The Blue Diaper Syndrome: Familial Hypercalcemia with Nephrocalcinosis and Indicanuria; a New Familial Disease, with Definition of the Metabolic Abnormality. *Am J Med* 1964 Dec;37:928–48. [PubMed: 14246093]
48. Saarela T, Simila S, Koivisto M. Hypercalcemia and nephrocalcinosis in patients with congenital lactase deficiency. *J Pediatr* 1995 Dec;127(6):920–3. [PubMed: 8523189]
49. Belmont JW, Reid B, Taylor W, Baker SS, Moore WH, Morriss MC, et al. Congenital sucrase-isomaltase deficiency presenting with failure to thrive, hypercalcemia, and nephrocalcinosis. *BMC Pediatr* 2002 Apr 25;2:4. [PubMed: 12014995]
50. Shoemaker L, Welch TR, Bergstrom W, Abrams SA, Yergey AL, Vieira N. Calcium kinetics in the hyperprostaglandin E syndrome. *Pediatr Res* 1993 Jan 1;33:92–6. [PubMed: 8433868]
51. Amirlak I, Dawson KP. Bartter syndrome: an overview. *QJM* 2000 Apr;93(4):207–15. [PubMed: 10787448]
52. Bettinelli A, Ciarmatori S, Cesareo L, Tedeschi S, Ruffa G, Appiani AC, et al. Phenotypic variability in Bartter syndrome type I. *Pediatr Nephrol* 2000 Sep;14(10-11):940–5. [PubMed: 10975303]
53. Vilain E, Le Merrer M, Lecointre C, Desangles F, Kay MA, Maroteaux P, et al. IMaGE, a new clinical association of intrauterine growth retardation, metaphyseal dysplasia, adrenal hypoplasia congenita, and genital anomalies. *J Clin Endocrinol Metab* 1999 Dec;84(12):4335–40. [PubMed: 10599684]

54. Kelly, A.; Levine, MA. Disorders of calcium, phosphate, parathyroid hormone and vitamin D. In: Kappy, MS.; Allen, DB.; Geffner, ME., editors. *Pediatric Practice: Endocrinology*. Springfield: Charles C Thomas Publisher, LTD; 2009. p. 191-256.
55. Kollars J, Zarroug AE, van Heerden J, Lteif A, Stavlo P, Suarez L, et al. Primary hyperparathyroidism in pediatric patients. *Pediatrics* 2005 Apr;115(4):974–80. [PubMed: 15805373]
56. Stalberg P, Carling T. Familial parathyroid tumors: diagnosis and management. *World J Surg* 2009 Nov;33(11):2234–43. [PubMed: 19184636]
57. Carpten JD, Robbins CM, Villablanca A, Forsberg L, Presciuttini S, Bailey-Wilson J, et al. HRPT2, encoding parafibromin, is mutated in hyperparathyroidism-jaw tumor syndrome. *Nat Genet* 2002 Dec;32(4):676–80. [PubMed: 12434154]
58. Shattuck TM, Valimaki S, Obara T, Gaz RD, Clark OH, Shoback D, et al. Somatic and germ-line mutations of the HRPT2 gene in sporadic parathyroid carcinoma. *N Engl J Med* 2003 Oct 30;349(18):1722–9. [PubMed: 14585940]
59. Bosch X. Hypercalcemia due to endogenous overproduction of active vitamin D in identical twins with cat-scratch disease. *JAMA* 1998 Feb 18;279(7):532–4. [PubMed: 9480364]
60. Monkawa T, Yoshida T, Hayashi M, Saruta T. Identification of 25-hydroxyvitamin D3 1alpha-hydroxylase gene expression in macrophages. *Kidney Int* 2000 Aug;58(2):559–68. [PubMed: 10916079]
61. Stewart AF, Adler M, Byers CM, Segre GV, Broadus AE. Calcium homeostasis in immobilization: an example of resorptive hypercalciuria. *N Engl J Med* 1982 May 13;306(19):1136–40. [PubMed: 6280047]
62. Vahntsevanos K, Kyrgidis A, Verrou E, Katodritou E, Triaridis S, Andreadis CG, et al. Longitudinal cohort study of risk factors in cancer patients of bisphosphonate-related osteonecrosis of the jaw. *J Clin Oncol* 2009 Nov 10;27(32):5356–62. [PubMed: 19805682]
63. Shoemaker LR. Expanding role of bisphosphonate therapy in children. *J Pediatr* 1999 Mar;134(3):264–7. [PubMed: 10064659]
64. Srivastava T, Alon US. Bisphosphonates: from grandparents to grandchildren. *Clin Pediatr (Phila)* 1999 Dec;38(12):687–702. [PubMed: 10618761]
65. Attard TM, Dhawan A, Kaufman SS, Collier DS, Langnas AN. Use of disodium pamidronate in children with hypercalcemia awaiting liver transplantation. *Pediatr Transplant* 1998 May;2(2):157–9. [PubMed: 10082449]
66. Kedlaya D, Brandstater ME, Lee JK. Immobilization hypercalcemia in incomplete paraplegia: successful treatment with pamidronate. *Arch Phys Med Rehabil* 1998 Feb;79(2):222–5. [PubMed: 9474008]
67. Platt C, Inward C, McGraw M, Dudley J, Tizard J, Burren C, et al. Middle-term use of Cinacalcet in paediatric dialysis patients. *Pediatr Nephrol* 2010 Jan;25(1):143–8. [PubMed: 19838738] . *An early report on the use of the calcimimetic




Figure 1. Hypercalcemia algorithm (54)

FeCa = Fractional (urinary) excretion of calcium

Adapted from: Kelly A, Levine MA. Disorders of calcium, phosphate, parathyroid hormone and vitamin D. In: Kappy MS, Allen DB, Geffner ME editor. Pediatric Practice: Endocrinology. Springfield: Charles C. Thomas Publisher, LTD; 2009. p. 191-256.

Table 1

Representative normal values for age for concentrations of serum total calcium

	Age (yr)	Serum total calcium (mg/dl)
Infants	0-0.25	8.8-11.3
	1-5	9.4-10.8
Children	6-12	9.4-10.3
Men	20	9.1-10.2
	50	8.9-10.0
	70	8.8-9.9
Women	20	8.8-10.0
	50	8.8-10.0
	70	8.8-10.0

Revised from reference (3).

Table 2

Differential diagnosis of hypercalcemia in neonates and infants (up to 2 years of age)

•	Iatrogenic
•	Phosphate Depletion
•	Premature infants on human milk or standard formula
•	Parenteral nutrition
•	Hyperparathyroidism
–	Congenital parathyroid hyperplasia
–	Maternal hypoparathyroidism
•	Inactivating mutations in Ca ²⁺ -sensing receptor gene
–	Familial hypocalciuric hypercalcemia (familial benign hypercalcemia)
–	Neonatal severe hyperparathyroidism
•	Jansen's metaphyseal chondrodysplasia
•	Persistent PTHrP
•	Hypervitaminosis D
•	Subcutaneous fat necrosis
•	Williams syndrome/Idiopathic infantile hypercalcemia
•	Other inborn metabolic disorders
–	Blue diaper syndrome
–	Lactase deficiency
–	Disaccharide intolerance
–	Bartter syndrome
•	Hypophosphatasia
•	IMAGe
•	Down syndrome
•	Severe congenital hypothyroidism
•	Maternal hypercalcemia
•	Vitamin A intoxication

Table 3

Laboratory values in differential diagnosis of hypercalcemia.

	Serum calcium	Serum phosphorus	Fractional excretion of calcium	PTH	PTHrP	25(OH)D	1,25(OH)2D
Familial hypocalciuric hypercalcemia	↑	N or ↓	<0.01	N or ↑	↓	N	↑
NSHPT	↑↑↑	↓	<0.01	↑↑	↓	N or ↓	↑
Subcutaneous fat necrosis	↑	↑	↑	↓	↓	N	↑↑
Williams syndrome	↑	↑	↑	↓	↓	N	N or ↑
Primary hyperparathyroidism	↑	↓	>0.01	↑	↓	N	↑
Humoral Malignancy	↑↑	↓	↑	↓	↑↑	N	N or ↑
Osteolytic malignancy	↑↑	↑	↑↑	↓	↓	N	↓
Granulomatous disease	↑	↑	↑↑	↓	↓	N	↑↑
Vitamin D intoxication	↑	↑	↑↑	↓	↓	↑↑	N or ↓
Immobilization	↑	↑	↑↑	↓	↓	N	↓

Table 4

Hypercalcemia in children (over 2 years of age) and adolescents

-
- Excessive calcium intake
 - Phosphate depletion
 - Parenteral nutrition
 - Hyperparathyroidism
 - Acquired primary
 - ◆ adenoma
 - ◆ multi-glandular
 - ◆ carcinoma
 - Genetic primary
 - ◆ autosomal dominant/récessive FHH
 - ◆ familial
 - ◆ MEN types I and IIa (IIb)
 - ◆ Hyperparathyroidism-Jaw tumor
 - Autonomous (tertiary)
 - Hypervitaminosis D
 - Excessive intake
 - Granulomatous diseases: cat scratch fever; sarcoidosis; tuberculosis; histoplasmosis; coccidiomycosis; leprosy; HIV
 - Chronic inflammatory disorders
 - Williams syndrome/ Idiopathic infantile hypercalcemia
 - Immobilization
 - Malignancy associated hypercalcemia
 - Primary bone tumors
 - Metastatic tumors with osteolysis
 - Tumors secreting PTHrP, prostaglandins, cytokines, growth factors
 - Hepatic disease
 - Hyperthyroidism
 - Adrenal insufficiency
 - Pheochromocytoma
 - Vasoactive intestinal polypeptide-secreting tumor
 - Drugs (thiazides, lithium, systemic retinoid derivatives, theophylline, acetosalicylic acid)
 - Milk alkali syndrome/ calcium gluttony
 - Renal tubular acidosis
-