

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

2-(5-Bromo-2-hydroxyphenyl)-1,2-dihydroquinazolin-4(3H)-one

Davar M. Boghaei,^{a*} Mohammad Mahdi Najafpour^a and Vickie McKee^b^aDepartment of Chemistry, Sharif University of Technology, PO Box 11155-8639, Tehran, Iran, and^bDepartment of Chemistry, Loughborough University, Leicestershire LE11 3TU, England

Correspondence e-mail: dboghaei@sharif.edu


Received 8 October 2008; accepted 30 October 2008

Key indicators: single-crystal X-ray study; $T = 150$ K; mean $\sigma(\text{C}-\text{C}) = 0.004$ Å; R factor = 0.040; wR factor = 0.099; data-to-parameter ratio = 17.9.

The asymmetric unit of the title compound, $\text{C}_{14}\text{H}_{11}\text{BrN}_2\text{O}_2$, contains two independent molecules connected into a dimer by intermolecular $\text{N}-\text{H}\cdots\text{O}$ hydrogen bonds involving the amine and carbonyl groups. The dimers are further connected by $\text{O}-\text{H}\cdots\text{O}$ hydrogen bonds, forming chains running parallel to the a axis, which are stabilized through $\pi-\pi$ stacking interactions, with a centroid-centroid distance of 3.679 (8) Å. The dihedral angle between the two aromatic rings is 89.2 (4)°.

Related literature

For general background to the chemistry of quinazolinone derivatives, see: Liu (2008); Goto *et al.* (1993); Mohri (2001). For hydrogen-bond motifs, see: Bernstein *et al.* (1995).



Experimental

Crystal data

 $\text{C}_{14}\text{H}_{11}\text{BrN}_2\text{O}_2$ $M_r = 319.16$ Triclinic, $P\bar{1}$ $a = 8.8392$ (5) Å $b = 11.2252$ (7) Å $c = 13.8817$ (8) Å $\alpha = 73.0392$ (9)° $\beta = 75.9620$ (9)° $\gamma = 85.0936$ (9)° $V = 1277.95$ (13) Å³ $Z = 4$ Mo $K\alpha$ radiation $\mu = 3.21$ mm⁻¹ $T = 150$ (2) K $0.21 \times 0.12 \times 0.07$ mm

Data collection

Bruker APEXII CCD diffractometer

Absorption correction: multi-scan

(SADABS; Sheldrick, 2003)

 $T_{\min} = 0.552$, $T_{\max} = 0.806$

13015 measured reflections

6129 independent reflections

4580 reflections with $I > 2\sigma(I)$ $R_{\text{int}} = 0.035$

Refinement

 $R[F^2 > 2\sigma(F^2)] = 0.040$ $wR(F^2) = 0.099$ $S = 1.02$

6129 reflections

343 parameters

H-atom parameters constrained

 $\Delta\rho_{\text{max}} = 0.64$ e Å⁻³ $\Delta\rho_{\text{min}} = -0.70$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$\text{N2A}-\text{H2NA}\cdots\text{O1B}$	0.95	1.97	2.897 (3)	165
$\text{N2B}-\text{H2NB}\cdots\text{O1A}$	0.91	2.05	2.914 (3)	157
$\text{O2A}-\text{H2OA}\cdots\text{O1A}^i$	0.85	1.90	2.701 (3)	157
$\text{O2B}-\text{H2OB}\cdots\text{O1B}^{ii}$	0.85	1.86	2.691 (3)	165

Symmetry codes: (i) $x - 1, y, z$; (ii) $x + 1, y, z$.

Data collection: APEX2 (Bruker, 2005); cell refinement: SAINT (Bruker, 2005); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: SHELXTL (Sheldrick, 2008); software used to prepare material for publication: SHELXTL.

We are grateful to the Research Council of Sharif University of Technology and Loughborough University for their financial support.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: RZ2254).

References

- Bernstein, J., Davis, R. E., Shimoni, L. & Chang, N.-L. (1995). *Angew. Chem. Int. Ed. Engl.* **34**, 1555-1573.
- Bruker (2005). APEX2 and SAINT. Bruker AXS Inc., Madison, Wisconsin, USA.
- Goto, S., Tsuboi, H. & Kagara, K. (1993). *Chem. Express*, **8**, 761-764.
- Liu, G. (2008). *Acta Cryst.* **E64**, o1677.
- Mohri, S. J. (2001). *Synth. Org. Chem. Jpn.* **59**, 514-515.
- Sheldrick, G. M. (2003). SADABS. Bruker AXS Inc., Madison, Wisconsin, USA.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112-122.

supplementary materials

Acta Cryst. (2009). E65, o193 [doi:10.1107/S1600536808035678]

2-(5-Bromo-2-hydroxyphenyl)-1,2-dihydroquinazolin-4(3H)-one

D. M. Boghaei, M. M. Najafpour and V. McKee

Comment

Quinazolinone derivatives are of interest because of their biological activity, and have been widely used as key compounds in medicinal drugs (Goto *et al.*, 1993; Mohri, 2001). We herein report the crystal structure of 1,2-dihydro-2-(5-bromo-2-hydroxybenzene)-4(3H)-quinazolinone.

The asymmetric unit of the title compound (Fig. 1) contains two crystallographically independent molecules, which are linked into a dimer by a pair of intermolecular N—H \cdots O hydrogen bonds (Table 1), generating a ring of graph set $R^2_2(8)$ (Bernstein *et al.*, 1995). Bond lengths and angles are within normal ranges. The dimers are further connected by O—H \cdots O hydrogen bonding interactions to form chains running parallel to the *a* axis (Figures 2). The chains are stabilized by π – π stacking interactions involving adjacent 5-bromo-2-hydroxybenzene rings, with a centroid-centroid separation of 3.679 (8) Å, a perpendicular interplanar distance of 3.561 (8) Å and a centroid \cdots centroid offset of 0.924 (6) Å.


Experimental


The title compound was synthesized by adding 5-bromo-2-hydroxybenzaldehyde (2 mmol, 402 mg) to a solution of 2-aminobenzamide (2 mmol, 272 mg) and manganese acetate (0.02 mmol, 4.90 mg) in ethanol (20 ml). The mixture was refluxed with stirring for 5 h. The resultant yellow solution was filtered. Yellow single crystals of the title compound suitable for *X*-ray structure determination were recrystallized from a mixture of water/ethanol (2:1 v/v) by slow evaporation of the solvents at room temperature over several days.

Refinement

All H atoms were placed in calculated positions and refined using the riding model approximation, with C—H = 0.95–1.0 Å, O—H = 0.85 Å, N—H = 0.89 Å and with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$. The isotropic thermal parameter of the hydrogen atoms bound to the N and O atoms was fixed at 0.04 Å².

Figures

 Fig. 1. The molecular structure of the title compound with displacement ellipsoids drawn at the 50% probability level. Hydrogen bonds are shown as dashed lines.

 Fig. 2. Crystal packing of the title compound viewed along the *a* axis with hydrogen atoms omitted for clarity. Hydrogen bonds are shown as dashed lines.

2-(5-Bromo-2-hydroxyphenyl)-1,2-dihydroquinazolin-4(3H)-one

Crystal data

$C_{14}H_{11}BrN_2O_2$	$Z = 4$
$M_r = 319.16$	$F_{000} = 640$
Triclinic, $P\bar{1}$	$D_x = 1.659 \text{ Mg m}^{-3}$
Hall symbol: -P 1	Mo $K\alpha$ radiation
$a = 8.8392 (5) \text{ \AA}$	$\lambda = 0.71073 \text{ \AA}$
$b = 11.2252 (7) \text{ \AA}$	Cell parameters from 3058 reflections
$c = 13.8817 (8) \text{ \AA}$	$\theta = 2.4\text{--}25.8^\circ$
$\alpha = 73.0392 (9)^\circ$	$\mu = 3.22 \text{ mm}^{-1}$
$\beta = 75.9620 (9)^\circ$	$T = 150 (2) \text{ K}$
$\gamma = 85.0936 (9)^\circ$	Block, yellow
$V = 1277.95 (13) \text{ \AA}^3$	$0.21 \times 0.12 \times 0.07 \text{ mm}$

Data collection

Bruker APEXII CCD diffractometer	6129 independent reflections
Radiation source: fine-focus sealed tube	4580 reflections with $I > 2\sigma(I)$
Monochromator: graphite	$R_{\text{int}} = 0.035$
$T = 150(2) \text{ K}$	$\theta_{\text{max}} = 28.0^\circ$
φ and ω scans	$\theta_{\text{min}} = 1.6^\circ$
Absorption correction: multi-scan (SADABS; Sheldrick, 2003)	$h = -11 \rightarrow 11$
$T_{\text{min}} = 0.552, T_{\text{max}} = 0.806$	$k = -14 \rightarrow 14$
13015 measured reflections	$l = -18 \rightarrow 18$

Refinement

Refinement on F^2	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.040$	H-atom parameters constrained
$wR(F^2) = 0.099$	$w = 1/[\sigma^2(F_o^2) + (0.0415P)^2 + 0.6583P]$
$S = 1.03$	where $P = (F_o^2 + 2F_c^2)/3$
6129 reflections	$(\Delta/\sigma)_{\text{max}} = 0.001$
343 parameters	$\Delta\rho_{\text{max}} = 0.64 \text{ e \AA}^{-3}$
Primary atom site location: structure-invariant direct methods	$\Delta\rho_{\text{min}} = -0.70 \text{ e \AA}^{-3}$
	Extinction correction: none

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against all reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on all data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
N1A	0.0154 (3)	-0.1619 (2)	0.82388 (19)	0.0233 (5)
C1A	0.1438 (3)	-0.2154 (3)	0.7715 (2)	0.0226 (6)
C2A	0.1414 (4)	-0.3350 (3)	0.7594 (2)	0.0274 (7)
H2A	0.0487	-0.3822	0.7874	0.033*
C3A	0.2724 (4)	-0.3839 (3)	0.7072 (2)	0.0308 (7)
H3A	0.2693	-0.4652	0.7000	0.037*
C4A	0.4098 (4)	-0.3167 (3)	0.6646 (2)	0.0301 (7)
H4A	0.4981	-0.3506	0.6262	0.036*
C5A	0.4164 (3)	-0.2011 (3)	0.6785 (2)	0.0260 (7)
H5A	0.5103	-0.1554	0.6508	0.031*
C6A	0.2848 (3)	-0.1506 (3)	0.7335 (2)	0.0221 (6)
C7A	0.2963 (3)	-0.0355 (3)	0.7617 (2)	0.0219 (6)
O1A	0.4227 (2)	0.0103 (2)	0.75507 (16)	0.0260 (5)
N2A	0.1608 (3)	0.0141 (2)	0.80201 (19)	0.0227 (5)
C8A	0.0102 (3)	-0.0267 (3)	0.7988 (2)	0.0224 (6)
H8A	-0.0707	-0.0030	0.8545	0.027*
C9A	-0.0345 (3)	0.0361 (3)	0.6958 (2)	0.0198 (6)
C10A	0.0667 (3)	0.1068 (3)	0.6088 (2)	0.0222 (6)
H10A	0.1705	0.1192	0.6113	0.027*
C11A	0.0160 (3)	0.1594 (3)	0.5183 (2)	0.0232 (6)
Br1A	0.15542 (4)	0.25729 (3)	0.40004 (2)	0.03233 (10)
C12A	-0.1353 (3)	0.1454 (3)	0.5130 (2)	0.0246 (6)
H12A	-0.1687	0.1827	0.4507	0.029*
C13A	-0.2376 (3)	0.0763 (3)	0.5999 (2)	0.0248 (6)
H13A	-0.3421	0.0664	0.5973	0.030*
C14A	-0.1883 (3)	0.0214 (3)	0.6905 (2)	0.0212 (6)
O2A	-0.2832 (2)	-0.0477 (2)	0.77902 (16)	0.0265 (5)
N1B	0.5589 (3)	0.1904 (2)	1.02282 (18)	0.0235 (5)
C1B	0.4498 (3)	0.2837 (3)	1.0340 (2)	0.0215 (6)
C2B	0.4523 (4)	0.3566 (3)	1.1007 (2)	0.0272 (7)
H2B	0.5320	0.3436	1.1381	0.033*
C3B	0.3386 (4)	0.4471 (3)	1.1116 (2)	0.0297 (7)
H3B	0.3396	0.4949	1.1577	0.036*

supplementary materials

C4B	0.2227 (4)	0.4691 (3)	1.0563 (2)	0.0306 (7)
H4B	0.1478	0.5341	1.0620	0.037*
C5B	0.2169 (3)	0.3961 (3)	0.9929 (2)	0.0266 (7)
H5B	0.1362	0.4095	0.9564	0.032*
C6B	0.3289 (3)	0.3025 (3)	0.9820 (2)	0.0209 (6)
C7B	0.3173 (3)	0.2169 (3)	0.9224 (2)	0.0224 (6)
O1B	0.1980 (2)	0.2107 (2)	0.89065 (17)	0.0307 (5)
N2B	0.4387 (3)	0.1394 (2)	0.90652 (19)	0.0224 (5)
C8B	0.5894 (3)	0.1544 (3)	0.9271 (2)	0.0217 (6)
H8B	0.6427	0.0710	0.9396	0.026*
C9B	0.6953 (3)	0.2433 (3)	0.8361 (2)	0.0203 (6)
C10B	0.6512 (3)	0.3015 (3)	0.7437 (2)	0.0217 (6)
H10B	0.5498	0.2894	0.7368	0.026*
C11B	0.7546 (3)	0.3770 (3)	0.6619 (2)	0.0246 (6)
Br1B	0.69147 (4)	0.45082 (3)	0.53632 (2)	0.03648 (11)
C12B	0.9020 (4)	0.3972 (3)	0.6699 (2)	0.0283 (7)
H12B	0.9714	0.4501	0.6133	0.034*
C13B	0.9479 (3)	0.3395 (3)	0.7613 (2)	0.0258 (6)
H13B	1.0494	0.3527	0.7675	0.031*
C14B	0.8457 (3)	0.2624 (3)	0.8440 (2)	0.0209 (6)
O2B	0.8846 (2)	0.2024 (2)	0.93607 (16)	0.0274 (5)
H1NA	-0.0747	-0.1968	0.8334	0.040*
H1NB	0.6366	0.1868	1.0581	0.040*
H2NA	0.1709	0.0883	0.8199	0.040*
H2NB	0.4285	0.0809	0.8750	0.040*
H2OA	-0.3770	-0.0516	0.7751	0.040*
H2OB	0.9813	0.2141	0.9279	0.040*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
N1A	0.0159 (11)	0.0269 (14)	0.0256 (13)	-0.0034 (10)	-0.0037 (10)	-0.0049 (11)
C1A	0.0204 (14)	0.0291 (16)	0.0183 (14)	0.0018 (12)	-0.0080 (11)	-0.0041 (12)
C2A	0.0309 (17)	0.0272 (17)	0.0253 (16)	-0.0028 (13)	-0.0102 (13)	-0.0054 (13)
C3A	0.0415 (19)	0.0265 (17)	0.0278 (16)	0.0022 (14)	-0.0123 (14)	-0.0098 (14)
C4A	0.0276 (16)	0.0382 (19)	0.0293 (17)	0.0080 (14)	-0.0100 (13)	-0.0163 (15)
C5A	0.0224 (15)	0.0331 (17)	0.0255 (15)	0.0019 (13)	-0.0102 (12)	-0.0094 (13)
C6A	0.0190 (14)	0.0276 (16)	0.0203 (14)	0.0012 (12)	-0.0084 (11)	-0.0049 (12)
C7A	0.0182 (14)	0.0290 (16)	0.0192 (14)	0.0000 (12)	-0.0076 (11)	-0.0052 (12)
O1A	0.0148 (10)	0.0353 (12)	0.0316 (12)	-0.0002 (9)	-0.0082 (9)	-0.0126 (10)
N2A	0.0144 (11)	0.0293 (14)	0.0281 (13)	-0.0006 (10)	-0.0059 (10)	-0.0130 (11)
C8A	0.0134 (13)	0.0323 (17)	0.0236 (15)	-0.0002 (12)	-0.0037 (11)	-0.0115 (13)
C9A	0.0163 (13)	0.0207 (14)	0.0250 (15)	0.0014 (11)	-0.0053 (11)	-0.0105 (12)
C10A	0.0150 (13)	0.0240 (15)	0.0288 (16)	-0.0027 (11)	-0.0044 (11)	-0.0093 (12)
C11A	0.0210 (14)	0.0206 (15)	0.0265 (15)	-0.0008 (11)	-0.0047 (12)	-0.0048 (12)
Br1A	0.02666 (17)	0.03157 (19)	0.03285 (18)	-0.00522 (13)	-0.00377 (13)	-0.00114 (14)
C12A	0.0247 (15)	0.0252 (16)	0.0261 (15)	0.0027 (12)	-0.0103 (12)	-0.0079 (13)
C13A	0.0159 (14)	0.0317 (17)	0.0311 (16)	0.0014 (12)	-0.0090 (12)	-0.0129 (13)

C14A	0.0159 (13)	0.0232 (15)	0.0266 (15)	-0.0002 (11)	-0.0030 (11)	-0.0118 (12)
O2A	0.0134 (10)	0.0361 (12)	0.0281 (11)	-0.0049 (9)	-0.0044 (8)	-0.0048 (9)
N1B	0.0178 (12)	0.0336 (14)	0.0213 (12)	0.0013 (10)	-0.0078 (10)	-0.0085 (11)
C1B	0.0179 (14)	0.0263 (15)	0.0170 (13)	-0.0048 (11)	-0.0011 (11)	-0.0023 (12)
C2B	0.0284 (16)	0.0340 (18)	0.0206 (15)	-0.0084 (13)	-0.0054 (12)	-0.0077 (13)
C3B	0.0310 (17)	0.0319 (18)	0.0263 (16)	-0.0092 (14)	0.0020 (13)	-0.0129 (14)
C4B	0.0252 (16)	0.0282 (17)	0.0370 (18)	-0.0018 (13)	0.0022 (14)	-0.0144 (14)
C5B	0.0189 (14)	0.0298 (17)	0.0309 (17)	-0.0025 (12)	-0.0052 (12)	-0.0082 (13)
C6B	0.0173 (13)	0.0248 (15)	0.0208 (14)	-0.0040 (11)	-0.0032 (11)	-0.0068 (12)
C7B	0.0165 (13)	0.0271 (16)	0.0240 (15)	-0.0051 (11)	-0.0039 (11)	-0.0071 (12)
O1B	0.0157 (10)	0.0404 (13)	0.0443 (13)	-0.0008 (9)	-0.0090 (9)	-0.0226 (11)
N2B	0.0167 (12)	0.0212 (13)	0.0343 (14)	-0.0003 (10)	-0.0080 (10)	-0.0133 (11)
C8B	0.0153 (13)	0.0279 (16)	0.0246 (15)	0.0021 (11)	-0.0078 (11)	-0.0094 (12)
C9B	0.0177 (13)	0.0233 (15)	0.0240 (15)	0.0020 (11)	-0.0070 (11)	-0.0117 (12)
C10B	0.0200 (14)	0.0252 (15)	0.0243 (15)	0.0030 (12)	-0.0069 (12)	-0.0133 (12)
C11B	0.0291 (16)	0.0263 (16)	0.0217 (15)	0.0064 (13)	-0.0096 (12)	-0.0108 (13)
Br1B	0.0462 (2)	0.0396 (2)	0.02454 (17)	0.00589 (16)	-0.01352 (15)	-0.00780 (14)
C12B	0.0254 (16)	0.0275 (17)	0.0284 (16)	-0.0011 (13)	0.0008 (13)	-0.0080 (13)
C13B	0.0167 (14)	0.0287 (16)	0.0337 (17)	0.0006 (12)	-0.0054 (12)	-0.0119 (14)
C14B	0.0186 (14)	0.0220 (15)	0.0247 (15)	0.0044 (11)	-0.0069 (12)	-0.0104 (12)
O2B	0.0153 (10)	0.0376 (13)	0.0288 (11)	0.0000 (9)	-0.0095 (8)	-0.0049 (10)

Geometric parameters (Å, °)

N1A—C1A	1.384 (4)	N1B—C1B	1.379 (4)
N1A—C8A	1.453 (4)	N1B—C8B	1.457 (4)
N1A—H1NA	0.8822	N1B—H1NB	0.9273
C1A—C2A	1.402 (4)	C1B—C6B	1.397 (4)
C1A—C6A	1.405 (4)	C1B—C2B	1.408 (4)
C2A—C3A	1.373 (4)	C2B—C3B	1.380 (4)
C2A—H2A	0.9500	C2B—H2B	0.9500
C3A—C4A	1.393 (5)	C3B—C4B	1.389 (5)
C3A—H3A	0.9500	C3B—H3B	0.9500
C4A—C5A	1.375 (4)	C4B—C5B	1.377 (4)
C4A—H4A	0.9500	C4B—H4B	0.9500
C5A—C6A	1.402 (4)	C5B—C6B	1.396 (4)
C5A—H5A	0.9500	C5B—H5B	0.9500
C6A—C7A	1.474 (4)	C6B—C7B	1.463 (4)
C7A—O1A	1.242 (3)	C7B—O1B	1.252 (3)
C7A—N2A	1.342 (4)	C7B—N2B	1.340 (4)
N2A—C8A	1.459 (3)	N2B—C8B	1.461 (3)
N2A—H2NA	0.9523	N2B—H2NB	0.9095
C8A—C9A	1.528 (4)	C8B—C9B	1.522 (4)
C8A—H8A	1.0000	C8B—H8B	1.0000
C9A—C10A	1.388 (4)	C9B—C10B	1.390 (4)
C9A—C14A	1.405 (4)	C9B—C14B	1.399 (4)
C10A—C11A	1.387 (4)	C10B—C11B	1.382 (4)
C10A—H10A	0.9500	C10B—H10B	0.9500
C11A—C12A	1.381 (4)	C11B—C12B	1.377 (4)

supplementary materials

C11A—Br1A	1.903 (3)	C11B—Br1B	1.894 (3)
C12A—C13A	1.386 (4)	C12B—C13B	1.385 (4)
C12A—H12A	0.9500	C12B—H12B	0.9500
C13A—C14A	1.386 (4)	C13B—C14B	1.391 (4)
C13A—H13A	0.9500	C13B—H13B	0.9500
C14A—O2A	1.368 (3)	C14B—O2B	1.370 (3)
O2A—H2OA	0.8493	O2B—H2OB	0.8511
C1A—N1A—C8A	116.9 (2)	C1B—N1B—C8B	117.7 (2)
C1A—N1A—H1NA	115.4	C1B—N1B—H1NB	113.3
C8A—N1A—H1NA	115.0	C8B—N1B—H1NB	121.5
N1A—C1A—C2A	122.7 (3)	N1B—C1B—C6B	119.3 (3)
N1A—C1A—C6A	118.9 (3)	N1B—C1B—C2B	121.6 (3)
C2A—C1A—C6A	118.3 (3)	C6B—C1B—C2B	119.0 (3)
C3A—C2A—C1A	120.2 (3)	C3B—C2B—C1B	119.8 (3)
C3A—C2A—H2A	119.9	C3B—C2B—H2B	120.1
C1A—C2A—H2A	119.9	C1B—C2B—H2B	120.1
C2A—C3A—C4A	121.3 (3)	C2B—C3B—C4B	121.0 (3)
C2A—C3A—H3A	119.3	C2B—C3B—H3B	119.5
C4A—C3A—H3A	119.3	C4B—C3B—H3B	119.5
C5A—C4A—C3A	119.4 (3)	C5B—C4B—C3B	119.6 (3)
C5A—C4A—H4A	120.3	C5B—C4B—H4B	120.2
C3A—C4A—H4A	120.3	C3B—C4B—H4B	120.2
C4A—C5A—C6A	120.0 (3)	C4B—C5B—C6B	120.5 (3)
C4A—C5A—H5A	120.0	C4B—C5B—H5B	119.8
C6A—C5A—H5A	120.0	C6B—C5B—H5B	119.8
C5A—C6A—C1A	120.5 (3)	C5B—C6B—C1B	120.1 (3)
C5A—C6A—C7A	120.5 (3)	C5B—C6B—C7B	121.3 (3)
C1A—C6A—C7A	118.7 (3)	C1B—C6B—C7B	118.5 (3)
O1A—C7A—N2A	120.9 (3)	O1B—C7B—N2B	120.1 (3)
O1A—C7A—C6A	123.0 (3)	O1B—C7B—C6B	122.8 (3)
N2A—C7A—C6A	115.9 (2)	N2B—C7B—C6B	117.0 (2)
C7A—N2A—C8A	122.3 (2)	C7B—N2B—C8B	122.6 (2)
C7A—N2A—H2NA	114.5	C7B—N2B—H2NB	117.5
C8A—N2A—H2NA	122.0	C8B—N2B—H2NB	119.5
N1A—C8A—N2A	107.8 (2)	N1B—C8B—N2B	107.5 (2)
N1A—C8A—C9A	113.3 (2)	N1B—C8B—C9B	113.8 (2)
N2A—C8A—C9A	112.4 (2)	N2B—C8B—C9B	112.8 (2)
N1A—C8A—H8A	107.7	N1B—C8B—H8B	107.5
N2A—C8A—H8A	107.7	N2B—C8B—H8B	107.5
C9A—C8A—H8A	107.7	C9B—C8B—H8B	107.5
C10A—C9A—C14A	118.8 (3)	C10B—C9B—C14B	118.7 (3)
C10A—C9A—C8A	124.0 (2)	C10B—C9B—C8B	122.6 (2)
C14A—C9A—C8A	117.2 (2)	C14B—C9B—C8B	118.6 (2)
C11A—C10A—C9A	119.9 (3)	C11B—C10B—C9B	120.2 (3)
C11A—C10A—H10A	120.0	C11B—C10B—H10B	119.9
C9A—C10A—H10A	120.0	C9B—C10B—H10B	119.9
C12A—C11A—C10A	121.4 (3)	C12B—C11B—C10B	121.2 (3)
C12A—C11A—Br1A	119.1 (2)	C12B—C11B—Br1B	120.0 (2)
C10A—C11A—Br1A	119.4 (2)	C10B—C11B—Br1B	118.7 (2)

C11A—C12A—C13A	119.1 (3)	C11B—C12B—C13B	119.3 (3)
C11A—C12A—H12A	120.5	C11B—C12B—H12B	120.4
C13A—C12A—H12A	120.5	C13B—C12B—H12B	120.4
C14A—C13A—C12A	120.3 (3)	C12B—C13B—C14B	120.2 (3)
C14A—C13A—H13A	119.8	C12B—C13B—H13B	119.9
C12A—C13A—H13A	119.8	C14B—C13B—H13B	119.9
O2A—C14A—C13A	123.3 (2)	O2B—C14B—C13B	122.9 (3)
O2A—C14A—C9A	116.2 (3)	O2B—C14B—C9B	116.7 (3)
C13A—C14A—C9A	120.5 (3)	C13B—C14B—C9B	120.4 (3)
C14A—O2A—H2OA	114.4	C14B—O2B—H2OB	107.1

Hydrogen-bond geometry (\AA , $^\circ$)

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
N2A—H2NA \cdots O1B	0.95	1.97	2.897 (3)	165.2
N2B—H2NB \cdots O1A	0.91	2.05	2.914 (3)	157.3
O2A—H2OA \cdots O1A ⁱ	0.85	1.90	2.701 (3)	156.7
O2B—H2OB \cdots O1B ⁱⁱ	0.85	1.86	2.691 (3)	165.4

Symmetry codes: (i) $x-1, y, z$; (ii) $x+1, y, z$.

Fig. 1




Fig. 2

