

(E)-3-(Anthracen-9-yl)-1-(4-bromophenyl)-prop-2-en-1-one¹

Thitipone Suwunwong,^a Suchada Chantrapromma,^{b*} Chatchanok Karalai,^b Paradorn Pakdeevanich^c and Hoong-Kun Fun^{d§}

^aDepartment of Chemistry and Center of Excellence for Innovation in Chemistry, Faculty of Science, Prince of Songkla University, Hat-Yai, Songkhla 90112, Thailand,

^bCrystal Materials Research Unit, Department of Chemistry, Faculty of Science, Prince of Songkla University, Hat-Yai, Songkhla 90112, Thailand, ^cDepartment of Physics, Faculty of Science, Prince of Songkla University, Hat-Yai, Songkhla 90112, Thailand, and ^dX-ray Crystallography Unit, School of Physics, Universiti Sains Malaysia, 11800 USM, Penang, Malaysia

Correspondence e-mail: suchada.c@psu.ac.th


Received 31 December 2008; accepted 26 January 2009

Key indicators: single-crystal X-ray study; $T = 100$ K; mean $\sigma(C-C) = 0.003 \text{ \AA}$; R factor = 0.032; wR factor = 0.076; data-to-parameter ratio = 21.5.

In the title molecule, $C_{23}H_{15}BrO$, the prop-2-en-1-one unit is planar and it makes dihedral angles of 20.9 (1) and 45.8 (1) $^\circ$, respectively, with the 4-bromophenyl ring and the anthracene ring system. The interplanar angle between the 4-bromophenyl ring and the anthracene ring system is 35.52 (7) $^\circ$. In the crystal structure, molecules are linked into dimers by C—H \cdots Br hydrogen bonds, and the dimers are linked into a zigzag network parallel to the bc plane by weak C—H \cdots O hydrogen bonds and C—H \cdots π interactions involving the central benzene ring of the anthracene ring system.

Related literature

For bond-length data, see: Allen *et al.* (1987). For related structures, see: Ng *et al.* (2006); Patil *et al.* (2006); Patil, Chantrapromma *et al.* (2007); Suwunwong *et al.* (2009). For background and applications of chalcones, see: Jung *et al.* (2008); Patil, Chantrapromma *et al.* (2007); Patil, Dharmaprabak *et al.* (2007); Patil & Dharmaprabak (2008); Prasad *et al.* (2008).



Experimental

Crystal data

$C_{23}H_{15}BrO$	$V = 1666.02 (6) \text{ \AA}^3$
$M_r = 387.25$	$Z = 4$
Monoclinic, $P2_1/c$	Mo $K\alpha$ radiation
$a = 5.3792 (1) \text{ \AA}$	$\mu = 2.47 \text{ mm}^{-1}$
$b = 19.1030 (4) \text{ \AA}$	$T = 100.0 (1) \text{ K}$
$c = 16.3005 (4) \text{ \AA}$	$0.57 \times 0.27 \times 0.15 \text{ mm}$
$\beta = 95.944 (1)^\circ$	

Data collection

Bruker SMART APEXII CCD area-detector diffractometer	29994 measured reflections
Absorption correction: multi-scan (<i>SADABS</i> ; Bruker, 2005)	4866 independent reflections
$T_{min} = 0.331$, $T_{max} = 0.714$	3803 reflections with $I > 2\sigma(I)$
	$R_{int} = 0.036$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.032$	226 parameters
$wR(F^2) = 0.076$	H-atom parameters constrained
$S = 1.02$	$\Delta\rho_{\max} = 0.53 \text{ e \AA}^{-3}$
4866 reflections	$\Delta\rho_{\min} = -0.54 \text{ e \AA}^{-3}$

Table 1

Hydrogen-bond geometry (\AA , $^\circ$).

$Cg1$ is the centroid of the C18–C23 ring.

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$C8-H8A\cdots O1^i$	0.93	2.42	3.308 (2)	159
$C13-H13A\cdots O1^{ii}$	0.93	2.57	3.288 (2)	135
$C21-H21A\cdots Br1^{iii}$	0.93	2.93	3.4722 (19)	119
$C9-H9A\cdots Cg1^{iv}$	0.93	2.83	3.4479 (18)	125

Symmetry codes: (i) $x - 1, y, z$; (ii) $-x + 2, -y + 1, -z + 2$; (iii) $-x, -y + 1, -z + 1$; (iv) $x + 1, y, z$.

Data collection: *APEX2* (Bruker, 2005); cell refinement: *APEX2*; data reduction: *SAINT* (Bruker, 2005); program(s) used to solve structure: *SHELXTL* (Sheldrick, 2008); program(s) used to refine structure: *SHELXTL*; molecular graphics: *SHELXTL*; software used to prepare material for publication: *SHELXTL* and *PLATON* (Spek, 2003).

Financial support from the Center of Excellence for Innovation in Chemistry (PERCH-CIC), Commission on Higher Education, Ministry of Education, Thailand, is gratefully acknowledged. The authors also thank the Thailand Research Fund (TRF) and the Prince of Songkla University for financial support, and the Universiti Sains Malaysia for the Research University Golden Goose grant No. 1001/PFIZIK/811012.

¹This paper is dedicated to the late Her Royal Highness Princess Galyani Vadhana Krom Luang Naradhiwas Rajanagarindra for her patronage of Science in Thailand.

[§]Additional correspondence author, e-mail: hkfun@usm.my.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: CI2756).

References

- Allen, F. H., Kennard, O., Watson, D. G., Brammer, L., Orpen, A. G. & Taylor, R. (1987). *J. Chem. Soc. Perkin Trans. 2*, pp. S1–19.
- Bruker (2005). *APEX2, SAINT and SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Jung, Y. J., Son, K. I., Oh, Y. E. & Noh, D. Y. (2008). *Polyhedron*, **27**, 861–867.
- Ng, S.-L., Shettigar, V., Razak, I. A., Fun, H.-K., Patil, P. S. & Dharmaprkash, S. M. (2006). *Acta Cryst. E***62**, o1570–o1572.
- Patil, P. S., Chantrapromma, S., Fun, H.-K., Dharmaprkash, S. M. & Babu, H. B. R. (2007). *Acta Cryst. E***63**, o2612.
- Patil, P. S. & Dharmaprkash, S. M. (2008). *Mater. Lett.* **62**, 451–453.
- Patil, P. S., Dharmaprkash, S. M., Ramakrishna, K., Fun, H. K., Sai Santosh Kumar, R. & Narayana Rao, D. (2007). *J. Cryst. Growth*, **303**, 520–524.
- Patil, P. S., Rosli, M. M., Fun, H.-K., Razak, I. A., Puranik, V. G. & Dharmaprkash, S. M. (2006). *Acta Cryst. E***62**, o4798–o4799.
- Prasad, Y. R., Kumar, P. R., Smile, D. J. & Babu, P. A. (2008). *Arkivoc*, **11**, 266–276.
- Sheldrick, G. M. (2008). *Acta Cryst. A***64**, 112–122.
- Spek, A. L. (2003). *J. Appl. Cryst.* **36**, 7–13.
- Suwunwong, T., Chantrapromma, S. & Fun, H.-K. (2009). *Acta Cryst. E***65**, o120.

supplementary materials

Acta Cryst. (2009). E65, o420-o421 [doi:10.1107/S1600536809003122]

(E)-3-(Anthracen-9-yl)-1-(4-bromophenyl)prop-2-en-1-one

T. Suwunwong, S. Chantrapromma, C. Karalai, P. Pakdeevanich and H.-K. Fun

Comment

Chalcones are compounds which have a wide range of applications covering from non-linear optical (Patil & Dharmaprakash, 2008) and electro-active fluorescent materials (Jung *et al.*, 2008) to materials with various biological activities (Prasad *et al.*, 2008). Our previous work (Patil Dharmaprakash *et al.*, 2007) has reported that 1-(4-bromophenyl)-3-(2,4,5-trimethoxyphenyl)-propanone shows efficient second-order nonlinear optical properties. The various interesting properties of chalcone derivatives lead us to synthesize the title chalcone derivative in order to study its photoluminescence and antimicrobial activities.

The molecule of the title chalcone derivative (Fig. 1) exists in an *E* configuration with respect to the C8=C9 double bond [1.333 (2) Å]. The anthracene ring system is planar, with atom C21 deviating a maximum of 0.147 (2) Å. The molecule is twisted as indicated by the interplanar angle between 4-bromophenyl ring and anthracene ring system of 35.52 (7)°, and torsion angles C5–C6–C7–C8 of 22.9 (1)° and C8–C9–C10–C23 of -50.2 (3)°. The pro-2-en-1-one unit (C7–C9/O1) is planar as evidenced by the torsion angle O1–C7–C8–C9 of 0.1 (3)°. The O1/C6–C9 plane makes dihedral angles of 20.9 (1)° and 45.8 (1)°, respectively, with the 4-bromophenyl ring and anthracene ring system. The bond distances show normal values (Allen *et al.*, 1987) and are comparable with those observed in related structures (Ng *et al.*, 2006; Patil *et al.*, 2006; Patil, Chantrapromma *et al.*, 2007; Suwunwong *et al.*, 2009).

In the crystal packing (Fig. 2), the molecules are linked into dimers by weak C—H···Br interactions (Table 1) and the dimers are further linked into a zigzag network parallel to the *bc* plane by weak C—H···O and C—H···π interactions (Table 1).

Experimental

The title compound was synthesized by the condensation of anthracene-9-carbaldehyde (0.01 mol) with 4-bromoacetophenone (0.01 mol) in ethanol (40 ml) in the presence of NaOH (10 ml, 10%). After stirring for 2 h, a yellow solid appeared and was then collected by filtration, washed with distilled water, dried and purified by repeated recrystallization from acetone. Yellow plate-shaped single crystals of the title compound suitable for *X*-ray structure determination were obtained by slow evaporation of an acetone solution at room temperature after several days.

Refinement

All H atoms were placed in calculated positions, with C–H = 0.93 Å, $U_{\text{iso}} = 1.2U_{\text{eq}}(\text{C})$. The highest residual electron density peak is located at 0.76 Å from Br1 and the deepest hole is located at 0.69 Å from Br1.

supplementary materials

Figures




Fig. 1. The molecular structure of the title compound, showing 60% probability displacement ellipsoids and the atom-numbering scheme.




Fig. 2. Part of the crystal packing of the title compound, viewed along the a axis, showing hydrogen-bonded (dashed lines) dimers.

(E)-3-(Anthracen-9-yl)-1-(4-bromophenyl)prop-2-en-1-one

Crystal data

C ₂₃ H ₁₅ BrO	$F_{000} = 784$
$M_r = 387.25$	$D_x = 1.544 \text{ Mg m}^{-3}$
Monoclinic, $P2_1/c$	Mo $K\alpha$ radiation
Hall symbol: -P 2ybc	$\lambda = 0.71073 \text{ \AA}$
$a = 5.3792 (1) \text{ \AA}$	Cell parameters from 4866 reflections
$b = 19.1030 (4) \text{ \AA}$	$\theta = 2.1\text{--}30.0^\circ$
$c = 16.3005 (4) \text{ \AA}$	$\mu = 2.47 \text{ mm}^{-1}$
$\beta = 95.944 (1)^\circ$	$T = 100.0 (1) \text{ K}$
$V = 1666.02 (6) \text{ \AA}^3$	Plate, yellow
$Z = 4$	$0.57 \times 0.27 \times 0.15 \text{ mm}$

Data collection

Bruker SMART APEXII CCD area-detector diffractometer	4866 independent reflections
Radiation source: fine-focus sealed tube	3803 reflections with $I > 2\sigma(I)$
Monochromator: graphite	$R_{\text{int}} = 0.036$
Detector resolution: 8.33 pixels mm^{-1}	$\theta_{\text{max}} = 30.0^\circ$
$T = 100.0(1) \text{ K}$	$\theta_{\text{min}} = 2.1^\circ$
ω scans	$h = -7 \rightarrow 7$
Absorption correction: multi-scan (SADABS; Bruker, 2005)	$k = -26 \rightarrow 26$
$T_{\text{min}} = 0.331$, $T_{\text{max}} = 0.714$	$l = -22 \rightarrow 22$
29994 measured reflections	

Refinement

Refinement on F^2	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.032$	H-atom parameters constrained
$wR(F^2) = 0.076$	$w = 1/[\sigma^2(F_o^2) + (0.0322P)^2 + 1.1607P]$ where $P = (F_o^2 + 2F_c^2)/3$
$S = 1.02$	$(\Delta/\sigma)_{\max} = 0.001$
4866 reflections	$\Delta\rho_{\max} = 0.53 \text{ e \AA}^{-3}$
226 parameters	$\Delta\rho_{\min} = -0.54 \text{ e \AA}^{-3}$
Primary atom site location: structure-invariant direct methods	Extinction correction: none

Special details

Experimental. The low-temperature data was collected with the Oxford Cyrosystem Cobra low-temperature attachment.

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Br1	0.32584 (4)	0.344013 (10)	0.411244 (11)	0.02530 (7)
O1	0.9073 (2)	0.51056 (7)	0.74450 (8)	0.0213 (3)
C1	0.7274 (4)	0.41439 (9)	0.62533 (11)	0.0201 (4)
H1A	0.8800	0.4043	0.6552	0.024*
C2	0.6526 (4)	0.37631 (10)	0.55484 (12)	0.0219 (4)
H2A	0.7530	0.3408	0.5374	0.026*
C3	0.4261 (4)	0.39203 (9)	0.51095 (10)	0.0180 (4)
C4	0.2711 (4)	0.44314 (10)	0.53719 (11)	0.0206 (4)
H4A	0.1176	0.4523	0.5075	0.025*
C5	0.3464 (3)	0.48075 (10)	0.60836 (10)	0.0184 (4)
H5A	0.2419	0.5149	0.6267	0.022*
C6	0.5781 (3)	0.46766 (9)	0.65244 (10)	0.0152 (3)
C7	0.6813 (3)	0.51052 (9)	0.72478 (10)	0.0159 (3)
C8	0.5083 (3)	0.55087 (9)	0.77158 (10)	0.0159 (3)
H8A	0.3374	0.5499	0.7555	0.019*
C9	0.5978 (3)	0.58858 (9)	0.83690 (10)	0.0161 (3)
H9A	0.7707	0.5920	0.8471	0.019*

supplementary materials

C10	0.4492 (3)	0.62533 (9)	0.89440 (10)	0.0158 (3)
C11	0.5151 (3)	0.61533 (9)	0.98020 (10)	0.0157 (3)
C12	0.7151 (4)	0.57046 (10)	1.01164 (11)	0.0195 (4)
H12A	0.8015	0.5452	0.9749	0.023*
C13	0.7822 (4)	0.56385 (10)	1.09428 (11)	0.0219 (4)
H13A	0.9149	0.5349	1.1132	0.026*
C14	0.6506 (4)	0.60082 (10)	1.15145 (11)	0.0220 (4)
H14A	0.7002	0.5970	1.2076	0.026*
C15	0.4527 (4)	0.64183 (10)	1.12478 (11)	0.0224 (4)
H15A	0.3652	0.6648	1.1631	0.027*
C16	0.3767 (3)	0.65030 (9)	1.03860 (10)	0.0169 (3)
C17	0.1715 (4)	0.69108 (9)	1.01020 (10)	0.0187 (4)
H17A	0.0769	0.7118	1.0482	0.022*
C18	0.1036 (3)	0.70178 (9)	0.92633 (10)	0.0167 (3)
C19	-0.1040 (4)	0.74534 (9)	0.89840 (11)	0.0196 (4)
H19A	-0.2054	0.7629	0.9365	0.023*
C20	-0.1560 (4)	0.76158 (9)	0.81740 (11)	0.0211 (4)
H20A	-0.2925	0.7898	0.8002	0.025*
C21	-0.0008 (4)	0.73530 (9)	0.75909 (11)	0.0205 (4)
H21A	-0.0309	0.7487	0.7041	0.025*
C22	0.1913 (3)	0.69077 (9)	0.78224 (10)	0.0184 (4)
H22A	0.2862	0.6730	0.7424	0.022*
C23	0.2504 (3)	0.67066 (9)	0.86689 (10)	0.0153 (3)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Br1	0.03907 (13)	0.02093 (10)	0.01575 (9)	-0.00663 (8)	0.00217 (7)	-0.00386 (7)
O1	0.0137 (7)	0.0297 (7)	0.0203 (6)	0.0009 (5)	0.0009 (5)	-0.0031 (5)
C1	0.0181 (10)	0.0199 (9)	0.0219 (9)	0.0037 (7)	0.0000 (7)	-0.0002 (7)
C2	0.0238 (10)	0.0180 (9)	0.0239 (9)	0.0060 (7)	0.0029 (7)	-0.0027 (7)
C3	0.0238 (10)	0.0159 (8)	0.0146 (8)	-0.0052 (7)	0.0032 (6)	-0.0018 (6)
C4	0.0170 (10)	0.0258 (9)	0.0182 (8)	-0.0003 (7)	-0.0010 (7)	-0.0016 (7)
C5	0.0158 (9)	0.0228 (9)	0.0168 (8)	0.0022 (7)	0.0020 (6)	-0.0031 (7)
C6	0.0155 (9)	0.0169 (8)	0.0134 (7)	-0.0013 (6)	0.0024 (6)	0.0007 (6)
C7	0.0156 (9)	0.0176 (8)	0.0148 (7)	-0.0011 (7)	0.0027 (6)	0.0005 (6)
C8	0.0119 (9)	0.0206 (8)	0.0155 (7)	-0.0010 (7)	0.0022 (6)	-0.0008 (6)
C9	0.0133 (9)	0.0179 (8)	0.0175 (8)	-0.0016 (7)	0.0032 (6)	0.0007 (6)
C10	0.0157 (9)	0.0163 (8)	0.0155 (8)	-0.0035 (7)	0.0024 (6)	-0.0019 (6)
C11	0.0163 (9)	0.0155 (8)	0.0154 (8)	-0.0034 (7)	0.0018 (6)	-0.0006 (6)
C12	0.0195 (10)	0.0211 (9)	0.0184 (8)	-0.0005 (7)	0.0034 (7)	-0.0008 (7)
C13	0.0209 (10)	0.0240 (9)	0.0202 (8)	-0.0003 (8)	-0.0003 (7)	0.0026 (7)
C14	0.0292 (11)	0.0223 (9)	0.0140 (8)	-0.0036 (8)	-0.0004 (7)	0.0005 (7)
C15	0.0307 (11)	0.0227 (9)	0.0141 (8)	-0.0017 (8)	0.0042 (7)	-0.0017 (7)
C16	0.0210 (9)	0.0144 (8)	0.0156 (8)	-0.0040 (7)	0.0036 (6)	-0.0015 (6)
C17	0.0225 (10)	0.0168 (8)	0.0175 (8)	-0.0003 (7)	0.0054 (7)	-0.0032 (6)
C18	0.0182 (9)	0.0142 (8)	0.0178 (8)	-0.0032 (7)	0.0021 (6)	-0.0015 (6)
C19	0.0193 (10)	0.0156 (8)	0.0244 (9)	-0.0002 (7)	0.0050 (7)	-0.0024 (7)

C20	0.0200 (10)	0.0160 (8)	0.0266 (9)	-0.0010 (7)	-0.0018 (7)	0.0006 (7)
C21	0.0216 (10)	0.0209 (9)	0.0182 (8)	-0.0033 (7)	-0.0016 (7)	0.0014 (7)
C22	0.0192 (10)	0.0191 (9)	0.0168 (8)	-0.0029 (7)	0.0020 (6)	-0.0020 (7)
C23	0.0155 (9)	0.0154 (8)	0.0150 (7)	-0.0048 (6)	0.0013 (6)	-0.0018 (6)

Geometric parameters (Å, °)

Br1—C3	1.8954 (17)	C12—C13	1.364 (2)
O1—C7	1.225 (2)	C12—H12A	0.93
C1—C2	1.384 (3)	C13—C14	1.416 (3)
C1—C6	1.396 (2)	C13—H13A	0.93
C1—H1A	0.93	C14—C15	1.356 (3)
C2—C3	1.380 (3)	C14—H14A	0.93
C2—H2A	0.93	C15—C16	1.431 (2)
C3—C4	1.380 (3)	C15—H15A	0.93
C4—C5	1.389 (2)	C16—C17	1.391 (3)
C4—H4A	0.93	C17—C18	1.393 (2)
C5—C6	1.395 (2)	C17—H17A	0.93
C5—H5A	0.93	C18—C19	1.429 (3)
C6—C7	1.495 (2)	C18—C23	1.440 (2)
C7—C8	1.480 (2)	C19—C20	1.357 (3)
C8—C9	1.333 (2)	C19—H19A	0.93
C8—H8A	0.93	C20—C21	1.420 (3)
C9—C10	1.472 (2)	C20—H20A	0.93
C9—H9A	0.93	C21—C22	1.361 (3)
C10—C23	1.413 (3)	C21—H21A	0.93
C10—C11	1.420 (2)	C22—C23	1.436 (2)
C11—C12	1.428 (3)	C22—H22A	0.93
C11—C16	1.433 (2)		
C2—C1—C6	121.23 (17)	C11—C12—H12A	119.3
C2—C1—H1A	119.4	C12—C13—C14	120.32 (18)
C6—C1—H1A	119.4	C12—C13—H13A	119.8
C3—C2—C1	118.75 (17)	C14—C13—H13A	119.8
C3—C2—H2A	120.6	C15—C14—C13	120.41 (17)
C1—C2—H2A	120.6	C15—C14—H14A	119.8
C4—C3—C2	121.49 (16)	C13—C14—H14A	119.8
C4—C3—Br1	118.73 (14)	C14—C15—C16	121.00 (17)
C2—C3—Br1	119.78 (14)	C14—C15—H15A	119.5
C3—C4—C5	119.44 (17)	C16—C15—H15A	119.5
C3—C4—H4A	120.3	C17—C16—C15	121.78 (16)
C5—C4—H4A	120.3	C17—C16—C11	119.28 (16)
C4—C5—C6	120.34 (17)	C15—C16—C11	118.94 (17)
C4—C5—H5A	119.8	C16—C17—C18	121.81 (16)
C6—C5—H5A	119.8	C16—C17—H17A	119.1
C5—C6—C1	118.69 (16)	C18—C17—H17A	119.1
C5—C6—C7	123.18 (16)	C17—C18—C19	120.98 (16)
C1—C6—C7	118.04 (16)	C17—C18—C23	119.57 (16)
O1—C7—C8	121.62 (16)	C19—C18—C23	119.41 (15)
O1—C7—C6	119.02 (15)	C20—C19—C18	121.24 (17)

supplementary materials

C8—C7—C6	119.35 (15)	C20—C19—H19A	119.4
C9—C8—C7	119.93 (16)	C18—C19—H19A	119.4
C9—C8—H8A	120.0	C19—C20—C21	119.62 (18)
C7—C8—H8A	120.0	C19—C20—H20A	120.2
C8—C9—C10	126.25 (17)	C21—C20—H20A	120.2
C8—C9—H9A	116.9	C22—C21—C20	121.16 (16)
C10—C9—H9A	116.9	C22—C21—H21A	119.4
C23—C10—C11	119.94 (15)	C20—C21—H21A	119.4
C23—C10—C9	122.24 (15)	C21—C22—C23	121.31 (17)
C11—C10—C9	117.80 (16)	C21—C22—H22A	119.3
C10—C11—C12	122.43 (16)	C23—C22—H22A	119.3
C10—C11—C16	119.84 (16)	C10—C23—C22	123.65 (16)
C12—C11—C16	117.72 (15)	C10—C23—C18	119.29 (15)
C13—C12—C11	121.49 (17)	C22—C23—C18	116.99 (16)
C13—C12—H12A	119.3		
C6—C1—C2—C3	0.3 (3)	C13—C14—C15—C16	-1.8 (3)
C1—C2—C3—C4	-1.9 (3)	C14—C15—C16—C17	178.78 (18)
C1—C2—C3—Br1	176.90 (14)	C14—C15—C16—C11	-0.8 (3)
C2—C3—C4—C5	1.4 (3)	C10—C11—C16—C17	3.0 (3)
Br1—C3—C4—C5	-177.43 (14)	C12—C11—C16—C17	-176.19 (16)
C3—C4—C5—C6	0.8 (3)	C10—C11—C16—C15	-177.41 (16)
C4—C5—C6—C1	-2.4 (3)	C12—C11—C16—C15	3.4 (2)
C4—C5—C6—C7	174.05 (17)	C15—C16—C17—C18	177.22 (17)
C2—C1—C6—C5	1.9 (3)	C11—C16—C17—C18	-3.2 (3)
C2—C1—C6—C7	-174.75 (17)	C16—C17—C18—C19	-178.36 (17)
C5—C6—C7—O1	-158.07 (17)	C16—C17—C18—C23	-0.7 (3)
C1—C6—C7—O1	18.4 (2)	C17—C18—C19—C20	173.37 (17)
C5—C6—C7—C8	22.9 (2)	C23—C18—C19—C20	-4.3 (3)
C1—C6—C7—C8	-160.59 (16)	C18—C19—C20—C21	-0.4 (3)
O1—C7—C8—C9	0.1 (3)	C19—C20—C21—C22	3.8 (3)
C6—C7—C8—C9	179.10 (16)	C20—C21—C22—C23	-2.3 (3)
C7—C8—C9—C10	-173.26 (16)	C11—C10—C23—C22	171.79 (16)
C8—C9—C10—C23	-50.2 (3)	C9—C10—C23—C22	-6.7 (3)
C8—C9—C10—C11	131.30 (19)	C11—C10—C23—C18	-5.1 (3)
C23—C10—C11—C12	-179.68 (17)	C9—C10—C23—C18	176.46 (16)
C9—C10—C11—C12	-1.1 (3)	C21—C22—C23—C10	-179.25 (18)
C23—C10—C11—C16	1.1 (3)	C21—C22—C23—C18	-2.3 (3)
C9—C10—C11—C16	179.69 (16)	C17—C18—C23—C10	4.9 (3)
C10—C11—C12—C13	177.21 (17)	C19—C18—C23—C10	-177.42 (16)
C16—C11—C12—C13	-3.6 (3)	C17—C18—C23—C22	-172.15 (16)
C11—C12—C13—C14	1.1 (3)	C19—C18—C23—C22	5.5 (2)
C12—C13—C14—C15	1.6 (3)		

Hydrogen-bond geometry (\AA , $^\circ$)

$D\text{—H}\cdots A$	$D\text{—H}$	$H\cdots A$	$D\cdots A$	$D\text{—H}\cdots A$
C8—H8A…O1 ⁱ	0.93	2.42	3.308 (2)	159
C13—H13A…O1 ⁱⁱ	0.93	2.57	3.288 (2)	135

supplementary materials

C21—H21A···Br1 ⁱⁱⁱ	0.93	2.93	3.4722 (19)	119
C9—H9A···Cg1 ^{iv}	0.93	2.83	3.4479 (18)	125

Symmetry codes: (i) $x-1, y, z$; (ii) $-x+2, -y+1, -z+2$; (iii) $-x, -y+1, -z+1$; (iv) $x+1, y, z$.

supplementary materials

Fig. 1




Fig. 2

