

4-(4-Octyloxybenzoyloxy)benzoic acid

Khushi Muhammad,^a M. Khawar Rauf,^a Masahiro Ebihara^b and Shahid Hameed^{a*}

^aDepartment of Chemistry, Quaid-i-Azam University, Islamabad 45320, Pakistan, and ^bDepartment of Chemistry, Faculty of Engineering, Gifu University, Yanagido, Gifu 501-1193, Japan

Correspondence e-mail: shameed@qau.edu.pk


Received 22 January 2009; accepted 27 January 2009

Key indicators: single-crystal X-ray study; $T = 123\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.003\text{ \AA}$; R factor = 0.081; wR factor = 0.138; data-to-parameter ratio = 17.6.

The title compound, $C_{22}\text{H}_{26}\text{O}_5$, is an important intermediate for the synthesis of side-chain ligands for polymeric liquid crystals. The octyl group is coplanar with the central C_6O moiety, where the maximum deviation of a C atom in the octyl group from the C_6O plane is 0.161 (5) \AA . The crystal structure is stabilized by intermolecular O—H···O hydrogen bonds.

Related literature

For studies of aromatic carboxylic acids and their applications, see: Naoum *et al.* (2008); Nazir *et al.* (2008a,b); Gabert *et al.* (2006); Aranzazu *et al.* (2006); Hussain *et al.* (2005); Shafiq *et al.* (2005); Ahmad *et al.* (2003); Ribeiro *et al.* (2008); Hameed & Rama (2004); For related structures, see: Muhammad *et al.* (2008); Hartung *et al.* (1997)



Experimental

Crystal data

$C_{22}\text{H}_{26}\text{O}_5$

$M_r = 370.43$

Monoclinic, $P2_1/n$

$a = 13.528 (8)\text{ \AA}$

$b = 7.245 (4)\text{ \AA}$

$c = 20.903 (12)\text{ \AA}$

$\beta = 111.407 (8)^\circ$

$V = 1907.5 (18)\text{ \AA}^3$

$Z = 4$

Mo $K\alpha$ radiation

$\mu = 0.09\text{ mm}^{-1}$

$T = 123\text{ K}$

$0.40 \times 0.30 \times 0.15\text{ mm}$

Data collection

Rigaku/MSC Mercury CCD diffractometer

Absorption correction: none
14669 measured reflections

4358 independent reflections

3870 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.048$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.081$

$wR(F^2) = 0.138$

$S = 1.31$

4358 reflections

247 parameters

H-atom parameters constrained

$\Delta\rho_{\text{max}} = 0.28\text{ e \AA}^{-3}$

$\Delta\rho_{\text{min}} = -0.19\text{ e \AA}^{-3}$

Table 1

Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
O3—H3O···O4 ⁱ	0.84	1.85	2.659 (3)	161
O4—H4O···O3 ⁱ	0.84	1.83	2.659 (3)	171

Symmetry code: (i) $-x + 2, -y + 1, -z + 1$.

Data collection: *CrystalClear* (Molecular Structure Corporation & Rigaku, 2001); cell refinement: *CrystalClear*; data reduction: *TEXSAN* (Molecular Structure Corporation & Rigaku, 2004); program(s) used to solve structure: *SIR97* (Altomare *et al.*, 1999); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEPII* (Johnson, 1976); software used to prepare material for publication: *SHELXL97* and *TEXSAN*.

MKR is grateful to the Higher Education Commission of Pakistan for financial support under the International Support Initiative Programme for a doctoral fellowship at Gifu University, Japan.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HG2473).

References

- Ahmad, H. B., Rama, N. H., Hussain, M., Hussain, M. T., Qasim, M. M., Hameed, S., Malana, M. A. & Malik, A. (2003). *Indian J. Chem. Sect. B*, **42**, 611–615.
- Altomare, A., Burla, M. C., Camalli, M., Cascarano, G. L., Giacovazzo, C., Guagliardi, A., Molterini, A. G. G., Polidori, G. & Spagna, R. (1999). *J. Appl. Cryst.*, **32**, 115–119.
- Aranzazu, M.-G., Ernesto, P. & Antonio, B. (2006). *Polymer*, **47**, 2080–2090.
- Gabert, A. J., Verpoegen, E., Hammond, P. T. & Schrock, R. R. (2006). *Macromolecules*, **39**, 3993–4000.
- Hameed, S. & Rama, N. H. (2004). *J. Chem. Soc. Pak.*, **26**, 157–162.
- Hartung, H., Hoffmann, F. & Weissflog, W. (1997). *J. Mol. Struct.*, **415**, 205–214.
- Hussain, M. T., Rama, N. H., Hameed, S., Malik, A. & Khan, K. M. (2005). *Nat. Prod. Res.*, **19**, 41–51.
- Johnson, C. K. (1976). *ORTEPII*. Report ORNL-5138. Oak Ridge National Laboratory, Tennessee, USA.
- Molecular Structure Corporation & Rigaku (2001). *CrystalClear*. MSC, The Woodlands, Texas, USA, and Rigaku Corporation, Tokyo, Japan.
- Molecular Structure Corporation & Rigaku (2004). *TEXSAN*. MSC, The Woodlands, Texas, USA, and Rigaku Corporation, Tokyo, Japan.
- Muhammad, K., Khawar Rauf, M., Ebihara, M. & Hameed, S. (2008). *Acta Cryst. E64*, o1251.
- Naoum, M. M., Fahmi, A. A. & Alaasar, M. A. (2008). *Mol. Cryst. Liq. Cryst.*, **487**, 74–91.
- Nazir, S., Khawar Rauf, M., Ebihara, M. & Hameed, S. (2008a). *Acta Cryst. E64*, o423.
- Nazir, S., Muhammad, K., Khawar Rauf, M., Ebihara, M. & Hameed, S. (2008b). *Acta Cryst. E64*, o1013.
- Ribeiro, G., Benadiba, M., Colquhoun, A. & Silva, D. D. (2008). *Polyhedron*, **27**, 1131–1137.
- Shafiq, Z., Arfan, M., Rama, N. H., Hameed, S., Abbas, G. & Hussain, M. T. (2005). *Turk. J. Chem.*, **29**, 321–325.
- Sheldrick, G. M. (2008). *Acta Cryst. A64*, 112–122.

supplementary materials

Acta Cryst. (2009). E65, o424 [doi:10.1107/S1600536809003298]

4-(4-Octyloxybenzoyloxy)benzoic acid

K. Muhammad, M. Khawar Rauf, M. Ebihara and S. Hameed

Comment

Aromatic carboxylic acids bearing different substituents have been investigated for their liquid crystalline properties (Naoum *et al.*, 2008; Nazir *et al.*, 2008*a,b*). Such acids have been utilized in the synthesis of both side-chain (Gabert *et al.*, 2006) and main-chain (Aranzazu *et al.*, 2006) liquid crystal polymers. In addition to their use as intermediates in the synthesis of a large number of organic compounds (Hussain *et al.*, 2005; Shafiq *et al.*, 2005; Ahmad *et al.*, 2003), carboxylic acids has also been used in the pharmaceutical industry (Ribeiro *et al.*, 2008; Hameed & Rama 2004). The title compound (I) was synthesized as an intermediate in the synthesis of side-chain liquid crystal polymers. In the present report, the crystal structure of (I) is being presented. Bond lengths and angles are within the normal ranges as given for benzoyloxybenzoic acids (Muhammad *et al.*, 2008; Hartung *et al.*, 1997). The C7—O1 and C7—O2 bond lengths are 1.205 (3) and 1.369 (3) Å, respectively, that reflect their double and single bond character. The very similar bond lengths of C14—O4 and C14—O3, 1.287 (3) and 1.261 (3) Å, are due to disorder of CO₂H moiety. The octyl group is coplanar with the central C₆O moiety where the *max* deviation of C atom in octyl group from the C₆O moiety is 0.161 (5) Å. Two molecules related by an inversion center form a dimer *via* two hydrogen bonds composed of two carboxyl groups as shown in Fig. 2.

Experimental

To a solution of 4-hydroxybenzaldehyde (0.032 moles) in 50 ml of triethylamine (TEA), was added an equivalent amount of 4-octyloxybenzoylchloride with stirring and the mixture heated at 60°C for 1 h. The excess TEA was removed *in vacuo* and the product, after recrystallization from hot ethanol, was subjected to KMnO₄ oxidation. The 4-(4-octyloxybenzoyloxy)benzaldehyde (0.025 moles) was dissolved in acetone (100 ml) and aqueous KMnO₄ (0.025 moles) was added dropwise at room temperature with stirring. The stirring was continued for three hours when the reaction mixture was filtered and the filtrate acidified using 6M HCl. The precipitated product was purified by recrystallization from acetone.

Refinement

The O-bound H atom was refined isotropically. All the other H atoms were placed in idealized positions and treated as riding atoms, with C—H distance in the range 0.95–0.99 Å and $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$ or $1.5U_{\text{eq}}(\text{C})$.

Figures




Fig. 1. Molecular structure of (I) showing atom-labelling scheme and displacement ellipsoids at the 30% probability level.

supplementary materials




Fig. 2. Showing hydrogen bonded molecules through O—H···O.

4-(4-Octyloxybenzoyloxy)benzoic acid

Crystal data

C ₂₂ H ₂₆ O ₅	$F_{000} = 792$
$M_r = 370.43$	$D_x = 1.290 \text{ Mg m}^{-3}$
Monoclinic, $P2_1/n$	Mo $K\alpha$ radiation
Hall symbol: -P 2yn	$\lambda = 0.71070 \text{ \AA}$
$a = 13.528 (8) \text{ \AA}$	Cell parameters from 4339 reflections
$b = 7.245 (4) \text{ \AA}$	$\theta = 3.0\text{--}27.5^\circ$
$c = 20.903 (12) \text{ \AA}$	$\mu = 0.09 \text{ mm}^{-1}$
$\beta = 111.407 (8)^\circ$	$T = 123 \text{ K}$
$V = 1907.5 (18) \text{ \AA}^3$	Chip, colourless
$Z = 4$	$0.40 \times 0.30 \times 0.15 \text{ mm}$

Data collection

Rigaku/MSC Mercury CCD diffractometer	4358 independent reflections
Radiation source: fine-focus sealed tube	3870 reflections with $I > 2\sigma(I)$
Monochromator: graphite	$R_{\text{int}} = 0.048$
Detector resolution: 14.62 pixels mm^{-1}	$\theta_{\text{max}} = 27.5^\circ$
$T = 123(2) \text{ K}$	$\theta_{\text{min}} = 3.2^\circ$
ω scans	$h = -17 \rightarrow 15$
Absorption correction: none	$k = -7 \rightarrow 9$
14669 measured reflections	$l = -25 \rightarrow 27$

Refinement

Refinement on F^2	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.081$	H-atom parameters constrained
$wR(F^2) = 0.138$	$w = 1/[\sigma^2(F_o^2) + (0.0263P)^2 + 1.5055P]$ where $P = (F_o^2 + 2F_c^2)/3$
$S = 1.31$	$(\Delta/\sigma)_{\max} < 0.001$
4358 reflections	$\Delta\rho_{\max} = 0.28 \text{ e \AA}^{-3}$
247 parameters	$\Delta\rho_{\min} = -0.19 \text{ e \AA}^{-3}$
Primary atom site location: structure-invariant direct methods	Extinction correction: none

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$	Occ. (<1)
C1	0.22298 (16)	0.5733 (3)	0.30577 (10)	0.0181 (4)	
C2	0.19154 (16)	0.5148 (3)	0.23723 (10)	0.0218 (5)	
H2	0.2435	0.4795	0.2189	0.026*	
C3	0.08502 (17)	0.5086 (3)	0.19644 (11)	0.0239 (5)	
H3	0.0638	0.4695	0.1500	0.029*	
C4	0.00858 (16)	0.5596 (3)	0.22337 (10)	0.0202 (5)	
C5	0.03873 (16)	0.6186 (3)	0.29139 (10)	0.0204 (5)	
H5	-0.0132	0.6530	0.3099	0.024*	
C6	0.14626 (17)	0.6261 (3)	0.33158 (10)	0.0211 (5)	
H6	0.1676	0.6683	0.3777	0.025*	
C7	0.33624 (16)	0.5860 (3)	0.35119 (10)	0.0189 (4)	
O1	0.36934 (12)	0.6558 (2)	0.40736 (7)	0.0245 (4)	
O2	0.40082 (11)	0.5073 (2)	0.32137 (7)	0.0231 (4)	
C8	0.51107 (16)	0.5074 (3)	0.35729 (10)	0.0198 (5)	
C9	0.55644 (17)	0.4315 (3)	0.42229 (11)	0.0218 (5)	
H9	0.5133	0.3826	0.4453	0.026*	
C10	0.66633 (16)	0.4284 (3)	0.45304 (11)	0.0201 (4)	
H10	0.6987	0.3783	0.4979	0.024*	

supplementary materials

C11	0.72990 (16)	0.4980 (3)	0.41892 (10)	0.0174 (4)	
C12	0.68191 (16)	0.5711 (3)	0.35287 (10)	0.0191 (4)	
H12	0.7246	0.6174	0.3291	0.023*	
C13	0.57211 (17)	0.5762 (3)	0.32203 (10)	0.0210 (5)	
H13	0.5392	0.6262	0.2772	0.025*	
C14	0.84738 (16)	0.4969 (3)	0.45311 (10)	0.0182 (4)	
O3	0.88859 (12)	0.4343 (2)	0.51339 (7)	0.0257 (4)	
H3O	0.9550	0.4364	0.5257	0.038*	0.50
O4	0.90241 (12)	0.5614 (2)	0.41929 (8)	0.0258 (4)	
H4O	0.9671	0.5599	0.4443	0.039*	0.50
O5	-0.09388 (11)	0.5459 (2)	0.17868 (7)	0.0249 (4)	
C15	-0.17837 (16)	0.5852 (3)	0.20264 (10)	0.0205 (5)	
H15A	-0.1738	0.7146	0.2188	0.025*	
H15B	-0.1743	0.5023	0.2412	0.025*	
C16	-0.28096 (16)	0.5541 (3)	0.14232 (10)	0.0196 (4)	
H16A	-0.2839	0.4239	0.1273	0.024*	
H16B	-0.2814	0.6332	0.1036	0.024*	
C17	-0.37919 (16)	0.5958 (3)	0.15880 (10)	0.0214 (5)	
H17A	-0.3764	0.7258	0.1740	0.026*	
H17B	-0.3792	0.5161	0.1973	0.026*	
C18	-0.48182 (16)	0.5646 (3)	0.09742 (10)	0.0212 (5)	
H18A	-0.4790	0.6383	0.0582	0.025*	
H18B	-0.4857	0.4329	0.0841	0.025*	
C19	-0.58294 (16)	0.6147 (3)	0.10966 (10)	0.0219 (5)	
H19A	-0.5865	0.5417	0.1488	0.026*	
H19B	-0.5806	0.7470	0.1221	0.026*	
C20	-0.68233 (16)	0.5782 (3)	0.04655 (10)	0.0205 (5)	
H20A	-0.6752	0.6432	0.0068	0.025*	
H20B	-0.6864	0.4443	0.0364	0.025*	
C21	-0.78576 (17)	0.6380 (4)	0.05333 (11)	0.0264 (5)	
H21A	-0.7825	0.7718	0.0635	0.032*	
H21B	-0.7944	0.5718	0.0924	0.032*	
C22	-0.88171 (17)	0.5991 (3)	-0.01178 (12)	0.0261 (5)	
H22A	-0.8708	0.6557	-0.0513	0.039*	
H22B	-0.9456	0.6510	-0.0070	0.039*	
H22C	-0.8904	0.4654	-0.0189	0.039*	

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C1	0.0150 (10)	0.0197 (11)	0.0193 (9)	-0.0018 (8)	0.0060 (8)	-0.0005 (8)
C2	0.0153 (10)	0.0294 (13)	0.0218 (10)	0.0000 (9)	0.0080 (8)	-0.0036 (9)
C3	0.0195 (11)	0.0332 (13)	0.0178 (10)	-0.0005 (10)	0.0055 (8)	-0.0049 (9)
C4	0.0153 (10)	0.0229 (12)	0.0202 (10)	-0.0004 (9)	0.0039 (8)	-0.0010 (9)
C5	0.0156 (10)	0.0266 (12)	0.0201 (10)	0.0003 (9)	0.0078 (8)	0.0002 (8)
C6	0.0190 (11)	0.0273 (12)	0.0170 (10)	0.0002 (9)	0.0066 (8)	-0.0008 (8)
C7	0.0176 (10)	0.0195 (11)	0.0204 (10)	-0.0012 (9)	0.0080 (8)	0.0012 (8)
O1	0.0180 (8)	0.0345 (10)	0.0195 (7)	-0.0007 (7)	0.0049 (6)	-0.0042 (7)

O2	0.0122 (7)	0.0339 (10)	0.0212 (7)	0.0011 (7)	0.0037 (6)	-0.0064 (7)
C8	0.0124 (10)	0.0240 (12)	0.0208 (10)	0.0011 (9)	0.0035 (8)	-0.0050 (9)
C9	0.0169 (10)	0.0244 (12)	0.0255 (10)	-0.0020 (9)	0.0095 (8)	-0.0002 (9)
C10	0.0174 (10)	0.0210 (11)	0.0210 (10)	0.0011 (9)	0.0059 (8)	0.0006 (9)
C11	0.0153 (10)	0.0159 (11)	0.0206 (9)	0.0005 (8)	0.0060 (8)	-0.0026 (8)
C12	0.0184 (10)	0.0219 (11)	0.0179 (9)	0.0005 (9)	0.0077 (8)	-0.0030 (8)
C13	0.0191 (10)	0.0266 (12)	0.0151 (9)	0.0021 (9)	0.0036 (8)	-0.0026 (8)
C14	0.0180 (10)	0.0172 (10)	0.0200 (9)	0.0004 (9)	0.0075 (8)	-0.0022 (8)
O3	0.0160 (7)	0.0338 (10)	0.0232 (8)	0.0005 (7)	0.0024 (6)	0.0036 (7)
O4	0.0140 (7)	0.0347 (10)	0.0291 (8)	-0.0015 (7)	0.0084 (6)	0.0029 (7)
O5	0.0130 (7)	0.0395 (10)	0.0206 (7)	0.0009 (7)	0.0042 (6)	-0.0046 (7)
C15	0.0158 (10)	0.0259 (12)	0.0203 (10)	0.0000 (9)	0.0071 (8)	0.0012 (9)
C16	0.0154 (10)	0.0211 (11)	0.0205 (10)	0.0005 (9)	0.0045 (8)	-0.0001 (8)
C17	0.0162 (10)	0.0266 (12)	0.0194 (10)	0.0008 (9)	0.0042 (8)	-0.0015 (9)
C18	0.0170 (10)	0.0253 (12)	0.0189 (10)	-0.0003 (9)	0.0038 (8)	-0.0007 (9)
C19	0.0173 (10)	0.0267 (13)	0.0201 (10)	-0.0004 (9)	0.0050 (8)	-0.0019 (9)
C20	0.0158 (10)	0.0248 (12)	0.0198 (10)	-0.0002 (9)	0.0050 (8)	0.0003 (9)
C21	0.0191 (11)	0.0376 (14)	0.0231 (11)	0.0044 (10)	0.0084 (9)	0.0004 (10)
C22	0.0152 (10)	0.0282 (13)	0.0333 (12)	0.0005 (9)	0.0068 (9)	-0.0001 (10)

Geometric parameters (Å, °)

C1—C6	1.387 (3)	O3—H3O	0.8400
C1—C2	1.403 (3)	O4—H4O	0.8400
C1—C7	1.482 (3)	O5—C15	1.434 (3)
C2—C3	1.382 (3)	C15—C16	1.513 (3)
C2—H2	0.9500	C15—H15A	0.9900
C3—C4	1.396 (3)	C15—H15B	0.9900
C3—H3	0.9500	C16—C17	1.520 (3)
C4—O5	1.363 (2)	C16—H16A	0.9900
C4—C5	1.396 (3)	C16—H16B	0.9900
C5—C6	1.390 (3)	C17—C18	1.525 (3)
C5—H5	0.9500	C17—H17A	0.9900
C6—H6	0.9500	C17—H17B	0.9900
C7—O1	1.205 (3)	C18—C19	1.525 (3)
C7—O2	1.369 (3)	C18—H18A	0.9900
O2—C8	1.404 (2)	C18—H18B	0.9900
C8—C9	1.384 (3)	C19—C20	1.524 (3)
C8—C13	1.385 (3)	C19—H19A	0.9900
C9—C10	1.388 (3)	C19—H19B	0.9900
C9—H9	0.9500	C20—C21	1.520 (3)
C10—C11	1.397 (3)	C20—H20A	0.9900
C10—H10	0.9500	C20—H20B	0.9900
C11—C12	1.398 (3)	C21—C22	1.526 (3)
C11—C14	1.486 (3)	C21—H21A	0.9900
C12—C13	1.387 (3)	C21—H21B	0.9900
C12—H12	0.9500	C22—H22A	0.9800
C13—H13	0.9500	C22—H22B	0.9800
C14—O3	1.261 (3)	C22—H22C	0.9800

supplementary materials

C14—O4	1.287 (3)		
C6—C1—C2	119.32 (19)	O5—C15—H15A	110.4
C6—C1—C7	118.64 (19)	C16—C15—H15A	110.4
C2—C1—C7	122.02 (19)	O5—C15—H15B	110.4
C3—C2—C1	119.9 (2)	C16—C15—H15B	110.4
C3—C2—H2	120.0	H15A—C15—H15B	108.6
C1—C2—H2	120.0	C15—C16—C17	113.15 (18)
C2—C3—C4	120.1 (2)	C15—C16—H16A	108.9
C2—C3—H3	119.9	C17—C16—H16A	108.9
C4—C3—H3	119.9	C15—C16—H16B	108.9
O5—C4—C5	124.50 (19)	C17—C16—H16B	108.9
O5—C4—C3	114.96 (19)	H16A—C16—H16B	107.8
C5—C4—C3	120.54 (19)	C16—C17—C18	112.50 (18)
C6—C5—C4	118.7 (2)	C16—C17—H17A	109.1
C6—C5—H5	120.7	C18—C17—H17A	109.1
C4—C5—H5	120.7	C16—C17—H17B	109.1
C1—C6—C5	121.39 (19)	C18—C17—H17B	109.1
C1—C6—H6	119.3	H17A—C17—H17B	107.8
C5—C6—H6	119.3	C19—C18—C17	114.88 (18)
O1—C7—O2	123.14 (19)	C19—C18—H18A	108.5
O1—C7—C1	125.5 (2)	C17—C18—H18A	108.5
O2—C7—C1	111.40 (17)	C19—C18—H18B	108.5
C7—O2—C8	119.08 (16)	C17—C18—H18B	108.5
C9—C8—C13	121.90 (19)	H18A—C18—H18B	107.5
C9—C8—O2	121.9 (2)	C20—C19—C18	112.06 (18)
C13—C8—O2	116.03 (18)	C20—C19—H19A	109.2
C8—C9—C10	118.5 (2)	C18—C19—H19A	109.2
C8—C9—H9	120.8	C20—C19—H19B	109.2
C10—C9—H9	120.8	C18—C19—H19B	109.2
C9—C10—C11	120.9 (2)	H19A—C19—H19B	107.9
C9—C10—H10	119.6	C21—C20—C19	114.91 (18)
C11—C10—H10	119.6	C21—C20—H20A	108.5
C10—C11—C12	119.38 (19)	C19—C20—H20A	108.5
C10—C11—C14	120.15 (19)	C21—C20—H20B	108.5
C12—C11—C14	120.46 (19)	C19—C20—H20B	108.5
C13—C12—C11	120.1 (2)	H20A—C20—H20B	107.5
C13—C12—H12	119.9	C20—C21—C22	112.13 (19)
C11—C12—H12	119.9	C20—C21—H21A	109.2
C8—C13—C12	119.24 (19)	C22—C21—H21A	109.2
C8—C13—H13	120.4	C20—C21—H21B	109.2
C12—C13—H13	120.4	C22—C21—H21B	109.2
O3—C14—O4	123.08 (19)	H21A—C21—H21B	107.9
O3—C14—C11	119.19 (19)	C21—C22—H22A	109.5
O4—C14—C11	117.73 (18)	C21—C22—H22B	109.5
C14—O3—H3O	109.5	H22A—C22—H22B	109.5
C14—O4—H4O	109.5	C21—C22—H22C	109.5
C4—O5—C15	119.13 (17)	H22A—C22—H22C	109.5
O5—C15—C16	106.55 (17)	H22B—C22—H22C	109.5

C6—C1—C2—C3	−0.6 (3)	C9—C10—C11—C12	−0.2 (3)
C7—C1—C2—C3	−178.8 (2)	C9—C10—C11—C14	178.9 (2)
C1—C2—C3—C4	−0.3 (4)	C10—C11—C12—C13	0.8 (3)
C2—C3—C4—O5	−179.2 (2)	C14—C11—C12—C13	−178.3 (2)
C2—C3—C4—C5	0.5 (4)	C9—C8—C13—C12	−0.8 (3)
O5—C4—C5—C6	179.9 (2)	O2—C8—C13—C12	−176.44 (19)
C3—C4—C5—C6	0.2 (3)	C11—C12—C13—C8	−0.3 (3)
C2—C1—C6—C5	1.4 (3)	C10—C11—C14—O3	−0.8 (3)
C7—C1—C6—C5	179.6 (2)	C12—C11—C14—O3	178.3 (2)
C4—C5—C6—C1	−1.2 (3)	C10—C11—C14—O4	179.5 (2)
C6—C1—C7—O1	−7.8 (4)	C12—C11—C14—O4	−1.4 (3)
C2—C1—C7—O1	170.4 (2)	C5—C4—O5—C15	−3.2 (3)
C6—C1—C7—O2	172.24 (19)	C3—C4—O5—C15	176.5 (2)
C2—C1—C7—O2	−9.6 (3)	C4—O5—C15—C16	−178.78 (19)
O1—C7—O2—C8	−0.1 (3)	O5—C15—C16—C17	−178.28 (18)
C1—C7—O2—C8	179.84 (18)	C15—C16—C17—C18	179.72 (19)
C7—O2—C8—C9	56.4 (3)	C16—C17—C18—C19	−176.7 (2)
C7—O2—C8—C13	−128.0 (2)	C17—C18—C19—C20	−179.4 (2)
C13—C8—C9—C10	1.4 (3)	C18—C19—C20—C21	−175.8 (2)
O2—C8—C9—C10	176.7 (2)	C19—C20—C21—C22	179.5 (2)
C8—C9—C10—C11	−0.8 (3)		

Hydrogen-bond geometry (\AA , °)

$D\cdots H$	$D—H$	$H\cdots A$	$D\cdots A$	$D—H\cdots A$
O3—H3O \cdots O4 ⁱ	0.84	1.85	2.659 (3)	161
O4—H4O \cdots O3 ⁱ	0.84	1.83	2.659 (3)	171

Symmetry codes: (i) $-x+2, -y+1, -z+1$.

supplementary materials

Fig. 1




Fig. 2

