

Published in final edited form as:

J Am Chem Soc. 2010 November 10; 132(44): 15512–15513. doi:10.1021/ja107329k.


Synthesis of Amino Acid-Derived Enaminones via Wolff Rearrangement Using Vinylogous Amides as Carbon Nucleophiles

Hajime Seki, Gunda I. Georg and

Department of Chemistry and Department of Medicinal Chemistry, Institute for Therapeutics Discovery and Development, College of Pharmacy, University of Minnesota, 717 Delaware Street SE, Minneapolis, Minnesota 55414

Abstract

Cyclic enaminones were synthesized in high yields from amino acids in two steps via Wolff rearrangement. The cyclization represents a rare 6-*exo-dig* cyclization involving a ketene as an electrophile. No racemization was observed during this reaction.



Piperidine, indolizidine and quinolizidine alkaloids (Figure 1) display important biological properties and therefore have been frequent targets for synthetic chemistry efforts.¹ It is well-known that monocyclic and bicyclic enaminones (2,3-dihydropyridin-4(1*H*)-ones) can serve as versatile intermediates for their synthesis.² The most common synthetic strategies (Scheme 1) to obtain enaminones in optically active form are: 1) the hetero Diels-Alder reaction, and 2) the nucleophilic addition to pyridinium salts utilizing chiral auxiliaries or chiral catalysts.^{2,3} In 2006 we reported a concise method that relies on the cyclization of amino acid-derived amino ynones to prepare enantioenriched enaminones under very mild conditions.⁴ In some instances, however, partial racemization of the reaction products was observed during these reactions.⁴ To address this issue a different disconnection, that retained the use of amino acids, was sought. We hypothesized that enaminones could be derived from diazoketones carrying a pendant enamine moiety that could trap the ketene generated by a Wolff rearrangement (Scheme 2).⁵

The Wolff rearrangement and its ketene intermediate have been studied extensively. Reactions involving the reactive ketene species are classified as either cycloaddition or nucleophilic addition reactions, exemplified by the Staudinger reaction and the Arndt-Eistert homologation.^{6,7} While a C–C bond is formed in cycloadditions, C–C bond formation in nucleophilic addition reactions of ketenes are less common. Although ketenes have been investigated in reactions with organolithium and magnesium reagents, other type of carbon nucleophiles have been studied less frequently,⁸ and typically involve intermolecular reactions. Only a few reports have appeared where a carbon nucleophile was employed in an intramolecular fashion.⁹ Herein, we detail our new synthetic strategy to prepare enaminones from amino acids via Wolff rearrangement using vinylogous amides as carbon nucleophiles. The cyclization represents a rare example of a 6-*exo-dig* cyclization involving a ketene as the electrophile.¹⁰

* georg@umn.edu .

Our efforts started with the synthesis of diazoketone **1** as a starting material for the reaction in order to test the feasibility of our hypothesis (Scheme 2). *N*-Benzylglycine readily underwent Michael addition to ethyl propiolate, and the acid group of the reaction product was subsequently converted into a diazo moiety. With diazoketone **1** in hand, we embarked on screening reaction conditions. A variety of silver salts, known to induce the Wolff rearrangement, as well as other reaction conditions were tested. We found that the use of halogenated solvents, PhCO₂Ag, CF₃CO₂Ag, and AcOAg readily facilitated the rearrangement to furnish the desired product in excellent yields. The presence of triethylamine, which is often used in Arndt-Eistert homologations as an additive, however diminished the yield. As a result of this study, we selected PhCO₂Ag as the catalyst and CH₂Cl₂ as the solvent for further studies of this method. With optimized reaction conditions in hand, we investigated the scope of the reaction (Table 1). First, diazoketones **3**, **5**, and **7** (entries 1–3) were synthesized from *N*-benzylglycine and the corresponding alkynes. Upon treatment with PhCO₂Ag, the desired enaminones **4**, **6**, and **8** were obtained in good yields. Next, α -substituted diazoketone **9** and **11** (entries 4–5) were prepared from *N*-methylalanine and *N*-methylleucine respectively, which underwent cyclization to afford β -alkyl enaminones **10** and **12**. 19F NMR analysis of the Mosher ester of (*S*)-**12** in comparison to the Mosher esters of (\pm)-**12** revealed that only a single diastereoisomer of the (*S*)-**12** Mosher ester had formed (see SI).¹¹ Diazoketones **13**, **15**, **17**, and **19** (entries 6–9) derived from cyclic amino acids afforded bicyclic enaminones **14**, **16**, **18**, **20** in good yields. These derivatives are known to be excellent intermediates for indolizidine, and quinolizidine syntheses.² Mosher ester derivatives of **14** and **16** showed single diastereoisomers by HPLC analysis (see SI).¹¹ (*2R,8aS*)-**18** was formed as a single diastereoisomer as determined by ¹H NMR.

In summary, we have developed a novel enantiospecific synthetic method to obtain monocyclic, bicyclic and tricyclic enaminones from amino acids in two steps. Due to the lack of racemization, a simple nucleophilic 6-*exo-dig* cyclization is a plausible mechanism, which is a favorable transformation according to Baldwin's rules.¹⁰ Further investigation regarding the synthetic utility of this methodology is ongoing.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

This work was supported by National Institutes of Health Grant GM081267 and the University of Minnesota through the Vince and McKnight Endowed Chairs. We thank Dr. Subhashree Rangarajan for mass spectrometry assistance.

References

- (1). For review: (a) Michael JP. Nat. Prod. Rep. 2008; 25:139. [PubMed: 18250900] (b) Chemler SA. Current Bioact. Compd. 2009; 5:2.
- (2). For review: (a) Comins DL, Joseph SP. Adv. Nitrogen Heterocycl. 1996; 2:251. (b) Joseph S, Comins DL. Curr. Opin. Drug Discovery Dev. 2002; 5:870. (c) Comins, DL.; O'Connor, S.; Alawar, RS. Comprehensive Heterocyclic Chemistry III. Alan, RK.; Christopher, AR.; Eric, FVS.; Richard, JKT., editors. Elsevier; Oxford: 2008. p. 41
- (3). (a) Danishefsky S, Kitahara T. J. Am. Chem. Soc. 1974; 96:7807. (b) Pfrengle W, Kunz H. J. Org. Chem. 1989; 54:4261. (c) Ishihara K, Miyata M, Hattori K, Tada T, Yamamoto H. J. Am. Chem. Soc. 1994; 116:10520. (d) Kobayashi S, Kusakabe K.-i. Komiyama S, Ishitani H. J. Org. Chem. 1999; 64:4220. (e) Yao S, Saaby S, Hazell RG, Jorgensen KA. Chem.--Eur. J. 2000; 6:2435. (f) Josephsohn NS, Snapper ML, Hoveyda AH. J. Am. Chem. Soc. 2003; 125:4018. [PubMed:

- 12670201] (g) Mancheno OG, Arrayas RG, Carretero JC. *J. Am. Chem. Soc.* 2004; 126:456. [PubMed: 14719929] (h) Yamashita Y, Mizuki Y, Kobayashi S. *Tetrahedron Lett.* 2005; 46:1803.
- (4). (a) Turunen BJ, Georg GI. *J. Am. Chem. Soc.* 2006; 128:8702. [PubMed: 16819843] (b) Niphakis MJ, Turunen BJ, Georg GI. *J. Org. Chem.* 2010; 75:6019. [PubMed: 20704319] (c) Niphakis MJ, Georg GI. *J. Org. Chem.* 2010 ASAP.
- (5). For Review: (a) Meier H, Zeller KP. *Angew. Chem.* 1975; 87:52. (b) Kirmse W. *Eur. J. Org. Chem.* 2002:2193. (c) Tidwell TT. *Eur. J. Org. Chem.* 2006:563.
- (6). For review: Georg, GI.; Ravikumar, VT. Stereocontrolled Ketene-Imine Cycloaddition Reactions. In: Georg, GI., editor. *The Organic Chemistry of β -Lactams*. Verlag Chemie; New York: 1993. p. 295
- (7). (a) Bachmann WE, Struve WS. *Org. React.* 1942:38–62. (b) Seikaly HR, Tidwell TT. *Tetrahedron.* 1986; 42:2587–613.
- (8). For ketene reactions with organo lithium/magnesium reagents, see: (a) Haener R, Laube T, Seebach D. *J. Am. Chem. Soc.* 1985; 107:5396. (b) Baigrie LM, Seikaly HR, Tidwell TT. *J. Am. Chem. Soc.* 1985; 107:5391. For reactions with other type of carbon nucleophiles, see: (c) Rathke MW, Sullivan DF. *Tetrahedron Lett.* 1973:1297. (d) Kita Y, Matsuda S, Kitagaki S, Tsuzuki Y, Akai S. *Synlett.* 1991:401. (e) Negri G, Kascheres C. *J. Heterocycl. Chem.* 2001; 38:109.
- (9). (a) Hickmott PW, Giasuddin Ahmed M, Ahmed SA, Wood S, Kapon M. *J. Chem. Soc., Perkin Trans.* 1985:2559. (b) Hickmott PW. *S. Afr. J. Chem.* 1989; 42:17. (c) Byeon C-H, Hart DJ, Lai C-S, Unch J. *Synlett.* 2000:119.
- (10). (a) Baldwin JE. *J. Chem. Soc., Chem. Commun.* 1976:734. (b) Johnson CD. *Acc. Chem. Res.* 1993; 26:476.
- (11). Dale JA, Dull DL, Mosher HS. *J. Org. Chem.* 1969; 34:2543. See SI for experimental details






Figure 1.
Examples of biologically active molecules containing piperidine, indolizidine and quinolizidine core structures



Scheme 1.
Various approaches for enaminone syntheses



entry ^a	catalyst	isolated yield (%)
1	Ag_2O	90
2	$\text{CF}_3\text{CO}_2\text{Ag}$	82
3	AcOAg^b	97
4	PhCO_2Ag	99









Scheme 2.











Optimization of reaction conditions

^aReaction conditions: diazoketone **1** in CH_2Cl_2 (0.2M), 24 h, dark^bSonication was used.

Table 1

Substrate scope of the reaction^a

entry	substrate	product	isolated yield (%)
1	 <p>3</p>	 <p>4</p>	80
2	 <p>5</p>	 <p>6</p>	88
3 ^b	 <p>7</p>	 <p>8</p>	93
4	 <p>9</p>	 <p>10</p>	93

entry	substrate	product	isolated yield (%)
5	 <p>11</p>	 <p>12</p>	76 ^c
6	 <p>13</p>	 <p>14</p>	93 ^c
7 ^d	 <p>15</p>	 <p>16</p>	86 ^c
8	 <p>17</p>	 <p>18</p>	79 ^e
9	 <p>19</p>	 <p>20</p>	81

^a Reaction conditions: diazoketone and PhCO₂Ag (10 mol%) in CH₂Cl₂ (0.2 M).

^b 20 mol% of PhCO₂Ag was used.

^c *ee* was determined by ¹⁹F NMR (Mosher ester derivatives); only one isomer was observed.

^d Ag₂O (10 mol%) and C₂H₄Cl₂ (0.2 M) were used.

^e *dr* was determined by ¹H NMR; only one isomer was observed.