

Published in final edited form as:

*Inorg Chem.* 2010 November 15; 49(22): 10238–10240. doi:10.1021/ic1018373.

## Coordination-Driven Self-Assembly of Truncated Tetrahedra Capable of Encapsulating 1,3,5-Triphenylbenzene


Yao-Rong Zheng<sup>†</sup>, Zhigang Zhao<sup>†</sup>, Hyunuk Kim<sup>‡</sup>, Ming Wang<sup>†</sup>, Koushik Ghosh<sup>†</sup>, J. Bryant Pollock<sup>†</sup>, Ki-Whan Chi<sup>‡</sup>, and Peter J. Stang<sup>†,\*</sup>

<sup>†</sup>Department of Chemistry, University of Utah, 315 South 1400 East, RM 2020, Salt Lake City, Utah 84112

<sup>‡</sup>Department of Chemistry, University of Ulsan, Ulsan 680-749, Republic of Korea

### Abstract

The design and synthesis of coordinative truncated tetrahedra is described. Coordination-driven self-assembly of a truncated tetrahedron was achieved using 90° organoplatinum acceptors and a hexapyridyl ligand with six-fold symmetry under mild conditions. This assembly can act as a host towards 1,3,5-triphenylbenzene. The truncated tetrahedral structures and the host-guest complex were identified using multinuclear (<sup>31</sup>P and <sup>1</sup>H) NMR spectroscopy, electrospray ionization (ESI) mass spectrometry, X-ray crystallography, and pulsed field gradient spin-echo (PGSE) NMR, along with computational simulations.



Coordination-driven self-assembly is a successful methodology for preparing three-dimensional (3-D) supramolecular structures.<sup>1</sup> During the past two decades, a variety of novel 3-D coordinative structures of high complexity, high symmetry, and well-defined size and shape, such as tetrahedra, cubes, double squares, cuboctahedra, adamantanoids, dodecahedra, and a sphere, have been developed.<sup>2</sup> By virtue of their robust structural backbone and well-defined 3-D cavity, 3-D coordinative supramolecules have been used in a variety of applications such as guest encapsulation, gas storage, catalysis, and drug delivery.<sup>3</sup> With further exo/endo functionalization,<sup>4</sup> 3-D supramolecules are endowed with a variety of novel properties, like 3-D dendrimers, nanoscale fluoro-droplet, and confined polymerization.<sup>5</sup> Among these 3-D supramolecules, because of their unique structural features and fascinating host-guest properties, tetrahedral structures represent one of the most widely studied systems, as witnessed by the impressive results of Raymond<sup>6</sup> and Fujita.<sup>7</sup> However, the design of tetrahedral structures is still limited.<sup>6–8</sup>

Homoleptic self-assembly of a hexapyridyl ligand and a 90° acceptor should form a tetrahedral structure, where the hexadentate ligand acts as faces and the organoplatinum acceptors are the connectors at the corners as shown in Scheme 1. We therefore carried out

\*To whom correspondence should be addressed. stang@chem.utah.edu.

**Supporting Information Available:** Experimental details for the synthesis and characterization of supramolecules **3** and the host-guest study. (<http://pubs.acs.org/>).

the self-assembly of the hexapyridyl donor **29** with the 90° organoplatinum acceptors **1**, resulting in a truncated tetrahedron in quantitative yield. The structures were characterized by multinuclear (<sup>31</sup>P and <sup>1</sup>H) NMR spectroscopy, electrospray ionization mass spectrometry (ESI-MS), pulsed field gradient spin-echo (PGSE) NMR, as well as X-ray crystallography. Furthermore, it was found that truncated tetrahedron **3b** is able to encapsulate 1,3,5-triphenylbenzene in an aqueous acetone solution. This host-guest complex was characterized by NMR spectroscopy and ESI-MS as well as computational simulations.

By mixing 90° organoplatinum acceptor **1** and hexapyridyl donor **29** in a 3:1 ratio in an acetone-*d*<sub>6</sub>/CD<sub>3</sub>NO<sub>2</sub> (*v/v* 7:3) solution for **3a** and an aqueous acetone solution (*v/v* 1:1) for **3b**, self-assembly of truncated tetrahedra **3** were obtained after 16 h of heating at 80 °C. The assemblies **3** can be quantitatively isolated via ion exchange with KPF<sub>6</sub>.

In the <sup>31</sup>P{<sup>1</sup>H} NMR spectra (Figure 1) of **3**, only one singlet at 0.92 ppm (**3a**) and -28.2 ppm (**3b**) with concomitant <sup>195</sup>Pt satellites can be found, respectively, for the coordinated platinum centers. Likewise, the <sup>1</sup>H NMR spectra (See Supporting Information) also exhibit sharp signals for the pyridyl protons of **3** ( $\delta_{\text{Py}\alpha\text{-H}}$ : 8.91 ppm and  $\delta_{\text{Py}\beta\text{-H}}$ : 7.44 ppm for **3a**;  $\delta_{\text{Py}\alpha\text{-H}}$ : 8.68 ppm and  $\delta_{\text{Py}\beta\text{-H}}$ : 7.48 ppm for **3b**), with approximately 0.1 ppm ( $\text{H}_{\text{Py-}\beta}$ ) and 0.2–4 ppm ( $\text{H}_{\text{Py-}\alpha}$ ) downfield shifts due to the loss of electron density upon coordination with the platinum centers. Signals corresponding to the phenyl protons on the donors are split into two sets of doublets, presumably caused by the difference between the exterior and interior of the cage structure. In the ESI mass spectra of **3** (see Supporting Information), peaks corresponding to [**3a** – 5PF<sub>6</sub>]<sup>5+</sup> and [**3a** – 6PF<sub>6</sub>]<sup>6+</sup> can be found at *m/z* = 2383.1 and *m/z* = 1961.9, as are those for **3b** at *m/z* = 2197.0 [**3b** – 5OTf]<sup>5+</sup> and *m/z* = 1806.2 [**3b** – 6OTf]<sup>6+</sup>. These signals are isotopically resolved and in agreement with their theoretical distributions.

The structure of **3** was unambiguously determined by X-ray diffraction analysis using synchrotron radiation. X-ray quality crystals of **3a** were obtained by slow diffusion of pentane into the acetone solution of **3a**. As shown in Figure 2 and S4 in the Supporting Information, the structure has a truncated tetrahedral shape with a diameter of 3.6 nm, and bears a 1.0 nm cavity core. The shortest Pt-Pt distance is 1.1 nm. The PF<sub>6</sub><sup>-</sup> anions are found surrounding the structure but not in the cavity, and the shortest Pt-F distance was measured to be 0.48 nm. As seen in Table S3 (see Supporting Information), the Pt atoms in the structure are coordinating with two PEt<sub>3</sub> and two nitrogen atoms from the pyridine moieties in a distorted square planar geometry. The mean values from the square planar Pt(II) is N-Pt-N 80.6°, P-Pt-P 98.2°, N-Pt-P 90.6°, Pt-P 2.28 Å, and Pt-N 2.09 Å, respectively.

PGSE NMR measurements were also used to characterize the structures in solution. Using the translational self-diffusion coefficient measured by PGSE NMR in conjunction with the Stokes-Einstein equation, the “effective size” of the overall assembly in acetone-*d*<sub>6</sub> was obtained: 3.35 ± 0.15 nm for **3a** and 3.23 ± 0.21 nm for **3b**, which are in good agreement with that of the crystal structure.

The host-guest properties of **3** were also studied. According to the symmetry and size of the host, 1,3,5-triphenylbenzene **4** was chosen for the investigation. The experiment was carried out by mixing **1b** and **2** in a 3:1 ratio with excess triphenylbenzene **4** in an aqueous acetone solution (*v/v* 1:1). After 16 h of heating at 70 °C, the encapsulated complex **3b**·**4**<sub>3</sub> was formed.

The singlet at -28.2 ppm with concomitant <sup>195</sup>Pt satellites in the <sup>31</sup>P{<sup>1</sup>H} NMR spectrum (see Supporting Information) and the identifiable peaks ( $\delta$  = 8.76 ppm  $\text{H}_{\text{Py-}\alpha\text{-3b}}$ ;  $\delta$  = 7.36 ppm  $\text{H}_{\text{Py-}\beta\text{-3b}}$ ;  $\delta$  = 6.94 ppm  $\text{H}_{\text{phenyl-3b}}$ ) in the <sup>1</sup>H NMR spectra (Figure 3b) show the formation of a truncated tetrahedron. By comparing the <sup>1</sup>H NMR spectra of **4** (acetone-*d*<sub>6</sub>),

**3b** (acetone-*d*<sub>6</sub>/D<sub>2</sub>O = 1:1), and **3b**·**4**<sub>3</sub> (acetone-*d*<sub>6</sub>/D<sub>2</sub>O = 1:1) in Figure 3, signals are found at 7.25 ppm ( $\Delta\delta = -0.6$  ppm, H<sub>Phenyl-4</sub>), 7.06 ppm ( $\Delta\delta = 0.24$  ppm, H<sub>Phenyl-3b</sub>), and 6.35 ppm ( $\Delta\delta = -1.0$ – $1.8$  ppm, H<sub>Phenyl-4</sub>), indicating that the triphenylbenzene **4** is encapsulated in the truncated tetrahedron **3b**, and integration of the peaks at 8.76 ppm (H<sub>Py- $\alpha$ -3b</sub>) and 6.35 ppm (H<sub>Phenyl-4</sub>) suggest that three guest molecules are encapsulated in each cage. In the ESI mass spectra (see Supporting Information), isotopically resolved signals at *m/z* = 2380.8 [**3b**·**4**<sub>3</sub> – 5OTf]<sup>5+</sup> and *m/z* = 1959.3 [**3b**·**4**<sub>3</sub> – 6OTf]<sup>6+</sup> confirm the complex **3b**·**4**<sub>3</sub>. Elemental analysis (see Supporting Information) of the isolated complex is also consistent with the composition of **3b**·**4**<sub>3</sub>.

While X-ray quality crystals for **3b**·**4**<sub>3</sub> were not obtained, a computational simulation was used to gain insight into the structural features of the encapsulated complex **3b**·**4**<sub>3</sub>. A 10 Å molecular dynamics simulation using a molecular mechanics force field (MMFF), 300K, in the gas phase was used to equilibrate each supramolecule, and the output of the simulation was then minimized to full convergence. As shown in Figure 4, in the model of **3b**·**4**<sub>3</sub>, three triphenylbenzene molecules **4** are stacked within the cavity of **3b**. The distance between these guests is about 4.3 Å, and that between the guest and the interior of the cage is 3.6–4.3 Å.

In conclusion, we report the facile synthesis of a new type of 3-D truncated tetrahedra via coordination-driven self-assembly, where the highly symmetrical hexapyridyl ligand acts as the faces and 90° organoplatinum acceptors are connectors at the edges. These truncated tetrahedra show a unique 3-D nanoscale pore, and preliminary studies indicate the nanocavity is able to encapsulate 1,3,5-triphenylbenzene.

## Supplementary Material

Refer to Web version on PubMed Central for supplementary material.


## Acknowledgments

P.J.S. thanks the NIH (GM-057052) for financial support. K.W.C. and P.J.S. gratefully acknowledge the WCU program (R33-2008-000-10003-0) of the Korean Ministry of Education, Science and Technology for the support of this work. X-ray diffraction experiments using synchrotron radiation were performed at the Pohang Accelerator Laboratory supported by MOST and POSTECH in Korea.


## References

- (a) Leininger S, Olenyuk B, Stang PJ. *Chem. Rev.* 2000; 100:853. [PubMed: 11749254] (b) Seidel SR, Stang PJ. *Acc. Chem. Res.* 2002; 35:972. [PubMed: 12437322] (c) Holliday BJ, Mirkin CA. *Angew. Chem., Int. Ed.* 2001; 40:2022. (d) Fujita M, Umemoto K, Yoshizawa M, Fujita N, Kusakawa T, Biradha K. *Chem. Commun.* 2001:509. (e) Fiedler D, Leung DH, Bergman RG, Raymond KN. *Acc. Chem. Res.* 2005; 38:351. (f) Fujita M, Tominaga M, Hori A, Therrien B. *Acc. Chem. Res.* 2005; 38:369. [PubMed: 15835883] (g) Ruben M, Rojo J, Romero-Salguero FJ, Uppadine LH, Lehn J-M. *Angew. Chem., Int. Ed.* 2004; 43:3644. (h) Severin K. *Chem. Commun.* 2006:3859. (i) Nitschke JR. *Acc. Chem. Res.* 2007; 40:103. [PubMed: 17309191] (j) Oliveri CG, Ulmann PA, Wiester MJ, Mirkin CA. *Acc. Chem. Res.* 2008; 41:1618. [PubMed: 18642933] (k) Ward MD. *Chem. Commun.* 2009; 30:4487. (l) Jin P, Dalgarno SJ, Atwood JL. *Coord. Chem. Rev.* 2010; 254:1760.
- (a) Fujita M, Oguro D, Miyazawa M, Oka H, Yamaguchi K, Ogura K. *Nature.* 1995; 378:469. (b) Parac TN, Caulder DL, Raymond KN. *J. Am. Chem. Soc.* 1998; 120:8003. (c) Johannessen SC, Brisbois RG, Fischer JP, Grieco PA, Counterman AE, Clemmer DE. *J. Am. Chem. Soc.* 2001; 123:3818. [PubMed: 11457114] (d) Roche S, Haslam C, Adams H, Heath SL, Thomas JA. *Chem. Commun.* 1998:1681. (e) Fujita M, Yu S-Y, Kusakawa T, Funaki H, Ogura K, Yamaguchi K. *Angew. Chem. Int. Ed. Engl.* 1998; 37:2082. (f) Olenyuk B, Whiteford JA, Fechtenkotter A, Stang


- PJ. Nature. 1999; 398:796. [PubMed: 10235260] (g) Schweiger M, Seidel SR, Schmitz M, Stang PJ. Org. Lett. 2000; 2:1255. [PubMed: 10810721] (h) Olenyuk B, Levin MD, Whiteford JA, Shield JE, Stang PJ. J. Am. Chem. Soc. 1999; 121:10434. (i) Suzuki K, Kawano M, Sato S, Fujita M. J. Am. Chem. Soc. 2007; 129:10652. [PubMed: 17685616] (j) Ghosh K, Hu J, White HS, Stang PJ. J. Am. Chem. Soc. 2009; 131:6695. [PubMed: 19397325] (k) Suzuki K, Tominaga M, Kawano M, Fujita M. Chem. Commun. 2009; 13:1638. (l) Zheng Y-R, Ghosh K, Yang H-B, Stang PJ. Inorg. Chem. 2010; 49:9007. (m) Sun Q-F, Iwasa J, Ogawa D, Ishido Y, Sato S, Ozeki T, Sei Y, Yamaguchi K, Fujita M. Science. 2010; 328:1144. [PubMed: 20430973]
3. (a) Rosi NL, Eckert J, Eddaoudi M, Vodak DT, Kim J, O'Keeffe M, Yaghi OM. Science. 2003; 300:1127. [PubMed: 12750515] (b) Trembleau, Laurent; Rebek, Julius, Jr. Science. 2003; 301:1219. [PubMed: 12947192] (c) Yoshizawa M, Tamura M, Fujita M. Science. 2006; 312:251. [PubMed: 16614218] (d) Pluth, MichaelD; Bergman, RobertG; Raymond, KennethN. Science. 2007; 316:85. [PubMed: 17412953]
4. Northrop BH, Yang H-B, Stang PJ. Chem. Commun. 2008:5896.
5. (a) Sato S, Iida J, Suzuki K, Kawano M, Ozeki T, Fujita M. Science. 2006; 313:1273. [PubMed: 16946067] (b) Kamiya N, Tominaga M, Sato S, Fujita M. J. Am. Chem. Soc. 2007; 129:3816. [PubMed: 17343384] (c) Murase T, Sato S, Fujita M. Angew. Chem., Int. Ed. 2007; 46:5133. (d) Zheng Y-R, Ghosh K, Yang H-B, Stang PJ. Inorg. Chem. 2010; 49:4747. [PubMed: 20443570]
6. (a) Pluth MD, Bergman RG, Raymond KN. Acc. Chem. Res. 2009; 42:1650. [PubMed: 19591461] (b) Pluth MD, Fiedler D, Mugridge JS, Bergman RG, Raymond KN. Proc. Natl. Acad. Sci. USA. 2009; 106:10438. [PubMed: 19181859] (c) Brown CJ, Bergman RG, Raymond KN. J. Am. Chem. Soc. 2009; 131:17530. [PubMed: 19950985] (d) Hastings CJ, Pluth MD, Bergman RG, Raymond KN. J. Am. Chem. Soc. 2010; 132:6938. [PubMed: 20443566]
7. (a) Yoshizawa M, Klosterman JK, Fujita M. Angew. Chem., Int. Ed. 2009; 48:3418. (b) Murase T, Horiuchi S, Fujita M. J. Am. Chem. Soc. 2010; 132:2866. [PubMed: 20155904] (c) Dolain C, Hatakeyama Y, Sawada T, Tashiro S, Fujita M. J. Am. Chem. Soc. 2010; 132:5644. (d) Horiuchi S, Nishioka Y, Murase T, Fujita M. Chem. Commun. 2010; 46:3460. (e) Murase T, Otsuka K, Fujita M. J. Am. Chem. Soc. 2010; 132:7864. [PubMed: 20481576]
8. (a) Leininger S, Fan J, Schmitz M, Stang PJ. Proc. Natl. Acad. Sci. USA. 2000; 97:1380. [PubMed: 10677470] (b) Schweiger M, Yamamoto T, Stang PJ, Blaeser D, Boese R. J. Org. Chem. 2005; 70:4861. [PubMed: 15932331]
9. (a) Rathore R, Burns CL, Guzei IA. J. Org. Chem. 2004; 69:1524. [PubMed: 14987006] (b) Hoffmann M, Kärnbratt J, Chuan MH, Herz LM, Albinsson B, Anderson HL. Angew. Chem., Int. Ed. 2008; 47:4993.
10. Caskey DC, Yamamoto T, Addicott C, Shoemaker RK, Vacek J, Hawkrigde AM, Muddiman DC, Kottas GS, Michl J, Stang PJ. J. Am. Chem. Soc. 2008; 130:7620. and references therein. [PubMed: 18491898]




**Figure 1.**  $^{31}\text{P}\{^1\text{H}\}$  NMR spectra (121.4 MHz) of truncated tetrahedra **3a** in acetone- $d_6$ / $\text{CD}_3\text{NO}_2$  (a) and **3b** in acetone- $d_6$ / $\text{D}_2\text{O}$  (b).



**Figure 2.** Crystal structure of truncated tetrahedron **3a** (Pt: green; C: grey; N: blue; P: Orange; protons, solvent, and  $\text{PF}_6^-$  omitted for clarity).





**Figure 3.** Partial  $^1\text{H}$  NMR spectra (300 MHz) of “free” **4** in acetone- $d_6$  (a), **3b** in acetone- $d_6/\text{D}_2\text{O}$  (b), and the encapsulated complex **3b·4<sub>3</sub>** in acetone- $d_6/\text{D}_2\text{O}$  (c).



**Figure 4.** Computational model (MMFF) of the encapsulated complex  $3b \cdot 4_3$  (For clarity, the three guest molecules are labeled as blue, green, and orange).





**Scheme 1.**  
Graphical representation of the [12 + 4] self-assembly of 90° organoplatinum acceptors **1** and hexapyridyl ligand **2** into a truncated tetrahedron **3**.