

## MINIREVIEW

### Acetic Acid Bacteria, Newly Emerging Symbionts of Insects<sup>∇</sup>

Elena Crotti,<sup>1</sup> Aurora Rizzi,<sup>1</sup> Bessem Chouaia,<sup>1</sup> Irene Ricci,<sup>2</sup> Guido Favia,<sup>2</sup> Alberto Alma,<sup>3</sup>  
Luciano Sacchi,<sup>4</sup> Kostas Bourtzis,<sup>5</sup> Mauro Mandrioli,<sup>6</sup> Ameer Cherif,<sup>7</sup>  
Claudio Bandi,<sup>8</sup> and Daniele Daffonchio<sup>1\*</sup>

*Dipartimento di Scienze e Tecnologie Alimentari e Microbiologiche (DiSTAM), Università degli Studi di Milano, 20133 Milan, Italy<sup>1</sup>;*  
*Dipartimento di Medicina Sperimentale e Sanità Pubblica (DMSSP), Università degli Studi di Camerino, 62032 Camerino, Italy<sup>2</sup>;*  
*Dipartimento di Valorizzazione e Protezione delle Risorse Agroforestali (DIVAPRA), Università degli Studi di Torino, 10095 Grugliasco, Italy<sup>3</sup>;*  
*Dipartimento di Biologia Animale (DBA), Università degli Studi di Pavia, 27100 Pavia, Italy<sup>4</sup>;*  
*Department of Environmental and Natural Resources Management, University of Ioannina, 30100 Agrinio, Greece<sup>5</sup>;*  
*Dipartimento di Biologia Animale (DBA), Università degli Studi di Modena e Reggio Emilia, 41125 Modena, Italy<sup>6</sup>;*  
*Laboratoire des Microorganismes et Biomolécules Actives (LMBA), Université de Tunis El Manar, Tunis, Tunisia<sup>7</sup>;*  
*Dipartimento di Patologia Animale, Igiene e Sanità Pubblica Veterinaria (DiPAV), Università degli Studi di Milano, 20133 Milan, Italy<sup>8</sup>*

Recent research in microbe-insect symbiosis has shown that acetic acid bacteria (AAB) establish symbiotic relationships with several insects of the orders Diptera, Hymenoptera, Hemiptera, and Homoptera, all relying on sugar-based diets, such as nectars, fruit sugars, or phloem sap. To date, the fruit flies *Drosophila melanogaster* and *Bactrocera oleae*, mosquitoes of the genera *Anopheles* and *Aedes*, the honey bee *Apis mellifera*, the leafhopper *Scaphoideus titanus*, and the mealybug *Saccharicoccus sacchari* have been found to be associated with the bacterial genera *Acetobacter*, *Gluconacetobacter*, *Gluconobacter*, *Asaia*, and *Saccharibacter* and the novel genus *Commensalibacter*. AAB establish symbiotic associations with the insect midgut, a niche characterized by the availability of diet-derived carbohydrates and oxygen and by an acidic pH, selective factors that support AAB growth. AAB have been shown to actively colonize different insect tissues and organs, such as the epithelia of male and female reproductive organs, the Malpighian tubules, and the salivary glands. This complex topology of the symbiosis indicates that AAB possess the keys for passing through body barriers, allowing them to migrate to different organs of the host. Recently, AAB involvement in the regulation of innate immune system homeostasis of *Drosophila* has been shown, indicating a functional role in host survival. All of these lines of evidence indicate that AAB can play different roles in insect biology, not being restricted to the feeding habit of the host. The close association of AAB and their insect hosts has been confirmed by the demonstration of multiple modes of transmission between individuals and to their progeny that include vertical and horizontal transmission routes, comprising a venereal one. Taken together, the data indicate that AAB represent novel secondary symbionts of insects.

Acetic acid bacteria (AAB), especially members of the genera *Acetobacter* and *Gluconacetobacter*, have a significant historical role in human activities, having been used over millennia for the production of vinegar for consumption and for medicinal purposes. AAB of the family *Acetobacteraceae* are ubiquitous and are known to be adapted to various sugar- and ethanol-rich environments. AAB are obligate aerobes, and most of them are unable to oxidize ethanol, sugars, and polyalcohols completely, accumulating large amounts of the corresponding oxidation products in their culture medium. Their oxidative capacity is commercially exploited not only for vinegar production but also in the manufacture of foods and chemical compounds, i.e., kombucha tea, cocoa, sorbose, gluconic acid, etc. (40).

AAB can be isolated from a variety of substrates and natural environments like plants, flowers, herbs, fruits, and fermented foods and beverages. Great attention has recently been directed toward symbiotic associations between AAB and insect hosts, which are emerging as a novel niche for AAB isolation. Interestingly, these prokaryotes have been reported to be associated with insects that rely on sugar-based diets, in particular those belonging to the orders Diptera, Hymenoptera, and Hemiptera (13).

Symbiotic associations can profoundly affect the evolutionary history, lifestyle, and physiology of organisms. Consequently, in recent years, the interactions between insects and microorganisms have received considerable attention (53). The remarkable adaptability of insects to very diverse terrestrial habitats, comprising those that are nutritionally limited or unbalanced, has contributed to their evolutionary success in different ecological environments. In many cases, microbial partners allow their insect hosts to specialize in nutrient-deficient food resources, giving them competitive advantages (24). For example, insects that feed exclusively on nutritionally poor

\* Corresponding author. Mailing address: Dipartimento di Scienze e Tecnologie Alimentari e Microbiologiche (DiSTAM), Università degli Studi di Milano, 20133 Milan, Italy. Phone: 39-0250316730. Fax: 39-0250316694. E-mail: daniele.daffonchio@unimi.it.

<sup>∇</sup> Published ahead of print on 17 September 2010.

diets such as plant sap, vertebrate blood, or woody material possess microbial endosymbionts. Arthropod-associated microorganisms play numerous different roles besides the nutritional aspects, influencing development, reproduction and speciation, defense against natural enemies, and immunity (14). For example, the aphid facultative endosymbiont *Hamiltonella defensa* increases the protection of its host from parasitoid wasps (56).

In this minireview, a description of the currently available information about AAB recovered from insects is presented. The main insect sources so far for AAB are bees, mosquitoes, fruit flies, and sugarcane mealybugs; it is, however, likely that the number of insect species found to harbor AAB will significantly increase in the near future. Since the beginning of the 20th century, it was well known that gluconobacters, environmental sugar-loving microbes, represent a significant component of the honeybee microflora (65), while only more recently AAB were found to also inhabit mosquitoes, fruit flies, and leafhoppers.

The family *Acetobacteraceae* comprises a large variety of species. Since the description of *Acetobacter* Beijerinck (7) and *Gluconobacter* Asai (1), the taxonomy of AAB has undergone many changes, especially in the last 30 years. These changes are due to the new species ascribed to the family, the reclassification of strains (67), and the development and introduction of novel technologies for microbial taxonomy. Early classification systems were mainly based on the analyses of morphological and biochemical characteristics. Nowadays, taxonomic classification is based on what is called "polyphasic taxonomy," in which independent approaches such as phenotypic, chemotaxonomic, and genotyping analyses are combined (10). According to this classification approach, the AAB of the family *Acetobacteraceae* are classified into 13 genera: *Acetobacter*, *Gluconobacter*, *Gluconacetobacter*, *Acidomonas*, *Asaia*, *Kozakia* (40), *Swaminathania* (45), *Saccharibacter* (37), *Neosasaia* (69), *Granulibacter* (33, 34), *Commensalibacter* (60), *Tanticharoenia* (68), and *Ameyamaea* (70).

#### WHICH INSECT HOSTS DO AAB COLONIZE?

Bees were the first insects from which AAB were recovered (65). Bees and the products of beekeeping have been extensively studied over the years, taking into account the microbiological perspective (31). Among the great variety of microorganisms that were identified and isolated, *Gluconobacter*, *Gluconacetobacter*, and *Acetobacter* are some of the predominant bacterial groups of the bee microbiota (4, 36, 50, 51). Another hymenopteran, the endoparasitoid wasp *Asobara tabida*, hosts several members of the *Acetobacteraceae* family, represented by *Acetobacter* (*A.*) *pasteurianus* and *Acidomonas* (*Ac.*) *methanolica*, as reported by Zouache et al. (71).

*Drosophila melanogaster*, the most widely used insect model organism, hosts a rich AAB microbiome (11, 12, 58, 61). Analyzing the bacterial community associated with laboratory-reared and wild-captured *D. melanogaster* by establishing 16S rRNA gene libraries, Cox and Gilmore (12) showed that *Acetobacter* represented one of the most abundant genera, with 29% of the clones analyzed. The following species were identified: *A. acetii*, *A. cerevisiae*, *A. pasteurianus*, *A. pomorum*, and *A. peroxydans*, together with several species of *Gluconobacter*

and *Gluconacetobacter*. In the olive fruit fly *Bactrocera oleae*, which is phylogenetically close to *Drosophila*, *Acetobacter* symbionts also dominate, with the single species *A. tropicalis* (42). Combining cultivation-dependent and -independent techniques with ultrastructural and microscopic analyses, the authors detected the symbionts in all laboratory and field-collected individuals originating from different locations in Greece. The investigation revealed that several *A. tropicalis* strains can coexist in a single fly.

A recently studied AAB-insect symbiosis is the association between the cultivable *Asaia* spp. and the pathogen-transmitting mosquitoes *Anopheles* (*An.*) *stephensi*, *An. maculipennis*, *An. gambiae*, and *Aedes aegypti* (13, 22). *Asaia* sp. was found as a dominant bacterium within the insect microbial community by the use of several techniques. *Asaia* sp. was also identified in the bacterial community of the leafhopper *Scaphoideus titanus*, the vector of flavescence dorée in grapes (13, 46). Assessed by quantitative PCR, 16S rRNA genes of *Asaia* sp. constituted 4.9% of the total bacterial 16S rRNA gene copies per leafhopper. Ashbolt and Inkerman (2) detected the presence of AAB in another leafhopper, *Perkinsiella saccharidica*, for which an AAB community of  $5 \times 10^3$  cells per individual was estimated.

Other insects in which *Asaia* sp. symbionts were detected by cultivation-independent techniques are the hymenopteran *Marietta leopardina*, a hyperparasitoid (47), and the lepidopteron *Pieris rapae*, the cabbage white butterfly (59).

The pink sugarcane mealybug *Saccharicoccus sacchari* is a common insect found in sugarcane-growing countries which is able to host several members of the family *Acetobacteraceae* (2). At least  $10^6$  AAB are typically carried by an adult female mealybug actively feeding on aerial storage, whereas fewer than  $10^4$  bacteria per insect are maintained in less active adults and individuals from underground mealybug populations. AAB isolates from *S. sacchari* were identified as *A. acetii*, *Gluconacetobacter* (*Ga.*) *diazotrophicus*, *Ga. liquefaciens*, and *Ga. sacchari* by Franke et al. (26). Further experiments, though, led the authors to the conclusion that AAB are only a relatively small proportion of the microbial community in *S. sacchari* (25, 27, 28). The presence of AAB in other mealybugs like *Planococcus* sp. and *Dysmicoccus brevis* was also demonstrated (2).

A list of AAB that colonize insect hosts is given in Table 1. It should be emphasized that all of these insects rely on sugar-based diets such as nectars, fruit sugars, or phloem sap.

#### WHICH INSECT ORGANS DO AAB INHABIT?

A major habitat of insect-associated microorganisms is the digestive system, due to the availability of nutrients that are degraded by both host enzymes and microbial activity (17). The microbiota in an insect gut is influenced by structural and physiological factors, as well as by the quality of the ingested food. However, AAB symbionts have been found associated not only with the insect interior but also with the insect surface, as in the case of *D. melanogaster* (58), underlining their ability to survive in unfavorable environments. Growing on the surface of *Drosophila*, *Acetobacter* sp. can have a smaller-than-normal size, as shown by scanning electron microscopy analysis (58). AAB are able to survive other harsh conditions represented, for instance, by starvation or an adverse environment, entering into a viable but not culturable (VBNC) state and reducing the size of the microbial

TABLE 1. AAB inhabit several insect hosts belonging to different orders like Diptera, Hymenoptera, Hemiptera, and Homoptera

AAB and insect host(s)	Reference(s)
<i>Acetobacter</i> sp.	
<i>D. melanogaster</i> (Diptera: Drosophilidae).....	11, 12, 58, 61
<i>A. mellifera</i> (Hymenoptera: Apidae).....	4, 50
<i>B. oleae</i> (Diptera: Tephritidae).....	42
<i>A. tabida</i> (Hymenoptera: Braconidae).....	71
<i>S. sacchari</i> (Homoptera: Pseudococcidae).....	2
<i>C. intestini</i>	
<i>D. melanogaster</i> (Diptera: Drosophilidae).....	60, 61
<i>Gluconobacter</i> sp.	
<i>A. mellifera</i> (Hymenoptera: Apidae).....	4, 51
<i>D. melanogaster</i> (Diptera: Drosophilidae).....	11, 12, 58, 60, 61
<i>S. sacchari</i> (Homoptera: Pseudococcidae).....	2
<i>Gluconacetobacter</i> sp.	
<i>D. melanogaster</i> (Diptera: Drosophilidae).....	11, 12, 61
<i>A. mellifera</i> (Hymenoptera: Apidae).....	4, 36, 50
<i>S. sacchari</i> (Homoptera: Pseudococcidae).....	2, 25–28
<i>Asaia</i> sp.	
<i>Anopheles</i> sp. (Diptera: Culicidae).....	13, 15, 22
<i>A. aegypti</i> (Diptera: Culicidae).....	13
<i>S. titanus</i> (Hemiptera: Cicadellidae).....	13, 46
<i>M. leopardiana</i> (Hymenoptera: Aphelinidae).....	47
<i>P. rapae</i> (Lepidoptera: Pieridae).....	59
<i>Saccharibacter floricola</i>	
<i>A. mellifera</i> (Hymenoptera: Apidae).....	51

cells (49). Consequently, AAB populations from various substrates can be underestimated when the evaluation is performed only by a culture-based approach. The application of culture-independent techniques such as epifluorescence staining and real-time PCR overcomes this problem (6).

Simple or complex microbial consortia can inhabit the insect gut. For instance, in *D. melanogaster*, 25 phylotypes with just a few dominant bacterial species, among which are AAB, are associated with the insect gut (12). The anterior hindgut region is the most densely inhabited part of the digestive system, due to the availability of partially digested food coming from the midgut, as well as the products excreted by the Malpighian tubules. It has been estimated that  $10^8$  to  $10^9$  bacterial cells per g of gut content are present in the honeybee gut (39, 57), while  $>10^6$  bacteria are typically recovered from an entire old *Drosophila* fly (58).

The insect gastrointestinal tract (GIT), more structured in the adult stage than in the larval one, is organized similarly in bees, mosquitoes, and fruit flies. The mouth is followed by the esophagus, the ventriculus (stomach), the intestine, the rectum, and the anus. *Drosophila* flies also possess an acidic crop, together with an alkaline ventriculus and a slightly acidic hindgut, while mosquitoes have three diverticula that arise near the terminal end of the esophagus. The ventral one, also called a crop, is acidic and able to enlarge into the abdomen. A sugar meal such as floral nectar is stored in the diverticula and passes slowly to the midgut for the digestion step (63). Bees possess a honey stomach, an enlargement of the esophagus in which nectar is stored during flight and which ends with the proventriculus. This structure avoids contamination of the nectar with the contents of the stomach, which follows the proventriculus (66). The adult bee gut is normally acidic in comparison to the

larval one. The pH is reported to vary from 4.8 to 7.0, depending on the acidity of the ingested pollen. Mohr and Tebbe (50) reported pH values of approximately 4.0 for larvae fed with worker jelly, as well as with nectar and honey.

In the insect digestive system, AAB find a suitable environment in which they flourish and reproduce, mainly due to the aerobic environment, the acidic pH, and the presence of diet-derived sugars. Moreover, AAB establish a tight association with insect epithelial cells. AAB are known to produce polysaccharidic matrices that are involved in microbial interactions. One of the polysaccharides produced by bacteria is cellulose, which, in animal pathogens, is implicated in biofilm formation, in multicellular behavior, in adhesion to animal cells, or in stress tolerance (5). Several members of the family *Acetobacteraceae* possess the ability to produce cellulose. *Ga. xylinus* is the best-known cellulose producer, but other examples can be found, such as *Ga. kombuchae*, *Ga. swingsii*, *Ga. rhaeticus*, *Ga. nataicola*, and some strains of *Ga. hansenii*, *Ga. europeus*, and *Ga. oboediens* (21, 44). The presence of a cellulose operon has also been reported for a strain of *Asaia bogorensis* isolated from tropical flowers (NCBI accession no. AB355706).

Investigations by transmission electron microscopy (TEM) of the microbiome associated with the epithelia of mosquitoes and other insects showed an extensive mass of extracellular polysaccharidic matrix around the AAB symbionts (13, 22, 42, 62). This matrix establishes tight contact between the microbial cells and the host epithelium, implying a role in microbial interaction with host surfaces. Crotti et al. (13) also discussed the possibility that this extracellular matrix could protect the bacterial cell from adverse conditions such as alkaline or acidic pH or high osmolarity. Indeed, several members of the family





FIG. 1. (A) *Asaia* sp. colonizes the gut epithelium of *Anopheles* sp., *Ae. aegypti*, and *S. titanus*, establishing a specific association with the insect epithelium mediated by an extracellular polysaccharidic matrix that surrounds *Asaia* sp. cells. Bar, 3.4  $\mu\text{m}$ . (B) *Asaia* sp. produces a thick pellicle when cultured in a glass tube without shaking for 10 days. (C) *Asaia* sp. has been found to be strictly associated with the surface of immature eggs in *S. titanus* and *An. gambiae*, as shown by FISH with *Asaia* sp.-specific fluorescent probes (*Asaia* sp.-specific probes for *S. titanus* are shown in yellow), suggesting egg-smearing transmission of *Asaia* sp. in its hosts. Bar, 400  $\mu\text{m}$ .

*Acetobacteraceae* (*Tanticharoenia*, *Acidomonas*, *Asaia*, *Saccharibacter*, *Swaminathania*, and *Neosaia* spp.) show the ability to grow at high osmolarity (30% glucose), whereas only weak growth under this condition is reported for *Kozakia* sp. (69, 70).

*Asaia* sp., a dominating bacterium in the microbiota associated with *Anopheles* sp. and *Ae. aegypti* mosquitoes and with the leafhopper *S. titanus*, was analyzed by TEM and in situ hybridization (ISH), proving its ability to produce a gelatinous matrix around the bacterial cells in contact with the host gut epithelium (13, 22, Fig. 1A). The capability of this strain to colonize and lodge in the gut system was also demonstrated using strains labeled with fluorescent proteins (13, 22). It was shown that the symbiont produces a thick pellicle when grown in glass tubes without shaking (Fig. 1B), a feature typically found in AAB. For instance, when AAB occur as spoilers in bottled wine, a distinctive ring of bacterial biomass is deposited in the neck of the bottle at the interface between the wine and the air headspace (6). *Asaia* sp. is intimately associated not only with the insect gut epithelium but also with the epithelia of the reproductive systems of both females and males, as well as with the salivary glands, as shown by TEM, ISH, and colonization experiments with *Asaia* sp. strains tagged with fluorescent proteins. In particular, in *S. titanus*, *Asaia* sp. was shown to be harbored in the testicles, intermixed within the spermatid bundles and in the Malpighian tubules associated with brochosomes, while in mosquitoes, the male epithelial genital duct was observed to be lined by microcolonies of *Asaia* sp. cells (13, 22). In *B. oleae*, *A. tropicalis* was found to be tightly entrapped in a brown gelatinous matrix within the peritrophic membrane (42), confirming that the main site of microbial colonization in *B. oleae* is the gut lumen inside the peritrophic matrix (48). Also, *A. tropicalis*, when grown in pure culture in a glass tube without shaking, forms a thick pellicle at the liquid-air interface.

#### TRANSMISSION ROUTES OF AAB SYMBIONTS IN INSECTS

In order to ensure bacterial transmission to different individuals and the next generation, insects have evolved a wide array of mechanisms. Among these are transovarian transmission (i.e., through the mother oocytes) and post-hatch mechanisms like “egg smearing” (i.e., eggs are contaminated by the symbionts on the surface, and thus the hatched larvae acquire the symbionts by consuming or probing the eggshell) and “coprophagy” (i.e., preadult stage individuals feed on adult excrement, reacquiring symbionts), although other unique modes of symbiont transmission exist (3, 41).

Most of the knowledge on the transmission routes followed by AAB in their hosts derives from studies on *Asaia* sp. symbionts (13, 15). Colonization experiments performed by feeding *An. stephensi*, *An. gambiae*, *Ae. aegypti*, and *S. titanus* with green fluorescent protein- or DsRed-labeled *Asaia* sp. showed a horizontal transmission route with rapid colonization of the gut, salivary glands, and reproductive organs. *Asaia* sp. can also be passed horizontally through feeding between insects that belong to phylogenetically distant species. For instance, it was shown that *Asaia* sp. from *Anopheles* spp. is able to cross colonize other sugar-feeding insects like *Ae. aegypti* and the hemipteran *S. titanus* (13).

These investigations showed that several routes occur simultaneously (15). Environmental acquisition appears to be an important way of transmission, aided by the ubiquitous nature of the symbiont, but transmission from the mother to the offspring and from the father to the mother and then to the offspring also occurs (15). An egg-smearing mode of vertical transmission of *Asaia* sp. in *S. titanus* was proven by fluorescence ISH (FISH) and colonization experiments with strains labeled with fluorescent proteins (Fig. 1C) (13). Dur-

ing ovarian egg development, an increasing ordered disposition of the superficial bacterial biomass is observable: from a more disperse pattern at the initial developmental stages, the bacterial cells are finally confined to the apical egg regions.

Transfer of *Asaia* sp. from larvae to pupae and from pupae to adults was also demonstrated (15). It is likely that during insect metamorphosis, *Asaia* sp. can be transtadially transmitted and escapes the reduction/removal of midgut bacteria that normally occurs during development, perhaps residing at other sites (and thus not being included in the meconium), or being resistant to the antimicrobial exuvial fluid that is ingested as part of the ecdysial process (52).

### THE INSECT BODY CAN BE AN APPROPRIATE NICHE FOR AAB

The presence of AAB in the insect digestive system and in other organs reflects the physiochemical properties of these habitats (such as the type of food ingested, redox conditions, and pH) and the metabolic responses and capabilities of these microorganisms (17).

The distribution of arthropod-associated AAB in the different insect species reflects their diverse preference of carbon sources acquired by insect nutrition. *Gluconobacter* sp., preferring sugars, is generally isolated from honeybees, while ethanol-loving *Acetobacter* sp. is mainly recovered from *Drosophila* and *Bactrocera* flies. Nectar and honey are basically sugar solutions composed of sucrose, glucose, and fructose (9, 64), thus being favorable substrates for *Gluconobacter* sp. On the other hand, *Drosophila* flies are attracted by fermenting fruits that are a more suitable environment for acetobacters. *Drosophila* flies are mostly fungivores, and their association with fruit is indirect in that they primarily eat yeasts that live in rotting fruit. *Drosophila* flies are also called vinegar flies because they are attracted to the acidic odor of fermentation. Moreover, insects from which AAB are usually recovered have sugar-rich diets that are unbalanced for nitrogen content, so they have to fulfill the nitrogen requirement from other sources, such as the atmosphere (55). Among AAB, several examples of N<sub>2</sub>-fixing bacteria have been reported, such as *Ga. diazotrophicus* (32, 66), *Ga. johannae*, *Ga. azotocaptans* (30), *Ga. kombuchae* (21), *A. peroxydans* (54), *A. nitrogenifigens* (20), and *Swaminathanian salitolerans* (45). However, further investigations are needed to evaluate if insect-associated AAB could contribute to insect nitrogen metabolism or recycling. The ability of several AAB strains to grow on media devoid of vitamins (16) also suggests the possibility that AAB in insects could synthesize vitamins or cofactors utilized by the host. *A. pasteurianus*, *A. peroxydans*, *A. estunensis*; *Ga. liquefaciens*, *As. bogorensis* (43), and *Asaia* sp. isolated from *An. stephensi* have been showed to be prototrophic with respect to vitamins. On the other hand, experimental removal of AAB from their insect hosts (as in the case of the *Asaia-Anopheles* system) did not appear to be detrimental to the host, underlining the secondary symbiont status of these bacteria.

The GIT and circulatory system of AAB-associated insects lack anoxic niches (8, 12). Studies on the gut microbiota associated with *Drosophila* or *Apis* spp. confirmed the absence of obligate anaerobic bacteria and the presence of aerobic, fac-

ultatively aerobic, or aerotolerant bacteria (12, 50). AAB have a respiratory metabolism that requires oxygen as the final electron acceptor, but they are also able to survive in environments with low oxygen availability (19, 38, 49). Hence, they can find a favorable environment in the insect gut.

Within the insect, AAB, as well as other symbiotic microorganisms, have to face the host innate immune system (35). According to the results obtained by Ryu et al. (61), the AAB microbiome is involved in modulating *Drosophila* immunity. The authors described a delicate equilibrium between the gut commensals and the fly innate immune system. The normal flora in the fly gut was sufficient to suppress the growth of pathogenic bacteria, maintaining the pathogenic commensal *Gluconobacter (G.) morbifer* at a low level. When the system was perturbed, an increased number of *G. morbifer* bacteria led to gut apoptosis. Hence, in healthy individuals, the immune system allows the dominance of two AAB strains (*A. pomorum* and *Commensalibacter intestini*), which suppress the proliferation of *G. morbifer* by competition. Another study gained insight into the interaction between the acetic acid bacterium *Asaia* sp. and the innate immune system of *An. gambiae* (18). When the expression of a host gene involved in the innate immune response (AgDscam) gene was silenced, *Asaia* sp., whose favorable site of colonization is the gut (22), was no longer controlled by the innate immune system and proliferated massively in the host hemolymph (18).

AAB are considered fastidious microorganisms due to their difficult isolation and cultivation on synthetic media, regardless of the great number of media proposed for their growth (6). Moreover, they have been shown to be capable of entering the VBNC state (6, 49). It was not possible to isolate *Asaia* sp. from *S. titanus* leafhoppers, in spite of the well-documented presence of the symbiont in the insect as assayed with several cultivation-independent techniques (13). The authors speculated that the failure to isolate *Asaia* sp. might have been due to the unfavorable environment represented by *S. titanus* in a not actively feeding state or due to the development of unknown nutritional requirements by the bacteria.

### CONCLUDING REMARKS AND PERSPECTIVES

For a long time, AAB have been considered environmental and ubiquitous bacteria. However, an increasing number of reports of stable associations of AAB with insects and the fact that they follow several transmission routes for their propagation in the next insect generation indicate that insect-associated AAB cannot be considered just environmental microorganisms but are also symbionts of the insect body, where they occupy a specific favorable niche. Phylogenetic studies of insect-associated AAB and those recovered from the environment do not show phylogenetic congruence of the insect-associated AAB (Fig. 2). This suggests that AAB have been acquired from the environment by their insect hosts recently (14). Moreover, not being essential for host survival, AAB have to be considered secondary symbionts of insects.

Clarification of the function(s) exerted by the bacteria in and for their hosts will be a major step toward understanding the bacterium-insect association. AAB are probably involved in many aspects of insect biology, such as (i) the host




FIG. 2. Phylogenetic positions of insect-associated AAB among the most representative members of family *Acetobacteraceae*. The tree shown is based on bacterial 16S rRNA gene sequences. AAB recovered from insects are in bold. Shaded species belong to clusters including environmental AAB, as well as those recovered from insects. Values at nodes are percentages of bootstrap replications calculated from 2,000 replicate trees. Accession numbers of reference sequences are in parentheses. The species used as the outgroup belongs to the *Gamma*proteobacteria taxon. The scale bar represents 10% sequence divergence.

metabolism by supplying nutrients or by oxidizing certain substrates, (ii) defense against harmful microorganisms by decreasing the gut pH or by competitive exclusion, (iii) contributing to the maintenance of host gut homeostasis

(vi), (iv) interference with cell-to-cell communication through the production of volatile compounds, and (v) maintaining an equilibrium of the microbial consortia by supplying metabolites to other microbes beneficial to the



host. Genome sequencing of the insect-associated AAB will be a particularly valuable tool in this context.

Not only are further studies needed to clarify the role of insect-associated AAB, but efforts should also be directed toward understanding AAB biological diversity and behavior in the host. An interesting topic could be the investigation of body invasion mechanisms adopted by AAB, whether they follow tissue tropism, as shown for the alphaproteobacterial symbiont *Wolbachia* sp. (29).

Symbiont biology receives increasing attention since insect symbionts can potentially be used to control vector-borne diseases or suppress insect pests. AAB could be used as agents for natural or paratransgenic symbiotic control. Bacteria like *Asaia* spp. or *A. tropicalis* possess features interesting in this context, like easy cultivability, preservability, transformability, and dominance and prevalence in the insect microbiome. Moreover, vertical and horizontal transmission routes and cross-colonizing capability could ensure the spread of these biological agents through host populations (23).

#### ACKNOWLEDGMENTS

We thank for financial support the European Union in the ambit of project BIODESERT (European Community's Seventh Framework Programme CSA-SA REGPOT-2008-2 under grant agreement 245746) and the Italian Ministry for Research (MIUR) in the ambit of project PRIN 2007 Caratterizzazione del microbiota associato a *Scaphoideus titanus* e *Hyalesthes obsoletus*, cicaline vettrici di fitoplasmi nella vite ed isolamento e studio della localizzazione di batteri acetici simbiotici. K.B. is grateful to the European Union (European Community's Seventh Framework Programme CSA-SA\_REGPROT-2007-1 under grant agreement 203590), the International Atomic Energy Agency, and the University of Ioannina, which have supported the research in his laboratory. C.B., K.B., A.C., E.C., and D.D. benefited from travel grants from Cost Action FA0701, Arthropod Symbiosis: from Fundamental Studies to Pest and Disease Management.

We thank Stefan Oehler for his comments.

#### REFERENCES

- Asai, T. 1935. Taxonomic studies on acetic acid bacteria and allied oxidative bacteria isolated from fruits. A new classification of the oxidative bacteria. *J. Agric. Chem. Soc. Jpn.* **11**:674–708. (In Japanese.)
- Ashbolt, N. J., and P. A. Inkeram. 1990. Acetic acid bacteria biota of the pink sugar cane mealybug, *Saccharococcus sacchari*, and its environs. *Appl. Environ. Microbiol.* **56**:707–712.
- Attardo, G. M., C. Lohs, A. Heddi, U. H. Alam, S. Yildirim, and S. Aksoy. 2008. Analysis of milk gland structure and function in *Glossina morsitans*: milk protein production, symbiont populations and fecundity. *J. Insect Physiol.* **54**:1236–1242.
- Babendreier, D., D. Joller, J. Romeis, F. Bigler, and F. Widmer. 2007. Bacterial community structures in honeybee intestines and their response to two insecticidal proteins. *FEMS Microbiol. Ecol.* **59**:600–610.
- Barak, J. D., C. E. Jahn, D. L. Gibson, and A. O. Charkowsky. 2007. The role of cellulose and O-antigen capsule in the colonization of plants by *Salmonella enterica*. *Mol. Plant Microbe Interact.* **20**:1083–1091.
- Bartowsky, E. J., and P. A. Henschke. 2008. Acetic acid bacteria spoilage of bottled red wine—a review. *Int. J. Food Microbiol.* **125**:60–70.
- Beijerinck, M. W. 1898. Über die Arten der Essigbakterien. *Zentralbl. Bakteriol. Naturwiss.* **4**:209–216.
- Bodenstein, D., K. W. Cooper, G. F. Ferris, A. Miller, D. F. Poulson, and B. P. Sonnenblick. 1950. *Biology of Drosophila*. John Wiley & Sons, Inc., New York, NY.
- Chalcoff, V. R., M. A. Aizen, and L. Galetto. 2006. Nectar concentration and composition of 26 species from the temperate forest of South America. *Ann. Bot.* **97**:413–421.
- Cleenwerck, I., and P. De Vos. 2008. Polyphasic taxonomy of acetic acid bacteria: an overview of the currently applied methodology. *Int. J. Food Microbiol.* **125**:2–14.
- Corby-Harris, V., A. C. Pontaroli, L. J. Shimkets, J. L. Bennetzen, K. E. Habel, and D. E. Promislow. 2007. Geographical distribution and diversity of bacteria associated with natural populations of *Drosophila melanogaster*. *Appl. Environ. Microbiol.* **73**:3470–3479.
- Cox, C., and M. Gilmore. 2007. Native microbial colonization of *Drosophila melanogaster* and its use as a model of *Enterococcus faecalis* pathogenesis. *Infect. Immun.* **75**:1565–1576.
- Crotti, E., C. Damiani, M. Pajoro, E. Gonella, A. Rizzi, I. Ricci, I. Negri, P. Scuppa, P. Rossi, P. Ballarini, N. Raddadi, M. Marzorati, L. Sacchi, E. Clementi, M. Genchi, M. Mandrioli, C. Bandi, G. Favia, A. Alma, and D. Daffonchio. 2009. *Asaia*, a versatile acetic acid bacterial symbiont, capable of cross-colonizing insects of phylogenetically-distant genera and orders. *Environ. Microbiol.* **11**:3252–3264.
- Dale, C., and N. Moran. 2006. Molecular interactions between bacterial symbionts and their hosts. *Cell* **126**:453–465.
- Damiani, C., I. Ricci, E. Crotti, P. Rossi, A. Rizzi, P. Scuppa, F. Esposito, C. Bandi, D. Daffonchio, and G. Favia. 2008. Paternal transmission of symbiotic bacteria in malaria vectors. *Curr. Biol.* **18**:R1087–R1088.
- De Ley, J., and J. Frateur. 1974. Genus *Acetobacter* Beijerinck, 1898, p. 276–278. In R. E. Buchanan and N. E. Gibbons (ed.), *Bergey's manual of determinative bacteriology*, 8th ed. The Williams & Wilkins Co., Baltimore, MD.
- Dillon, R. J., and V. M. Dillon. 2004. The gut bacteria of insects: non-pathogenic interactions. *Annu. Rev. Entomol.* **49**:71–92.
- Dong, Y., H. E. Taylor, and G. Dimopoulos. 2006. AgDscam, a hypervariable immunoglobulin domain-containing receptor of the *Anopheles gambiae* innate immune system. *PLoS Biol.* **4**:1137–1146.
- Du Toit, W. J., and I. S. Pretorius. 2002. The occurrence, control and esoteric effect of acetic acid bacteria in winemaking. *Ann. Microbiol.* **52**:155–179.
- Dutta, D., and R. Gachhui. 2006. Novel nitrogen-fixing *Acetobacter nitrogenifigens* sp. nov., isolated from Kombucha tea. *Int. J. Syst. Evol. Microbiol.* **56**:1899–1903.
- Dutta, D., and R. Gachhui. 2007. Nitrogen-fixing and cellulose-producing *Gluconacetobacter kombuchae* sp. nov., isolated from Kombucha tea. *Int. J. Syst. Evol. Microbiol.* **57**:353–357.
- Favia, G., I. Ricci, C. Damiani, N. Raddadi, E. Crotti, M. Marzorati, A. Rizzi, R. Urso, L. Brusetti, S. Borin, D. Mora, P. Scuppa, L. Pasqualini, E. Clementi, M. Genchi, S. Corona, I. Negri, G. Grandi, A. Alma, L. Kramer, F. Esposito, C. Bandi, L. Sacchi, and D. Daffonchio. 2007. Bacteria of the genus *Asaia* stably associate with *Anopheles stephensi*, an Asian malarial mosquito vector. *Proc. Natl. Acad. Sci. U. S. A.* **104**:9047–9051.
- Favia, G., I. Ricci, M. Marzorati, I. Negri, A. Alma, L. Sacchi, C. Bandi, and D. Daffonchio. 2008. Bacteria of the genus *Asaia*: a potential paratransgenic weapon against malaria. *Adv. Exp. Med. Biol.* **627**:49–59.
- Feldhaar, H., and R. Gross. 2009. Insects as hosts for mutualistic bacteria. *Int. J. Med. Microbiol.* **299**:1–8.
- Franke, I. H., M. Fegan, C. Hayward, G. Leonard, and L. I. Sly. 2000. Molecular detection of *Gluconacetobacter sacchari* associated with the pink sugarcane mealybug *Saccharicoccus sacchari* (Cockerell) and the sugarcane leaf sheath microenvironment by FISH and PCR. *FEMS Microbiol. Ecol.* **31**:61–71.
- Franke, I. H., M. Fegan, C. Hayward, G. Leonard, E. Stackebrandt, and L. I. Sly. 1999. Description of *Gluconacetobacter sacchari* sp. nov., a new species of acetic acid bacterium isolated from the leaf sheath of sugar cane and from the pink sugar-cane mealy bug. *Int. J. Syst. Bacteriol.* **49**:1681–1693.
- Franke-Whittle, I. H., M. G. O'Shea, G. J. Leonard, and L. I. Sly. 2004. Molecular investigation of the microbial populations of the pink sugarcane mealybug, *Saccharicoccus sacchari*. *Ann. Microbiol.* **54**:455–470.
- Franke-Whittle, I. H., M. G. O'Shea, G. J. Leonard, and L. I. Sly. 2005. Design, development, and use of molecular primers and probes for the detection of *Gluconacetobacter* species in the pink sugarcane mealybug. *Microb. Ecol.* **50**:128–139.
- Frydman, H. M., J. M. Li, D. N. Robson, and E. Wieschaus. 2006. Somatic stem cell niche tropism in *Wolbachia*. *Nature* **441**:509–512.
- Fuentes-Ramírez, L. E., R. Bustillos-Cristales, A. Tapia-Hernandez, T. Jimenez-Salgado, E. T. Wang, E. Martínez-Romero, and J. Caballero-Melgado. 2001. Novel nitrogen-fixing acetic acid bacteria, *Gluconacetobacter johannae* sp. nov. and *Gluconacetobacter azotocaptans* sp. nov., associated with coffee plants. *Int. J. Syst. Evol. Microbiol.* **51**:1305–1314.
- Gilliam, M. 1997. Identification and roles of non-pathogenic microflora associated with honey bees. *FEMS Microbiol. Lett.* **155**:1–10.
- Gillis, M., K. Kersters, B. Hoste, D. Janssens, R. M. Kroppenstedt, M. P. Stephan, K. R. S. Teixeira, J. Dobreiner, and J. De Ley. 1989. *Acetobacter diazotrophicus* sp. nov., a nitrogen-fixing acetic acid bacterium associated with sugarcane. *Int. J. Syst. Bacteriol.* **39**:361–364.
- Greenberg, D. E., L. Ding, A. M. Zelazny, F. Stock, A. Wong, V. L. Anderson, G. Miller, D. E. Kleiner, A. R. Tenorio, L. Brinster, D. W. Dorward, P. R. Murray, and S. M. Holland. 2006. A novel bacterium associated with lymphadenitis in a patient with chronic granulomatous disease. *PLoS Pathog.* **2**:e28.
- Greenberg, D. E., S. F. Porcella, F. Stock, A. Wong, P. S. Conville, P. R. Murray, S. M. Holland, and A. M. Zelazny. 2006. *Granulibacter bethesdensis* gen. nov., sp. nov., a distinctive pathogenic acetic acid bacterium in the family Acetobacteraceae. *Int. J. Syst. Evol. Microbiol.* **56**:2609–2616.

35. Gross, R., F. Vavre, A. Heddi, G. D. Hurst, E. Zchori-Fein, and K. Bourtzis. 2009. Immunity and symbiosis. *Mol. Microbiol.* **73**:751–759.
36. Jeyaprakash, A., M. A. Hoy, and M. H. Allsopp. 2003. Bacterial diversity in worker adults of *Apis mellifera capensis* and *Apis mellifera scutellata* (Insecta: Hymenoptera) assessed using 16S rRNA sequences. *J. Invertebr. Pathol.* **84**:96–103.
37. Jojima, Y., Y. Mihara, S. Suzuki, K. Yokozeki, S. Yamanaka, and R. Fudou. 2004. *Saccharibacter floricola* gen. nov., sp. nov., a novel osmophilic acetic acid bacterium isolated from pollen. *Int. J. Syst. Evol. Microbiol.* **54**:2263–2267.
38. Joyeux, A., S. Lafon-Lafourcade, and P. Ribereau-Gayon. 1984. Evolution of acetic acid bacteria during fermentation and storage of wine. *Appl. Environ. Microbiol.* **48**:153–156.
39. Kacaniova, M., R. Chlebo, M. Kopernicky, and A. Trakovicka. 2004. Microflora of the honeybee gastrointestinal tract. *Folia Microbiol.* **49**:169–171.
40. Kersters, K., P. Lisdiyanti, K. Komagata, and J. Swings. 2006. The family Acetobacteraceae: the genera *Acetobacter*, *Acidomonas*, *Asaia*, *Gluconacetobacter*, *Gluconobacter*, and *Kozakia*, p. 163–200. In M. Dworkin, S. Falkow, E. Rosenberg, K.-H. Schleifer, and E. Stackebrandt (ed.), *The prokaryotes*, 3rd ed., vol. 5. Springer, New York, NY.
41. Kikuchi, Y., and T. Fukatsu. 2009. Insect-bacterium mutualism without vertical transmission, p. 143–161. In K. Bourtzis and T. A. Miller (ed.), *Symbiosis*, vol. 3. CRC Press, Boca Raton, FL.
42. Kounatidis, I., E. Crotti, P. Sapountzis, L. Sacchi, A. Rizzi, B. Chouaia, C. Bandi, A. Alma, D. Daffonchio, P. Mavragani-Tsipidou, and K. Bourtzis. 2009. *Acetobacter tropicalis* is a major symbiont in the olive fruit fly *Bactrocera oleae*. *Appl. Environ. Microbiol.* **75**:3281–3288.
43. Lisdiyanti, P., H. Kawasaki, T. Seki, Y. Yamada, T. Uchimura, and K. Komagata. 2000. Systematic study of the genus *Acetobacter* with descriptions of *Acetobacter indonesiensis* sp. nov., *Acetobacter tropicalis* sp. nov., *Acetobacter orleanensis* (Henneberg 1906) comb. nov., *Acetobacter lovaniensis* (Frater 1950) comb. nov., and *Acetobacter estunensis* (Carr 1958) comb. nov. *J. Gen. Appl. Microbiol.* **46**:147–165.
44. Lisdiyanti, P., R. R. Navarro, T. Uchimura, and K. Komagata. 2006. Re-classification of *Gluconacetobacter hansenii* strains and proposals of *Gluconacetobacter saccharivorans* sp. nov. and *Gluconacetobacter nataicola* sp. nov. *Int. J. Syst. Evol. Microbiol.* **56**:2101–2111.
45. Loganathan, P., and S. Nair. 2004. *Swaminathania salitolerans* gen. nov., sp. nov., a salt-tolerant, nitrogen-fixing and phosphate-solubilizing bacterium from wild rice (*Porteresia coarctata* Tateoka). *Int. J. Syst. Evol. Microbiol.* **54**:1185–1190.
46. Marzorati, M., A. Alma, L. Sacchi, M. Pajoro, S. Palermo, L. Brusetti, N. Raddadi, A. Balloi, R. Tedeschi, E. Clementi, S. Corona, F. Quaglino, P. A. Bianco, T. Beninati, C. Bandi, and D. Daffonchio. 2006. A novel *Bacteroidetes* symbiont is localized in *Scaphoideus titanus*, the insect vector of flavescence dorée in *Vitis vinifera*. *Appl. Environ. Microbiol.* **72**:1467–1475.
47. Matalon, Y., N. Katzir, Y. Gottlieb, V. Portnoy, and E. Zchori-Fein. 2007. *Cardinium* in *Plagiomerus diaspidis* (Hymenoptera: Encyrtidae). *J. Invertebr. Pathol.* **96**:106–108.
48. Mazzon, L., A. Piscedda, M. Simonato, I. Martinez-Sañudo, A. Squartini, and V. Girolami. 2008. Presence of specific symbiotic bacteria in flies of the subfamily Tephritinae (Diptera Tephritidae) and their phylogenetic relationships: proposal of 'Candidatus Stammerula tephritidis'. *Int. J. Syst. Evol. Microbiol.* **58**:1277–1287.
49. Millet, V., and A. Lonvaud-Funel. 2000. The viable but non-culturable state of wine micro-organisms during storage. *Lett. Appl. Microbiol.* **30**:136–141.
50. Mohr, K. I., and C. C. Tebbe. 2006. Diversity and phylotype consistency of bacteria in the guts of three bee species (Apoidea) at an oilseed rape field. *Environ. Microbiol.* **8**:258–272.
51. Mohr, K. I., and C. C. Tebbe. 2007. Field study results on the probability and risk of a horizontal gene transfer from transgenic herbicide-resistant oilseed rape pollen to gut bacteria of bees. *Appl. Microbiol. Biotechnol.* **75**:573–582.
52. Moll, R. M., W. S. Romoser, M. C. Modrzakowski, A. C. Moncayo, and K. Lerdthusnee. 2001. Meconial peritrophic membranes and the fate of midgut bacteria during mosquito (Diptera: Culicidae) metamorphosis. *J. Med. Entomol.* **38**:29–32.
53. Moran, N. A. 2006. Symbiosis. *Curr. Biol.* **16**:R866–R871.
54. Muthukumarasamy, R., I. Cleenwerck, G. Revathi, M. Vadivelu, D. Janssens, B. Hoste, K. U. Gum, K. D. Park, C. Y. Son, T. Sa, and J. Caballero-Mellado. 2005. Natural association of *Gluconacetobacter diazotrophicus* and diazotrophic *Acetobacter peroxydans* with wetland rice. *Syst. Appl. Microbiol.* **28**:277–286.
55. Nardi, J. B., R. I. Mackie, and J. O. Dawson. 2002. Could microbial symbionts of arthropod guts contribute significantly to nitrogen fixation in terrestrial ecosystems? *J. Insect Physiol.* **48**:751–763.
56. Oliver, K. M., J. A. Russell, N. A. Moran, and M. S. Hunter. 2003. Facultative bacterial symbionts in aphids confer resistance to parasitic wasps. *Proc. Natl. Acad. Sci. U. S. A.* **100**:1803–1807.
57. Rada, V., M. Machova, J. Huk, M. Marounek, and D. Duskova. 1997. Microflora in the honeybee digestive tract: counts, characteristics and sensitivity to veterinary drugs. *Apidologie* **28**:357–365.
58. Ren, C., P. Webster, S. E. Finkel, and J. Tower. 2007. Increased internal and external bacterial load during *Drosophila* aging without life-span trade-off. *Cell Metab.* **6**:144–152.
59. Robinson, C. J., P. Schloss, Y. Ramon, K. Raffa, and J. Handelsman. 2010. Robustness of the bacterial community in the cabbage white butterfly larval midgut. *Microb. Ecol.* **59**:199–211.
60. Roh, S. W., Y.-D. Nam, H.-W. Chang, K.-H. Kim, M.-S. Kim, J.-H. Ryu, S.-H. Kim, W.-J. Lee, and J.-W. Bae. 2008. Characterization of two novel commensal bacteria involved with innate immune homeostasis in *Drosophila melanogaster*. *Appl. Environ. Microbiol.* **74**:6171–6177.
61. Ryu, J.-H., S.-H. Kim, H.-Y. Lee, J. Y. Bai, Y.-D. Nam, J.-W. Bae, D. G. Lee, S. C. Shin, E.-M. Ha, and W.-J. Lee. 2008. Innate immune homeostasis by the homeobox gene *Caudal* and commensal-gut mutualism in *Drosophila*. *Science* **319**:777–782.
62. Sacchi, L., M. Genchi, E. Clementi, E. Bigliardi, A. M. Avanzati, M. Pajoro, I. Negri, M. Marzorati, E. Gonella, A. Alma, D. Daffonchio, and C. Bandi. 2008. Multiple symbiosis in the leafhopper *Scaphoideus titanus* (Hemiptera: Cicadellidae): details of transovarial transmission of *Cardinium* sp. and yeast-like endosymbionts. *Tissue Cell* **40**:231–242.
63. Thompson, M. T. 1905. Alimentary canal of the mosquito. *Proc. Boston Soc. Nat. Hist.* **32**:145–202.
64. White, J. W., Jr., M. L. Riethof, M. H. Subers, and I. Kushnir. 1962. Composition of American honeys. Technical bulletin 1261. Agricultural Research Service, U.S. Department of Agriculture, Washington, DC.
65. White, P. B. 1921. The normal bacterial flora of the bee. *J. Pathol. Bacteriol.* **24**:64–78.
66. Winston, M. L. 1987. The biology of the honey bee. Harvard University Press, Cambridge, MA.
67. Yamada, Y., K.-I. Hoshino, and T. Ishikawa. 1997. The phylogeny of acetic acid bacteria based on the partial sequences of 16S ribosomal RNA: the elevation of the subgenus *Gluconoacetobacter* to the generic level. *Biosci. Biotechnol. Biochem.* **61**:1244–1251.
68. Yukphan, P., T. Malimas, Y. Muramatsu, M. Takahashi, S. Tanasupawat, Y. Nakagawa, K. W. Suzuki, and Y. Yamada. 2008. *Tanticharoenia sakaeratisensis* gen. nov., sp. nov., a new osmotolerant acetic acid bacterium in the alpha-Proteobacteria. *Biosci. Biotechnol. Biochem.* **72**:672–676.
69. Yukphan, P., T. Malimas, W. Potacharoen, S. Tanasupawat, M. Tanticharoen, and Y. Yamada. 2005. *Neoasaia Chiangmaiensis* gen. nov., sp. nov., a novel osmotolerant acetic acid bacterium in the alpha-Proteobacteria. *J. Gen. Appl. Microbiol.* **51**:301–311.
70. Yukphan, P., T. Malimas, Y. Muramatsu, M. Takahashi, M. Kaneyasu, W. Potacharoen, S. Tanasupawat, Y. Nakagawa, K. Hamana, Y. Tahara, K. Suzuki, M. Tanticharoen, and Y. Yamada. 2009. *Ameayamaea Chiangmaiensis* gen. nov., sp. nov., an acetic acid bacterium in the alpha-Proteobacteria. *Biosci. Biotechnol. Biochem.* **73**:2156–2162.
71. Zouache, K., D. Voronin, V. Tran-Van, and P. Mavingui. 2009. Composition of bacterial communities associated with natural and laboratory populations of *Asobara tabida* infected with *Wolbachia*. *Appl. Environ. Microbiol.* **75**:3755–3764.