



Published in final edited form as:

*J Am Chem Soc.* 2009 October 21; 131(41): 14620–14621. doi:10.1021/ja906186f.

## PCR with an Expanded Genetic Alphabet

Denis A. Malyshev<sup>†</sup>, Young Jun Seo<sup>†</sup>, Phillip Ordoukhanian<sup>‡</sup>, and Floyd E. Romesberg<sup>†,\*</sup>

<sup>†</sup>Department of Chemistry, The Scripps Research Institute, 10550 North Torrey Pines Road, La Jolla, CA, 92037, USA.

<sup>‡</sup>Center for Protein and Nucleic Acid Research, The Scripps Research Institute, 10550 North Torrey Pines Road, La Jolla, CA, 92037, USA.

### Abstract



Expansion of the genetic alphabet with a third base pair would lay the foundation of a semi-synthetic organism with an expanded genetic code and also have immediate *in vitro* applications. Previously, the unnatural base pairs formed between **d5SICS** and either **dNaM** or **dMMO2** were shown to be well replicated by DNA polymerases under steady-state conditions and also transcribed by T7 RNA polymerase efficiently in either direction. We now demonstrate that DNA containing either the **d5SICS-dNaM** or **d5SICS-dMMO2** unnatural base pair may be PCR amplified with fidelities and efficiencies that approach those of fully natural DNA. These results further demonstrate that the determinants of a functional unnatural base pair may be designed into predominantly hydrophobic nucleobases with no structural similarity to the natural purines or pyrimidines. Importantly, the results reveal that the unnatural base pairs may function within an expanded genetic alphabet and make possible many *in vitro* applications.

The genetic alphabet is constrained by the efficient polymerase-mediated replication of DNA and RNA containing the two natural base pairs. In addition to laying the foundation for a semi-synthetic organism with an expanded genetic code, an efficiently and selectively replicated and transcribed unnatural base pair would dramatically increase the potential of the already ubiquitous *in vitro* methodologies based on DNA and RNA, and their sequence specific amplification. PCR amplification of DNA containing unnatural base pairs was first reported by Benner and co-workers, using pairs with orthogonal H-bonding complementarity,<sup>1</sup> followed by Hirao and co-workers,<sup>2</sup> using a pair formed between substituted pyrimidine and pyrrole nucleotide analogs. These studies represent landmarks in the effort to expand the genetic alphabet, but the unnatural pairs used are limited by strong sequence dependencies and/or inefficient transcription into RNA. This may not limit their use in some *in vitro* applications, but does reduce their generality and preclude their eventual use *in vivo* as part of a semi-synthetic organism.

We have focused on developing unnatural base pairs formed between predominantly hydrophobic nucleobases that have no structural homology to the natural nucleobases, and that pair based on hydrophobic and packing forces. Screening of a library of nucleotides, followed by hit optimization, identified the pair formed between **d5SICS** and **dMMO2** (Figure 1), which is relatively well recognized by different DNA polymerases under steady-state, single

floyd@scripps.edu.

**Supporting Information Available:** Details of template synthesis and PCR analysis. This information is available free of charge via the Internet at <http://www.pubs.acs.org>.

nucleotide incorporation conditions.<sup>3a</sup> Further optimization identified dNaM, which pairs with d5SICS to form an unnatural pair that is even better replicated under the same steady state conditions.<sup>3b</sup> Importantly, d5SICS-dMMO2, and especially d5SICS-dNaM, are also efficiently transcribed in both directions,<sup>3c</sup> suggesting that they might have immediate practical applications. Here, we examine whether these unnatural base pairs, which are much less natural-like than those previously examined, are also sufficiently well amplified for practical use.

Nucleosides were synthesized and converted to the corresponding triphosphates or phosphoramidites, and the phosphoramidites were incorporated into DNA duplexes D1–D6 using automated DNA synthesis (Figure 1 and Supporting Information). The duplexes are 134 to 149 nucleotides in length with a single, centrally positioned d5SICS-dMMO2 (D1) or d5SICS-dNaM (D2–D6) pair. D1 is similar to the duplex used by Hirao *et al.*,<sup>2</sup> while D2–D5 systematically vary the flanking dG-dC base pairs to probe for sequence dependencies among sequences that are relatively challenging to PCR amplify.<sup>4</sup> D6 contains randomized nucleotides to further explore sequence-specific effects and D7 is identical to D1 but with the unnatural base pair replaced by a natural dA-dT. Using a gel-based assay, we first explored PCR amplification of D1 using exonuclease proficient DeepVent DNA polymerase in the presence and absence of d5SICSTP and dNaMTP (note that this results in the replacement of dMMO2 with dNaM during the first round of replication). Promisingly, PCR conditions were easily identified where DNA was amplified only when both natural and unnatural triphosphates were present (Supporting Information).

To better characterize amplification, we determined the yields of PCR product after 14 cycles, starting with 1 ng of D1–D6, dNTPs, d5SICSTP, and dNaMTP (Table 1). The 424-fold amplification of D1 was the highest and compares favorably with the 556-fold amplification of the control D7. Amplification levels of the other duplexes were slightly lower, likely due to their GC-content, but remained greater than 100-fold, except for D3 and D5, which were amplified 74- and 35-fold, respectively. The lower efficiency observed with D3 and D5 is not surprising considering that they position the unnatural base pair within a dG:dC run, which is particularly challenging to amplify by PCR.<sup>4</sup> While amplification levels may be increased with additional rounds of PCR (see below), the data suggest that DNA containing the unnatural base pairs may be amplified with an efficiency that is sufficient for *in vitro* applications.

To better characterize fidelity, the amplicons were sequenced (Table 1). In most cases, standard sequencing reactions (ABI 3730 DNA Analyzer) lacking unnatural triphosphates terminated at the unnatural nucleotide. In these cases, the fidelities (*i.e.* the percentage of unnatural base pair retention per doubling) are reported as the average determined by sequencing both amplified strands (Supporting Information). For D5, significant read-through was observed in one direction. However, because read-through was also observed with chemically synthesized control strands, we conclude that it results from sequencing and not from PCR amplification. In this case, the reported fidelity was determined from one strand context. Remarkably, in four of the six sequence contexts examined the average fidelity is at least 99%. It is slightly lower in D3 (98.5%) and D5 (98.0%), which again likely results from their particularly difficult, GC-rich sequence context. Despite the expected challenges associated with replicating the GC-rich sequences, the 99.5% fidelity observed with the random sequence of D6 suggests that in general, most sequences are compatible with high fidelity amplification of the unnatural base pair.

Many *in vitro* applications rely on the efficient amplification of minuscule quantities of a template. To further explore the potential utility of the unnatural base pair, we examined PCR amplification via quantitative real time PCR with decreasing amounts of D1 and D6 (Table 1 and Figure 2). After 30 cycles of PCR, we found that D1 amplification remains efficient (2.7

$\times 10^6$ ) and accurate (99.8% fidelity) even with only 100 fg template. D6 was examined down to 1 pg, where amplification also remained efficient ( $1.9 \times 10^4$ ) and accurate (98.2% fidelity). The efficient and high fidelity amplification of D6 is particularly noteworthy given its randomized sequence. Moreover, sequencing revealed no major differences before and after amplification of D6 (Supporting Information), further suggesting that amplification is general and not strongly sequence-dependent.

To explore the determinants of efficient unnatural base pair amplification, we compared the amplification of DNA using **d5SICSTP** and **dMMO2TP** with that described above using **d5SICSTP** and **dNaMTP**. We examined amplification with duplexes D1 and D5, which correspond to the best and worst sequence contexts, respectively, for **d5SICS-dNaM** amplification, as well as with D6 (Table 1). The two unnatural pairs are replicated with similar efficiencies and fidelities in D1, reinforcing the idea that their incorporation into DNA does not rely solely on hydrophobicity, since **dNaM** is much more hydrophobic than **dMMO2**, and also that the intrabase packing interactions, which are similar in the two base pairs, make an important contribution. However, the data also reveal that **d5SICS-dMMO2** is replicated with lower fidelity and efficiency in D5 and D6, suggesting that its recognition is more sequence context dependent than that of **d5SICS-dNaM**.

We next examined amplification with two other thermostable polymerases commonly used for PCR, *Taq* and Phusion<sup>TM</sup>, using templates D1, D5, or D6, and dNTPs as well as **d5SICSTP** and **dNaMTP** (Table 1). Relative to DeepVent, *Taq* recognizes the unnatural base pair in D1 and D5 with similar efficiency and fidelity, but the fidelity is somewhat reduced with D6. Relative to DeepVent, Phusion<sup>TM</sup> polymerase amplifies **d5SICS-dNaM** in D1 with generally similar efficiency and fidelity, but in this case the fidelities are somewhat lower with both D5 and D6. The data suggest that the different polymerases behave similarly, except with increasing GC content, where divergent behavior is also observed with fully natural sequences,<sup>4b</sup> further demonstrating that the determinants of replication are contained within the base pairs and are not specific for a given polymerase.

We have now demonstrated not only that **d5SICS-dNaM** and **d5SICS-dMMO2** are efficiently transcribed in either direction,<sup>3c</sup> but also that they may be amplified by PCR, and are thus the first pairs to fulfill all of the primary requirements of a fully functional unnatural base pair. From a theoretical perspective, the data reveal that all of the properties required of a functional unnatural base pair may be optimized within predominantly hydrophobic nucleobases that bear no homology to the natural nucleobases. From a practical perspective, the data suggest that **d5SICS-dMMO2** and especially **d5SICS-dNaM** are sufficiently optimized for use as part of an *in vitro* expanded genetic alphabet. This should enable a variety of unprecedented applications,<sup>5</sup> which are currently being explored.

## Supplementary Material

Refer to Web version on PubMed Central for supplementary material.


## Acknowledgments

Funding provided by NIH GM060005.


## References

1. (a) Sismour AM, Lutz S, Park JH, J LM, Boyer PL, Hughes SH, Benner SA. *Nucleic Acids Res* 2004;32:728–735. [PubMed: 14757837] (b) Yang Z, Sismour AM, Sheng P, Puskar NL, Benner SA. *Nucleic Acids Res* 2007;35:4238–4249. [PubMed: 17576683]

2. (a) Hirao I, Kimoto M, Mitsui T, Fujiwara T, Kawai R, Sato A, Harada Y, Yokoyama S. *Nat. Methods* 2006;3:729–735. [PubMed: 16929319] (b) Kimoto M, Kawai R, Mitsui T, Yokoyama S, Hirao I. *Nucleic Acids Res* 2009;37:e14. [PubMed: 19073696]
3. (a) Leconte AM, Hwang GT, Matsuda S, Capek P, Hari Y, Romesberg FE. *J. Am. Chem. Soc* 2008;130:2336–2343. [PubMed: 18217762] (b) Seo YJ, Hwang GT, Ordoukhanian P, Romesberg FE. *J. Am. Chem. Soc* 2009;131:3246–3252. [PubMed: 19256568] (c) Seo YJ, Matsuda S, Romesberg FE. *J. Am. Chem. Soc* 2009;131:5046–5047. [PubMed: 19351201]
4. (a) Hansen, LL.; Justensen, J. *PCR Primer: A Laboratory Manual*. Diffenbach, CW.; Dveksler, GS., editors. Woodbury, NY: Cold Spring Harbor Laboratory Press; 2003. p. 43-62. (b) Arezi B, Xing W, Sorge JA, Hogrefe HH. *Anal. Biochem* 2003;321:226–235. [PubMed: 14511688]
5. Klussmann, S., editor. *The Aptamer Handbook: Functional Oligonucleotides and Their Applications*. Weinheim: Wiley-VCH; 2006.



**Figure 1.**  
Unnatural base pairs (with sugar and phosphate backbone omitted for clarity) and duplexes employed in this study.



**Figure 2.** Quantitative real time PCR analysis. (A) 30 cycle amplification of (from left to right) 1 ng, 100 pg, 10 pg, 1 pg, and 0.1 pg of D1. The rightmost curve corresponds to no template (negative control) where signal results from primer-dimer formation. Each curve represents the average of two independent experiments. (B) 30 cycle amplification of 1 pg of D6. See Supporting Information for details.

**Table 1**Efficiencies and fidelities of PCR amplification. <sup>a</sup>

Template	dXTPs incorporated	Enzyme	Amplification	Fidelity <sup>b</sup>
D1	dNaM, d5SICS	DeepVent	424	99.7
D2	dNaM, d5SICS	DeepVent	118	99.0
D3	dNaM, d5SICS	DeepVent	74	98.5
D4	dNaM, d5SICS	DeepVent	150	99.2
D5	dNaM, d5SICS	DeepVent	35	98.0
D6	dNaM, d5SICS	DeepVent	121	99.5
D7	-	DeepVent	556	-
D1 <sup>c</sup>	dNaM, d5SICS	DeepVent	$2.7 \times 10^6$	99.8
D6 <sup>d</sup>	dNaM, d5SICS	DeepVent	$1.9 \times 10^4$	98.2
D1	dMMO2, d5SICS	DeepVent	224	99.4
D5	dMMO2, d5SICS	DeepVent	25	97.1
D6	dMMO2, d5SICS	DeepVent	52	92.9
D1	dNaM, d5SICS	Taq	159	98.7
D5	dNaM, d5SICS	Taq	83	99.1
D6	dNaM, d5SICS	Taq	104	92.7
D1	dNaM, d5SICS	Phusion	257	99.7
D5	dNaM, d5SICS	Phusion	28	85.7
D6	dNaM, d5SICS	Phusion	82	95.9

<sup>a</sup>Conditions: 1 ng DNA template; dNTPs/dXTP = 600/400  $\mu$ M, 6 mM MgSO<sub>4</sub>, 0.03 U/ $\mu$ L enzyme, 8 min extension, 14 cycles.

<sup>b</sup>Calculated as average fidelity for unnatural base pair replication in both directions, except with template D5, where it was calculated in one direction (see text).

<sup>c</sup>100 fg DNA template, 30 cycles.

<sup>d</sup>1 pg DNA template, 30 cycles.