

2,4-Dichlorobenzaldehyde

Ricardo Cabello, Maksymilian Chruszcz and Wladek Minor*

University of Virginia, Department of Molecular Physiology & Biological Physics,
 1340 Jefferson Park Avenue, Charlottesville, VA 22908, USA
 Correspondence e-mail: wladek@wonka.med.virginia.edu


Received 4 December 2009; accepted 16 December 2009

Key indicators: single-crystal X-ray study; $T = 100\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.001\text{ \AA}$;
 R factor = 0.036; wR factor = 0.114; data-to-parameter ratio = 34.9.

In the crystal structure of the title compound, $\text{C}_7\text{H}_4\text{Cl}_2\text{O}$, the molecules form a network of weak $\text{C}-\text{H}\cdots\text{O}$ interactions involving the aldehyde O atom and the *ortho*-H atom on the benzene ring together with $\text{C}-\text{H}\cdots\text{O}$ interactions between the formyl groups. Together, these connect the molecules into $(10\bar{1})$ layers, which are stabilized additionally by $\pi-\pi$ stacking interactions of the benzene rings [centroid–centroid distance = $3.772(1)\text{ \AA}$]. The aldehyde group is twisted relative to the benzene ring by $7.94(13)^\circ$.

Related literature

For applications of the title compound, see: Katagi (1988); Wang *et al.* (2004). For a related structure, see: Gawlicka-Chruszcz *et al.* (2006).



Experimental

Crystal data

$\text{C}_7\text{H}_4\text{Cl}_2\text{O}$	$V = 693.2(3)\text{ \AA}^3$
$M_r = 175.01$	$Z = 4$
Monoclinic, $P2_1/n$	Mo $K\alpha$ radiation
$a = 13.100(1)\text{ \AA}$	$\mu = 0.85\text{ mm}^{-1}$
$b = 3.772(1)\text{ \AA}$	$T = 100\text{ K}$
$c = 15.332(1)\text{ \AA}$	$0.40 \times 0.10 \times 0.10\text{ mm}$
$\beta = 113.797(2)^\circ$	

Data collection

Rigaku R-AXIS RAPID diffractometer	6924 measured reflections
Absorption correction: multi-scan (Otwinowski <i>et al.</i> , 2003)	3737 independent reflections
$T_{\min} = 0.90$, $T_{\max} = 0.92$	3221 reflections with $I > 2\sigma(I)$
	$R_{\text{int}} = 0.063$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.036$	107 parameters
$wR(F^2) = 0.114$	All H-atom parameters refined
$S = 1.10$	$\Delta\rho_{\max} = 0.67\text{ e \AA}^{-3}$
3737 reflections	$\Delta\rho_{\min} = -0.41\text{ e \AA}^{-3}$

Table 1
 Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
C7—H7 \cdots O1 ⁱ	0.946 (17)	2.533 (17)	3.4289 (11)	158.0 (14)
C6—H6 \cdots O1 ⁱⁱ	0.950 (19)	2.512 (17)	3.2774 (11)	137.8 (12)

Symmetry codes: (i) $-x + \frac{3}{2}, y + \frac{1}{2}, -z + \frac{3}{2}$; (ii) $-x + 1, -y - 1, -z + 1$.

Data collection: *HKL-2000* (Otwinowski & Minor, 1997); cell refinement: *HKL-2000*; data reduction: *HKL-2000*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008) and *HKL-3000SM* (Minor *et al.*, 2006); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008) and *HKL-3000SM*; molecular graphics: *HKL-3000SM*, *ORTEPIII* (Burnett & Johnson, 1996), *ORTEP-3* (Farrugia, 1997), *Mercury* (Macrae *et al.*, 2006) and *POV-RAY* (The *POV-RAY* Team, 2004); software used to prepare material for publication: *HKL-3000SM*.

This work was supported by contract GI11496 from HKL Research, Inc.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: GK2246).

References

- Burnett, M. N. & Johnson, C. K. (1996). *ORTEPIII*. Report ORNL-6895. Oak Ridge National Laboratory, Tennessee, USA.
- Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
- Gawlicka-Chruszcz, A., Zheng, H., Hyacinth, M., Cymborowski, M., Sabat, M. & Minor, W. (2006). *Z. Kristallogr. New Cryst. Struct.* **221**, 545–546.
- Katagi, T. (1988). *J. Agric. Food Chem.* **36**, 344–349.
- Macrae, C. F., Edgington, P. R., McCabe, P., Pidcock, E., Shields, G. P., Taylor, R., Towler, M. & van de Streek, J. (2006). *J. Appl. Cryst.* **39**, 453–457.
- Minor, W., Cymborowski, M., Otwinowski, Z. & Chruszcz, M. (2006). *Acta Cryst. D* **62**, 859–866.
- Otwinowski, Z., Borek, D., Majewski, W. & Minor, W. (2003). *Acta Cryst. A* **59**, 228–234.
- Otwinowski, Z. & Minor, W. (1997). *Methods in Enzymology*, Vol. 276, *Macromolecular Crystallography*, Part A, edited by C. W. Carter Jr & R. M. Sweet, pp. 307–326. New York: Academic Press.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- The POV-RAY Team (2004). *POV-RAY*. <http://www.povray.org/download/>.
- Wang, S.-X., Tan, Z.-C., Di, Y.-Y., Xu, F., Zhang, H.-T., Sun, L.-X. & Zhang, T. (2004). *J. Chem. Thermodyn.* **36**, 393–399.

supplementary materials

Acta Cryst. (2010). E66, o243 [doi:10.1107/S160053680905435X]

2,4-Dichlorobenzaldehyde

R. Cabello, M. Chruscz and W. Minor

Comment

2,4-Dichlorobenzaldehyde is primarily used in the preparation of dyes, insecticides, herbicides, antiseptics and disinfectants (Wang *et al.*, 2004). It is also used as an intermediate of organic synthesis of fungicide diniconazole (Katagi, 1988).

In the crystal structure of 2,4-dichlorobenzaldehyde (Fig. 1), the aldehyde group is twisted relative to the benzene ring with torsion angles C6—C1—C7—O1 and C2—C1—C7—O1 being -7.94 (13)° and 170.86 (9)°. These torsion angles are significantly smaller in comparison to the corresponding angles in 2,6-dichlorobenzaldehyde (Gawlicka-Chruscz *et al.*, 2006) which are -27.3° and 152.6° respectively. Significantly bigger twist of the aldehyde group in the case of 2,6-dichlorobenzaldehyde is caused by presence of the chlorine atoms in ortho positions.

The change of the position of chlorine atom causes that interactions in which chlorine atoms are involved in 2,4-dichlorobenzaldehyde and 2,6-dichlorobenzaldehyde differ significantly. In the case of 2,6-dichlorobenzaldehyde Cl2 was involved in weak interaction with hydrogen atom from neighboring benzene ring, while in 2,4-dichlorobenzaldehyde structure such interactions are not observed for any of the chlorine atoms. However, in the case of 2,4-dichlorobenzaldehyde, the chlorine atoms from neighboring molecules form short contacts with Cl1···Cl2 ($1/2 + x, 1/2 - y, 1/2 + z$) distance being 3.442 Å (Fig. 2).

The weak O···H—C interactions (Table 1) between the aldehyde oxygen and the benzene hydrogen atoms connect molecules to form layers, which are additionally stabilized by stacking of benzene rings (Fig. 2). The oxygen atom from the aldehyde group plays a central role in the formation of weak interactions, and O1···H6—C6 ($1 - x, -1 - y, 1 - z$) and O1···H7—C7 ($1.5 - x, -1/2 + y, 1.5 - z$) distances are 2.51 Å and 2.53 Å respectively.

Experimental

2,4-dichlorobenzaldehyde was purchased from ALDRICH (99% purity, lot 08722CD). The compound was provided in crystalline form.

Refinement

All hydrogen atoms were localized using the difference density Fourier map. Their positions and isotropic displacement parameters were refined.

supplementary materials

Figures




Fig. 1. The asymmetric unit of the reported structure. Displacement ellipsoids are drawn at the 50% probability level and hydrogen atoms are drawn as grey spheres of an arbitrary radius.




Fig. 2. The molecular packing of 2,4-dichlorobenzaldehyde. Weak interactions, in which the oxygen atom participates, are shown as blue, dashed lines.

2,4-Dichlorobenzaldehyde

Crystal data

C ₇ H ₄ Cl ₂ O	<i>F</i> (000) = 352
<i>M_r</i> = 175.01	<i>D_x</i> = 1.677 Mg m ⁻³
Monoclinic, <i>P2₁/n</i>	Mo <i>K<alpha< i=""> radiation, λ = 0.71074 Å</alpha<></i>
Hall symbol: -P 2yn	Cell parameters from 31891 reflections
<i>a</i> = 13.100 (1) Å	θ = 1.0–37.8°
<i>b</i> = 3.772 (1) Å	μ = 0.85 mm ⁻¹
<i>c</i> = 15.332 (1) Å	<i>T</i> = 100 K
β = 113.797 (2)°	Block, colorless
<i>V</i> = 693.2 (3) Å ³	0.40 × 0.10 × 0.10 mm
<i>Z</i> = 4	

Data collection

Rigaku R-AXIS RAPID diffractometer	3737 independent reflections
Radiation source: fine-focus sealed tube graphite	3221 reflections with $I > 2\sigma(I)$
Detector resolution: 10 pixels mm ⁻¹	$R_{\text{int}} = 0.063$
ω scans	$\theta_{\text{max}} = 37.8^\circ$, $\theta_{\text{min}} = 1.0^\circ$
Absorption correction: multi-scan (Otwinowski <i>et al.</i> , 2003)	$h = -22 \rightarrow 22$
$T_{\text{min}} = 0.90$, $T_{\text{max}} = 0.92$	$k = -6 \rightarrow 6$
	$l = -26 \rightarrow 24$

6924 measured reflections

Refinement

Refinement on F^2	Primary atom site location: structure-invariant direct methods
Least-squares matrix: full	Secondary atom site location: difference Fourier map
$R[F^2 > 2\sigma(F^2)] = 0.036$	Hydrogen site location: difference Fourier map
$wR(F^2) = 0.114$	All H-atom parameters refined
$S = 1.10$	$w = 1/[\sigma^2(F_o^2) + (0.0708P)^2 + 0.0197P]$ where $P = (F_o^2 + 2F_c^2)/3$
3737 reflections	$(\Delta/\sigma)_{\text{max}} = 0.001$
107 parameters	$\Delta\rho_{\text{max}} = 0.67 \text{ e \AA}^{-3}$
0 restraints	$\Delta\rho_{\text{min}} = -0.41 \text{ e \AA}^{-3}$

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
Cl2	0.138241 (15)	0.17792 (6)	0.579292 (15)	0.02422 (7)
Cl1	0.558151 (16)	0.15135 (6)	0.840216 (14)	0.02526 (7)
C1	0.49878 (6)	-0.1166 (2)	0.66074 (6)	0.01968 (13)
C3	0.35318 (6)	0.1385 (2)	0.70157 (6)	0.02006 (14)
C2	0.46410 (6)	0.0484 (2)	0.72565 (5)	0.01980 (13)
C6	0.41880 (6)	-0.1866 (2)	0.56890 (6)	0.02060 (14)
C5	0.30765 (6)	-0.0964 (2)	0.54240 (6)	0.02080 (13)
C4	0.27652 (6)	0.0634 (2)	0.60987 (5)	0.01992 (13)
O1	0.64561 (5)	-0.4059 (2)	0.63526 (5)	0.02975 (14)
C7	0.61622 (6)	-0.2245 (2)	0.68671 (6)	0.02340 (14)
H3	0.3319 (13)	0.260 (4)	0.7450 (12)	0.038 (3)*
H5	0.2576 (13)	-0.150 (4)	0.4813 (13)	0.042 (4)*
H6	0.4423 (13)	-0.293 (5)	0.5239 (12)	0.046 (4)*
H7	0.6686 (14)	-0.131 (4)	0.7449 (13)	0.041 (4)*

supplementary materials

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Cl2	0.01697 (10)	0.02944 (12)	0.02558 (11)	0.00285 (6)	0.00789 (8)	0.00236 (6)
Cl1	0.02130 (11)	0.02926 (12)	0.02104 (11)	-0.00059 (6)	0.00422 (8)	-0.00518 (6)
C1	0.0165 (3)	0.0207 (3)	0.0213 (3)	-0.0005 (2)	0.0071 (2)	-0.0007 (2)
C3	0.0190 (3)	0.0209 (3)	0.0206 (3)	0.0003 (2)	0.0083 (3)	-0.0003 (2)
C2	0.0182 (3)	0.0203 (3)	0.0195 (3)	-0.0014 (2)	0.0062 (2)	-0.0017 (2)
C6	0.0189 (3)	0.0226 (3)	0.0206 (3)	-0.0002 (2)	0.0082 (2)	-0.0008 (2)
C5	0.0185 (3)	0.0229 (3)	0.0195 (3)	-0.0007 (2)	0.0061 (2)	-0.0008 (2)
C4	0.0168 (3)	0.0209 (3)	0.0215 (3)	0.0001 (2)	0.0071 (2)	0.0016 (2)
O1	0.0214 (3)	0.0378 (3)	0.0304 (3)	0.0045 (2)	0.0108 (2)	-0.0046 (3)
C7	0.0178 (3)	0.0266 (3)	0.0251 (3)	-0.0001 (3)	0.0080 (3)	-0.0013 (3)

Geometric parameters (\AA , $^\circ$)


Cl2—C4	1.7327 (7)	C3—H3	0.939 (17)
Cl1—C2	1.7343 (8)	C6—C5	1.3869 (11)
C1—C2	1.3961 (11)	C6—H6	0.950 (19)
C1—C6	1.3999 (11)	C5—C4	1.3930 (11)
C1—C7	1.4820 (11)	C5—H5	0.923 (18)
C3—C4	1.3877 (11)	O1—C7	1.2180 (11)
C3—C2	1.3893 (11)	C7—H7	0.946 (17)
C2—C1—C6	118.32 (7)	C1—C6—H6	118.6 (9)
C2—C1—C7	122.14 (7)	C6—C5—C4	118.43 (7)
C6—C1—C7	119.53 (7)	C6—C5—H5	118.4 (10)
C4—C3—C2	118.11 (7)	C4—C5—H5	123.2 (10)
C4—C3—H3	121.2 (10)	C3—C4—C5	122.11 (7)
C2—C3—H3	120.6 (10)	C3—C4—Cl2	118.26 (6)
C3—C2—C1	121.73 (7)	C5—C4—Cl2	119.62 (6)
C3—C2—Cl1	116.99 (6)	O1—C7—C1	123.05 (8)
C1—C2—Cl1	121.28 (6)	O1—C7—H7	121.4 (10)
C5—C6—C1	121.30 (7)	C1—C7—H7	115.5 (10)
C5—C6—H6	120.0 (9)		
C4—C3—C2—C1	-0.72 (12)	C1—C6—C5—C4	-0.68 (12)
C4—C3—C2—Cl1	179.11 (6)	C2—C3—C4—C5	-0.19 (12)
C6—C1—C2—C3	0.90 (12)	C2—C3—C4—Cl2	-179.59 (6)
C7—C1—C2—C3	-177.92 (8)	C6—C5—C4—C3	0.88 (12)
C6—C1—C2—Cl1	-178.92 (6)	C6—C5—C4—Cl2	-179.73 (6)
C7—C1—C2—Cl1	2.26 (11)	C2—C1—C7—O1	170.86 (9)
C2—C1—C6—C5	-0.18 (12)	C6—C1—C7—O1	-7.94 (13)
C7—C1—C6—C5	178.67 (7)		

Hydrogen-bond geometry (\AA , $^\circ$)

$D—H\cdots A$	$D—H$	$H\cdots A$	$D\cdots A$	$D—H\cdots A$
C7—H7—O1 ⁱ	0.946 (17)	2.533 (17)	3.4289 (11)	158.0 (14)

C6—H6···O1ⁱⁱ 0.950 (19) 2.512 (17) 3.2774 (11) 137.8 (12)
Symmetry codes: (i) $-x+3/2, y+1/2, -z+3/2$; (ii) $-x+1, -y-1, -z+1$.

Fig. 1



supplementary materials

Fig. 2

