

INFECTION

The BEACHES Study: health effects and exposures from non-point source microbial contaminants in subtropical recreational marine waters

Jay M Fleisher,^{1,2*} Lora E Fleming,^{2,3} Helena M Solo-Gabriele,^{2,4} Jonathan K Kish,² Christopher D Sinigalliano,^{2,5} Lisa Plano,^{2,3} Samir M Elmir,^{2,6} John D Wang,² Kelly Withum,² Tomoyuki Shibata,^{2,4,5} Maribeth L Gidley,^{2,5} Amir Abdelzaher,^{2,4} Guoqing He,^{2,4} Cristina Ortega,^{2,4} Xiaofang Zhu,² Mary Wright,^{2,4} Julie Hollenbeck² and Lorraine C Backer⁷

¹Nova Southeastern University COM/MPH, Fort Lauderdale, FL, USA, ²NSF NIEHS Oceans and Human Health Center, Rosenstiel School, University of Miami, Miami, FL, USA, ³Miller School of Medicine, University of Miami School of Medicine, Miami, FL, USA, ⁴College of Engineering, University of Miami, Coral Gables, FL, USA, ⁵NOAA Atlantic Oceanographic and Meteorological Laboratory, Miami, FL, USA, ⁶Miami Dade County Public Health Department, Miami, FL, USA and ⁷National Center for Environmental Health, Centers for Disease Control and Prevention, Atlanta, GA, USA

*Corresponding author. Nova Southeastern University COM/MPH, 3200 South University Drive, Fort Lauderdale, FL 33328, USA. E-mail: jmfleish@nova.edu

Accepted 14 April 2010

Background Microbial water-quality indicators, in high concentrations in sewage, are used to determine whether water is safe for recreational purposes. Recently, the use of these indicators to regulate recreational water bodies, particularly in sub/tropical recreational marine waters without known sources of sewage, has been questioned. The objectives of this study were to evaluate the risk to humans from exposure to subtropical recreational marine waters with no known point source, and the possible relationship between microbe densities and reported symptoms in human subjects with random-exposure assignment and intensive individual microbial monitoring in this environment.

Methods A total of 1303 adult regular bathers were randomly assigned to bather and non-bather groups, with subsequent follow-up for reported illness, in conjunction with extensive environmental sampling of indicator organisms (enterococci).

Results Bathing was 1.76 times more likely to report gastrointestinal illness [95% confidence interval (CI) 0.94–3.30; $P = 0.07$]; 4.46 times more likely to report acute febrile respiratory illness (95% CI 0.99–20.90; $P = 0.051$) and 5.91 times more likely to report a skin illness (95% CI 2.76–12.63; $P < 0.0001$) relative to non-bathers. Evidence of a dose–response relationship was found between skin illnesses and increasing enterococci exposure among bathers [1.46 times (95% CI 0.97–2.21; $P = 0.07$) per increasing \log_{10} unit of enterococci exposure], but not for gastrointestinal or respiratory illnesses.

Conclusions This study indicated that bathers may be at increased risk of several illnesses relative to non-bathers, even in the absence of any known

source of domestic sewage impacting the recreational marine waters. There was no dose–response relationship between gastroenteritis and increasing exposure to enterococci, even though many current water-monitoring standards use gastroenteritis as the major outcome illness.

Keywords Gastrointestinal illness, respiratory illness, skin illness, indicator organisms, enterococci, recreational water quality

Introduction

Traditionally, analogous to drinking water, monitoring of microbial water quality for coastal waters used for recreational purposes has been regulated by measuring the concentrations of indicator microbes. These microbes are those typically found in human faeces in high concentrations, and are not pathogenic. Thus, an elevated concentration of these indicator microbes in coastal waters should indicate that these waters have been contaminated by human sewage, and are unsafe for recreational use.^{1–5}

Recently, the use of indicator microbes to regulate the recreational use of coastal waters has come into question, particularly in subtropical marine environments characterized by no known source of human sewage (i.e. non-point source). Studies conducted in subtropical areas have shown collectively that, in the absence of any known sources of human faecal materials, indicator microbes are consistently present, and may be recovered in high concentrations in the environment.^{6–14} It is thought that indicator microbes may re-grow due to climate conditions conducive for re-growth, and thus may not be representative of any pathogens that might be present. However, the regulator is presented with a perplexing situation where the possibility of micro-organisms or pathogens within urban run-off might be responsible for an increased risk of illness among bathers. In addition, within subtropical waters, it remains unclear which indicator microbe(s) should be utilized, and once the data are obtained, how these data should be interpreted.^{5,15}

Several epidemiological studies have found that bathing in temperate recreational waters with known point sources of faecal contamination (such as domestic sewage or storm-drain runoff) has been associated with an increased risk for transmission of infectious diseases (including gastroenteritis, and febrile respiratory, skin, eye and ear illnesses).^{3,16–26} The few epidemiological studies conducted in subtropical environments have shown no statistically significant relationship between human health and current US Environmental Protection Agency (EPA) recommended indicator microbes for evaluating beach water quality in subtropical regions.^{2,8,12,27} The objective of this study, the first prospective randomized exposure study in non-point source subtropical

recreational marine waters, was to evaluate the risks to human health in recreational marine environment with no known source of domestic sewage, and the use of indicator organisms to assess these exposures and risks.

Methods

The design of the epidemiological study was based on past studies at the study site and on the work of Fleisher *et al.*^{11,14,16–19,28–30} The investigators conducted a prospective randomized exposure study with participant exposure randomly assigned to either marine recreational waters or to beach-only exposure. The epidemiological study data-collection activities were conducted over 15 individual study days beginning 15 December 2007 and ending 21 June 2008.

Epidemiological study protocol

Local adult residents (≥ 18 years of age) who reported regular bathing in recreational marine waters were recruited to participate using word of mouth, email and local publicity as recruitment tools. After recruitment, the participants were screened for current illnesses and symptoms, possible alternative risk factors for the study illnesses, and a brief health history; they were then given an appointment for the study beach-exposure day. Members of the same household or family members were recruited, but subject to the same randomization procedures as all other study participants. Possible secondary spread of infection among family members was ascertained by asking ‘Did any member of your family become ill before or after recruitment into the study?’, with control in subsequent statistical analyses.

On the study beach-exposure day, participants read and signed the consent form. At this point, participants were again interviewed briefly about any current illnesses and symptoms, food consumption and beach exposure since the baseline recruitment interview. Study subjects were then randomly assigned to either (exposed) bather group or (unexposed) non-bather group. The bathers were assigned to the bathing station where staff members supervised the exposure activity of each bather, including the time, location, unusual activities and duration each

individual bather spent in the water. Bathers were required to spend 15 min in knee-deep water (due to the relative shallowness of the study site), and to immerse their head three times completely under water. Using ropes, a 30–40-m stretch of beach was subdivided into 5-m intervals forming six to eight bathing exposure zones, with exposure of any individual bather restricted to their own individual swim zone that was 5 m wide. Each subject was instructed to take their own water sample at 5-min intervals near the surface before their head immersion, as well as provided with an appropriate individual water-sampling container. Staff members instructed participants to thoroughly rinse the collection container before filling completely with the marine water, as well as the avoidance of microbial contamination of the collection container by the participant. When the subjects left the water, they gave their individual water samples to the environmental research study staff for microbial analysis processing (described below). No bather was allowed to enter the water more than once during the actual study exposure. The participants in the randomized non-bather group were restricted to sitting on chairs on plastic sheeting in a covered roped-off area distant from water and sand exposure for 15 min.

After participating in the beach-day protocol, all participants were given a US\$50 gift card, and an appointment to complete an extensive follow-up phone questionnaire 7 days after the beach visit. After completing the follow-up questionnaire, study participants received an additional US \$25 gift card to compensate them for participating in the study.

Informed consent documents and study questionnaires were provided in English and Spanish, and approved by the Florida Department of Health and University of Miami Human Subjects Committees. The questionnaires (as well as the study design) were adapted from those of Dr David Kay and Dr Jay M. Fleisher from the prospective randomized exposure studies in the UK and Europe.^{16–19,22} Adaptations included some language changes to 'American' English, as well as logistical changes (in particular, instead of follow-up at 7 days and 3 weeks, due to resources and logistics, these follow-ups were combined into a single follow-up questionnaire given 7 days post exposure via telephone).

Disease endpoint criteria

The disease endpoints of interest in this study were self-reported symptoms consistent with gastrointestinal illness, acute febrile respiratory illness [i.e. International Classification of Diseases 9: 461–466, 480], eye or ear infections and skin infections occurring within 7 days of beach exposure.

The following definitions were used to derive disease endpoint categories prior to data analysis based on the results of the seventh-day follow-up questionnaire. 'Gastrointestinal illness' was defined as a

report of the following symptoms: all cases of vomiting or diarrhoea, or all reported cases of indigestion or nausea accompanied by a fever; 'diarrhoea' defined as having three or more runny stools within a 24-h period; 'acute febrile respiratory illness' defined as report of at least one of each of the symptoms listed in each of the three categories: (i) fever; (ii) headache and/or body-aches and/or unusual fatigue and/or loss of appetite; and (iii) sore throat and/or runny nose and/or dry or productive cough; 'skin illness' defined as report of at least one of the following symptoms: (i) skin rash; (ii) skin ulcer/sore; or (iii) itching/irritation; 'ear illness' defined as report of an ear infection (sore/discharge); 'eye illness' as report of an eye infection on the follow-up questionnaire. These illness definitions were compatible with past studies carried out by the investigators in Europe, and are roughly compatible with the 'highly creditable illness' definitions used in the majority of the US recreational water cohort studies.^{3,16–19,22}

Environmental sampling and microbial assays

The study site had been well characterized by the investigators previously in terms of the indicator organisms in the water and sediments as being a non-point source subtropical beach with periodic use by people, dogs and birds as well as seasonal heavy rains.^{14,28–33}

As described above, while each study participant was in the water, each subject collected their own environmental water sample from the demarcated bathing lanes. All samples were collected in the morning between 8 am and 12 pm on each sampling day concomitant with the epidemiological study. As described above, water samples were collected in knee-deep water by each study participant using 5 litres sterilized plastic containers; sampling points from the shoreline ranged from 10 to 40 m depending on tidal stage corresponding with the knee height.

Samples taken by each study participant were assayed for enterococci by membrane filtration using the method recommended by the US EPA.^{34,35}

Randomization, epidemiological study database and statistical analyses

Block randomization into bather and non-bather groups was used to maximize the probability of achieving an equal number of study participants in the two groups under study. Block size consisted of a random ordering of blocks of two, four and six study participants per individual block. It should be noted that the random ordering of block size eliminates any chance of study personnel discovering the blocking pattern. Participants were only informed of their randomized status after completing the pre-exposure day interview at the beach. The sample size was calculated to obtain a difference in gastrointestinal illness between bathers and non-bathers of 5/100 at $\alpha=0.05$ and $\beta=0.80$.

The statistical software packages Statistical Analysis Software (SAS), and where necessary StatXact, were used to perform all statistical analyses. Initial analyses evaluated the effects of randomization for all demographic variables to insure equal distribution of demographic risk factors among exposed (i.e. bathers) and the unexposed (i.e. non-bathers). Subsequent basic analyses were geared to answer two questions: (i) 'Was there an excess of the illnesses under study among bathers vs non-bathers?'; and (ii) 'Was there a dose-response relationship between indicator organism density and the incidence of the outcome illnesses studied among bathers?'. Participants reporting a particular illness on the study day were excluded from the statistical analysis of that particular illness. Statistical analyses included univariate analysis using chi-square and, where expected cell size was <5 , the Fisher's exact test (two-tailed). Unconditional logistic regression modelling was then used to control for possible confounders/non-water-related risk factors among bathers vs non-bathers. Non-water-related risk factors or possible confounders for all outcome illness under analysis were evaluated separately in univariate analyses (Table 1). Inclusion criterion for the potential confounders was $P < 0.15$, with a model retention criterion of $P < 0.05$. A backward stepwise elimination strategy was employed.

Results

Of 1341 subjects who participated in the 15 beach study days, 38 (2.9%) were lost to follow-up, resulting in a final total study cohort of 1303 with a follow-up of 97.1%. Of note, 80% of the 1303 subjects were successfully followed up within 7 days of exposure, and the remaining 20% within 3 weeks. After randomization, there were 652 bathers and 651 non-bathers. Randomization of subjects into bather and non-bather exposure for the purposes of the

study was successful in terms of equal distribution of demographics with no significant differences (Table 2).

The distribution of enterococci exposure among bathers was: for all 668 bathers, the mean enterococci exposure was 71 enterococci/100 ml [standard deviation (SD) 244]; the median = 19 enterococci/100 ml; the minimum exposure below the detection limit of 2 enterococci/100 ml; and the maximum enterococci exposure = 3320 enterococci/100 ml. We recorded the number of days between the exposure study day for each individual participant and the day of onset of their symptoms. For gastrointestinal illnesses, the average time for bathers to the onset of symptoms was 6.10 (SD 2.81) days and for non-bathers 7.78 (SD 3.15) days ($P=0.16$); for acute febrile illness, bathers 5.18 (SD 2.44) days and for non-bathers 12 (SD 2.82) days ($P=0.004$); for skin illnesses, bathers 4.36 (SD 2.56) days and for non-bathers 5.55 (SD 2.17) ($P=0.19$); for eye illnesses, bathers 5.75 (SD 2.56) days and for non-bathers 9.20 (SD 7.19) days, ($P=0.41$); and for ear illnesses, bathers 3.67 (SD 1.75) days and for non-bathers 9.67 (SD 0.58) days, ($P=0.0008$). For every illness outcome, the un-exposed non-bathers reported a longer time to onset than the exposed bathers.

Bathers vs non-bathers

Bathers reported more gastrointestinal illness ($P=0.08$), respiratory illness ($P=0.04$) and skin illness ($P=0.0001$) relative to non-bathers (Table 3). After controlling for non-water-related risk factors/possible confounders for each of these illnesses using multiple logistic regression analysis, subsequent to 7 days of follow-up from beach exposure, bathers were 1.79 times [odds ratio 1.79; 95% confidence interval (CI) 0.94–3.43; $P=0.07$] more likely to report gastrointestinal illness relative to non-bathers; bathers were 4.46 times (95% CI 0.99–20.97; $P=0.051$) more likely to report acute febrile

Table 1. Non-water-related risk factors and possible confounders evaluated for both bathers and non-bathers

Risk factor/possible confounder

- Age
 - Gender
 - History of significant illness: gastrointestinal, respiratory, skin, eye, ear, migraines, stress
 - Use of medications within 4 weeks of exposure day
 - Illness within 4 weeks of exposure day
 - Consumption of the following foods from 3 days before to 7 days after exposure day: ethanol, mayonnaise, chicken, eggs, ice cream, salad, hamburgers, hot dogs, raw milk, meat pies, seafood, purchased sandwiches
 - Illness in household after exposure day
 - Additional bathing after exposure day
 - Various measures of risk perception
-

Source: Fleisher *et al.*¹⁸ and Kay *et al.*²²

respiratory illness; and bathers were 5.31 times (95% CI 2.58–10.96; $P=0.0001$) more likely to report skin illness. No substantial confounding was observed between any of these three illnesses and the non-water-related risk factor/confounders (Table 4). No differences between bathers and non-bathers for ear and eye illnesses were observed (Table 3).

Expressed in terms of attributable risk, the excess risk for gastroenteritis was 3.17/100 ($P=0.01$); for acute febrile respiratory illness, the excess risk was 1.28/100 ($P=0.04$); and for skin illness, the excess risk was 6.03/100 ($P < 0.0001$).

Bathers only

Enterococci by membrane filtration showed a dose–response relationship with skin illness. Unconditional multiple logistic regression modelling was used to compute the probability of bathers acquiring a skin illness with increasing enterococci exposure rose while controlling for any of the non-water-related risk factors/confounders shown in Table 1. Covariates in which the P -value was <0.15 in the univariate analysis comparing bathers reporting skin illnesses with bathers who did not report such

illness were included in the model, while model retention was set at $P < 0.05$.

Enterococci exposure was modelled as a continuous variable (data not shown). Enterococci exposures were transformed via \log_{10} prior to inclusion into the model. The results showed that the risk of a skin illness increased by 1.46 times (0.97–2.21; $P=0.07$) per increasing \log_{10} unit of enterococci exposure, while controlling for gender and for the question ‘Have you heard anything about chemical pollution and bathing waters?’ as a measure of risk perception ($P=0.04$ and 0.02, respectively). A plot of the probability of bathers contracting skin illness with increasing enterococci exposure (after \log_{10} transformation) with 95% CIs is illustrated in Figure 1. With regard to gastroenteritis, the same method of analysis was used, but demonstrated no relationship with the increasing levels of enterococci (1.39 per increasing \log_{10} unit of enterococci exposure (0.74–2.61; $P=0.31$), and thus no dose–response relationship).

Discussion

This study was the first randomized exposure study of bathers with individual exposure assessment in subtropical recreational marine waters with no known source of domestic sewage. The results demonstrated bathers to be at increased risk of reported gastrointestinal, acute febrile respiratory and skin illnesses relative to non-bathers. In addition, there was evidence for a dose–response relationship between bather exposure to increasing levels of enterococci and an increased risk of reporting skin illness.

The fact that we found no dose–response relationship between respiratory or gastrointestinal illnesses and enterococci exposure remains unexplained, and may be a result of the study sample size. Based on our prior studies, some contribution to each individual participant’s enterococci exposure may have come from their own bacterial shedding; however, our prior research has indicated that this is not the major source of enterococci exposure in non-point source subtropical recreational marine waters.^{29,32,33} It should be noted that the actual pathogens responsible for all of these illnesses remain unknown; they may

Table 2. Demographics of the study population

Variable	Bather	Non-bather
Bather status, N (%)	652 (50.04)	651 (49.96)
Mean age (\pm SE)	32.20 \pm 12.64	32.50 \pm 13.40
Female, N (%)	370 (49.27)	381 (50.73)
Hispanic ethnicity, N (%)	243 (48.70)	256 (51.30)
Occupation, N (%)		
Office	141 (47.80)	154 (52.20)
Catering/leisure	9 (64.29)	5 (35.71)
Agriculture	5 (50)	5 (50)
Student/school/university	285 (49.83)	287 (50.17)
Factory	7 (87.50)	1 (12.50)
Caring for others	26 (41.94)	36 (58.06)
Building/construction	23 (62.16)	14 (37.84)
Other	154 (51.16)	147 (48.84)

SE, standard error.

Table 3. Crude rates of the outcome illnesses assessed by bather vs non-bather groups


	Total analysed ^a	Bathers, N (%)	Non-bathers, N (%)	P -value ^b
Gastroenteritis	1239	31 (4.75)	18 (2.90)	0.08
Acute febrile respiratory illness	1240	12 (1.94)	4 (0.64)	0.04
Skin	1253	47 (7.47)	9 (1.44)	0.0001
Eye	1299	4 (0.61)	5 (0.77)	0.75
Ear	1300	6 (0.92)	3 (0.46)	0.51

^aNumbers vary due to exclusion of participants reporting particular symptoms on exposure day.

^bStatistical testing by chi-square, or (where any expected cell size was <5), Fisher’s exact test.

Table 4. Results of unconditional multiple logistic regression analyses by reported illness: bathers vs non-bathers

Risk factor	Chi-square	P-value	Odds ratio (95% CI)
Gastrointestinal illness			
Gastrointestinal symptoms	3.17	0.07	1.79 (0.94–3.43)
Excessive tiredness 3 weeks prior to initial interview, lasting <24 h	7.21	0.007	4.65 (1.51–14.27)
Eat salad after exposure day	4.54	0.03	0.75 (0.57–0.98)
Illness in household within 7 days of beach exposure	7.91	0.005	0.49 (0.30–0.81)
Eat ice-cream since exposure day	3.50	0.06	0.79 (0.62–1.01)
Respiratory illness			
Report of respiratory symptoms	3.80	0.051	4.46 (0.99–20.97)
Illness in household within 7 days of beach exposure	7.81	0.005	2.03 (1.24–3.34)
Skin illness			
Skin symptoms	20.96	0.0001	5.31 (2.58–10.96)
Illness in household within 7 days of beach exposure	8.33	0.004	0.49 (0.30–0.80)

**Figure 1.** Dose–response (with 95% CIs) between the probability of skin illnesses and increasing enterococci levels (bathers only). Maximum enterococci measured = 3.52 log₁₀ CFU, colony forming units

vary from study site to study site, and thus, the apparent lack of correlation with enterococci exposure reported herein. Although ‘urban runoff’ is a probable source of pathogen(s), we cannot rule out other micro-organisms that are naturally present in subtropical marine waters and not the result of contamination by man and other animals. If this is the case, we would not expect a correlation with any indicator organism. Since this study was the first randomized prospective exposure of bathers performed in subtropical recreational marine waters with no known point source, additional studies need to be performed to confirm and explore our results further.

This is the only study of the association between concentrations of microbial water contamination and subsequent illnesses in which the bathers collected

their own water samples at the same time they were being ‘exposed’. This should provide the most accurate estimate of individual exposure to water-borne microbial indicators, including enterococci, to date. This should aid to minimize the misclassification of exposures associated with non-randomized cohort designs, which would likely be non-differential, and thus result in an underestimation of the risk. It must, however, be emphasized that the bather cohort used in our study consisted of regular adult bathers. Care must therefore be used when interpreting these findings, especially with respect to susceptible subpopulations such as small children or anyone with a compromised immune system.^{15,36}

In addition to individual exposure assessment, another strength of our study was that, unlike the designs of almost all other published epidemiological studies which used a prospective cohort design, the basic unit of measurement was the individual bather rather than the average rates of illness among many bathers on different study days. This use of an aggregated measure of individual exposures (i.e. rates or rate differences) as the basic unit of measurement in all previous prospective cohort designs would lead to significant misclassification of exposure bias.¹⁶ The design of the study reported herein would minimize this source of misclassification of exposure bias. In addition, the randomization of individual study participants who reported bathing regularly in recreational marine waters into exposed and unexposed groups solely for the purposes of the study further avoided another bias possibly inherent in the prospective cohort design: the hypothesized phenomena that persons who report regular non-bathing might constitute a less healthy group relative to regular bathers, leading to a self-selection bias.^{17,37,38} In the presence of such selection bias (unavoidable in previous prospective cohort studies), the comparison of

bathers with non-bathers might lead to results of questionable validity.

In a regulatory context, the results we report can be transformed into the probability of illness, attributable risk and other measures of risk, thus avoiding the use of rates as the basic unit of measurement published in previous prospective cohort designs, which inevitably introduce misclassification of exposure. Thus, the methods used in this study will supply the regulator with more accurate estimates of risk upon which to build criteria in the future.

The implications of the findings of this study are far reaching. Our findings suggest that there is an increased risk to bathers even when using marine recreational waters with no known source of domestic sewage. We also observed a correlational dose–response association between skin illnesses and increasing enterococci exposure. These findings might indicate that microbes from non-point sources are causing increased risk from pathogens introduced by urban run-off, and thus from a regulatory viewpoint,²⁰ the notion of zero additional risk to bathers could be challenged.¹⁵ Further epidemiological and laboratory studies will be necessary to confirm this hypothesis.

Funding

National Center for Environmental Health (NCEH), Centers for Disease Control and Prevention (CDC); Florida Dept of Health (FL DOH) through monies from the Florida Dept of Environmental Protection (FL DEP);

the EPA Internship Program; the National Science Foundation (NSF) and the National Institute of Environmental Health Sciences (NIEHS) Oceans and Human Health Center at the University of Miami Rosenstiel School (NSF 0CE0432368) and (NIEHS P50 ES12736) and NSF REU in Oceans and Human Health and the NSF SGER (NSF SGER 0743987) in Oceans and Human Health.

Acknowledgements

The researchers would like to dedicate this research to the memory of Ms Seana Campbell, a very talented, hardworking and creative young researcher who enriched all people whose lives she touched, and who died too young. The researchers would like to thank the following collaborators at the following Institutions: NOAA Southeast Miami Lab; Miami-Dade County Department of Health; Miami Sea Aquarium, University of Florida; BCS Laboratories; University of South Florida; NOAA Charleston Hollings Laboratory; Woods Hole Oceanographic Institute and the University of Missouri. The researchers would also like to thank the many students of University of Miami and Florida International University, other student researchers and other researchers for their assistance in the performance of this study.

Conflict of interest: None declared.

KEY MESSAGES

- The objectives of this study were to evaluate the risk to humans from exposure to subtropical recreational marine waters with no known point source of microbial pollution.
- This study indicated that bathers may be at increased risk of several illnesses relative to non-bathers, even in the absence of any known source of domestic sewage impacting the recreational marine waters.
- Interestingly, no dose–response relationship between gastroenteritis and increasing exposure to enterococci was detected, given that many current water monitoring standards use gastroenteritis as the major outcome illness and enterococcus as the recommended indicator organism for monitoring purposes.

References

- 1 Cabelli VJ, Dufour AP, McCabe LJ, Levin MA. Swimming associated gastroenteritis and water quality. *Am J Epidemiol* 1982;**115**:606–16.
- 2 Fujioka RS, Byappanahalli MN. *Draft of Final Report, Tropical Indicator Workshop*. Washington, DC: USEPA Office of Water, 2001.
- 3 Wade TJ, Pai N, Eisenberg JN, Colford JM Jr. Do U.S. Environmental Protection Agency water quality guidelines for recreational waters prevent gastrointestinal illness? A systematic review and meta-analysis. *Environ Health Perspect* 2003;**111**:1102–9.
- 4 Craun GF, Calderon RL, Wade TJ. Assessing waterborne risks: an introduction. *J Water Health* 2004;**4**:3–18.
- 5 Boehm AB, Ashbolt NJ, Colford JM *et al*. A sea change ahead for recreational water quality criteria. *J Water Health* 2009;**7**:9–20.
- 6 Toranzos GA. Current and possible alternate indicators of fecal contamination in tropical waters: a short review. *Environ Toxicol Water Quality* 1991;**6**:121–30.
- 7 Calderon RL, Mood EW, Dufour AP. Health effects of swimmers and nonpoint sources of contaminated water. *Int J Environ Health Res* 1991;**1**:21–31.
- 8 Dwight RH, Baker DB, Semenza JC, Olson BH. Health effects associated with recreational coastal water

- use: urban versus rural California. *Am J Public Health* 2004;**94**:565–67.
- ⁹ Solo-Gabriele HM, Wolfert MA, Desmarais TR, Palmer CJ. Sources of *Escherichia coli* in a coastal subtropical environment. *Applied Environ Microbiol* 2000;**66**:230–37.
 - ¹⁰ Desmarais TR, Solo-Gabriele HM, Palmer CJ. Influence of soil on fecal indicator organisms in a tidally influenced subtropical environment. *Applied Environ Microbiol* 2002;**68**:1165–72.
 - ¹¹ Shibata T, Solo Gabriele HM, Fleming LE, Elmir S. Monitoring marine recreational water quality using multiple microbial indicators in an urban tropical environment. *Water Res* 2004;**38**:3119–31.
 - ¹² Colford JM Jr, Wade TJ, Schiff KC *et al*. Water quality indicators and the risk of illness at beaches with non-point sources of fecal contamination. *Epidemiology* 2007;**18**:27–35.
 - ¹³ Yamahara KM, Layton BA, Santoro AE, Boehm AB. Beach sands along the California coast are diffuse sources of fecal bacteria to coastal waters. *Environ Sci Technol* 2007;**41**:4515–21.
 - ¹⁴ Wright ME. *Evaluation of enterococci, an indicator microbe, and the sources that impact the water quality at a subtropical non-point source recreational beach*. Masters Thesis in Engineering. Coral Gables, FL: University of Miami, 2008.
 - ¹⁵ US Environmental Protection Agency (US EPA). *Report of the Experts Scientific Workshop on Critical Research Needs for the Development of New or Revised Recreational Water Quality Criteria*. Washington, DC. EPA 823-R-07-006, <http://www.epa.gov/waterscience/criteria/recreation/>, 2007 (1 May 2010, date last accessed).
 - ¹⁶ Fleisher JM. A re-analysis of the data supporting U.S. federal bacteriological water quality criteria governing marine recreational waters. *Water Pollution Control Federation J* 1991;**63**:259–65.
 - ¹⁷ Fleisher JM, Jones F, Kay D, Stanwell-Smith R, Wyer M, Morano R. Water and non-water risk factors for gastroenteritis among bathers exposed to sewage-contaminated marine waters. *Int J Epidemiol* 1993;**22**:698–708.
 - ¹⁸ Fleisher JM, Kay D, Salmon RL, Jones F, Wyer MD, Godfree AF. Marine waters contaminated with domestic sewage: nonenteric illness associated with bather exposure in the United Kingdom. *Am J Public Health* 1996;**86**:1228–34.
 - ¹⁹ Fleisher JM, Kay D, Wyer MD, Godfree AF. Estimates of the severity of illnesses associated with bathing in marine recreational waters contaminated with domestic sewage. *Int J Epidemiol* 1998;**27**:722–26.
 - ²⁰ Fujioka RS, Roll K, Morens D. *A Pilot Epidemiological Study of Health Risks Associated with Swimming at Kuhio Beach*. Honolulu, Hawaii: Hawaii Water Resources Research Center, 1994.
 - ²¹ Haile RW, Witte JS, Gold M *et al*. The health effects of swimming in ocean water contaminated by storm drain runoff. *Epidemiology* 1999;**10**:355–63.
 - ²² Kay D, Fleisher JM, Salmon RL *et al*. Predicting likelihood of gastroenteritis from sea bathing: results from randomised exposure. *Lancet* 1994;**344**:905–9.
 - ²³ Kueh CS, Tam TY, Lee T. Epidemiological study of swimming associated illnesses relating to bathing beach water quality. *Water SciTech* 1995;**31**:1–4.
 - ²⁴ Prieto MD, Lopez B, Juanes JA, Revilla JA, Llorca J, Delgado Rodriguez M. Recreation in coastal waters: health risks associated with bathing in sea water. *J Epidemiol Community Health* 2001;**55**:442–47.
 - ²⁵ Pruss A. Review of epidemiological studies on health effects from exposure to recreational water. *Int J Epidemiol* 1998;**27**:1–9.
 - ²⁶ Shuval H. Estimating the global burden of thalassogenic diseases: human infectious diseases caused by wastewater pollution of the marine environment. *J Water Health* 2003;**1**:53–64.
 - ²⁷ Fleming LE, Solo Gabriele H, Elmir S *et al*. A pilot study of microbial contamination of subtropical recreational waters. *Fl J Env Health* 2004;29–33.
 - ²⁸ Elmir SM. *Development of a water quality model which incorporates non-point microbial sources*. PhD Thesis. Coral Gables, FL: University of Miami, 2006.
 - ²⁹ Elmir SM, Wright ME, Solo-Gabriele HM *et al*. Quantitative evaluation of bacteria released by bathers in marine water. *Water Res* 2007;**41**:3–10, (with supplement available online).
 - ³⁰ Fleming LE, Solo Gabriel HM, Fleisher JM *et al*. *Final Report on the Pilot Epidemiologic Assessment of Microbial Indicators for Monitoring Recreational Water Quality in Marine Sub/Tropical Environments*. Miami, FL: The NSF NIEHS Oceans and Human Health Center, Rosenstiel School of Marine and Atmospheric Sciences, University of Miami, 2008.
 - ³¹ Solo Gabriele HM, Abdelzaher A, Wright M *et al*. *Pathogen Monitoring of a South Florida Beach Which Has Been Characterized by Elevated Microbe Levels*. Tallahassee, FL: Final Report to the Florida Department of Health, 2008.
 - ³² Elmir SM, Shibata T, Solo Gabriele HM *et al*. Quantitative evaluation of enterococci and Bacteroidales released by adults and toddlers in marine water. *Water Research* 2009;**43**:4610–16.
 - ³³ Wright ME, Solo Gabriele HM, Elmir S, Fleming LE. Microbial load from animal feces at a recreational beach. *Mar Pollut Bull* 2009;**58**:1649–56.
 - ³⁴ Sinigalliano CD, Fleisher JH, Gidley ML *et al*. Traditional and molecular analyses for fecal indicator Bacteria in non-point source subtropical recreational marine waters. *Water Research* in press.
 - ³⁵ US Environmental Protection Agency (US EPA). *Improved Enumeration Methods for the Recreational Water Quality Indicators: Enterococci and Escherichia coli*. U.S. Environmental Protection Agency. Washington, DC: Office of Science and Technology, 2000.
 - ³⁶ Wade TJ, Calderon RL, Brenner KP *et al*. High sensitivity of children to swimming-associated gastrointestinal illness: results using a rapid assay of recreational water quality. *Epidemiology* 2008;**19**:375–83.
 - ³⁷ Fleisher JM, Kay D. Risk perception bias, self-reporting of illness, and the validity of reported results in an epidemiologic study of recreational water associated illnesses. *Mar Pollut Bull* 2006;**52**:264–68.
 - ³⁸ Kay D, Bartram J, Prüss A *et al*. Derivation of numerical values for the World Health Organization guidelines for recreational waters. *Water Res* 2004;**38**:1296–304.