

ARTICLE

Expression of *SNURF–SNRPN* upstream transcripts and epigenetic regulatory genes during human spermatogenesis

Michaela Wawrzik¹, Andrej-Nikolai Spiess², Ralf Herrmann³, Karin Buiting¹ and Bernhard Horsthemke^{*,1}

¹Institut für Humangenetik, Universitätsklinikum Essen, Essen, Germany; ²Abteilung für Andrologie, Universitätsklinikum Hamburg-Eppendorf, Hamburg-Eppendorf, Germany; ³Zentrum für Kinderheilkunde, Universitätsklinikum Essen, Essen, Germany

The imprinted domain in human 15q11-q13 is controlled by a bipartite imprinting centre (IC), which overlaps the 5' part of the paternally expressed *SNURF–SNRPN* gene. We have recently described two novel genes upstream of *SNURF–SNRPN* (*PWRN1* and *PWRN2*), which are biallelically expressed in the testis. We have now found that *PWRN1* represents an alternative 5' part of *SNURF–SNRPN*, and that its expression in the brain is imprinted. To determine when the locus is activated during spermatogenesis and which factors are involved in this process, we have mined gene-expression data of testicular biopsies from men with different types of spermatogenic failure. Whereas *PWRN1–SNURF–SNRPN* and *PWRN2* are expressed in post-meiotic germ cells only, a hitherto undetected *SNURF–SNRPN* upstream transcript is expressed already at meiosis. Several epigenetic factors (eg, *MBD1* and *MBD2* isoforms, *MBD3L1*, *SUVH39H2*, *BRDT*, and *EZH2*) are upregulated at specific stages of spermatogenesis, suggesting that they play an important role in the epigenetic reprogramming during spermatogenesis.

European Journal of Human Genetics (2009) 17, 1463–1470; doi:10.1038/ejhg.2009.83; published online 27 May 2009

Keywords: epigenetics; imprinting; spermatogenesis; gene expression

Introduction

Human chromosome 15q11-q13 harbors a 2-Mb imprinted domain, which is regulated by a bipartite imprinting centre (for a review see Horsthemke and Wagstaff¹). One element of the imprinting centre (IC) is defined by the shortest region of deletion overlap in patients with Prader–Willi syndrome and a paternal imprinting defect (PWS-SRO). The other element is defined by the shortest region of deletion overlap in patients with Angelman syndrome and a maternal imprinting defect (AS-SRO). The two elements are ~30 kb apart (see Figure 1a).

The IC overlaps the 5' part of the paternally expressed *SNURF–SNRPN* gene, which has at least three alternative start sites and many alternative exons.^{2,3} In 2000 we identified *C15orf2*, which is a 7.5 kb intronless gene located ~300 kb upstream of the major *SNURF–SNRPN* start site.⁴ *C15orf2* has a 3.5 kb open reading frame encoding an 1156 amino-acid protein of unknown function. Recently we have identified two novel genes (*PWRN1* and *PWRN2*), which are located centromeric to *C15orf2*⁵ and which do not appear to have any protein-coding potential. *PWRN1*, *PWRN2* and *C15orf2* are mainly expressed in the testis, where expression is biallelic. *PWRN1* and *C15orf2* are also expressed in the brain. The finding of monoallelic expression in the brain suggested that these genes are subject to genomic imprinting.⁵ Here we show that *PWRN1* is part of the *SNURF–SNRPN* transcription unit and that expression in the brain is from the paternal allele only.

*Correspondence: Professor B Horsthemke, Institut für Humangenetik, Universitätsklinikum Essen, Hufelandstr 55, Essen D-45122, Germany. Tel: +49 201 723 4556; Fax: +49 201 723 4556; E-mail: bernhard.horsthemke@uni-due.de
Received 5 March 2009; revised 17 April 2009; accepted 23 April 2009; published online 27 May 2009




Figure 1 Physical map. (a) Schematic overview of the ~800 kb chromosomal region from *PWRN2* to *SNURF-SNRPN*. Not drawn to scale. The start and orientation of transcription are indicated by arrows. AS-SRO and PWS-SRO, the two elements of the bipartite imprinting centre. Exons are indicated as black boxes. U6, U7, and U8 are unpublished *SNURF-SNRPN* upstream exons (U6, Chr15: 22.717.397–22.717.495; U7, Chr.15: 22.725.258–22.725.314; U8, Chr.15: 22.746.815–22.746.930). (b) Published EST clones and mRNA fragments. (c) RT-PCR clones generated in this study.

The mechanism by which the IC controls imprinting in 15q11-q13 is unknown. The overlap of the IC with the 5' part of the *SNURF-SNRPN* locus suggests that the transcripts may play a role in this process, although there is no experimental evidence for this to date. Based on the phenotypic consequences of microdeletions affecting the PWS-SRO element of the IC, the AS-SRO element of the IC, or both, we have proposed that the maternal pattern of epigenetic modification, with CpG methylation of *MKRN3*, *NDN* and *SNURF-SNRPN* promoters and silencing of those genes, is the default state of the imprinted domain, that the PWS-SRO is unconditionally required for a chromosome to have the paternal pattern of epigenetic modification and gene expression, and that the AS-SRO is required for a chromosome to have the maternal pattern, if the chromosome has an intact PWS-SRO.¹ In other words, the PWS-SRO acts to activate the paternal copy of the imprinted domain or keep this copy active,⁶ and the AS-SRO counteracts this activity.

As the AS-SRO contains two *SNURF-SNRPN* upstream exons, we and others have proposed that these transcripts may play a role in methylating the PWS-SRO and silencing the region in the female germline.^{2,7} This hypothesis is supported by the work of Chotalia *et al.*,⁸ who found that transcription is required for the establishment of germline-methylation marks at imprinted genes. On the other hand, transgene experiments in which the 880 bp AS-SRO was found to be sufficient for silencing the *SNURF-SNRPN* promoter seem to argue against this hypothesis.⁹ Alternatively, the upstream transcripts may play a role in activating the locus, and the AS-SRO may contain a silencer, which represses transcription in the maternal germline. In any case, it is clear that *trans*-acting factors must be involved.

To lay the foundation for answering these questions, we have investigated the expression patterns of the *SNURF-SNRPN* upstream transcripts, as well as of 20 genes encoding epigenetic factors. Owing to the limited availability of normal human testis samples and technical problems associated with RNA *in situ* hybridization of this material, we have mined gene-expression data of testicular biopsies from men with different types of spermatogenic failure (complete tubular atrophy, Sertoli-cell only syndrome, arrest before meiosis, arrest at meiosis, arrest at round spermatids, and uniform hypospermatogenesis) and with full spermatogenesis.¹⁰ We show that these data can be used to infer gene-expression levels in spermatogonia, spermatocytes and round spermatids.

Materials and methods

Preparation of DNA and RNA

Total RNA and DNA from the fetal brain were extracted using the RNeasy Mini Kit and the QIAamp DNA Mini Kit (Qiagen, Hilden, Germany) according to the manufacturer's recommendations.

Exon connection PCR and RT-PCR

Exon connection PCR was performed on human cDNA from the testis and the fetal brain (Marathon-Ready cDNA, Clontech, Heidelberg, Germany) with the following primers: *SNRPN*_Exon23 (5'-TCCTAACAACTACTATAAGCTTGG-3') + *SNRPN*_4_rw (5'-GATGCTTCACGGTTTTCACA-3') (annealing temperature 58°C) and *SNRPN*_4_fw (5'-CCCA CAAGTAGGGTTGACCA-3') + *SNRPN*_U5 (5'-CAGATTCTTCTGCCGCAAAC-3') (annealing temperature 58°C).

RT-PCR for characterization of mRNA BC035402 was performed on human testis cDNA (Marathon-Ready cDNA, Clontech) with the following primers: BC035402_fw1

(5'-TTAACCTGCGTTTTCCCGTA-3') + BC04ex_rw (5'-AAACACAACACCCCAAGCAT-3') (annealing temperature 60°C) and BC035402_fw1 (5'-TTAACCTGCGTTTTCCCGTA-3') + BC04exnest_rw (5'-TGTTGCAAGGAACGAGACTG-3') (annealing temperature 60°C).

PCR products were re-amplified, subcloned into the pGEM-T Easy vector (Promega, Mannheim, Germany) and sequenced with specific primers. In the case of multiple PCR products, the products were excised from the gel, purified with the MinElute Gel Extraction Kit (Qiagen) and sequenced directly or sequenced with specific primers after subcloning into the pGEM-T Easy vector.

Sequence analysis

Sequencing reactions were carried out with fluorescence-tagged dideoxynucleotides (BIG Dye kit v1.1, Applied Biosystems, Foster City, CA, USA) and the *Taq* cycle sequencing procedure. Sequences were analysed on an ABI 3100 DNA sequencer.

Allele-specific expression analysis

DNA from the fetal brain (cerebrum) and corresponding parental DNA samples from blood were analysed for an expressed SNP (rs12908526) in exon 26 of *PWRN1* by PCR. For sequencing, the following primers were used: Ex26SNPF (5'-GGAACAAGTTTATTACACACAGA-3') and Ex26SNPR (5'-TGTC AACAGGGAAAGTGGA-3') (annealing temperature 58°C). RNA from the same fetal brain was reverse transcribed to cDNA with random hexamers and also analysed for SNP rs12908526. To check the integrity of the RNA and to exclude contamination with the genomic DNA, we used primers for the β -actin gene as a control.⁵

Expression profiles

The expression data of spermatogenesis genes, genes encoding DNA and histone-binding and -modifying proteins, as well as the *SNURF-SNRPN* upstream transcripts were extracted from the dataset published by Spiess *et al.*¹⁰ To corroborate the findings, which were obtained with Affymetrix GeneChip microarrays (Affymetrix, Santa Clara, CA, USA), we included expression data acquired with a completely different platform based on single-oligonucleotide probes (Codewalk system, Applied Microarray Inc., Tempe, AZ, USA) in the analysis.¹¹ This was done by building an intersection of the two platforms using HUGO gene abbreviations. Hierarchical cluster analysis was carried out with the mean expression values from the different stages of spermatogenic impairment and clustered with the TIGR MeV software v4.3 (The TIGR Consortium, <http://www.tm4.org/mev.html>) using Manhattan distance and average linkage. Genes that were common to both platforms (GeneChip and Codewalk), and which displayed similar expression patterns, were denoted with one asterisk on the heatmap and two asterisks if expression patterns were not concordant.

Results

PWRN1 is part of the IC-*SNURF-SNRPN* transcription unit

A database search (UCSC browser hg18, <http://genome.ucsc.edu/cgi-bin/hgGateway>) for expressed sequences upstream of *SNURF-SNRPN* revealed the presence of two overlapping spliced EST clones, DB525036 and BI825340. DB525036 consists of five exons and BI825340 consists of four exons. BI825340 shares two exons with DB525036, but contains two additional exons not present in EST DB525036 (see Figure 1b). By detailed sequence comparison we found that the most upstream exon of EST DB525036 represents exon 23 of *PWRN1*. *PWRN1* has no protein-coding potential and is subject to alternative splicing and polyadenylation.⁵ As DB525036 lacks exons 24–26 of *PWRN1*, but has additional four exons downstream to exon 26, it represents a novel splice variant, which extends more telomeric. Interestingly, the novel exons flank *C15orf2*, which lies now within an intron of *PWRN1*. By exon connection PCR we were able to verify the novel exons in fetal brain cDNA and could also identify one additional exon (see RT1 in Figure 1c). The most telomeric novel exon maps only 285 bp proximal to *UIB*, which represents one of the two known alternative start sites of the *SNURF-SNRPN* upstream transcripts. *PWRN1* transcripts display the same complex splicing pattern as previously observed for the upstream transcripts of *SNURF-SNRPN*, thus it was possible that the *PWRN1* exons represent an additional part of the complex IC-*SNURF-SNRPN* transcription transcripts. To find out whether this is true we performed exon connection PCR on cDNA of the fetal brain and the testis with primers that anneal in *SNURF-SNRPN* exons U2 and U5 and in the last but one novel downstream exon of *PWRN1*. Indeed, we could connect *PWRN1* with the upstream exon U5 in both tissues (see RT3 and RT4 in Figure 1c). Furthermore, we found in testis cDNA a novel exon between U3 and U4 (see RT4 in Figure 1c).

PWRN1 expression is imprinted in the brain

We have recently reported that *PWRN1* shows monoallelic expression in the fetal brain. However, as parental DNA samples were not available at that time, we could not distinguish whether expression is restricted to the paternal or maternal allele. In the meantime, we were able to obtain DNA and RNA samples from fetal brain and parental DNA samples. Analysis of an expressed single-nucleotide repeat (SNP, rs12908526, A/T) in exon 26 of *PWRN1* showed heterozygosity (A/T) in fetal brain DNA (Figure 2). The mother was homozygous for the A allele and the father was homozygous for the T allele. In the fetal brain RNA only the T allele was detected, indicating that monoallelic expression is paternal. These findings suggest that expression is imprinted.




Figure 2 Paternal expression of *PWRN1*. Sequence analysis of DNA samples from mother, father, and fetus (upper three panels) around a single-nucleotide polymorphism in exon 26 (rs12908526). The fetus is heterozygous, but expresses only the paternal (T) allele (lower panel).

Expression profiles

The finding of biallelic expression of the *SNURF-SNRPN* upstream transcripts in the testis raised three questions: (i) Are these transcripts expressed in somatic cells or in germ cells? (ii) If they are expressed in germ cells, at which stage of spermatogenesis are they expressed? (iii) Which epigenetic factors regulate the expression? To address these questions we have mined gene-expression data of testicular biopsies from men with different types of spermatogenic failure as determined with Affymetrix GeneChips.¹⁰ The arrest types used in this study differ essentially in their germ-cell composition so that data can be extracted by filtering genes that pertain to those germ-cell types that appear from a higher impaired pathology to a lower one. In detail, this would be the following (using the abbreviations defined in the legend to Figure 3)

CTA => SCO: appearance of Sertoli cells

SCO => PMA: appearance of spermatogonia (pre-meiotic arrest)

PMA => MA: appearance of primary/secondary spermatocytes (meiotic arrest)

MA => RS: appearance of round spermatids

RS => HYS: appearance of low amounts of elongated spermatids and testicular spermatozoa

To determine whether these data can be used to determine the upregulation of genes at specific stages of spermatogenesis, we examined several spermatogenesis genes. The synaptonemal complex protein 3 (*SYCP3*) is involved in the pairing of homologous chromosomes during meiosis I. As shown in Figure 3a, the *SYCP3* gene is expressed at very low levels in the testis lacking germ cells (CTA and SCO). Low expression was observed in men with spermatogenic arrest before meiosis (PMA), and high expression was observed in men with spermatogenic arrest at meiosis (MA) and at the round spermatid stage (RS), as well as in hypo- (HYS) and normospermic (FS) men. Transition protein 1 (*TNP1*) as well as the protamines *PRM1* and *PRM2* are involved in the packaging of DNA after meiosis into the sperm nucleus. As expected, these genes are expressed at high levels only in men with spermatogenic arrest at the round spermatid stage, as well as in hypo- and normospermic men.

These data show that the expression profiles of testicular biopsies from men with spermatogenic failure can be used to determine the stage-specific upregulation of genes during spermatogenesis. Therefore, we used this dataset to analyze the developmental expression profile of the *SNURF-SNRPN* upstream transcripts and genes encoding DNA and histone-binding and -modifying proteins. Several genes are represented by two or more probe sets on the microarray, and in some cases the different probe sets detect different spliceforms (isoforms).

Using hierarchical cluster analysis (Figure 3b), we identified four clusters of genes. Cluster 1 contains genes with little difference in expression (*DNMT2*, *DNMT3A*, *DNMT3B*, *DNMT3L*, *MBD3*, *MBD3L2*, *MBD4*, *MBD5*, *MBD6*, *MECP2*, *SUV39H1*, *CTCF*, and isoforms of *MBD1* and *MBD2*). Some of these genes tend to be expressed at a lower level in men with late spermatogenic arrest or normal spermatogenesis, indicating that the expression is mainly in somatic cells. We did not find any stage-specific differences in the expression of histone deacetylases (HDCA) either (not shown). Cluster 2 contains only one gene (*DNMT3L*), which is below the level of detection.

Cluster 3 contains genes that are mainly expressed in germ cells and upregulated at late meiosis or after meiosis. This cluster includes the spermatogenesis genes *TNP1*, *PRM1* and *PRM2*, as well as *MBD1*, *MBD3L1*, *BRDT*, and several IC-*SNURF-SNRPN* transcripts.

The remaining genes form a somewhat heterogeneous cluster (Cluster 4). Most of these genes are upregulated




Figure 3 Heatmap of the mean expression values of selected genes in each group of men as determined on the Affymetrix GeneChip platform. Types of spermatogenic failure (number of individuals): CTA, complete tubular atrophy ($n=1$); SCO, Sertoli-cell only ($n=7$); PMA, premeiotic arrest of spermatogenesis ($n=3$); MA, meiotic arrest of spermatogenesis ($n=5$); RS, presence of round spermatids ($n=4$); HYS, hypospermatogenesis ($n=7$); FS, full spermatogenesis ($n=8$). Dark blue indicates no expression ($\log_2=2$), yellow a high expression ($\log_2=10$). (a) Expression of four spermatogenesis genes. (b) Hierarchical cluster analysis reveals four clusters (1–4). Note that several genes are represented by different probe sets. The expression pattern of genes marked with one asterisk could be confirmed by a different microarray platform based on single-oligonucleotide probes (Codelink, see Materials and methods). The two asterisks denote different patterns between the two platforms.

before or at early meiosis. This cluster includes the spermatogenesis gene *SYCP3*, as well as *DNMT1*, *MBD1*, *MBD2*, *MBD4*, *MDB6*, *SUV39H2*, *EZH2*, *CTCF*, and one *IC-SNURF-SNRPN* transcript. Among the epigenetic genes of this cluster, *MBD2*, *EZH2*, *SUV39H2* and *CTCF* transcripts are clearly germ cell-specific and upregulated before meiosis. A detailed analysis of the histone methyltransferase genes *SUV39H1*, *SUV39H2* and *EZH2* is shown in Figure 4a.

To corroborate these findings, genes displayed in the heatmap of Figure 3 were interrogated on a dataset from a completely different, single-oligonucleotide microarray platform¹¹ by using the intersection obtained from HUGO annotations. This cross-platform comparison validates the expression changes described here and obviates the need to

validate single genes, as has been shown in a study in which 93% of genes common to two microarray platforms could be confirmed in their differential expression by quantitative real-time PCR¹². Similar to the results described there, we also observed a high concordance in the gene-expression profiles obtained from the two platforms (16 out of 18; see asterisks in Figure 3).

MBD1, *MBD2* and *IC-SNURF-SNRPN* are represented by different probe sets, which revealed different patterns of expression. The detailed expression pattern and the location of the *MBD1* and *MBD2* probe sets are shown in Figure 4b and Figure 4c, respectively. This analysis shows that the testis-specific isoforms of *MBD1* and *MBD2* use alternative 3' exons. The premeiotic form of *MBD2* is a truncated form.




Figure 4 Detailed diagrams of gene-expression values in individual biopsies. Each position on the x-axis represents one individual. The background of the expression is between $\log_2 = 3-4$. Abbreviations are as in Figure 3. (a) Expression profile of *SUV39H1* (green), *SUV39H2* (magenta), and *EZH2* (blue). (b) Expression of the *MBD1* isoforms (a, magenta; b, blue). Below the diagram is an overview of the different splice forms as given in the UCSC browser hg18. The two probe sets detect alternative 3' exons. (c) Expression of the short (blue) and the long isoform (magenta) of *MBD2*. The location of the probe sets is indicated below the diagram. (d) Expression of *PWRN2* (blue), *PWRN1* (magenta), and the mRNA BC035402 (green), which maps between the *SNURF-SNRPN* upstream exons U1B* and U1A. The location of the probe sets is indicated below the diagram.

As shown in Figure 4d, *PWRN1* and *PWRN2* are upregulated after meiosis. In contrast, the transcript containing the mRNA BC035402 is upregulated at meiosis. This mRNA represents no known upstream exon of *SNURF–SNRPN*, and we failed to link it to any of the known upstream exons in the testis, the fetal brain and leukocyte cDNA. Although the mRNA is colinear with genomic DNA, we identified two spliced alternative isoforms by RT-PCR of testis cDNA (data not shown).

Discussion

As shown by Spiess *et al.*,¹⁰ the microarray-based analysis of mRNA levels in testicular biopsies from fertile men and from men with different types of spermatogenic failure is highly reliable: in 19 of 20 genes tested, the expression patterns could be confirmed by quantitative RT-PCR.¹⁰ For validation of the gene-expression data described here, we used a second, independent dataset of human spermatogenesis.¹¹ Except for *EZH2* and *MBD1*, concordant expression profiles were found. The apparent discrepancy regarding the latter two genes may, at least in the case of *MBD1*, be because of different probes on the two array platforms and the presence of different splice forms (see below). The validity of our data is further supported by the finding that the expression values of different individuals within each disease group are very similar, with the exception of hypospermatogenesis (see Figure 4). The latter group is likely to be etiologically heterogeneous, and the expression patterns may help to classify these patients further. Last, but not least, in those cases in which mouse data are available, the gene-expression profiles described here are in complete concordance with the gene-expression profiles observed in the mouse (see below).

Here we have used the data to investigate the stage-specific upregulation of the *SNURF–SNRPN* locus as well as epigenetic regulator genes during male germ-cell development. The expression patterns of the spermatogenesis genes *SYCP3*, *TNP1*, *PRM1* and *PRM2* showed that this is a valid approach. It should be noted, however, that this approach has two limitations. (i) It is not possible to analyze genes expressed in germ cells and somatic cells, unless there is a germ-cell-specific isoform, because the presence of somatic transcripts in the biopsies masks possible differences in the abundance of germ-cell-specific transcripts. (ii) It is not possible to study the stage-specific downregulation of genes, because each biopsy contains cells at different stages of the germ-cell development. For example, biopsies from men with arrest at the round spermatid stage contain all stages of spermatogonia and spermatocytes. Thus, a gene that is active in spermatocytes but silenced in round spermatids will show up in these biopsies, as well as in biopsies from men with hypospermatogenesis and full spermatogenesis.

We have recently reported the identification of two novel genes upstream of *SNURF–SNRPN*.⁵ Here we show that *PWRN1* is not an independent gene, but an alternative 5' part of the *SNURF–SNRPN* gene. Similar to other transcripts made from this locus, expression in brain is imprinted and from the paternal chromosome only. It was unclear why *PWRN1* and *PWRN2*, which is transcribed from the opposite strand, are biallelically expressed in the testis. Here we show that both transcripts are made in post-meiotic germ cells, that is, from the haploid genome after the establishment of the paternal imprint. If they play a role in imprinting, they may not have a role in establishing the paternal imprint, but in maintaining the paternal allele in an open chromatin configuration.

In contrast to the *PWRN1* and *PWRN2* transcripts, the transcript containing the mRNA BC035402 is upregulated at meiosis. We have not been able to link this transcript to other *SNURF–SNRPN* upstream exons, and thus the identity of this transcript remains elusive.

To determine which epigenetic factors might be involved in upregulating the upstream transcripts and how this relates to the timing of epigenetic reprogramming during spermatogenesis,¹³ we have analysed the expression patterns of several epigenetic regulator genes. Our results are in agreement with and extend previous findings in the mouse and/or humans. We confirm in humans that CTCFL is a germ-cell-specific factor that is upregulated before meiosis. As shown in the mouse by Loukinov *et al.*,¹⁴ CTCFL is expressed in a mutually exclusive manner with CTCF during male germ-cell development, and the erasure of methylation marks during male germ-line development is associated with dramatic up-regulation of CTCFL and down-regulation of CTCF expression. This makes CTCFL a prime candidate protein for the epigenetic reprogramming factor acting in the male germ line.

SUV39H2, which encodes a testis-specific histone H3 lysine 9 methyltransferase related to *SUV39H1*, is upregulated during meiosis. Immunolocalization of the murine Suv39h2 protein during spermatogenesis by O'Carroll *et al.*,¹⁵ indicated enriched distribution at the heterochromatin from the leptotene to the round spermatid stage.

BRDT is a testis-specific member of the distinctive BET sub-family of bromodomain motif-containing proteins, a motif that binds acetylated lysines and is implicated in chromatin remodeling. As shown by Shang *et al.*,¹⁶ in the mouse, its expression is restricted to the germ line, specifically to pachytene and diplotene spermatocytes and early spermatids. Our findings are consistent with observations in the mouse.

MBD3L1 is a methyl-cytosine binding transcriptional repressor. As reported earlier by Jiang *et al.*¹⁷ and suggested by our findings, the expression of MBD3L1 is observed only in round spermatids, suggesting a role for the gene product in the postmeiotic stages of male germ-cell development.

We confirm that the *MBD2* gene encodes at least two isoforms of a methyl-CpG binding domain protein, one of which is testis-specific.¹⁸ It was unknown, however, when the testis-specific isoform is expressed. Here we show that it is expressed very early during spermatogenesis, before the meiosis. The function of *MBD2* has been a matter of dispute. Bhattacharya *et al.*¹⁹ have suggested that it acts as a DNA demethylase and is thus a transcriptional activator. However, no other group has been able to reproduce this finding. In fact, Ng *et al.*²⁰ have found strong evidence that *MBD2* acts as transcriptional repressor. Both isoforms contain the methyl-CpG binding domain. It is tempting to speculate that the short, testis-specific isoform lacks repressor activity and demethylates DNA. If so, our findings suggest that it may be involved in erasing methylation patterns in primordial germ cells or spermatogonia.

We show for the first time that the male germ line expresses an *MBD1* isoform, which is derived from an alternative 3' exon. In contrast to the above-mentioned isoform of *MBD2*, it is not shorter than the somatic isoform and is expressed somewhat later than *MBD2*. We also show for the first time that *EZH2*, which encodes a Polycomb group protein involved in methylating histone H3 at lysine 27, is expressed very early during spermatogenesis.

In summary, we have found that several epigenetic factors (*MBD1* and *MBD2* isoforms, *MBD3L1*, *SUVH39H2*, *BRDT*, and *EZH2*) are upregulated at specific stages of human spermatogenesis, suggesting that they play an important role in the epigenetic reprogramming of the genome during spermatogenesis. More studies are necessary to elucidate all the *cis*- and *trans*-acting factors involved in activating the paternal copy of 15q11-q13.

Acknowledgements

We thank Professor C Kirchhoff for helpful discussions and the Deutsche Forschungsgemeinschaft for financial support.

Ethical statement

This study was approved by the local ethic committee.

References

- Horsthemke B, Wagstaff J: Mechanisms of imprinting of the Prader-Willi/Angelman region. *Am J Med Genet A* 2008; **146A**: 2041–2052.
- Dittrich B, Buiting K, Korn B *et al*: Imprint switching on human chromosome 15 may involve alternative transcripts of the *SNRPN* gene. *Nat Genet* 1996; **14**: 163–170.
- Färber C, Dittrich B, Buiting K, Horsthemke B: The chromosome 15 imprinting centre (IC) region has undergone multiple duplication events and contains an upstream exon of *SNRPN* that is deleted in all Angelman syndrome patients with an IC microdeletion. *Hum Mol Genet* 1999; **8**: 337–343.
- Färber C, Gross S, Neesen J, Buiting K, Horsthemke B: Identification of a testis-specific gene (*C15orf2*) in the Prader-Willi syndrome region on chromosome 15. *Genomics* 2000; **65**: 174–183.
- Buiting K, Nazlican H, Galetzka D, Wawrzik M, Gross S, Horsthemke B: *C15orf2* and a novel noncoding transcript from the Prader-Willi/Angelman syndrome region show monoallelic expression in fetal brain. *Genomics* 2007; **89**: 588–595.
- El-Maarri O, Buiting K, Peery EG *et al*: Maternal methylation imprints on human chromosome 15 are established during or after fertilization. *Nat Genet* 2001; **27**: 341–344.
- Mapendano CK, Kishino T, Miyazaki K *et al*: Expression of the *Snurf-Snrpn* IC transcript in the oocyte and its putative role in the imprinting establishment of the mouse 7C imprinting domain. *J Hum Genet* 2006; **51**: 236–243.
- Chotalia M, Smallwood SA, Ruf N *et al*: Transcription is required for establishment of germline methylation marks at imprinted genes. *Genes Dev* 2009; **23**: 105–117.
- Shemer R, Hershko AY, Perk J *et al*: The imprinting box of the Prader-Willi/Angelman syndrome domain. *Nat Genet* 2000; **26**: 440–443.
- Spieß AN, Feig C, Schulze W *et al*: Cross-platform gene expression signature of human spermatogenic failure reveals inflammatory-like response. *Hum Reprod* 2007; **22**: 2936–2946.
- Feig C, Kirchhoff C, Ivell R, Naether O, Schulze W, Spieß AN: A new paradigm for profiling testicular gene expression during normal and disturbed human spermatogenesis. *Mol Hum Reprod* 2007; **13**: 33–43.
- Bosotti R, Locatelli G, Healy S *et al*: Cross platform microarray analysis for robust identification of differentially expressed genes. *BMC Bioinformatics* 2007; **8**: 1–55.
- Sasaki H, Matsui Y: Epigenetic events in mammalian germ-cell development: reprogramming and beyond. *Nat Rev Genet* 2008; **9**: 129–140.
- Loukinov DI, Pugacheva E, Vatolin S *et al*: BORIS, a novel male germ-line-specific protein associated with epigenetic reprogramming events, shares the same 11-zinc-finger domain with CTCF, the insulator protein involved in reading imprinting marks in the soma. *Proc Natl Acad Sci USA* 2002; **99**: 6806–6811.
- O'Carroll D, Scherthan H, Peters AH *et al*: Isolation and characterization of *Suv39h2*, a second histone H3 methyltransferase gene that displays testis-specific expression. *Mol Cell Biol* 2000; **20**: 9423–9433.
- Shang E, Salazar G, Crowley TE, Wang X, Lopez RA, Wolgemuth DJ: Identification of unique, differentiation stage-specific patterns of expression of the bromodomain-containing genes *Brd2*, *Brd3*, *Brd4*, and *Brd1* in the mouse testis. *Gene Expr Patterns* 2004; **4**: 513–519.
- Jiang CL, Jin SG, Lee DH *et al*: *MBD3L1* and *MBD3L2*, two new proteins homologous to the methyl-CpG-binding proteins *MBD2* and *MBD3*: characterization of *MBD3L1* as a testis-specific transcriptional repressor. *Genomics* 2002; **80**: 621–629.
- Hendrich B, Bird A: Identification and characterization of a family of mammalian methyl-CpG binding proteins. *Mol Cell Biol* 1998; **18**: 6538–6547.
- Bhattacharya SK, Ramchandani S, Cervoni N, Szyf M: A mammalian protein with specific demethylase activity for mCpG DNA. *Nature* 1999; **397**: 579–583.
- Ng HH, Zhang Y, Hendrich B *et al*: *MBD2* is a transcriptional repressor belonging to the MeCP1 histone deacetylase complex. *Nat Genet* 1999; **23**: 58–61.