

ARTICLE

Identification of *ANKRD11* and *ZNF778* as candidate genes for autism and variable cognitive impairment in the novel 16q24.3 microdeletion syndrome

Marjolein H Willemsen^{*1}, Bridget A Fernandez², Carlos A Bacino³, Erica Gerkes⁴, Arjan PM de Brouwer¹, Rolph Pfundt¹, Birgit Sikkema-Raddatz⁴, Stephen W Scherer⁵, Christian R Marshall⁵, Lorraine Potocki³, Hans van Bokhoven¹ and Tjitske Kleefstra¹

The clinical use of array comparative genomic hybridization in the evaluation of patients with multiple congenital anomalies and/or mental retardation has recently led to the discovery of a number of novel microdeletion and microduplication syndromes. We present four male patients with overlapping molecularly defined *de novo* microdeletions of 16q24.3. The clinical features observed in these patients include facial dysmorphisms comprising prominent forehead, large ears, smooth philtrum, pointed chin and wide mouth, variable cognitive impairment, autism spectrum disorder, structural anomalies of the brain, seizures and neonatal thrombocytopenia. Although deletions vary in size, the common region of overlap is only 90 kb and comprises two known genes, *Ankyrin Repeat Domain 11 (ANKRD11)* (MIM 611192) and *Zinc Finger 778 (ZNF778)*, and is located approximately 10 kb distally to *Cadherin 15 (CDH15)* (MIM 114019). This region is not found as a copy number variation in controls. We propose that these patients represent a novel and distinctive microdeletion syndrome, characterized by autism spectrum disorder, variable cognitive impairment, facial dysmorphisms and brain abnormalities. We suggest that haploinsufficiency of *ANKRD11* and/or *ZNF778* contribute to this phenotype and speculate that further investigation of non-deletion patients who have features suggestive of this 16q24.3 microdeletion syndrome might uncover other mutations in one or both of these genes.

European Journal of Human Genetics (2010) 18, 429–435; doi:10.1038/ejhg.2009.192; published online 18 November 2009

Keywords: 16q24.3 microdeletion; *ANKRD11*; *ZNF778*; cognitive impairment; autism

INTRODUCTION

Whole-genome scanning technologies such as array comparative genomic hybridization (array CGH) and single-nucleotide polymorphism oligonucleotide arrays (SNP array) have enabled the detection of submicroscopic chromosomal aberrations, which previously escaped detection by routine cytogenetic and molecular cytogenetic techniques.^{1–3} These methods have proven invaluable in the elucidation of genomic regions associated with mental retardation and/or congenital anomalies.^{4–8} Several clinically distinct microdeletion and microduplication syndromes have been reported on the basis of data derived from these techniques, such as the 17q21.31 microdeletion syndrome⁵ and the 1q41–1q42 microdeletion syndrome,⁹ as well as microduplication syndromes involving 7q11.23^{10,11} and 17p11.2.¹² The phenotypic characteristics of microdeletion syndromes can be caused by haploinsufficiency of single genes, for example, *TCF4* (MIM 602272) in Pitt–Hopkins syndrome,¹³ *EHMT1* (MIM 607001) in the 9q34.3 subtelomeric deletion syndrome¹⁴ and either *CREBBP* (MIM 600140) or *EP300* (MIM 602700) in Rubinstein–Taybi syndrome.^{15,16} The application of genome-wide array technologies with an increasing density of probes has led to the identification and

localization of several genes associated with developmental disorders or abnormal brain development,⁸ including *CHD7* (MIM 608892) in CHARGE syndrome¹⁷ and *FOXG1* (MIM 164874) in congenital Rett syndrome.¹⁸ Aberrations of chromosome 16q with clinical relevance have rarely been reported. Before this report, interstitial deletions restricted to band 16q24.3 have not been described. There are only a few reports in medical literature with regard to patients with larger, cytogenetically visible deletions comprising this region, mostly because of an unbalanced complex chromosomal rearrangement.^{19–29} In this study, we aimed to characterize the clinical and molecular features of four patients with submicroscopic interstitial 16q24.3 microdeletions ascertained by genome-wide array analysis and to determine the shortest region of overlap (SRO) to identify candidate genes responsible for their overlapping phenotype.

PATIENTS AND METHODS

Patient 1

Patient 1 was ascertained at the age of 22 years when SNP array was performed because of mental retardation, autism spectrum disorder, dysmorphic features and congenital anomalies. He was born after an uncomplicated pregnancy,

¹Department of Human Genetics, Radboud University Nijmegen Medical Centre, Nijmegen, The Netherlands; ²Disciplines of Genetics and Medicine, Memorial University of Newfoundland and Provincial Medical Genetic Program, Eastern Health, St John's, Newfoundland and Labrador, Canada; ³Department of Molecular and Human Genetics, Baylor College of Medicine, Texas Children's Hospital, Houston, TX, USA; ⁴Department of Genetics, University Medical Centre Groningen, Groningen, The Netherlands; ⁵The Centre for Applied Genomics, The Hospital for Sick Children and University of Toronto, Toronto, Ontario, Canada
*Correspondence: Dr MH Willemsen, 849 Department of Human Genetics, Radboud University Nijmegen Medical Centre, PO Box 9101, Nijmegen 6500 HB, The Netherlands. Tel: +31 24 3613 946; Fax: +31 24 3668 753; E-mail: m.willemsen@antrg.umcn.nl

Received 30 June 2009; revised 2 September 2009; accepted 22 September 2009; published online 18 November 2009

induced at 36-week gestation because of maternal preeclampsia. Growth parameters were appropriate for gestational age, with a birth weight between the 50th and 75th percentile. Head circumference was not measured at birth. He was diagnosed with congenital hip dysplasia, but had no other difficulties in the neonatal period. His psychomotor development was delayed. He walked independently at the age of 2.5 years and began using single words after the age of 4 years. At the age of 3 years, he was referred to a pediatric neurologist because of his developmental delay and epilepsy (absence and generalized epilepsy). Cerebral imaging with a computer tomography scan (CT-scan) showed colpocephaly, hypoplasia of the corpus callosum and heterotopia. Vision and audiological evaluations were normal. Routine chromosome analysis showed a normal male karyotype. At the age of 22 years, he was referred for reevaluation. He was moderately mentally retarded and showed behavior consistent with autism spectrum disorder. A Dutch formal test for autism spectrum disorders (AVZ-R 'Autisme- en Verwante stoornissenschaal-Z- Revisie', 1999) showed borderline results for the diagnosis of an autism spectrum disorder. His epilepsy was well under control and medical treatment was gradually stopped. On physical examination, he exhibited dysmorphic features, with a high forehead, bipitonal narrowing, long palpebral fissures, large ears, smooth philtrum, wide mouth and micrognathia (Figure 1c and d). He had kyphoscoliosis and strabismus. His height was 175 cm (3–10th percentile), weight was 64 kg (50th percentile) and head circumference was 59.5 cm (90th percentile). Cardiac ultrasound revealed no abnormalities. Bone densitometry showed normal bone density. Genetic diagnostic evaluation with subtelomeric multiplex ligation-dependent probe amplification (MLPA, SALSA MLPA Kit P036, MRC Holland, Amsterdam, The Netherlands) was normal.

Patient 2

A cursory analysis of patient 2 has been included in a large Canadian autism cohort study,³⁰ and here we provide a more detailed clinical and molecular description. Patient 2 was ascertained at the age of 3 years and 3 months when SNP array analysis was performed because of developmental delay, autism spectrum disorder and dysmorphic features. He was born at term after an uncomplicated pregnancy and delivered by cesarean section on maternal indication. He was born with strikingly outstanding and asymmetric ears. On day 3, he developed low platelets. It was documented as having been caused by alloimmune idiopathic thrombocytopenic purpura (ITP) because of his mother's ITP. After treatment, his platelets remained within the normal range. Psychomotor development was delayed. At 21 months of age, he was unable to walk independently and used three single words with meaning. He was diagnosed with autism by formal tests at the age of 2 years by ADOS-1 (Autism Diagnostic Observation Schedule) and at the age of 3 years and 4 months by ADI-R (Autism Diagnostic Interview Revised). Psychometric testing at the age of 6 years and 5 months showed average results for nonverbal intelligence; The Naglieri Nonverbal Ability test score was at the 50th percentile and Test of nonverbal intelligence gave a nonverbal IQ of 100. Preschool Language Scale 4 (PLS-4) showed that auditory comprehension was severely delayed (1st percentile) and that expressive communication was moderately delayed (4th percentile). On physical examination, he had a height of 94 cm (10–25th percentile) and weight of 13.5 kg (10–25th percentile) and a head circumference of 49.5 cm (50th percentile). Facial dysmorphic features included frontal bossing, long palpebral fissures, a wide mouth with full lips, long and smooth philtrum, anteverted nares, pointed chin and large ears (ears originally outstanding/cupped and asymmetric, now after surgical correction) (Figure 1e). Diagnostic studies, including routine G-banded chromosome analysis, FISH for the Williams syndrome microdeletion and molecular analysis for fragile X syndrome, were negative. CT-scan of the brain was normal and examination by an ophthalmologist did not show any abnormalities.

Patient 3

Patient 3 was ascertained at the age of 6 years and 3 months when array CGH analysis was performed because of mental retardation, dysmorphic features and congenital anomalies. He was born at term by vaginal delivery after an uncomplicated pregnancy and his size was appropriate for gestational age. At delivery, he was noted to have a skin rash and hepatosplenomegaly with thrombocytopenia. TORCH (toxoplasmosis, rubella, cytomegalovirus, herpes)

infection was excluded and organomegaly and thrombocytopenia were resolved. He suffered a mild intracranial hemorrhage secondary to thrombocytopenia, and was diagnosed with a ventricular septal defect (VSD), patent foramen ovale (PFO), cleft mitral valve and left cryptorchidism in the newborn period. VSD and PFO were repaired at the age of 3 months. Routine G-banded chromosome analysis was normal. At the time of genetics consultation at the age of 6 years and 3 months, he was admitted to hospital for an acute parvovirus myocarditis and dilated cardiomyopathy with severely depressed biventricular function. Further evaluation revealed a history of developmental delay (he walked at the age of 3 years, and spoke his first words at 18–20 months), mental retardation and febrile seizures. On physical examination, he had a height on the 5–12th percentile and a head circumference on the 10th percentile. Dysmorphic features that were observed included a triangular face with a high forehead and frontal bossing, arched eyebrows, large ears, smooth philtrum, wide mouth, pointed chin and high palate (Figure 1f and g). MRI of the brain revealed a posterior fossa arachnoid cyst, thinned corpus callosum, periventricular heterotopias, optic nerve hypoplasia and evidence for an old left hemorrhagic parietal infarct. The heterotopias were unilaterally localized along the lateral margin of the trigone of the right lateral ventricle extending along the temporal horn of the right lateral ventricle. Ophthalmic examination revealed high myopia, intermittent exotropia, horizontal nystagmus, and possible right amblyopia. Audiogram revealed moderate-to-severe bilateral mixed sensorineural and conductive hearing impairment. His complete blood count showed mild macrocytosis (MCV 92.4 fL; normal 76–90), with normal homocysteine and folate values. The cardiomyopathy and macrocytosis resolved, yet he developed complex partial seizures at the age of 7½ years, with EEG showing a seizure focus of the left temporal region. He continued to make progress in school. He meets criteria for autism under the ADOS.

Patient 4

Patient 4 was ascertained at the age of 8 years and 10 months when array CGH analysis was performed because of psychomotor retardation, features of autism spectrum disorder and dysmorphic features. He was born at term after an uneventful pregnancy by cesarean section because of breech presentation. His birth weight was 3850 g (50–90th percentile), but birth length and head circumference are unknown. In the first few weeks, there were feeding problems, probably caused by poor sucking. There was no hypotonia. Psychomotor development was slightly delayed with the child being able to sit independently at 12 months of age and able to walk independently at the age of 18 months. Language development was delayed. Throughout childhood, there were periods of obstipation. There were frequent ear infections. Hearing was tested normal. There is severe bilateral astigmatism. At the age of 7 years and 2 months, verbal IQ was 81 and performal IQ was 67, as assessed by the WISC III intelligence test. He was diagnosed with an Autism Spectrum Disorder (ASD)/Pervasive Developmental Disorder Not Otherwise Specified (PDD–NOS) by a child psychologist and psychiatrist. There are no signs of epilepsy. On physical examination, he showed dysmorphic features with a high and broad forehead, mildly upslanting palpebral fissures, a double hair whorl, large ears, a preauricular tag and bilateral fusion of the central and lateral incisors (Figure 1h and i). His height was 126.3 cm (10–25th percentile), weight was 23.7 kg (5–25th percentile) and head circumference was 52.2 cm (50–75th percentile). Cardiac examination was normal. The left hand showed a single transverse palmar crease. Thumbs were slightly proximally placed. X-rays of the hands showed pseudo-epiphyses of the second metacarpals. Routine G-banded chromosome analysis showed a normal male karyotype. Molecular analysis for fragile X syndrome was negative. Complete blood count showed no abnormalities at the age of 8 years.

Methods

DNA was obtained from peripheral blood leukocytes and isolated according to standard procedures.

In patient 1 and 2 and both parental couples, a 500K SNP array analysis with the combined two-chip Affymetrix *Nspl1* and *Sty1* GeneChip Human Mapping Commercial was performed according to the standard Affymetrix GeneChip protocol (Affymetrix Inc., Santa Clara, CA, USA). For CNV validation in patient 2, multiple SYBR Green-based qPCR assays were used (primer



Figure 1 Facial profiles of patients 1–4. (a–c) Patient 1 at several ages, from 6 years (a) to 22 years (b and c). (d–f) Patient 2 at the age of 3 years (d) and at the age of 8 years (e and f). (g and h) Patient 3 at the age of 7 years. (i and j) Patient 4 at the age of 8 years and 10 months. Facial features comprise high forehead/frontal bossing and large ears (all patients), broad mouth, long smooth philtrum and pointed chin (patient 1–3). Note the change in facial phenotype in patient 1 at adult age, showing a long and oval face with a full upturned nose, retrognathia and a pronounced groove in the chin.

sequences available upon request) to measure relative copy number in the patient, parents and controls between this chromosome 16q24.3 and a control region (*FOXP2*). The same approach was used to validate a *de novo* 500 kb CNV gain at 3p14.2 near the common fragile site *FRA3B* in patient 2. Subsequently, patient 2 was also tested on the Affymetrix 6.0 array.

In patient 3 and his parents, a 244K oligo array analysis, according to the Agilent protocol (Agilent Technologies Inc., Santa Clara, CA, USA), was performed.

In patient 4 and his parents, a 105K oligo array according to the Agilent protocol was performed (design ID 019015, Agilent Technologies Inc.).

RESULTS

Clinical features

A summary of the clinical features of the four patients with a chromosome 16q24.3 microdeletion is shown in Table 1. All patients show a prominent forehead and large ears. Moreover, patients 1–3 share other distinct features (Figure 1a–g), including long smooth philtrum, broad mouth and pointed chin. These were not observed in

patient 4 (Figure 1h and i). In addition to variable cognitive impairment, ranging from moderate mental retardation to normal nonverbal IQ with moderate-to-severe speech delay, all four patients had either autistic features (patient 1) or met the diagnostic criteria for an ASD (patients 2–4). Interestingly, brain abnormalities, including structural anomalies and neuronal migration disorders, were noticed in two of the three patients who underwent brain imaging (patient 1 and 3). Epilepsy and (transient) thrombocytopenia were each noticed in two of the four patients. Nonspecific ocular problems were seen in three of the four patients. One patient, patient 3, with the largest deleted region, had a ventricular septal defect and cleft mitral valve.

Molecular findings

The *de novo* CNV results observed in the four patients are summarized in Table 1 and the overlapping deletions at chromosome 16q24.3 are schematically shown in Figure 2. All deletions were mapped according to the USCS genome browser build May 2004.

In patient 1, an interstitial 378 kb loss was identified (87.65–88.03 Mb): 46, XY, arr snp 16q24.3 (SNP_A-1895824→SNP_A-2058988)×1 dn.

In patient 2, an interstitial loss of 265 kb (87.80–88.06 Mb): 46, XY, 16q24.3 (SNP_A-2223520→SNP_A-4274603)×1 dn was identified. A detailed analysis of the qPCR data in patient 2 revealed that the deletion does not extend to the last exon of *CDH15*. According to the assays, coordinates 87 788 807–87 788 909 (Build 36) are not deleted (results not shown). Patient 2 also carried a 500 kb *de novo* gain at 3p14.2. An interstitial loss of 2.07 Mb in chromosome band 16q24.2q24.3 was identified in patient 3 (86.06–88.13 Mb) (probe A_16_P40717088→probe A_16_P20559418). In patient 4, an interstitial loss of 1.1 Mb in chromosome band 16q24.2q24.3 was detected (86.79–87.89 Mb) (probe A_14_P119277→A_14_P111955).

In all four cases, analysis of both parents showed normal copy numbers for the deleted regions, suggesting a *de novo* origin of the deletions. These four chromosome 16q24.3 deletion cases are the only cases that have been detected in a total cohort of more than 3000

patients with developmental delay and/or autism or congenital anomalies. The same region was interrogated over 1000 population controls³¹ and found not to be copy number variable. *The Database of Genomics Variants*³² shows one CNV (Variation_4018) at this site,³³ involving the first two exons of *ANKRD11* but not *ZNF778*, but it is not found in other studies.

DISCUSSION

We present detailed clinical and molecular features of four patients with microdeletions within the chromosomal band 16q24.3. The four patients have autistic features or a diagnosis of ASD, variable cognitive impairment and facial dysmorphism, and share a 90 kb overlapping region comprising two annotated genes, *Ankyrin Repeat Domain 11* (*ANKRD11*) and *zinc finger 778* (*ZNF778*) (Figure 2). The level of cognitive impairment that was observed in these four patients is considerably variable, ranging from moderate mental retardation in patient 1 and 3 to a normal nonverbal IQ with moderate-to-severe

Table 1 Clinical features of the four presented patients with a 16q24.3 microdeletion

Patient	Patient 1	Patient 2	Patient 3	Patient 4	Total
Size of 16q24 deletion	378 kb	265 kb	2.07 Mb	1.1 Mb	90 kb overlap
Mb positions	87.65–88.03	87.80–88.06	86.06–88.13	86.79–87.89 Mb	SRO: 87.80–87.89 Mb
Age at examination	22 years	3 years, 3 months	6 years, 3 months	8 years, 10 months	Mean age 10 years
Cognitive impairment	Moderate mental retardation	Normal nonverbal IQ with moderate-severe speech delay	Moderate mental retardation	Borderline-normal verbal IQ mildly impaired nonverbal IQ	4/4
Autism spectrum disorder	Borderline	+	+	+	4/4
Height	3–10th percentile	10–25th percentile	5–12th percentile	10–25th percentile	All (low) normal
Head circumference	90th percentile	50th percentile	10th percentile	50–75th percentile	All normal
<i>Facial characteristics</i>					
High forehead	+	–	+	+	3/4
Frontal bossing	–	+	+	–	2/4
Bitemporal narrowing	+	–	–	–	1/4
Long oval face	+	–	–	–	1/4
Long palpebral fissures	+	+	–	–	2/4
Arched eyebrows	–	–	+	–	1/4
Large ears	+	+	+	+	4/4
Smooth philtrum	+	+	+	–	3/4
Broad mouth	+	+	+	–	3/4
Pointed chin	+	+	+	–	3/4
Micrognathia	+	+	–	–	2/4
High palate	+	–	+	–	2/4
Congenital heart defect	–	–	VSD, PFO and cleft mitral valve with severe mitral regurgitation	NT	1/2
Structural brain malformation	Corpus callosum hypoplasia, colpocephaly	–	Corpus callosum hypoplasia, dilated ventricles, optic nerve hypoplasia	NT	2/3
Neuronal migration disorder	Heterotopias	Normal unenhanced CT cerebrum	Perventricular heterotopias (unilateral)	NT	2/3
Seizures	Absences and generalized	–	Partial complex	–	2/4
Ocular problems	Strabismus	–	High myopia, astigmatism, horizontal nystagmus	Severe bilateral astigmatism	3/4
Hearing loss	–	–	Mixed sensorineural and conductive hearing loss	–	1/4
Skeletal anomalies	Kyphoscoliosis	–	–	–	1/4
Genital anomalies	–	–	Unilateral cryptorchidism	–	1/4
Hematologic disorder	–	Neonatal thrombopenia (resolved)	Thrombopenia, macrocytosis (resolved)	–	2/4




Figure 2 Schematic overview of deleted regions on chromosome 16q24.3 in the presented patients. All deletions were mapped according to the USCS genome browser build May 2004. The relative positions of the genes of interest are indicated. The region of 90 kb overlap is demarcated with a gray zone. The deletions in patient 3 and 4 extend beyond the figure (Mb positions of proximal delineations: 86.06 and 86.79, respectively). Arrows indicate the direction in which the genes are transcribed. SRO: shortest region of overlap.

speech delay in patient 2. Patient 2 showed a remarkable discrepancy between his verbal and nonverbal capacities in disadvantage of speech/auditory development. A reverse discrepancy in IQ levels, however less extreme, was reported in patient 4 who had a borderline-to-normal verbal IQ of 81 and a mildly impaired nonverbal IQ of 67.

Remarkably, both patients 1 and 3 have structural brain abnormalities, including hypoplasia of corpus callosum, colpocephaly/dilated ventricles, as well as periventricular heterotopias that might be causative for the epilepsy these two patients experienced. They share a deletion interval comprising *ANKRD11*, *ZNF778* and *CDH15*, suggesting that haploinsufficiency for one or more of these genes is involved in neuronal migration and causes the gray matter heterotopias in patient 1 and 3. The localization of the heterotopias in patient 1 was likely bilateral, but no further details were available by cerebral imaging for this patient. The heterotopias in patient 3, however, were unilaterally localized, which raises the possibility that this migrational defect was caused by an (*in utero*) acquired event rather than by a genetic defect. Both patients 2 and 3 had temporary thrombocytopenia, although the neonatal thrombocytopenia in patient 2 was attributed to his mother's ITP. Nonspecific ocular problems were noted in three of the four patients. Features that were only observed in patient 3 (with the largest deleted region) were a congenital heart defect, unilateral cryptorchidism and sensorineural hearing impairment. Patient 2 also carried a 500 kb *de novo* gain at 3p14.2 at the *FRA3B* (MIM 601153) locus, which may also contribute to his phenotype; however, patient 2 had no other specific features.

By molecular characterization of these four overlapping deletions in 16q24.3, two likely candidate genes for the observed autism phenotype with cognitive impairment were present in the SRO. A third gene, *Cadherin 15* (*CDH15*), is located just proximal to the SRO and is

involved in the deletion observed in patients 1, 3 and 4 (Figure 2). *CDH15* does not seem to be involved in the deletion in patient 2. The possibility that a position effect or disruption of gene regulatory elements could also create a functional null allele for *CDH15* in this patient is not excluded. The potential involvement of *CDH15* is of particular interest, a nonsynonymous variants of this gene have been identified in a cohort of mentally disabled patients.³⁴ However, the group of nonaffected controls in this study was relatively small and the segregation of variants in unaffected family members has not been tested consistently nor was it in agreement with the presumed pathogenicity of the detected variant. Of interest, a recently published genome-wide association study among cases with ASD identified common genetic variants on 5p14.1 that are associated with susceptibility to ASD. The nearby genes, *CDH9* (MIM 609974) and *CDH10* (MIM 604555), are also members of the Cadherin gene family involved in neuronal cell adhesion. This implicates a role of *CDH9* and *CDH10* and possibly of other members of the Cadherin family in the pathogenesis of ASD by affecting neuronal cell adhesion.³⁵

Haploinsufficiency for either or both *ANKRD11* and *ZNF778* is an attractive explanation for the syndromic autism phenotype, with variable cognitive impairment observed in the four patients. *ANKRD11* encodes a member of the family of ankyrin repeat containing cofactors that interacts with p160 nuclear receptor coactivators and histone deacetylases corepressors and inhibits ligand-dependent transcriptional activation of target genes by nuclear receptors.³⁶ Therefore, haploinsufficiency of *ANKRD11* may disrupt transcription regulation. Disruption of epigenetic processes has been shown to have a role in causing mental retardation. Examples are disruption of a methyl-CpG-binding protein caused by mutations in *MECP2* (MIM 30005) in Rett syndrome, mutations in *EMHT1* in the 9q subtelomere

deletion syndrome and mutations in either *CREBBP* or *EP300* in the Rubinstein–Taybi syndrome. In these cases, mental retardation might be caused by epigenetic effects on the expression of genes that are not themselves mutated.³⁷

Interestingly, Barber *et al*³⁸ described a mouse mutant, named *Yoda*, with a missense mutation in the *Ankrd11* gene. Homozygous knockout *Yoda* mice are not viable and die during embryogenesis. This suggests that the gene has a crucial function in embryonic development. A major feature of heterozygous mutant mice is reduced bone mineral density. They also show cranio-facial abnormalities, such as shortened snouts, wider skulls, deformed nasal bones and failure of the interfrontal suture to close. Body size is reduced and with aging many *Yoda* mice developed kyphosis.³⁸ Mild kyphoscoliosis was seen only in the oldest patient (patient 1); however, an examination of bone density in this patient showed normal results. On the basis of *in silico* data, *ANKRD11* is widely expressed, including in several brain regions (amygdala, parietal lobe, occipital lobe and cerebellum) and the heart (UCSC Genome Browser, Microarray Expression Data, Unigene).

The other gene localized in the SRO, *ZNF778*, encodes a member of the extensive family of KRAB-domain zinc finger proteins. On the basis of available *in silico* data, this gene is also expressed in several tissues, including the brain and heart (Unigene). KRAB-domain zinc finger proteins are found in transcription regulatory complexes, which are directed to the regulatory elements of target genes through the C2H2 zinc finger domains that recognize specific DNA binding sites. Moreover, several human zinc finger genes have been associated with MR: *ZNF41* (MIM 314995), *ZNF81* (MIM 314998) and *ZNF674*

(MIM 300573) with X-linked MR^{39–41} and *ZNF385B* (MIM 612344) with the 2q31.2 deletion syndrome.⁴¹ Therefore, haploinsufficiency of *ZNF778* might also, or exclusively, be involved in the phenotype observed in the reported patients.

Table 2 provides an overview of these and other interesting genes that are involved in nonoverlapping deleted regions. Proposed functions, phenotype in knockout mice (if available), previously reported cases and the possible clinical correlation of haploinsufficiency of these genes with the observed phenotype in our patients are indicated. Interestingly, the deletion interval of patients 2 and 3 with transient thrombocytopenia comprises the gene *ZFPM1* (*FOG1*) (MIM 601950), which is involved in the regulation of the expression of genes during erythroid and megakaryocytic differentiation. On the basis of available data, we were unable to recognize a candidate gene for the congenital heart defect observed in patient 3.

In conclusion, we hypothesize that *ANKRD11* and *ZNF778* are strong candidate genes for the observed core phenotype of this novel 16q24.3 microdeletion syndrome comprising autism spectrum disorder, variable cognitive impairment, facial dysmorphism and brain abnormalities, although a role of one or more of the other genes in the deleted regions of individual patients is not excluded. Moreover, the involvement of *ANKRD11* and *ZNF778* warrants further studies searching for *ANKRD11* and/or *ZNF778* mutations in patients with a similar phenotype, but without 16q24 abnormalities by molecular karyotyping. Follow-up of known patients might give answers to the possible occurrence of other phenotypic features. Moreover, the identification of new cases with overlapping deletions is needed to

Table 2 Genes of interest in the entire reported 16q24.3 deletion interval

Gene (Mb positions)	Function	Mouse null	Mouse heterozygous	Previously reported cases	Possible clinical correlate
<i>ANKRD11</i> (87.86–88.08)	Nuclear receptor-mediated transcriptional regulation (inhibitory) ^{36,38}	Lethal ³⁸	Reduced bone mineral density craniofacial abnormalities reduced body size ³⁸	Marshall <i>et al</i> identified a 16q24.3 deletion encompassing <i>ANKRD11</i> in an ASD case. ³⁰	Kyphoscoliosis in patient 1, however he has a normal bone density. Syndromic autism phenotype with variable cognitive impairment in patients 1, 2, 3 and 4
<i>ZNF778</i> (87.81–87.82)	Encodes a member of the KRAB domain containing zinc finger protein family, suggesting involvement in transcription regulation. ^{39,42}	No data available	No data available	Several other human zinc finger genes have been associated with mental retardation. ^{39,42}	Syndromic autism phenotype with variable cognitive impairment in patients 1, 2, 3 and 4
<i>CDH15</i> (87.77–87.79)	Cell–cell adhesion in skeletal muscle and human brain. ^{34,43}	No data available	No data available	Bhalla <i>et al</i> identified 4 nonsynonymous variants of <i>CDH15</i> in a cohort of patients with mental retardation. ³⁴	Cognitive impairment in patient 1, 3 and 4
<i>ZFPM1</i> (<i>FOG1</i>) (87.05–87.13)	Co-regulator of the transcriptional activator <i>GATA1</i> , which regulates the expression of certain genes during erythroid and megakaryocytic differentiation. ⁴⁴	<i>GATA-1</i> knockout mice are not viable. Mice with selective loss of <i>GATA-1</i> expression in megakaryocytes have thrombopenia.	No data available	Freson <i>et al</i> demonstrated a disturbed interaction between mutants of <i>GATA1</i> and <i>FOG1</i> in patients with macrothrombocytopenia and dyserythropoiesis. ⁴⁴	Neonatal thrombocytopenia in patient 2 and 3 unexplained macrocytosis in patient 3
<i>CBFA2T3</i> (MIM 603870) (87.47–87.57)	Regulates the proliferation and the differentiation of erythroid progenitors	No data available	No data available	—	Unexplained macrocytosis and neonatal thrombocytopenia in patient 3
<i>CDT1</i> (87.39–87.40)	Transcriptional regulator (initiation) ⁴⁵	No data available	No data available	—	?

extend phenotypic knowledge and to confirm or even further narrow down the critical region of involved genes.

CONFLICT OF INTEREST

The authors declare no conflict of interest.

ACKNOWLEDGEMENTS

We thank the parents of patients who participated in this study. This work was supported by grants from the Consortium VG Oost-Nederland (to TK and MHW) and the Genome Canada/Ontario Genomics Institute (to SWS and BF). CRM is supported by the SickKids Foundation and the National Alliance for Research on Schizophrenia and Depression (NARSAD). SWS holds the GlaxoSmithKline-CIHR Pathfinder Chair in Genetics and Genomics at the University of Toronto and Hospital for Sick Children. Patient consent was obtained from the patient's parents for publishing patient pictures in Figure 1. We also thank Dr Zhinshuo Ou and the Kleberg Cytogenetics Laboratory at Baylor College of Medicine.

RESOURCES

Database of Genomic Variants <http://projects.tcag.ca/variation/>; Decipher^{v41} Database of Chromosomal Imbalance and Phenotype in Humans using Ensembl Resources <https://decipher.sanger.ac.uk/application/>; European Cytogeneticists Association Register of Unbalanced Chromosome Aberrations (ECARUCA) <http://ecaruca.net>; Ensembl genome browser <http://www.ensembl.org>; Unigene <http://www.ncbi.nlm.nih.gov/UniGene>; USC Genome Browser (Assembly May 2004): <http://genome.ucsc.edu>; UniProt KB, Assembly July 2008 <http://www.uniprot.org>.

- 1 Feuk L, Carson AR, Scherer SW: Structural variation in the human genome. *Nat Rev Genet* 2006; **7**: 85–97.
- 2 Veltman JA: Genomic microarrays in clinical diagnosis. *Curr Opin Pediatr* 2006; **18**: 598–603.
- 3 Zahir F, Friedman JM: The Impact of array genomic hybridization on mental retardation research: a review of current technologies and their clinical utility. *Clin Genet* 2007; **72**: 271–287.
- 4 Friedman JM, Baross A, Delaney AD *et al*: Oligonucleotide microarray analysis of genomic imbalance in children with mental retardation. *Am J Hum Genet* 2006; **79**: 500–513.
- 5 Koolen DA, Sharp AJ, Hurst JA *et al*: Clinical and molecular delineation of the 17q21.31 microdeletion syndrome. *J Med Genet* 2008; **45**: 710–720.
- 6 Slavotinek AM: Novel microdeletion syndromes detected by chromosome microarrays. *Hum Genet* 2008; **124**: 1–17.
- 7 De Vries BBA, Pfundt R, Leisink M *et al*: Diagnostic genome profiling in mental retardation. *Am J Hum Genet* 2005; **77**: 606–616.
- 8 Fan Y, Jayakar P, Zhu H *et al*: Detection of pathogenic gene copy number variations in patients with mental retardation by genomewide oligonucleotide array comparative genomic hybridization. *Hum Mut* 2007; **28**: 1124–1132.
- 9 Shaffer LG, Theisen A, Bejjani BA *et al*: The discovery of microdeletion syndromes in the post-genomic era: review of the methodology and characterization of a new 1q41q42 microdeletion syndrome. *Genet Med* 2007; **9**: 607–616.
- 10 Somerville MJ, Mervis CB, Young EJ *et al*: Severe expressive-language delay related to duplication of the Williams-Beuren locus. *N Engl J Med* 2005; **353**: 1694–1701.
- 11 Berg JS, Brunetti-Pierri N, Peters SU *et al*: Speech delay and autism spectrum behaviors are frequently associated with duplication of the 7q11.23 Williams-Beuren syndrome region. *Genet Med* 2007; **9**: 427–441.
- 12 Potocki L, Bi W, Treadwell-Deering D *et al*: Characterization of Potocki-Lupski syndrome (dup(17)(p11.2p11.2)) and delineation of a dosage-sensitive critical interval that can convey an autism phenotype. *Am J Hum Genet* 2007; **80**: 633–649.
- 13 Zweier C, Peippo MM, Hoyer J *et al*: Haploinsufficiency of *TCF4* causes syndromal mental retardation with intermittent hyperventilation (Pitt-Hopkins Syndrome). *A J Hum Genet* 2007; **80**: 994–1001.
- 14 Kleefstra T, Brunner HG, Amiel J *et al*: Loss-of-function mutations in *Euchromatin Histone Methyl Transferase 1 (EHMT1)* cause the 9q34 subtelomeric deletion syndrome. *Am J Hum Genet* 2006; **79**: 370–377.
- 15 Petrij F, Giles RH, Dauwerse HG *et al*: Rubinstein-Taybi syndrome caused by mutations in the transcriptional co-activator CBP. *Nature* 1995; **376**: 348–351.
- 16 Roelfsema JH, White SJ, Ariyurek Y *et al*: Genetic heterogeneity in Rubinstein-Taybi syndrome: mutations in both the CBP and EP300 genes cause disease. *Am J Hum Genet* 2005; **76**: 572–580.
- 17 Vissers LE, van Ravenswaaij CM, Admiraal R *et al*: Mutations in a new member of chromodomain gene family cause CHARGE syndrome. *Nat Genet* 2004; **36**: 955–957.
- 18 Ariani F, Hayek G, Rondinella D *et al*: FOXP1 is responsible for the congenital variant of Rett syndrome. *Am J Hum Genet* 2008; **83**: 89–93.
- 19 Conte RA, Kleyman SM, Kharode C, Verma RS: Delineation of a ring chromosome 16 by the FISH-technique: a case report with review. *Clin Genet* 1997; **51**: 196–199.
- 20 Ionasescu V, Patil S, Hart M, Rhead W, Smith W: Multiple congenital anomalies syndrome with myopathy in chromosome 16 abnormality. *Am J Med Genet* 1987; **26**: 189–194.
- 21 Ohdo S, Madokoro H, Sonoda T, Ohba K: Sibs lacking characteristic features of duplication of distal 17q. *J Med Genet* 1989; **26**: 465–468.
- 22 Rack KA, Harris PC, MacCarthy AB *et al*: Characterization of three *de novo* derivative chromosomes 16 by 'reverse chromosome painting' and molecular analysis. *Am J Hum Genet* 1993; **52**: 987–997.
- 23 Ravnani JB, Tepperberg JH, Papenhausen P *et al*: Subtelomere FISH analysis of 11 688 cases: an evaluation of the frequency and pattern of subtelomere rearrangements in individuals with developmental disabilities. *J Med Genet* 2006; **43**: 478–489.
- 24 Rossi E, Piccini F, Zollino M *et al*: Cryptic telomeric rearrangements in subjects with mental retardation associated with dysmorphism and congenital malformations. *J Med Genet* 2001; **38**: 417–420.
- 25 Schinzel A: Catalogue of unbalanced chromosome aberrations in man; in de Gruyter (eds): Berlin, 2nd edn, 2001, pp 73–677.
- 26 Soltan HC, Jung JH, Pyatt Z, Singh R: Partial trisomy 9q resulting from a familial translocation t(9;16)(q32;q24). *Clin Genet* 1984; **25**: 449–454.
- 27 Werner W, Kraft S, Callen DF, Bartsch O, Hinkel GK: A small deletion of 16q23.1→16q24.2 in a boy with iris coloboma and minor anomalies. *Am J Med Genet* 1997; **70**: 371–376.
- 28 Yamamoto T, Dowa Y, Ueda H *et al*: Tetralogy of Fallot Associated With Pulmonary Atresia and Major Aortopulmonary Collateral Arteries in a Patient With Interstitial Deletion of 16q21–q22.1. *Am J Med Genet Part A* 2008; **146A**: 1575–1580.
- 29 De Vries BBA, Winter R, Schinzel A, van Ravenswaaij-Arts C: Telomeres: a diagnosis at the end of the chromosomes. *J Med Genet* 2003; **40**: 385–398.
- 30 Marshall CR, Noor A, Vincent JB *et al*: Structural Variation of Chromosomes in Autism Spectrum Disorder. *Am J Hum Genet* 2008; **82**: 477–488.
- 31 Pinto D, Marshall C, Feuk L, Scherer SW: Copy-number variation in control population cohorts. *Hum Mol Genet* 2007; **16**: R168–R173.
- 32 Iafrate AJ, Feuk L, Rivera MN *et al*: Detection of large-scale variation in the human genome. *Nat Genet* 2004; **36**: 949–951.
- 33 Redon R, Ishikawa S, Fitch KR *et al*: Global variation in copy number in the human genome. *Nature* 2006; **444**: 444–454.
- 34 Bhalla K, Luo Y, Buchan T *et al*: Alterations in CDH15 and KIRREL3 in patients with mild to severe intellectual disability. *Am J Hum Genet* 2008; **83**: 703–713.
- 35 Wang K, Zhang H, Ma D *et al*: Common genetic variants on 5p14.1 associate with autism spectrum disorders. *Nature* 2009; **459**: 528–533.
- 36 Zhang A, Li C, Don Chen J: Characterization of transcriptional regulatory domains of ankyrin repeat cofactor-1. *Biochem and Biophys Research Comm* 2007; **358**: 1034–1040.
- 37 Grant ME: The epigenetic origins of mental retardation. DNA methylation and mental retardation. Loss of dynamic histone regulation underlies mental retardation. *Clin Genet* 2008; **73**: 528–534.
- 38 Barberic I, Perry MJ, Dear TN *et al*: An ENU-induced mutation in the *Ankrd11* gene results in an osteopenia-like phenotype in the mouse mutant Yoda. *Physiol. Genomics* 2008; **32**: 311–321.
- 39 Shoichet SA, Hoffmann K, Menzel C *et al*: Mutations in the ZNF41 gene are associated with cognitive deficits: identification of a new candidate for X-linked mental retardation. *Am J Hum Genet* 2003; **73**: 1341–1354.
- 40 Kleefstra T, Yntema HG, Oudakker AR *et al*: Zinc finger 81 (ZNF81) mutations associated with X-linked mental retardation. *J Med Genet* 2004; **41**: 394–399.
- 41 Lugtenberg D, Yntema HG, Banning MJG *et al*: ZNF674: a new Kuppel-associated box-containing zinc-finger gene involved in nonsyndromic X-linked mental retardation. *Am J Hum Genet* 2006; **78**: 265–278.
- 42 Monfort S, Rosello M, Orellana C *et al*: Detection of known and novel genomic rearrangements by array based comparative genomic hybridisation: deletion of ZNF533 and duplication of CHARGE syndrome genes. *J Med Genet* 2008; **45**: 432–437.
- 43 Shimoyama Y, Tsujimoto G, Kitajima M, Natori M: Identification of three human type-II classic cadherins and frequent heterophilic interactions between different subclasses of type-II classic cadherins. *Biochem J* 2000; **349**: 159–167.
- 44 Freson K, Thys C, Wittewrongel C, Vermeylen J, Hoylaerts MF, Van Geet C: Molecular cloning and characterization of the GATA1 cofactor human FOG1 and assessment of its binding to GATA1 proteins carrying D218 substitutions. *Hum Genet* 2003; **112**: 42–49.
- 45 Rialland M, Sola F, Santocanale C: Essential role of human CDT1 in DNA replication and chromatin licensing. *J Cell Sci* 2002; **115**: 1435–1440.