

Published in final edited form as:

J Am Chem Soc. 2010 November 24; 132(46): 16324–16326. doi:10.1021/ja106571g.

NosA Catalyzing Carboxyl-Terminal Amide Formation in Nosiheptide Maturation via an Enamine Dealkylation on the Serine-Extended Precursor Peptide

Yi Yu[‡], Heng Guo[‡], Qi Zhang[‡], Lian Duan[‡], Ying Ding[‡], Rijng Liao[‡], Chun Lei[‡], Ben Shen[§], and Wen Liu[‡]

State Key Laboratory of Bioorganic and Natural Products Chemistry, Shanghai Institute of Organic Chemistry, Chinese Academy of Sciences, 345 Lingling Rd., Shanghai 200032, China, and Division of Pharmaceutical Sciences, School of Pharmacy, University of Wisconsin-Madison, Madison, Wisconsin 53705, USA

Wen Liu: wliu@mail.sioc.ac.cn

Abstract

The carboxyl-terminal amide group has been often found in many bioactive peptide natural products, including nosiheptide belonging to the over 80 entity-containing thiopeptide family. Upon functional characterization of a novel protein NosA in nosiheptide biosynthesis, herein we report an unusual C-terminal amide forming strategy in general for maturing certain amide-terminated thiopeptides by processing their precursor peptides featuring a serine extension. NosA acts on an intermediate bearing a bis-dehydroalanine tail, and catalyzes an enamide dealkylation to remove the acrylate unit originating from the extended serine residue.

The amidation catalyzed by asparagine synthetase-like proteins usually utilizes the γ -carboxamide of Gln to *in situ* generate a nascent ammonia, attack of which to the activated carboxylate results in amide formation with Glu, AMP and PPI as the coproducts (Scheme 1A).¹ For many peptide natural products, carboxyl-terminal amides can be generated endogenously (Scheme 1B), as shown in the biosynthesis of ribosomally synthesized animal hormones and non-ribosomally microbial metabolites such as melithiazols.² Their amide moieties arise from the oxidative cleavage of C-terminal Gly-extended peptides accompanying with glyoxylate production,³ during which the α -hydroxylation of Gly followed by dealkylation is divalent ion or flavin-dependent. Distinct from these well established strategies, we here report a new way to furnish the C-terminal amide in general for maturing certain thiopeptides, based on *in vivo* and *in vitro* characterization of a novel enzyme NosA that acts on the Ser-extended precursor peptide in nosiheptide (**1**) biosynthesis.

Nosiheptide is one of the parent compounds in the thiopeptide family that contains over 80 clinically interesting entities.⁴ The characteristic macrocyclic core of **1** consists of a hydroxypyridine central to multiple thiazoles, dehydroamino acids, and an amide-terminated peptide chain (Figure 1). We have recently unveiled that **1** has a ribosomal origin as other

Correspondence to: Wen Liu, wliu@mail.sioc.ac.cn.

[‡]Shanghai Institute of Organic Chemistry

[§]University of Wisconsin-Madison

Supporting Information **Available:** Experimental procedures for gene inactivation and complementation, characterization of reactions and compounds, protein expression and analysis, sequence analysis, and bioassay; supplementary tables; and supplementary figures. These materials are available free of charge via the internet at <http://pubs.acs.org>.

members newly to be characterized biosynthetically,^{5,6} showing complex posttranslational modifications on the precursor peptide NosM. The structural peptide of NosM, consistent with the amino acids constituting the peptide backbone, possesses an extended Ser residue proven to be the nitrogen source of the terminal amide.⁷ Functional assignment of biosynthetic genes in the *nos* cluster allows for identifying a hypothetic gene, *nosA*, which encodes a 151-aa protein likely involved in the late-stage tailoring to give **1**. Whether or not NosA catalyzes C-terminal amide formation remains elusive, given that all NosA homologues in the database are annotated as unknown proteins.

We first validated the role of NosA in **1** maturation by gene inactivation in the producing strain *Streptomyces actuosus*. To exclude the potential effects on downstream gene expression, *nosA* mutant strain SL4008 was generated by in-frame deletion. SL4008 completely lost the ability to produce **1**; however, it produced a distinct compound (**2**) that exhibits the UV absorption pattern quite similar to **1** (Figure 2, II). *In trans* expression of *nosA* in SL4008 led to the complete conversion of **2** to **1** in the resulting recombinant strain SL4010 (Figure 2, III), confirming the indispensability of *nosA* in **1** biosynthesis. For structural elucidation, **2** was purified, and then subjected to spectroscopic analysis (Figures S1-S3). HR-ESI-MS analysis established the molecular formula as C₅₄H₄₅N₁₃O₁₄S₆Na (*m/z* [M + Na]⁺ 1314.1467, 1314.1431 calculated), and extensive MS/MS analyses suggested the only difference of **2** from **1** in the C-terminal functionality of the peptidyl side chain. The ¹H, ¹³C, and 2D NMR spectra (including HSQC and HMBC correlations) finally established that **2** is a new biosynthetic intermediate of **1** that bears a bis-dehydroalanine tail (Figure 1). These findings strongly supported the involvement of *nosA* in amide formation by removing the terminal acrylate unit of **2** to give **1**.

The counterparts of *nosA* were also found for biosynthesizing some amidated thiopeptides (Figures S4 and S5), as *nocA* (64% identity) for nocathiacin and *tpdK* (34% identity) for GE2270A.6d,f. Similar to NosM in **1** biosynthesis, their precursor peptides contain one (Ser for NocM) or two (Ser-Ala for TpdA) extended amino acid(s) at the C-terminus in addition to the sequence for forming the thiopeptide backbone. We therefore carried out heterologous complementation to examine if the C α -N cleavage on the dehydroalanine unit (derived from Ser) is common for maturing these thiopeptides. *nocA* was introduced into SL4008, yielding the recombinant strain SL4009 for product detection. Remarkably, about 50% of **2** was converted to **1** (Figure 2, IV), ascertaining the functional identity of *nocA* to *nosA*. This interchangeability indicates a strategy in general to afford a C-terminal amide by processing precursor peptides that feature a Ser residue extension.

To characterize NosA as a novel protein for C-terminal amide formation, we next explored its cofactor dependence and subsequent chemical conversion. NosA was overproduced in *Escherichia coli* BL21(DE3) in a 6 \times His tagged form, and then purified to homogeneity for *in vitro* studies (Figure S6A). The resulting protein was subjected to MALDI-TOF MS analysis, exhibiting an M⁺ ion at *m/z* = 18,302.405 (Figure S6B). Taking the calculated molecular weight 18,318.46 and UV-absorptions into account, it is unlikely that any cofactor binds to NosA. In the presence of NosA, the conversion of **2** to **1** efficiently took place (Figure 2, VII), requiring no any cofactor. To probe the catalytic chemistry, 2,4-dinitrophenylhydrazine was added in NosA-catalyzed reaction for derivatization, leading to identification of the co-product pyruvate (**3**, corresponding to the derivative 2-(2,4-dinitrophenylhydrazono)-propanoic acid) (Figure S7). These results validated that NosA acts on **2**, catalyzing the amide moiety formation via a C α -N bond cleavage to afford **1** by eliminating the terminal acrylate moiety. The characteristic thiopeptide framework might be necessary for substrate recognition, since 2-acetamidoacrylic acid, a mimic to the peptidyl tail of **2**, can not be turned over by NosA to generate **3**.

For analyzing NosA-catalyzed reaction *in vitro*, we evaluated the stability of the substrate **2** before the condition optimization. **2** was stable at pH from 3.0 to 8.0 (Figure S8). By contrast, the incubation at pH 9.0 resulted in a time-dependent transformation of **2** to a major distinct compound (**X**, HR-ESI-MS m/z $[M + H]^+$ calcd. 1310.1717 for $C_{54}H_{48}N_{13}O_{15}S_6$, found 1310.1765), which, with the 18-Da increase in molecular weight, could be a hydrated or side ring-hydrolyzed product of **2** (Figure 2, IX, and Figure S9). **X** can be efficiently converted by NosA to **Y** (HR-ESI-MS m/z $[M + H]^+$ calcd. 1240.1662 for $C_{51}H_{46}N_{13}O_{13}S_6$, found 1240.1725), a putative acrylate-removed analogue (Figure 2, X), excluding the possibility of H_2O addition onto the terminal dehydroalanine unit. The mixtures containing **X** and **Y**, respectively, were therefore subjected to sulfhydryl derivatization by adding of 5,5'-dithio-bis(2-nitrobenzoic acid) (DTNB) (Figure S10), showing the generation of their corresponding derivatives **X'** (ESI-MS m/z $[M + H]^+$ calcd. 1507.1, found 1506.6) and **Y'** (ESI-MS m/z $[M + H]^+$ calcd. 1437.1, found 1436.8). This indicated the cleavage of the thioester bond by hydrolysis at high pH to give a free sulfhydryl group. Further, MS/MS analysis supported the deduced structures of **X** and **Y** (Figures S11 and S12), both of which are indole side ring-opened. The finding that NosA tolerates **X** as the substrate is consistent with the hypothesis that the main thiopeptide core plays a key structural role in the enzyme action, as a similar modification for maturation could be involved in the biosynthesis of the monocyclic thiopeptide GE2270A.6d

Thus, the pH dependence of NosA-catalyzed conversion was investigated by using **2** as the substrate in the reaction buffers varying from 4.0 to 8.5, displaying an optimal activity at pH 8.0 (Figure S13). The addition of EDTA into the reaction mixture did not afford the significant change in enzymatic activity, suggesting the metal ion-independence of NosA. Under the optimized condition, the steady-state kinetic parameters were measured, showing a conversion with a k_m value of $93.7 \pm 0.7 \mu M$ for **2**, and a k_{cat} value of $1200 \pm 30 \text{ min}^{-1}$ (Figure S14).

We thereby proposed that NosA-catalyzed reaction may proceed via an enamide dealkylation: 1) specific tautomerization of the terminal dehydroalanine unit of **2** to the corresponding methyl imine is followed by nucleophilic attack of H_2O to generate the terminally hydrated intermediate, and 2) subsequent cleavage of the $C\alpha$ -N bond yields **1** with pyruvate production (Scheme 1C). Consistent with previous studies in chemical synthesis of thiostrepton,⁸ the reaction can be base-initiated, whereas acidic condition makes against tautomerization of enamide to the labile imine and led to the dramatic decrease in NosA activity (Figure S13). The NosA chemistry is non-oxidative, and takes advantage of the terminal enamide moiety generated from the extended Ser by the dehydratase pair NosE and NosD that act at the early-stage posttranslational modifications on the precursor peptide NosM.6 This strategy is clearly distinct from that for maturation of the thiopeptides originating from the non-Ser-extended precursor peptides, such as thiostrepton and siomycin (Figure S5),^{7b-d} which requires an amidotransferase (TsrT/TsrC or SioC belonging to the asparagine synthetase family) to amidate the free carboxylic acid (Scheme 1A). Though terminal amide formation is often of importance to bioactivity,⁹ **2** showed the activity comparable to **1** against the test strain *Bacillus subtilis* (Table S2).

In conclusion, we have characterized that NosA in a cofactor-independent manner catalyzes an enamide dealkylation that may be common for maturing the thiopeptides nosiheptide, nocaithiacin and GE2270A, by processing the Ser-extended precursor peptide to furnish the terminal amide moiety with pyruvate production. The NosA chemistry is mechanistically distinct from known C-terminal amide forming proteins. Taking advantage of the substrate tolerance, it can be exploited by combinatorial biosynthesis for structural diversity in thiopeptide natural products.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

We thank Prof. Heinz G. Floss, University of Washington, for providing of *S. actuosus* ATCC 25421 and his pioneer works on nosiheptide biosynthesis. This work was supported in part by grants from NIH of USA (CA094426 to B.S.); and NNSF (20832009, 30525001 and 20921091), MST (2009ZX09501-008), “973 program” (2010CB833200), CAS (KJCX2-YW-H08 and KSCX2-YW-G-06), and STCSM (09QH1402700) of China (all to W.L.).

References

1. Walsh, CT. Enzymatic Reaction Mechanism. W H Freeman and Company; San Francisco, US: 1977.
2. (a) Kulathila R, Merkler KA, Merkler DJ. Nat Prod Rep. 1999; 16:145–154. [PubMed: 10331284] (b) Weinig S, Hecht HJ, Mahmud T, Müller R. Chem Biol. 2003; 10:939–952. [PubMed: 14583260]
3. (a) Bradbury AF, Finnie MDA, Smyth DG. Nature. 1982; 298:686–688. [PubMed: 7099265] (b) Eipper BA, Mains RE, Glembotski CC. Proc Natl Acad Sci U S A. 1983; 80:5144–5148. [PubMed: 6576381]
4. Bagley MC, Dale JW, Merritt EA, Xiong X. Chem Rev. 2005; 105:685–714. [PubMed: 15700961]
5. Yu Y, Duan L, Zhang Q, Liao R, Ding Y, Pan H, Wendt-Pienkowski E, Tang G, Shen B, Liu W. ACS Chem Biol. 2009; 4:855–864. [PubMed: 19678698]
6. (a) Brown LCW, Acker MG, Clardy J, Walsh CT, Fischbach MA. Proc Natl Acad Sci U S A. 2009; 106:2549–2553. [PubMed: 19196969] (b) Liao R, Duan L, Lei C, Pan H, Ding Y, Zhang Q, Chen D, Shen B, Yu Y, Liu W. Chem Biol. 2009; 16:141–147. [PubMed: 19246004] (c) Kelly WL, Pan L, Li C. J Am Chem Soc. 2009; 131:4327–4334. [PubMed: 19265401] (d) Morris RP, Leeds JA, Naegeli HU, Oberer L, Memmert K, Weber E, LaMarche MJ, Parker CN, Burrer N, Esterow S, Hein AE, Schmitt EK, Krastel P. J Am Chem Soc. 2009; 131:5946–5955. [PubMed: 19338336] (e) Wang J, Yu Y, Tang K, Liu W, Deng Z, He X, Huang X. Appl Environ Microbiol. 2010; 76:2335–2344. [PubMed: 20154110] (f) Ding Y, Yu Y, Pan H, Guo H, Li Y, Liu W. Mol Biosyst. 2010; 6:1180–1185. [PubMed: 20473441]
7. Mocek U, Knaggs AR, Tsuchiya R, Nguyen T, Beale JM, Floss HG. J Am Chem Soc. 1993; 115:7557–7568.
8. (a) Nicolaou KC, Zak M, Safina BS, Estrada AA, Lee SH, Nevalainen M. J Am Chem Soc. 2005; 127:11176–11183. [PubMed: 16076225] (b) Schoof S, Baumann S, Ellinger B, Arndt HD. ChemBioChem. 2009; 10:242–245. [PubMed: 19072817]
9. Merkler DJ. Enzyme Microb Technol. 1994; 15:450–456. [PubMed: 7764886]

VDA AHLSDLDIDALEISEFLDESRLDESEV VAKVMSA-**SCTTCECCSCSS*** NosM




Figure 1.
Precursor peptide NosM and structures of **1**, and **2**. The structural peptide is shown in bold. Asterisk indicates the extended Ser.





Figure 2.

Validation of NosA as a terminal amide-forming protein. *In vivo* **1** or **2** production in *S. actuosus* strains, including the wild type (I), *nosA* mutant strain SL4008 (II), SL4010, a SL4008 derivative carrying *nosA* (III); and SL4009, a SL4008 derivative carrying *nocA* (IV). Using standard **1** (V) and purified **2** (VI) as the controls, *in vitro* conversion of **2** to **1** catalyzed by active NosA (VII) and NosA inactivated by heating (VIII); transformation of **2** to **X** by incubation at pH 9.0 for 12 hrs in the absence of NosA (IX); and transformation of the mixture of **2** and **X** respectively to **1** and **Y** in the presence of NosA (X).



Scheme 1. Variable mechanisms for carboxyl-terminal amide formation