



Published in final edited form as:

Dev Psychopathol. 2009 ; 21(4): 1263–1291. doi:10.1017/S0954579409990150.

Extending Extant Models of the Pathogenesis of Borderline Personality Disorder to Childhood Borderline Personality Symptoms: The Roles of Affective Dysfunction, Disinhibition, and Self- and Emotion-Regulation Deficits

Kim L. Gratz^{1,2}, Matthew T. Tull², Elizabeth K. Reynolds³, Courtney L. Bagge², Robert D. Latzman², Stacey B. Daughters⁴, and C. W. Lejuez³

²Department of Psychiatry and Human Behavior, University of Mississippi Medical Center, Jackson, MS

³Center for Addictions, Personality, and Emotion Research, and the Department of Psychology, University of Maryland, College Park, MD

⁴Department of Public and Community Health, University of Maryland, College Park, MD

Abstract

Although research has been conducted on the course, consequences, and correlates of borderline personality disorder (BPD), little is known about its emergence in childhood, and no studies have examined the extent to which theoretical models of the pathogenesis of BPD in adults are applicable to the correlates of borderline personality symptoms in children. The goal of this study was to examine the interrelationships between two BPD-relevant personality traits (affective dysfunction and disinhibition), self- and emotion regulation deficits, and childhood borderline personality symptoms among 263 children aged 9 to 13. We predicted that affective dysfunction, disinhibition, and their interaction would be associated with childhood borderline personality symptoms, and that self- and emotion-regulation deficits would mediate these relationships. Results provided support for the roles of both affective dysfunction and disinhibition (in the form of sensation seeking) in childhood borderline personality symptoms, as well as their hypothesized interaction. Further, both self- and emotion-regulation deficits partially mediated the relationship between affective dysfunction and childhood borderline personality symptoms. Finally, results provided evidence of different gender-based pathways to childhood borderline personality symptoms, suggesting that models of BPD among adults are more relevant to understanding the factors associated with borderline personality symptoms among girls than boys.

Keywords

borderline personality disorder; childhood disorders; emotion dysregulation; sensation seeking; emotional vulnerability

Borderline personality disorder (BPD) is a serious mental health problem with great public health significance. Although BPD is found at rates of 1–2% in the general population (Lenzenweger, Lane, Loranger, & Kessler, 2007; Skodol et al., 2002), individuals with BPD represent approximately 15% of clinical populations (Skodol et al., 2002; Widiger &

¹Correspondence concerning this article should be addressed to Kim L. Gratz, Department of Psychiatry and Human Behavior, University of Mississippi Medical Center, Jackson, MS 39216, USA; Phone: (601) 815-6450; Fax: (601) 984-4489; KLGratz@aol.com..

Weissman, 1991) and are major consumers of health care resources (Skodol et al., 2005; Zanarini, Frankenburg, Khera, & Bleichmar, 2001). Further, BPD is associated with severe functional impairment (Skodol et al., 2002, 2005), high rates of co-occurring psychiatric disorders (Skodol et al., 2002; Zanarini et al., 1998a, 1998b), and elevated risk for completed suicide (occurring at a rate of 10%; Work Group on Borderline Personality Disorder, 2001).

Although research has been conducted on the prognosis, course, consequences, and correlates of BPD, little is known about the emergence and early manifestations of this disorder (Crick, Murray-Close, & Woods, 2005; Paris, 2005). However, given evidence that personality develops from an early age (Hartup & van Lieshout, 1995) and that adults with BPD report having experienced many borderline personality symptoms in childhood (Reich & Zanarini, 2001; Zanarini et al., 2006), early manifestations of borderline personality pathology can likely be observed in childhood and warrant empirical attention. Indeed, not only would research on the early manifestations of borderline pathology have important implications for the development of secondary prevention programs, studies examining the risk factors associated with the emergence of borderline pathology in childhood may increase our understanding of the pathogenesis of BPD *per se*.

Despite the obvious clinical relevance of this research, few studies have examined the factors associated with the emergence of borderline pathology in children. One likely reason for the paucity of research in this area is that the identification of personality disorders in youth was discouraged historically (due to concerns about the validity of personality disorder diagnoses prior to adulthood). Further, given that past research on childhood borderline pathology has been fraught with the use of inconsistent terms, differing conceptualizations and operationalizations of childhood borderline pathology, and a lack of empirically-validated measures, the progression of research in this area has been limited.

Conceptualizations of Childhood Borderline Pathology

The earliest descriptions of borderline pathology in children encompassed diverse clinical criteria (Ekstein & Wallerstein, 1954; Freud, 1969; Frijling-Schreuder, 1969). The term “borderline” was first used by Ekstein and Wallerstein (1954) to refer to a group of children characterized primarily by unpredictability and rapid fluctuations in ego functioning and interpersonal relationships. These children were considered to be on the border between neurosis and psychosis. The term borderline has also been used to refer to children on the border of receiving a diagnosis of an organic disorder (Kernberg, 1983). Other early conceptualizations of borderline children emphasized the role of micro-psychotic symptoms, loneliness, and intense anxiety (Frijling-Schreuder, 1969; Rosenfeld & Sprince, 1963). Extending this research, the early 1980's brought an interest in identifying specific criteria for the diagnosis of borderline disorders in childhood (Bemporad, Smith, Hanson, & Cicchetti, 1982; Pine, 1983; Vela, Gottlieb, & Gottlieb, 1983). Some of the first sets of diagnostic criteria were based on reviews of the extant clinical literature and/or clinical observations of “borderline children” and demonstrated considerable overlap with one another, including criteria pertaining to excessive and severe anxiety, impulse control problems, disturbed interpersonal relationships, disturbances in cognitive processes (including quasi-psychotic phenomena), and marked fluctuations in functioning (Bemporad et al., 1982, Vela et al., 1983; see also Pine, 1983).

These early conceptualizations of childhood borderline pathology describe a heterogeneous population (likely encompassing children with both borderline personality and schizotypal personality pathologies; Petti & Vela, 1990) and have been criticized for showing little resemblance to adult BPD (Gualtieri, Koriath, & Van Bourgondien, 1983; Palombo, 1982). Thus, it is unclear to what extent childhood borderline disorders (as defined above) reflect

the emergence of adult BPD pathology. Indeed, more recently, this clinical presentation has been termed “multiple complex developmental disorder” (Towbin, Dykens, Pearson, & Cohen, 1993), “multidimensionally impaired disorder” (Kumra et al., 1998), and “emoto-cognitive dys-social disorder” (Ad-Dab'Bagh & Greenfield, 2001), in light of the apparent lack of continuity between these conceptualizations of childhood borderline disorder and adult BPD.

Bearing a stronger resemblance to adult BPD, more recent conceptualizations of childhood borderline pathology reflect efforts to adapt adult BPD criteria for use with children (e.g., Goldman, D'Angelo, DeMaso, & Mezzacappa, 1992; Goldman, D'Angelo, & DeMaso 1993; Guzder, Paris, Zerkowitz, & Marchessault, 1996; Guzder, Paris, Zerkowitz, & Feldman, 1999; Greenman, Gunderson, Cane, & Saltzman, 1986; Paris, Zerkowitz, Guzder, Joseph, & Feldman, 1999). These approaches use modified criteria corresponding to adult BPD diagnostic criteria to diagnose BPD in children. Nevertheless, despite the clearer overlap between these conceptualizations of childhood borderline pathology and adult BPD (as well as their greater reliability and predictive validity, compared to earlier conceptualizations), the practice of diagnosing BPD in childhood has been criticized for assuming a stability in functioning and developmental trajectory that likely vary (Coolidge, 2005; Crick et al., 2005).

In contrast to these approaches, developmental researchers emphasize the importance of assessing childhood borderline pathology dimensionally (vs. categorically) and within a developmental psychopathology framework. Such an approach allows for an examination of the full range of borderline personality symptom severity and dose-response relationships, and does not assume the presence of pathology (as problem behavior is conceptualized as falling on a continuum ranging from none to severe). Further, a developmental psychopathology perspective suggests the importance of examining the ways in which various biological, psychological, and social-contextual factors interact to predict normal and abnormal development across the life course (Cicchetti & Toth, 2009), emphasizing the likelihood of varying pathways to both adaptive and maladaptive outcomes. As such, the same constellation of symptoms in childhood may lead to a variety of outcomes (i.e., multifinality; Cicchetti & Rogosch, 1996, 2002), and one outcome may emerge as a result of different risk factors or developmental processes (i.e., equifinality; Cicchetti & Rogosch, 1996, 2002).

As applied here, a developmental psychopathology approach suggests that childhood borderline personality symptoms may have many different trajectories, and should not be considered indicative of an enduring disorder (as these symptoms will not necessarily evolve on one course to the development of BPD in adulthood). Rather, consistent with the concept of multifinality, children who display early signs and symptoms of borderline personality pathology are expected to take different pathways depending on the risk and protective factors they encounter (Cicchetti & Lynch, 1993). For example, the presence of borderline personality symptoms in children may lead to a variety of outcomes, ranging from the development of personality disorders or other forms of pathology in adolescence and adulthood to more adaptive outcomes (reflecting the absence of any enduring pathology). Furthermore, the emergence of borderline pathology in childhood may, for some children, be indicative of risk for the later development of BPD (as it is likely that this disorder has its origins earlier in development). Of course, it is also important to note that the presence of borderline personality symptoms in childhood likely represent only one pathway through which BPD may develop, with other difficulties in childhood also potentially leading to BPD in adulthood. Nonetheless, given evidence of a connection between childhood and adulthood forms of other disorders (see Rutter, 1996), examining the presence of childhood

borderline personality symptoms is a useful first step in examining the emergence of borderline personality pathology earlier in the lifespan.

Consistent with this approach, Crick et al. (2005) modified a dimensional measure of BPD features in adults for use with children aged 9 and older. This measure provides an assessment of childhood borderline personality features that represent developmentally-appropriate manifestations of the corresponding adult BPD features. The researchers argue that “although BPD is not clearly defined in childhood, some children do exhibit features characteristic of adult borderline pathology, reflecting the emergence of borderline pathology across development” (Crick et al., 2005, pg. 1052). Likewise, Coolidge (2005) argues for the importance of assessing age-appropriate manifestations of borderline personality symptoms dimensionally among children, suggesting that although it may be neither desirable nor possible to diagnose children with BPD, research on the emergence and early manifestations of borderline personality symptoms among children has important clinical and research implications.

It is this approach to the conceptualization and assessment of childhood borderline pathology that is used here. Specifically, this study examines the emergence of borderline personality pathology among children, referred to here as childhood borderline personality symptoms. According to the conceptualization used here, although borderline personality symptoms may be expressed differently in childhood than adulthood (e.g., with the BPD relationship instability criterion pertaining to friends in childhood and intimate relationships in adulthood, or the anger criterion corresponding to temper tantrums in childhood and physical violence in adulthood), the presence of these symptoms in childhood may reflect the emergence of borderline personality pathology. Thus, although childhood borderline personality symptoms are expected to take variable pathways into adulthood (ranging from resilience to pathology), these symptoms may be indicative of risk for later BPD, and, as such, warrant empirical attention. In particular, research on the factors associated with childhood borderline personality symptoms may provide important information regarding the development and pathogenesis of borderline pathology in general, as well as facilitate the identification of children at risk for the development of borderline personality symptoms (potentially paving the way for prevention efforts).

Theories of the Pathogenesis of Borderline Personality Pathology

Extant theoretical models of the pathogenesis of BPD suggest that this disorder is best accounted for within the context of a diathesis-stress model, resulting from the interaction of environmental stressors and trait vulnerabilities (Linehan, 1993; Paris, 1997; Zanarini & Frankenburg, 1997). With regard to the former, researchers have highlighted the role of adverse childhood experiences in the development of BPD, in particular childhood maltreatment (including sexual, physical, and emotional abuse [Bornovalova, Gratz, Delany-Brumsey, Paulson, & Lejuez, 2006; Gibb, Wheeler, Alloy, & Abramson, 2001; Trull, 2001; Zanarini et al., 1997; Zweig-Frank & Paris, 1991], and emotional and physical neglect [Gunderson & Englund, 1981; Johnson, Smailes, Cohen, Brown, & Bernstein, 2000; Zanarini et al., 1997]). These experiences alone are not sufficient to account for the development of BPD, however, and are thought to lead to BPD only in the context of an underlying vulnerability. Specifically, consistent with its categorization as a personality disorder, theories of BPD implicate certain personality traits in the vulnerability for this disorder, with affective dysfunction and disinhibition in particular identified as the “core” traits underlying BPD (Linehan, 1993; Livesley, Jang, & Vernon, 1998; Nigg, Silk, Stavro, & Miller, 2005; Siever & Davis, 1991; Skodol et al., 2002). Moreover, although these traits are not considered to be specific to BPD in and of themselves, researchers have suggested that it is the interaction of affective dysfunction and disinhibition that likely distinguishes

BPD from other disorders (Depue & Lenzenweger, 2001; Nigg et al., 2005; Paris, 2005; Siever & Davis, 1991; Silverman et al., 1991; Trull, 2001).

Consistent with the emphasis on the roles of affective dysfunction (including emotional intensity, reactivity, and lability) and disinhibition (including impulsivity, sensation-seeking, and risk-taking) in BPD, these higher-order traits have received the most attention from researchers, and their relationship to BPD among adult patients is well-documented. For example, research provides evidence for a strong association between affective dysfunction and BPD. This affective dysfunction is thought to encompass several lower-order emotion-related traits, including anxiousness and affective lability (Livesley et al., 1998), emotional intensity, reactivity, and sensitivity (Linehan, 1993), and affective instability (i.e., marked, reactive shifts in mood; Siever & Davis, 1991; Skodol et al., 2002), all of which have been found to be heightened in adult patients with BPD. Specifically, research indicates that adult patients with BPD: (a) report heightened affective instability (Bornovalova et al., 2006; Henry et al., 2001; Koenigsberg et al., 2002) and affect intensity/reactivity (Henry et al., 2001; Koenigsberg et al., 2002); (b) evidence heightened sensitivity to emotional stimuli (Donegan et al., 2003; Herpertz et al., 2001); and (c) exhibit elevated symptoms of anxiety (Snyder & Pitts, 1988).

Research likewise provides evidence for a relationship between disinhibition and BPD. Specifically, behavioral dyscontrol or impulsiveness (in the form of deliberate self-harm, suicidal behaviors, substance use, and risky sexual behavior, among others) is one of the central defining features of BPD (see Paris, 2005; Skodol et al., 2002), and is thought to stem from one of several lower-order disinhibition-related traits, including sensation seeking, risk-taking, novelty seeking, and impulsivity (see Verheul, 2006). Indeed, research provides evidence for an association between each of these lower-order traits and BPD, with studies finding that adults with BPD report heightened levels of: (a) sensation seeking (Reist, Haier, DeMet, & Chicz-DeMet, 1990); (b) risk-taking (Dowson et al., 2004; Kilpatrick et al., 2007); (c) novelty seeking (Pukrop, 2002); and (d) impulsivity (Bornovalova et al., 2006; Henry et al., 2001; Hochhausen, Lorenz, & Newman, 2002). Further, Aluja, Cuevas, Garcia, and Garcia (2007) found that sensation seeking was one of the traits most strongly predictive of BPD pathology among adults.

Of course, the relationship between these personality traits (i.e., affective dysfunction and disinhibition) and BPD is not considered to be direct. Instead, researchers have suggested that these traits increase the risk for BPD through their relationship with deficits in self- and emotion-regulation, both of which are considered to be core mechanisms underlying the development of BPD (Linehan, 1993; Ryan, 2005). Self-regulation is broadly defined as the capacity to plan and execute control over one's experience (Baumeister, 1998), and refers to the ability to adaptively regulate all aspects of one's experience, particularly impulses and behaviors (represented by the construct of ego-control within the developmental literature; Block & Block, 1980). Although maladaptive self-regulation can take the form of both over-regulation (including inflexible responding and non-awareness of one's internal experience) and under-regulation (reflecting a lack of adequate regulatory capacities and including such difficulties as impulsive behaviors; Ryan, 2005), the deficits in self-regulation considered to be most relevant to BPD are those related to under-regulation. Indeed, studies of young adults have found that self-regulation is negatively associated with BPD-related pathology ($r = -.65$; Tangney, Baumeister, & Boone, 2004).

More specifically, *emotion* regulation is considered to play an integral role in the development of BPD (Putnam & Silk, 2005) and to be one of the central defining features of this disorder (Linehan, 1993). Indeed, Linehan (1993) suggests that deficits in emotion regulation mediate the relationship between affective dysfunction and BPD. Unlike affective

dysfunction, emotion regulation does not refer to the nature or quality of one's emotional responses. Instead, deficits in emotion regulation involve maladaptive ways of responding to one's emotions (regardless of their intensity, reactivity, or frequency), including difficulties modulating emotional arousal, difficulties controlling behaviors in the face of emotional distress, and deficits in the functional use of emotions as information. As such, emotion regulation deficits are distinguished from trait affective dysfunction, as the presence of this trait vulnerability (in and of itself) does not preclude adaptive regulation. Nonetheless, consistent with theories suggesting the mediating role of emotion regulation deficits in the relationship between trait affective dysfunction and BPD (Linehan, 1993), evidence suggests that affective dysfunction increases the risk for emotion regulation difficulties (Flett, Blankstein, & Obertynski, 1996; Gratz, Tull, Baruch, Bornovalova, & Lejuez, 2008), as intense/reactive emotions are more difficult to regulate (Flett et al., 1996). Consistent with the theoretical emphasis on the role of emotion regulation in BPD, research provides evidence for a relationship between emotion regulation deficits and BPD in adulthood (Bornovalova et al., 2008; Gratz, Rosenthal, Tull, Lejuez, & Gunderson, 2006; Levine, Marziali, & Hood, 1997). Further, recent evidence provides support for the mediating role of emotion dysregulation in the relationship between affective dysfunction and BPD among adults, with Gratz et al. (2008) finding that emotion dysregulation fully mediated the relationship between negative affect intensity/reactivity and BPD symptoms.

Unfortunately, despite the complexity of the theorized interrelationships of these traits and developmental processes in the development of BPD, no studies have examined the ways in which affective dysfunction and disinhibition interact to predict BPD, nor have studies examined the mediating role of self- and emotion-regulation deficits in the relationship between these traits and BPD. Furthermore, no research has examined if these personality traits and mediators are associated with the emergence of borderline personality pathology in childhood. However, findings that the environmental and neuropsychological factors associated with a BPD diagnosis in childhood are the same as those associated with BPD in adults (Goldman et al., 1992; Guzder et al., 1996, 1999; Paris et al., 1999; Zelkowitz, Paris, Gudzer & Feldman, 2001) suggest that the traits and mediators theorized to be central to the pathogenesis of BPD in adults may likewise be associated with childhood borderline personality symptoms. At the very least, examining the applicability of extant models of the pathogenesis of BPD among adults to the emergence of borderline pathology in childhood may be a useful starting point.

Factors Associated with Borderline Personality Pathology in Childhood

Although almost no research to date has examined the personality traits associated with childhood borderline personality symptoms, researchers have suggested that the trait vulnerabilities of affective dysfunction and disinhibition likely underlie the development of borderline personality pathology in children (Guzder et al., 1999). Further, studies of the likely behavioral manifestations of these traits provide suggestive evidence for the role of both affective dysfunction and disinhibition in childhood borderline personality pathology.

With regard to the relationship between affective dysfunction and childhood borderline personality pathology, Crick et al. (2005) found that one aspect of affective dysfunction (i.e., emotional sensitivity) uniquely predicted borderline personality features in a community sample of children over time. As for the potential manifestations of affective dysfunction, studies of primarily male psychiatric patients aged 6–12 have found higher levels of anxiety and depression among patients with a diagnosis of BPD (vs. those without a BPD diagnosis; Greenman, Gunderson, Cane, & Saltzman, 1986; Guzder et al., 1999). Although this use of BPD diagnoses differs from the dimensional approach to the conceptualization and operationalization of childhood borderline personality symptoms used here, these findings provide suggestive support for a relationship between affective dysfunction (or, more

specifically, psychopathological manifestations of affective dysfunction) and childhood borderline personality pathology.

As for the relationship between disinhibition and childhood borderline personality pathology, Paris et al. (1999) found that the presence of a BPD diagnosis among a sample of primarily male psychiatric patients aged 6–12 was associated with heightened levels of disinhibition (in the form of impulsivity). However, as with the aforementioned research on affective dysfunction, most research on the relationship between disinhibition and childhood borderline personality pathology has focused on behavioral manifestations of disinhibition, particularly aggressive and delinquent behavior. These studies have found that a diagnosis of BPD among samples of primarily male psychiatric patients aged 6–12 is associated with heightened levels of aggressive and delinquent behavior (Greenman et al., 1986; Guzder et al., 1999). Further, utilizing a dimensional measure of borderline personality features in childhood, Crick et al. (2005) found that both physical and relational aggression predicted borderline personality features in children over time (with changes in relational aggression in particular being *uniquely* associated with changes in borderline personality features).

As was the case with the adult BPD literature, however, researchers have not examined explicitly the interaction of affective dysfunction and disinhibition in the risk for childhood borderline personality symptoms. Further, in regard to the potential mediating role of self- and emotion-regulation deficits in the relationship between these personality traits (and their interaction) and childhood borderline personality symptoms, no studies to date have examined the relationship between self- and emotion-regulation deficits and childhood borderline personality symptoms. However, researchers in the area of developmental psychology have long suggested that these capacities are integral to normative development (Block & Block, 1980; Cole, Michel, & Teti, 1994; Fox, 1994), and research on the correlates of self- and emotion-regulation among children provides suggestive evidence for the importance of these phenomena to borderline personality symptoms in childhood (as deficits in each have been found to be associated with a range of negative borderline personality-relevant outcomes). For example, research indicates that ego-undercontrol is associated with socially-inappropriate behavior during late childhood (Eisenberg et al., 2003) and engagement in high-risk behaviors during adolescence (Block, Block, & Keyes, 1988). Moreover, prospective studies of the predictors of interpersonal difficulties during early adolescence have found that ego-undercontrol predicts aggressive behavior (Asendorpf & van Aken, 1999; Hart, Keller, Edelstein, & Hofmann, 1998). Finally, research indicates that emotion regulation deficits are associated with behavioral problems (Shields & Cicchetti, 1998).

As such, preliminary evidence suggests a potential relationship between deficits in self- and emotion-regulation and borderline personality-relevant pathology among children and adolescents, although the extent to which these deficits are associated with childhood borderline personality symptoms *per se* remains unclear. Further, the extent to which deficits in these capacities mediate the relationship between affective dysfunction and disinhibition and childhood borderline personality symptoms is unknown. Nonetheless, lending support to the conceptualization of self- and emotion-regulation deficits as mediators of the relationship between the traits of interest and childhood borderline personality symptoms, research suggests that both affective dysfunction and disinhibition interfere with the development of self- and emotion-regulation capacities throughout childhood (Calkins & Johnson, 1998; Carlson & Wang, 2007; Eisenberg et al., 1997; Santucci et al., 2008). For example, Eisenberg et al. (1997) found that parent and teacher ratings of children's negative emotionality (i.e., emotional intensity and reactivity) were negatively associated with ratings of children's ego-control, and Calkins and Johnson (1998) found that emotional reactivity to a frustrating task was negatively associated with the use of

adaptive emotion regulation strategies among toddlers. As such, these findings (in combination with the aforementioned findings that both the traits and proposed mediators predict borderline personality pathology and/or borderline personality-relevant outcomes) suggest that self- and emotion regulation deficits may mediate the relationship between these traits and borderline personality symptoms in childhood.

Gender-based Equifinality in the Development of Childhood Borderline Personality Symptoms

The principle of equifinality suggests that children may take different pathways to the development of borderline personality symptoms, with the risk factors for these symptoms differing across children. One factor that may arguably be of particular relevance to investigations of equifinality within developmental psychopathology is gender, with researchers proposing different gender-based pathways to psychopathology (e.g., depression; Gjerde & Block, 1996; see also Duggal, Carlson, Sroufe, & Egeland, 2001). In general, research on gender-based pathways to psychopathology highlights the importance of examining the risk factors associated with various forms of psychopathology among females and males separately.

The concept of gender-based equifinality may be particularly relevant to an outcome such as borderline personality pathology, which has long been studied primarily among women. Yet, despite the long-held assumption that BPD is much more prevalent among women than men, a growing body of research within community samples indicates no significant gender differences in the rates of BPD among adolescents and adults (Bernstein et al., 1993; Lenzenweger et al., 2007; Torgersen, Kringlen, & Cramer, 2001). Further, research suggests that the clinical presentation of BPD may be similar across gender (Johnson et al., 2003). However, despite evidence that BPD may be just as relevant to men as women, the majority of the research on BPD has involved primarily female samples, and most theories of the pathogenesis of BPD are based on empirical and clinical literature on women. Thus, it is unclear to what extent these models are applicable to the emergence of borderline personality pathology among males.

Indeed, preliminary evidence suggests that there may be gender differences in the risk factors for borderline personality pathology, as research indicates that the factors associated with both BPD (Paris, Zweig-Frank, & Guzder, 1994a, 1994b) and specific BPD-related behaviors (e.g., deliberate self-harm; Gratz, 2006; Gratz & Chapman, 2007; Gratz, Conrad, & Roemer, 2002) differ across gender (with extant theoretical models generally having greater relevance for women; Gratz et al., 2002). Thus, findings suggest the importance of exploring the factors associated with borderline personality pathology across gender. Given that there is virtually no research on the role of gender in childhood borderline personality pathology (with no known studies exploring differences in the risk factors associated with childhood borderline personality pathology among girls and boys, and research on gender differences in levels/rates of borderline personality pathology producing mixed results [see Crick et al., 2005; Greenman et al 1986; Guzder et al. 1996; Paris, 2003]), studies exploring the factors associated with childhood borderline personality symptoms among girls and boys separately may be particularly important.

The Current Study

The goal of the current study was to examine the relationship between two core BPD-relevant personality traits, self- and emotion regulation deficits, and childhood borderline personality symptoms among a sample of children aged 9 to 13. In regard to the personality traits, we were interested in examining affective dysfunction and disinhibition (as well as their interaction). Consistent with the literature on affective dysfunction in BPD among

adults, we defined affective dysfunction to include several lower-order BPD-relevant emotion-related traits, including anxiousness, affective lability, emotional intensity, and emotional reactivity. As for the particular dimensions of disinhibition of interest in this study, we focused on sensation-seeking and risk-taking, both of which have been shown to be more strongly related to borderline personality-relevant impulsive behaviors (including substance use, risky sexual behavior, and aggressive delinquent behaviors) among youth than trait impulsivity (see Lejuez, Aklin, Bornoalova, & Moolchan, 2005; Lejuez et al., 2007). Consistent with past findings (Goldman et al., 1992; Guzder et al., 1996, 1999; Paris et al., 1999), we hypothesized that models of the pathogenesis of BPD in adults would be applicable to the correlates of borderline personality symptoms in children, such that affective dysfunction, disinhibition, and their interaction would be associated with childhood borderline personality symptoms, and self- and emotion-regulation deficits would mediate these relationships. Further, to explore whether the factors associated with childhood borderline personality symptoms differ for girls and boys, post-hoc analyses examined the factors associated with borderline personality symptoms among girls and boys separately. Although this approach does not allow for a direct test of gender differences in the factors associated with childhood borderline personality symptoms, it was considered a useful first step in exploring the factors that emerge as most relevant for each gender.

Method

Participants

Participant dyads (children and their parents/legal guardians) were recruited through flyers, mailings, and media outreach in the greater Washington, DC metropolitan area, as well as study advertisements distributed to area schools, libraries, and Boys and Girls Clubs. Child-caregiver dyads were eligible for participation if the child was 9 to 13 years of age and both the primary caregiver and child were fluent in English. Data were collected from 263 children and their primary caregivers (87.3% mothers, 6.3% fathers, and 6.3% legal guardians). The child participants ranged in age from 9 to 13 ($M = 11.29$, $SD = 1.04$), and 45% ($n = 118$) were female. With regard to the children's racial/ethnic background, 49% were White, 35% were Black/African-American, 2% were Latino, 1% were Asian, and 13% were of another or unidentified racial/ethnic background. With regard to the educational background of the parents, 7% of the mothers and 14% of the fathers had completed high school or received a GED, 27% of the mothers and 24% of the fathers had attended at least some college or technical school, 30% of the mothers and 20% of the fathers had graduated college, and 32% of the mothers and 31% of the fathers had received an advanced degree. Median family income was \$85,000 per year.

Measures

Caregiver-report measures—The Coolidge Personality and Neuropsychological Inventory for Children (CPNI; Coolidge, 2005) is a 200-item, caregiver-as-respondent measure of DSM-IV Axis I and II symptoms and related difficulties among children and adolescents. Caregivers are asked to rate the extent to which each item accurately describes their child on a 4-point Likert scale (ranging from 1 “This is strongly false for my child; My child is not at all like this.” to 4 “This is strongly true for my child; I would say this describes my child very well.”). The CPNI was normed on a sample of 780 ethnically diverse children ranging in age from 5 to 17 (Coolidge, 2005), and its psychometric properties are well-established (Coolidge, Segal, Stewart, & Ellett, 2000; Coolidge & Thede, 2000; Coolidge, Thede, & Jang, 2001; Coolidge, Thede, Stewart, & Segal, 2002). Of particular interest to the current study were the subscales assessing borderline personality symptoms and the personality trait of affective dysfunction.

The borderline personality subscale includes 9 items adapted from the DSM-IV criteria for adult BPD to be age-appropriate for children (Coolidge, 2005). To ensure that developmentally-appropriate behaviors are not erroneously considered indicative of the emergence of borderline personality pathology, the CPNI requires that parents compare their child's behavior to the behavior of other children of the same age. Further, parents are instructed to endorse only those behaviors that have been typical of their child over an extended period of time (i.e., for several months). Example items include, "My child tries very hard to avoid being alone or feeling abandoned," "My child has threatened or tried to commit suicide or has hurt himself/herself on purpose," and "My child makes friends quickly but soon after seems to hate them" (for a full copy of this measure, see: <http://web.uccs.edu/fcoolidge/cpni/default.htm>). Consistent with the conceptualization of childhood borderline personality symptoms used in this study, the CPNI takes a dimensional approach to the assessment of borderline personality symptoms among children.

The borderline personality subscale has been found to have good test-retest reliability across a 4–6 week period ($r = .67$; Coolidge et al., 2002), as well as good construct and concurrent validity (see Coolidge et al., 2000). Specifically, in support of the construct validity of the borderline personality subscale, children with elevated levels of borderline personality symptoms have been found to exhibit co-occurring difficulties commonly found among adults with BPD, including executive functioning deficits (Coolidge et al., 2000), impulsiveness (Kristensen & Torgersen, 2007), social anxiety (Kristensen & Torgersen, 2007), and ADHD (Coolidge et al., 2000). Further, in-line with evidence that BPD (as assessed by the SCID) is heritable among adults (with Torgersen et al. [2000] finding a heritability estimate of .69 for subthreshold BPD and .80 for definite BPD), Coolidge et al. (2001) found that elevated borderline personality symptoms as assessed by the CPNI had a heritability of .76 among 112 child twin pairs (70 monozygotic and 42 dizygotic) averaging 9 years of age. For the present study, items were summed to create a total borderline personality symptom score. Internal consistency in this sample was adequate ($\alpha = .78$).

The affective dysfunction subscale of the CPNI was used to assess the trait of affective dysfunction. Although the original subscale consists of 10 items, one of these items overlaps with an item on the borderline personality subscale (i.e., the item used to assess the emotional lability criterion of borderline personality, "My child's moods change quickly."). Therefore, this item was excluded from the affective dysfunction subscale, which was created by summing scores on the remaining 9 items (none of which overlapped with the borderline personality subscale). Sample items of this subscale include "My child's emotions seem to shift rapidly and seem to be shallow" and "My child is too touchy or easily annoyed." The affective dysfunction subscale has been found to have good test-retest reliability over a 4–6 week period ($r = .86$; Coolidge et al., 2002). Scores on the affective dysfunction items were summed to create an overall subscale score, with higher scores indicating greater affective dysfunction. Internal consistency in the current sample was good ($\alpha = .83$).

The *Emotion Regulation Checklist* (ERC; Shields & Cicchetti, 1997) is a 24-item caregiver-report questionnaire that assesses children's abilities to regulate their emotions adaptively. Consistent with the conceptualization of emotion regulation used here, the ERC is based on the conceptual definition of emotion regulation as the ability to modulate emotional arousal so as to engage effectively with the environment (Shields & Cicchetti, 1997). Caregivers use a 4-point Likert scale (1 = *rarely/never*; 4 = *almost always*) to indicate the frequency with which their child exhibits a variety of emotion regulation-related behaviors (e.g., "Can manage excitement" and "Shows the kinds of negative feelings you would expect (anger, fear, frustration, distress) when other kids are mean, aggressive or intrusive towards him/her"). The ERC has been found to be associated with other measures of childhood emotion

regulation and problem behaviors among well-adjusted and maltreated samples of children aged 6 to 12 (Shields & Cicchetti, 1997, 1998).

Previous research has demonstrated that the ERC assesses two separate factors, Lability/Negativity and Emotion Regulation (Shields & Cicchetti, 1997). Unlike the Lability/Negativity subscale (which assesses emotional arousal and reactivity; e.g., “Exhibits wide mood swings”), the Emotion Regulation subscale of the ERC assesses the ability to act effectively in the context of emotional arousal (including modulating emotional arousal and controlling behaviors) and emotional awareness/understanding (e.g., “Can say when s/he is feeling sad, angry or mad, fearful or afraid;” Shields & Cicchetti, 1997, 1998). Given the focus of this study on children's self- and emotion-regulation abilities, only the second factor pertaining to emotion regulation was used in this study. Items composing this factor were summed to create a total score of emotion regulation, with higher scores reflecting more adaptive emotion regulation. Internal consistency within this sample was adequate ($\alpha = .70$).

The *Ego-Control Measure* (ECM; Eisenberg et al., 1997) is an 18-item questionnaire adapted from Block and Block's (1980) well-validated California Child Q-Sort to provide a faster caregiver-as-respondent measure of a child's level of self-regulation or ego-control. As measured here, ego-control is distinct from emotion regulation, being more closely related to the construct of reactive control (i.e., the capacity to inhibit or constrain impulsive behaviors and desires; Block & Block, 1980; Eisenberg et al., 2003). Example items include “My child starts to act immature when he/she faces difficult problems or is under stress” (reverse-coded) and “My child is inhibited and constricted (e.g., holds things in; has a hard time expressing himself/herself; is a little uptight).” Caregivers rate each item using a 9-point Likert scale reflecting the extent to which the item describes their child (1 = *most un-descriptive*, 9 = *most descriptive*). This measure has been found to be associated with resiliency (Eisenberg et al., 1997, 2003) and social functioning (Eisenberg et al., 1997, 2003) among children aged 7 and older.

Consistent with past research, items on this scale were summed to create a total ego-control score, with high scores reflecting ego-overcontrol and low scores reflecting ego-undercontrol. Although this scoring procedure results in a variable for which both high and low scores represent maladjustment, studies using this measure have examined ego-control as a linear variable (Eisenberg et al., 1997, 2003), consistent with findings that the quadratic ego-control term is not significant when controlling for the linear ego-control term (Eisenberg et al., 1997, 2003). Internal consistency within this sample was adequate ($\alpha = .78$).

Finally, caregivers completed a basic demographics questionnaire that asked for information regarding their educational background and annual family income, as well as their child's age, gender, and racial/ethnic background.

Child-report measures—As mentioned previously, we were interested in assessing two dimensions of trait disinhibition found to be associated with BPD-relevant impulsive behaviors among youth: risk-taking propensity and sensation seeking. With regard to the former, we used the *Balloon Analogue Risk Task – Youth Version* (BART-Y; Lejuez et al., 2007) to provide a behavioral assessment of risk-taking propensity. In this behavioral task, participants inflate a computer generated balloon that will explode at some point. Each pump of the balloon accrues one point in a temporary bank. Participants have the opportunity to stop pumping the balloon at any time prior to an explosion and allocate the accrued points to a permanent prize meter; if a balloon is pumped past its explosion point, then all points accrued for that balloon are lost. After a balloon explodes or points are allocated to the permanent prize meter, a new balloon appears. Upon completion of 30

balloon trials, the position of the prize meter determines the final prize, with markings indicating small, medium, large, and bonus prize.

Standardized instructions were given to each participant prior to beginning the task. These instructions included the total number of balloons and the fact that points in the prize meter would be exchangeable for prizes immediately following the task. Moreover, participants were informed that: “It is your choice to determine how much to pump up the balloon, but be aware that at some point the balloon will explode” and that “the explosion point varies across each of the thirty balloons, ranging from the first pump to enough pumps to make the balloon fill the entire computer screen.” Participants were given no other information about the probability underlying the explosion point for each balloon. Consistent with previous studies using the BART-Y (Lejuez et al., 2007), the average number of pumps on balloons that did not explode was used as an index of risk-taking propensity.

Second, we used the *Brief Sensation Seeking Scale* (BSSS; Hoyle, Stephenson, Palmgreen, Lorch, & Donohew, 2002) to assess sensation seeking. The BSSS is an 8-item self-report measure designed for use with child and adolescent samples. Example items include, “I would love to have new and exciting experiences, even if they are illegal” and “I like wild parties.” Participants are asked to rate the extent to which each item accurately describes their experience using a 5-point Likert scale (1 = *strongly disagree*; 5 = *strongly agree*). The BSSS has been found to be associated with other well-established measures of disinhibition (see Stephenson, Hoyle, Palmgreen, & Slater, 2003), and is predictive of impulsive behaviors such as substance use (Hoyle et al., 2002; Stephenson et al., 2003). Items were summed to create a total sensation seeking score. Internal consistency within this sample was adequate ($\alpha = .72$).

The *Revised Child Anxiety and Depression Scale* (RCADS; Chorpita, Yim, Moffitt, Umemoto, & Francis, 2000) was used to assess symptoms of anxiety disorders and major depression. Participants are asked to rate the frequency with which they have experienced each item on a 4-point Likert scale (0 = *never*; 3 = *always*). The RCADS has demonstrated convergent validity with other well-established measures of childhood anxiety and depression within both nonclinical (Chorpita et al., 2000) and clinical (Chorpita, Moffitt, & Gray, 2005) samples. Items were summed to create a total score, with higher scores reflecting greater depression and anxiety symptom severity. The RCADS was included in this study as a potential covariate to examine whether affective dysfunction, disinhibition, and self- and emotion regulation deficits are associated with childhood borderline personality symptoms above and beyond psychopathology in general (thereby enabling a more conservative test of the study's hypotheses). Internal consistency within this sample was good ($\alpha = .94$).

A shortened version of the Centers for Disease Control and Prevention Youth Risk Behavior Surveillance System (CDC, 2001) was used to assess delinquent behaviors (a form of externalizing psychopathology). Children were asked to indicate the past year frequency of seven different delinquent behaviors, including fighting, gambling, stealing, and carrying a weapon. Consistent with past research, items were summed to create a total delinquency score (Lejuez, Aclin, Zvolensky, & Pedulla, 2003; see also Feinberg, Greenberg, Osgood, Sartorius, & Bontempo, 2007). Like the RCADS, this measure was included as a potential covariate to examine whether the traits and proposed mediators of interest are associated with childhood borderline personality symptoms above and beyond psychopathology in general.

Procedure

All procedures were reviewed and approved by the University of Maryland Institutional Review Board (IRB). Study advertisements instructed interested caregivers to call the University of Maryland for further details about the study. Upon calling, caregivers were informed that the purpose of the study was to examine youth risk-taking behaviors. Eligible participants (i.e., those who were fluent in English and whose children were 9–13 years of age) were scheduled for an assessment session at the University of Maryland. Participants' travel costs to and from the University were reimbursed.

Upon arrival at the assessment session, a more detailed description of the study procedures was provided and the caregivers and children signed informed consent/assent forms, respectively. The child and caregiver were then accompanied to separate rooms to complete the assessments. Standardized instructions for completing the self-report questionnaires were read aloud to each member of the dyad separately, and participants were encouraged to ask the researchers questions regarding the content or response format of the questionnaires. After completing the questionnaires, the children were provided with instructions for completing the BART-Y and encouraged to ask the researcher any questions they had about the behavioral task.

Once the participants had completed the assessments, they were debriefed and provided with their reimbursement. In particular, children were able to choose their prize (based upon their performance on the BART-Y), and parents were provided with \$25 for their time.

Results

Identification of Covariates

Preliminary analyses were conducted to explore the impact of demographic factors (including age, ethnic/racial background, gender, and family income) and measures of general psychopathology (including depression and anxiety symptoms and delinquent behaviors) on the dependent variable and proposed mediators, in order to identify potential covariates for later analyses (see Tabachnick & Fidell, 1996). Neither the dependent variable nor the proposed mediators were significantly associated with any of the demographic variables ($r_s \leq .11$; $p_s > .05$). However, the composite measure of depression and anxiety symptom severity was significantly associated with both childhood borderline personality symptoms ($r = .14$; $p < .05$) and ego-control ($r = -.21$; $p < .01$), and the measure of delinquent behaviors was significantly associated with ego-control ($r = -.17$; $p < .01$). Thus, both of these measures of general psychopathology (representing both internalizing and externalizing symptoms) were included as covariates in subsequent analyses.

Preliminary Analyses

Table 1 presents descriptive data for the primary variables of interest, as well as the correlations between these variables. Results of the correlational analyses indicate that childhood borderline personality symptoms were significantly associated with the traits of affective dysfunction and sensation-seeking, and significantly negatively associated with the proposed mediators of emotion regulation and ego-control.¹ Further, both emotion regulation and ego-control were significantly negatively associated with affective

¹Prior to conducting correlational analyses, the presence of a quadratic relationship between ego-control and borderline personality symptoms was explored. Results of a regression analysis predicting childhood borderline personality symptoms indicated that the quadratic ego-control term (i.e., the interaction of ego-control by ego-control) was not significant when controlling for the linear ego-control term (and other variables of interest) among the full sample and subsamples by gender ($\beta_s < .07$, $p_s > .10$), consistent with findings from other studies using this measure (e.g., Eisenberg et al., 1997, 2003). Thus, results support the use of the ego-control variable as a linear variable.

dysfunction, and ego-control was significantly negatively associated with sensation seeking. Given that scores on the BART-Y (i.e., average number of pumps on unexploded balloons) were not significantly associated with any of the other variables, this variable was excluded from subsequent analyses.

Primary Analyses

A series of hierarchical regression analyses was conducted to test the proposed mediational model. According to Baron and Kenny (1986), support for the mediational model will be provided if: (a) the personality traits of interest (and/or their interaction) significantly predict childhood borderline personality symptoms, (b) the personality traits (and/or their interaction) significantly predict emotion regulation and ego-control, (c) emotion regulation and ego-control significantly predict childhood borderline personality symptoms, and (d) the personality traits (and their interaction) do not remain significant predictors of childhood borderline personality symptoms once emotion regulation and ego-control are entered into the equation as independent variables.

First, to examine if the personality traits (and/or their interaction) predict childhood borderline personality symptoms, a hierarchical regression analysis was conducted with childhood borderline personality symptoms as the dependent variable, the covariates of depression and anxiety symptom severity and delinquent behaviors entered in the first step of the equation, the traits of affective dysfunction and sensation seeking entered in the second step of the equation, and the interaction of affective dysfunction and sensation seeking entered in the third step of the equation. In line with Aiken and West (1991), predictor variables embedded within interaction terms were standardized prior to analyses. The overall model was significant, accounting for 54% of the variance in childhood borderline personality symptoms, $F(5, 257) = 62.95, p < .01$ (see Table 2). As predicted, the personality traits significantly predicted childhood borderline personality symptoms (above and beyond the covariates; $F_{\Delta}[2, 258] = 145.11, p < .01$). Further, both affective dysfunction and sensation seeking accounted for a significant amount of unique variance in childhood borderline personality symptoms (see Table 2).

Moreover, the interaction of affective dysfunction and sensation seeking accounted for a significant amount of additional variance in borderline personality symptoms above and beyond the main effects of these traits ($F_{\Delta}[1, 257] = 5.44, p < .05$). This significant interaction was explored following methods outlined by Aiken and West (1991). First, the regression lines were plotted at high (one SD above the mean), medium (the mean) and low (one SD below the mean) levels of sensation seeking and affective dysfunction, and then follow-up tests were conducted to examine whether the slopes of the regression lines differed significantly from zero. These tests revealed that the relationship between sensation seeking and borderline personality symptoms increased in magnitude as affective dysfunction moved from low ($b = .13, ns$), to medium ($b = .51, p < .05$), to high ($b = .90, p < .001$; Figure 1).

Next, two hierarchical regression analyses were conducted to examine if the personality traits (and/or their interaction) predict the proposed mediators (controlling for the same covariates described above). Results indicate that the personality traits significantly predicted both emotion regulation ($R^2_{\Delta} = .12, F_{\Delta}[2, 258] = 17.14, p < .001$) and ego-control ($R^2_{\Delta} = .11, F_{\Delta}[2, 258] = 17.50, p < .001$; see Table 2). However, only affective dysfunction accounted for a significant amount of unique variance in ego-control and emotion regulation (above and beyond the covariates and sensation seeking; see Table 2). Further, the interaction of affective dysfunction and sensation seeking did not account for a significant amount of additional variance in emotion regulation ($F_{\Delta}[1, 257] = .05, p > .10$) or ego-

control ($F_{\Delta}[1, 257] = 2.88, p > .05$), above and beyond the main effects of these traits (see Table 2).

Next, to examine if emotion regulation and ego-control mediate the relationship between affective dysfunction (the only personality trait found to be uniquely associated with either mediator) and childhood borderline personality symptoms, a hierarchical regression analysis was conducted with borderline personality symptoms as the dependent variable, the covariates entered in the first step of the equation, affective dysfunction and sensation seeking entered in the second step of the equation, the interaction of these traits entered in the third step of the equation, and emotion regulation and ego-control entered in the final step of the equation. Although emotion regulation and ego-control emerged as significant unique predictors of childhood borderline personality symptoms (accounting for an additional 8% of the variance in these symptoms; $F_{\Delta}[2, 255] = 28.73, p < .001$), affective dysfunction remained a significant predictor of borderline personality symptoms with the proposed mediators in the equation (see Table 3). Thus, results suggest that emotion regulation and ego-control do not fully mediate the relationship between affective dysfunction and childhood borderline personality symptoms. However, providing evidence for partial mediation, computation of Goodman (I) equations indicated that the indirect effect of affective dysfunction on childhood borderline personality symptoms through its effects on both emotion regulation ($z = 2.52, p < .05$) and ego-control ($z = 4.51, p < .001$) was significant. Thus, results suggest that two ways in which affective dysfunction may contribute to borderline personality symptoms in childhood is through its negative effects on emotion regulation and ego-control.

Finally, in order to examine if the synergistic influence of affective dysfunction and sensation seeking relates to borderline personality symptoms indirectly through deficits in emotion regulation and/or ego-control, we tested for the presence of mediated moderation (as outlined by Muller, Judd, and Yzerbyt, 2005). As described by Muller et al. (2005), the first condition for mediated moderation is the presence of a significant interaction between the independent variable and moderating variable in the prediction of the dependent variable. The second condition may be met in one of two ways (2a or 2b), with each way having two necessary subcomponents (*i* and *ii*). The 2a condition is met by the presence of a significant interaction effect between the independent variable and moderating variable on the mediator (*i*; see Table 4, equation 2 or 3) and a main effect of the mediating variable on the dependent variable (when controlling for the interactions between the moderating variable and independent variable and between the moderating variable and mediating variable; *ii*, see Table 4, equation 4). The 2b condition is met by the presence of an independent variable main effect on the mediating variable (*i*; see Table 4, equation 2 or 3) and a significant interaction between the moderating variable and mediating variable on the dependent variable (when controlling for the interaction between the moderating variable and independent variable on the dependent variable; *ii*, see Table 4, equation 4). Finally, the third condition for mediated moderation is met when the interaction effect of the independent variable and moderating variable on the dependent variable decreases in magnitude after entering the mediating variable into the equation.

As shown in Table 4, findings provided no evidence of mediated moderation, suggesting that the effect of the interaction of affective dysfunction and sensation seeking on childhood borderline personality symptoms is not mediated by ego-control or emotion regulation deficits.

Post-hoc Analyses

To explore whether the factors associated with childhood borderline personality symptoms differ for girls and boys, the analyses described above were conducted again among girls and boys separately.

Girls—Results indicate that the personality traits significantly predicted borderline personality symptoms among girls, $F_{\Delta}(2, 113) = 49.92, p < .001$, with both affective dysfunction and sensation seeking accounting for a significant amount of unique variance in borderline personality symptoms (see Table 5). Moreover, the interaction of affective dysfunction and sensation seeking accounted for a significant amount of additional variance in borderline personality symptoms above and beyond the main effects of these traits ($F_{\Delta}[1, 112] = 7.22, p < .01$; see Table 5). Consistent with findings for the sample as a whole, analyses examining the nature of this interaction effect revealed that the relationship between sensation seeking and borderline personality symptoms increased in magnitude as affective dysfunction moved from low ($b = .09, ns$), to medium ($b = .72, p < .05$), to high ($b = 1.36, p < .001$).

As for the relationships between the personality traits (and their interaction) and proposed mediators among girls, results indicate that the personality traits significantly predicted both emotion regulation ($R^2_{\Delta} = .08, F_{\Delta}[2, 113] = 4.74, p < .05$) and ego-control ($R^2_{\Delta} = .11, F_{\Delta}[2, 113] = 7.32, p < .01$), above and beyond the covariates. However, only affective dysfunction accounted for a significant amount of unique variance in emotion regulation and ego-control ($\beta_s = -.29$ and $-.33$, respectively, $ps < .01$); sensation seeking was not uniquely associated with emotion regulation ($\beta = .06, p > .10$) or ego-control ($\beta = -.04, p > .10$). Moreover, contrary to the findings for the sample as a whole, the interaction of affective dysfunction and sensation seeking accounted for a significant amount of additional variance in ego-control among girls ($R^2_{\Delta} = .03, F_{\Delta}[1, 112] = 4.33, p < .05$). The interaction term was not significantly associated with emotion regulation ($R^2_{\Delta} = .00, F_{\Delta}[1, 112] = .43, p > .10$).

With regard to the mediating roles of emotion regulation and ego-control in borderline personality symptoms among girls, results of a hierarchical regression analysis indicate that both emotion regulation and ego-control emerged as unique predictors of borderline personality symptoms among girls (see Table 5, Step 4), accounting for an additional 10% of the variance in these symptoms ($F_{\Delta}[2, 110] = 15.30, p < .001$). Further, although affective dysfunction remained a significant predictor of borderline personality symptoms with the proposed mediators in the equation, findings that the indirect effects of affective dysfunction on childhood borderline personality symptoms through its effects on both emotion regulation ($z = 2.23, p < .05$) and ego-control ($z = 2.98, p < .01$) were significant provide evidence for partial mediation.

As described above, we also conducted a series of regression analyses to test for the presence of mediated moderation among girls. As shown in Table 4 (lower panel), findings provide support for the presence of mediated moderation among girls, indicating that the sensation seeking X affective dysfunction interaction on borderline personality symptoms is mediated by deficits in ego-control (but not by deficits in emotion regulation). Specifically, the 2_{ai} condition for mediated moderation was met by the findings of a significant interaction between sensation seeking and affective dysfunction in the prediction of ego-control (see Table 4, lower panel, equation 2), with post-hoc analyses revealing that the relationship between sensation seeking and ego-control deficits (in the form of under-control) decreased in magnitude as affective dysfunction moved from high ($b = -.20, p = .09$), to medium ($b = -.03, p > .10$), to low ($b = .13, p > .10$). Further, the other conditions of mediated moderation were met by findings of a significant main effect of ego-control on borderline personality symptoms (when controlling for the interactions between affective

dysfunction and sensation seeking and between affective dysfunction and ego-control; see Table 4, lower panel, equation 4), as well as findings that the sensation seeking X affective dysfunction interaction effect on borderline personality symptoms reduced in magnitude from equation 1 ($b = .64$) to equation 4 ($b = .33$), and was no longer significant in equation 4.

Boys—Results indicate that the personality traits significantly predicted borderline personality symptoms among boys, above and beyond the covariates, $F_{\Delta}(2, 140) = 101.17, p < .001$ (see Table 5). However, only affective dysfunction accounted for a significant amount of unique variance in borderline personality symptoms among boys (see Table 5, Step 2). Moreover, the interaction of affective dysfunction and sensation seeking did not account for a significant amount of additional variance in borderline personality symptoms above and beyond the main effects of these traits ($F_{\Delta}[1, 139] = .16, p > .10$; see Table 5, Step 3).

With regard to the relationships between the personality traits and proposed mediators among boys, results indicate that the personality traits significantly predicted both emotion regulation ($R^2_{\Delta} = .18, F_{\Delta}[2, 140] = 15.17, p < .001$) and ego-control ($R^2_{\Delta} = .12, F_{\Delta}[2, 140] = 9.70, p < .001$), above and beyond the covariates. Further, both affective dysfunction and sensation seeking accounted for a significant amount of unique variance in emotion regulation ($\beta_s = -.33$ and $-.26$ for affective dysfunction and sensation seeking, respectively, $p_s < .01$) and ego-control ($\beta_s = -.29$ and $-.18$ for affective dysfunction and sensation seeking, respectively, $p_s < .05$). However, as with the findings for borderline personality symptoms described above, the interaction of affective dysfunction and sensation seeking failed to account for additional variance in emotion regulation ($R^2_{\Delta} = .01, F_{\Delta}[1, 139] = 1.51, p > .10$) or ego-control ($R^2_{\Delta} = .00, F_{\Delta}[1, 139] = .22, p > .10$) among boys, above and beyond the main effects of these traits.

With regard to the mediating roles of emotion regulation and ego-control in childhood borderline personality symptoms among boys, results of a hierarchical regression analysis indicate that the proposed mediators accounted for an additional 8% of the variance in borderline personality symptoms when controlling for the personality traits (and their interaction) and the covariates ($F_{\Delta}[2, 137] = 16.43, p < .001$). However, only ego-control emerged as a unique predictor of borderline personality symptoms among boys (see Table 5, Step 4). Finally, findings indicate that ego-control partially mediates the relationship between affective dysfunction and borderline personality symptoms among boys. Specifically, although affective dysfunction remained a significant predictor of borderline personality symptoms with ego-control in the equation, computation of the Goodman (I) equation indicated that the indirect effect of affective dysfunction on childhood borderline personality symptoms through its effect on ego-control was significant ($z = 3.25, p < .01$). Given that the interaction of affective dysfunction and sensation seeking was not significantly associated with borderline personality symptoms among boys, further analyses examining mediated moderation were not conducted for boys.

Discussion

Recent findings highlighting the early developmental origins of BPD have underscored the need to examine the emergence of borderline personality pathology in children (Crick et al., 2005; Paris, 2005). Although childhood borderline personality symptoms are expected to take variable pathways into adulthood (ranging from resilience to pathology), the presence of these symptoms may, for some children, be indicative of risk for later BPD. Therefore, studies examining the factors associated with childhood borderline personality symptoms may facilitate the identification of children at risk for the later development of borderline

personality pathology, as well as provide a better understanding of the pathogenesis of BPD *per se*. The goal of the current study was to examine the relationships between two core BPD-relevant personality traits (affective dysfunction and disinhibition), self- and emotion-regulation deficits, and childhood borderline personality symptoms among a sample of children aged 9 to 13.

Consistent with past research (Goldman et al., 1992; Guzder et al., 1996; Zerkowicz et al., 2001), results provided support for the applicability of extant models of the pathogenesis of BPD among adults to the emergence of borderline personality pathology among children. In particular, findings suggest that the same traits and mediators theorized to be central to the pathogenesis of BPD in adulthood are likewise associated with childhood borderline personality symptoms. Specifically, evidence was provided for the unique roles of both sensation seeking (one dimension of the trait of disinhibition) and affective dysfunction in childhood borderline personality symptoms. Moreover, consistent with extant theoretical models of the development of BPD (e.g., Paris, 2005), results provided support for the hypothesized interaction of affective dysfunction and sensation seeking in childhood borderline personality symptoms. Specifically, although sensation seeking was related to borderline personality symptoms among children with at least average levels of trait affective dysfunction, it was not related to borderline personality symptoms among children with lower than normal levels of affective dysfunction.

Support was also provided for the role of both self- and emotion-regulation deficits in childhood borderline personality symptoms. Each of these proposed mediators was uniquely associated with borderline personality symptoms, suggesting that deficits in the ability to modulate emotional arousal and control impulsive behaviors (within and outside of the context of emotional distress) are associated with the emergence of borderline personality pathology in children. Further, results suggest that affective dysfunction may increase the risk for deficits in each of these areas, interfering with the development of self- and emotion-regulation capacities and partially accounting for the relationship between affective dysfunction and childhood borderline personality symptoms. Contrary to expectations, neither of these putative mediators explained the main effect of sensation seeking on borderline personality symptoms. As such, results suggest that the relationship between sensation seeking and childhood borderline personality symptoms may be less about the inability to control behaviors, inhibit impulses, and/or modulate emotional arousal, and more about a proclivity for or interest in novel or stimulating stimuli. For example, sensation seeking may increase the risk for borderline personality symptoms by creating an interest in novel sensory experiences (and thereby motivating exploratory behaviors; Collins, Litman, & Spielberger, 2004) or resulting in an underestimation of the risk associated with specific behaviors or activities (Rosenbloom, 2003). Future research is needed to continue to identify the potential mechanisms and developmental processes that may underlie the relationship between this aspect of disinhibition and childhood borderline personality symptoms.

Interestingly, results indicated no gender differences in levels of childhood borderline personality symptoms among our community sample. Although findings of comparable levels of borderline personality symptoms among girls and boys are consistent with extant research indicating comparable rates of BPD across gender among community samples of adolescents and adults (Bernstein et al., 1993; Lenzenweger et al., 2007; Torgersen et al., 2001), they differ from the findings of another recent study of children in the community (which reported higher levels of borderline personality features among girls versus boys; Crick et al., 2005). Given the similarity in sample characteristics between the present study and Crick et al.'s (2005) study (i.e., community youth ranging from 9–13 years of age in the present study and 4th to 6th grade in Crick et al.'s study), further research is needed to

examine whether, to what extent, and for whom levels of childhood borderline personality symptoms differ by gender.

Despite findings of comparable levels of borderline personality symptoms among girls and boys, results provided preliminary evidence of different gender-based pathways to borderline personality symptoms in childhood, suggesting that the model examined here may be more applicable to girls. Specifically, results provided support for the roles of both affective dysfunction and sensation seeking (as well as their interaction) in borderline personality symptoms among girls. Further, findings provided support for the mediating roles of emotion regulation and ego-control, indicating that deficits in each of these areas partially mediated the relationship between affective dysfunction and borderline personality symptoms, and that deficits in ego-control in particular (in the form of ego-undercontrol) mediated the relationship between the interactive effect of affective dysfunction and sensation seeking on borderline personality symptoms. Specifically, findings indicate that sensation seeking predicted ego-control deficits only among girls with elevated levels of affective dysfunction, but ego-control deficits were associated with borderline personality symptoms among girls in general (regardless of their levels of the personality traits of interest).

Contrary to the findings for girls, results provided limited support for the relevance of the proposed etiological model for boys, as only affective dysfunction and ego-control deficits emerged as unique predictors of borderline personality symptoms among boys. Further, although findings that the relationship between affective dysfunction and borderline personality symptoms among boys is partially mediated by ego-control deficits is consistent with the aforementioned findings for girls, the absence of relationships between affective dysfunction and emotion regulation deficits and between emotion regulation and borderline personality symptoms is surprising. Moreover, despite the emphasis on the role of disinhibition (both in and of itself and in interaction with affective dysfunction; Depue & Lenzenweger, 2001; Siever & Davis, 1991; Silverman et al., 1991; Trull, 2001) in BPD, findings provided no support for the role of sensation seeking in borderline personality symptoms among boys.

Of course, findings that different factors emerged as relevant within the models conducted among girls and boys separately do not provide direct evidence for the presence of different gender-based pathways to borderline personality symptoms. Indeed, only by testing the extent to which gender interacts with the factors examined here can we examine directly whether the factors associated with childhood borderline symptoms differ significantly for males and females. Nonetheless, results of this study provide preliminary evidence that extant models of the pathogenesis of BPD may be less applicable to the emergence of borderline personality pathology among boys, consistent with past research showing that models of BPD and related pathology among adults (developed primarily on the basis of clinical and empirical literature on women; see, e.g., Linehan, 1993) are generally less applicable to men (see Gratz & Chapman, 2007; Gratz et al., 2002). Given increasing evidence that borderline personality and related pathology (e.g., self-harm) occur at comparable rates among male and female youth and adults (see Bernstein et al., 1993; Lenzenweger et al., 2007; Gratz, 2001; Muehlenkamp & Gutierrez, 2004), findings highlight the need to develop models of the pathogenesis of these difficulties among males specifically.

Altogether, findings from this study provide evidence for the role of particular personality traits (i.e., affective dysfunction and sensation seeking) in borderline personality symptoms among girls and boys aged 9–13, suggesting that these trait vulnerabilities (either alone or in conjunction with one another) may interfere with the development of self- and emotion-

regulation capacities, thereby increasing the risk for borderline personality symptoms. Yet, although the results of this study highlight the importance of trait vulnerabilities to borderline personality symptoms among children, it is important to note that the presence of these traits alone is not sufficient to predict the emergence of borderline personality pathology (Paris, 2003). Instead, risk and protective factors across a number of developmental-ecological domains must be considered. For example, environmental influences likely determine whether and to what extent individuals with these trait vulnerabilities go on to develop borderline personality symptoms. Further, environmental influences during childhood and adolescence are likely to influence whether children who display early manifestations of borderline personality pathology actually go on to develop BPD as young adults. In all likelihood (consistent with the principle of multifinality), many vulnerable or at-risk individuals (including those evidencing childhood borderline personality symptoms) will not develop BPD. However, given that children with heightened levels of affective dysfunction and sensation seeking (as well as emerging borderline pathology *per se*) may be at greater risk for the later development of BPD than individuals without these trait vulnerabilities or early borderline personality symptoms, research on the traits and mediators associated with borderline personality symptoms in children has important implications for prevention and early intervention efforts.

Although interesting, the results of this study are preliminary and must be evaluated in light of the study's limitations. First, this study used correlational data and a cross-sectional design to examine questions pertaining to the development of borderline personality symptoms in childhood. As such, it is possible that the direction or temporal order of these relationships differs from our predictions. Prospective, longitudinal studies are needed to address these limitations and more fully examine these relationships. In addition, although we attempted to utilize both self-report and behavioral measures, our behavioral measure of risk-taking was not found to be associated with childhood borderline personality symptoms or any other variables examined here. Given limitations associated with the exclusive use of self-report measures, future studies would benefit from a more expanded use of behavioral, biological, and/or psychophysiological measures of the key constructs examined here.

An additional limitation is the reliance on caregiver reports of children's affective dysfunction, self- and emotion regulation capacities, and childhood borderline personality symptoms. It is possible that some caregivers may not have had access to the personal experiences of their children, thereby limiting their ability to accurately report on certain aspects of key constructs, such as fears of abandonment, emotional awareness, or the child's self-efficacy in regulating emotions. Further, given that most of the constructs of interest were assessed through caregiver-report only (with children completing assessments of only the covariates and disinhibition-related variables), it is possible that method variance (i.e., caregiver- vs. child-report measures) may explain the differential strength of the relationships between affective dysfunction and borderline personality symptoms on the one hand (both of which were assessed with caregiver-report measures) and disinhibition and borderline personality symptoms on the other (the former which was assessed through two child-report measures). Given that we did not include both caregiver- and child-report measures of the same constructs, there is no way to determine to what extent reporter bias influenced our findings. However, providing support for our use of a caregiver-report measure of childhood borderline personality symptoms specifically, researchers have suggested that caregivers are better informants than youth themselves on issues where normative comparisons are needed (see Cohen & Crawford, 2005), as in the case of the assessment of borderline personality symptoms here (Coolidge, 2005). Nonetheless, it will be important for future studies to combine caregiver-report measures with child self-report measures of key constructs. Such an approach will clarify the strength of the interrelationships between these constructs.

It is also important to note that although we found evidence for the unique and interactive effects of sensation seeking and affective dysfunction on childhood borderline personality symptoms, we did not assess whether the combination of these traits distinguishes borderline personality symptoms from other forms of pathology, as has been theorized (Paris, 2005; Silverman et al., 1991). Although findings of a significant relationship between these traits and borderline personality symptoms when controlling for both anxiety and depression symptom severity and delinquent behaviors provide evidence for a unique relationship between these traits and childhood borderline personality symptoms above and beyond general internalizing and externalizing psychopathology, future studies should examine the specificity of the model examined here to childhood borderline personality symptoms. Moreover, future research should examine the relationships between childhood borderline personality symptoms and other risk factors thought to be relevant to BPD, including impulsivity (the dimension of disinhibition that has received the most attention with regard to BPD). Research examining all aspects of disinhibition will help clarify the nature and extent of the relationship between this trait and childhood borderline personality symptoms.

Further, future studies should examine the extent to which various BPD-relevant personality traits interact with environmental risk and protective factors to predict both childhood borderline personality symptoms and the varying adaptive and maladaptive trajectories of these symptoms. In particular, given findings that the environmental stressors associated with a BPD diagnosis in childhood are the same as those associated with BPD in adults (i.e., abuse and neglect; Goldman et al., 1992; Guzder et al., 1996, 1999; Zelkowitz et al., 2001), future studies should examine the ways in which maltreatment experiences and trait vulnerabilities interact to influence self- and emotion regulation deficits and, consequently, borderline personality symptoms. Further, given evidence of a relationship between insecure attachment and BPD (e.g., Barone, 2003; Sack, Sperling, Fagen, & Foelsch, 1996; Sperling, Sharp, & Fishler, 1991), future studies should examine if insecure parental attachment interacts with or compounds trait vulnerabilities and childhood stressors and, conversely, if secure attachment serves as a protective factor in the face of other vulnerabilities or risk factors. In particular, given evidence that secure attachment protects against the negative consequences of childhood abuse (e.g., Aspelmeier, Elliott, & Smith, 2007), research should examine if secure attachment to a caregiver protects against the development of borderline personality symptoms among at-risk children and/or increases the likelihood of resilience among children displaying borderline personality symptoms. Moreover, it is likely that different domains of risk (e.g., trait vs. environmental) are differentially related to particular borderline personality symptoms, with traits such as affective dysfunction being more strongly associated with some of the symptoms related to emotion dysregulation, and environmental factors playing a stronger role in symptoms related to self- or cognitive-dysregulation (such as identity disturbance and dissociation). As such, not only will research on the interrelations of traits, environmental stressors, and attachment provide a better understanding of the varying adaptive and maladaptive outcomes associated with childhood borderline personality symptoms, it may also increase our understanding of the emergence of all symptoms of borderline personality pathology (as well as the different pathways to different symptoms).

Finally, it will be important for future research to continue to explore the influence of pertinent demographic factors on the nature, extent, and correlates of childhood borderline personality symptoms. In particular, findings that different factors emerged as relevant to borderline personality symptoms among girls and boys highlight the importance of continuing to examine gender differences (and similarities) in the risk factors associated with childhood borderline personality symptoms. Further, future research should examine the extent to which other demographic factors (such as age) influence the presentation and correlates of borderline personality symptoms. Interestingly, despite examining a sample of

children ranging in age from 9 to 13, age was not significantly associated with levels of childhood borderline personality symptoms. However, it is possible that borderline personality symptoms may increase after the age of 13, or that age may interact with gender to influence levels of borderline personality symptoms in childhood (see Crick et al., 2005 for preliminary findings to this effect). Thus, research should continue to explore the ages at which borderline personality symptoms peak, and the extent to which this is influenced by other factors such as gender.

Despite limitations, the present study provides preliminary support for the applicability of extant models of the pathogenesis of BPD in adults to borderline personality symptoms in childhood (particularly among girls). In particular, this study provides evidence for the roles of sensation-seeking, affective dysfunction, and self- and emotion-regulation deficits in childhood borderline personality symptoms, and suggests the possibility of different gender-based pathways to borderline personality symptoms in childhood. In addition to furthering our understanding of the complex interrelationships of personality traits and childhood developmental processes in the development of childhood borderline personality symptoms, the results of this study have important clinical implications. Specifically, findings suggest the utility of teaching emotionally vulnerable girls and boys skills for effectively managing their emotions and behaviors (including strategies for regulating their emotions and controlling impulsive behaviors, as well as more adaptive ways of responding to their emotions). Given that the pervasiveness of personality pathology and related difficulties is negatively associated with treatment outcome, earlier intervention efforts may confer long-term benefits and be more efficacious.

Acknowledgments

This work was supported by National Institute on Drug Abuse Grant R01 DA18647, awarded to C. W. Lejuez.

References

- Ad-Dab'Bagh Y, Greenfield B. Multiple complex developmental disorder: The "multiple and complex" evolution of the "childhood borderline syndrome" construct. *Journal of the American Academy of Child and Adolescent Psychiatry*. 2001; 40:954–964. [PubMed: 11501696]
- Aiken, LS.; West, SG. *Multiple regression: Testing and interpreting interactions*. Sage Publishing; Thousand Oaks, CA: 1991.
- Aluja A, Cuevas L, Garcia LF, Garcia O. Zuckerman's personality model predicts MCMI-III personality disorders. *Personality and Individual Differences*. 2007; 42:1311–1321.
- Asendorpf JB, van Aken MA. Resilient, overcontrolled, and undercontrolled personality prototypes in childhood: Replicability, predictive power, and the trait-type issue. *Journal of Abnormal and Social Psychology*. 1999; 77:815–832.
- Aspelmeier JE, Elliot AN, Smith CH. Childhood sexual abuse, attachment, and trauma symptoms in college females: The moderating role of attachment. *Child Abuse and Neglect*. 2007; 31:549–566. [PubMed: 17391758]
- Baron RM, Kenny DA. The moderator-mediator variable distinction in social psychological research: Conceptual, strategic, and statistical considerations. *Journal of Personality and Social Psychology*. 1986; 51:1173–1182. [PubMed: 3806354]
- Barone L. Developmental protective and risk factors in borderline personality disorder: A study using the Adult Attachment Interview. *Attachment and Human Development*. 2003; 5:64–77. [PubMed: 12745829]
- Baumeister, RF. The self. In: Gilbert, DT.; Fiske, ST.; Gardner, L., editors. *The handbook of social psychology*. Vol. 1. McGraw-Hill; New York, NY: 1998. p. 680-740.
- Bemporad JR, Smith HF, Hanson G, Cicchetti D. Borderline syndromes in childhood: Criteria for diagnosis. *American Journal of Psychiatry*. 1982; 139:596–602. [PubMed: 7072844]

- Bernstein DP, Cohen P, Velez CN, Schwab-Stone M, Siever LJ, Shinsato L. Prevalence and stability of the DSM-III-R personality disorders in a community-based survey of adolescents. *American Journal of Psychiatry*. 1993; 150:1237–1243. [PubMed: 8328570]
- Block, JH.; Block, J. The role of ego-control and ego-resiliency in the organization of behavior. In: Collins, WA., editor. *Development of cognition, affect, and social relations. The Minnesota Symposia on Child Psychology*. Vol. Vol. 13. Erlbaum; Hillsdale, NJ: 1980. p. 39-101.
- Block J, Block JH, Keyes S. Longitudinally foretelling drug usage in adolescence: Early childhood personality and environmental precursors. *Child Development*. 1988; 59:336–355. [PubMed: 3359859]
- Bornoalova MA, Gratz KL, Daughters SB, Bettina N, Delany-Brumsey A, Lynch TR, et al. A multimodal assessment of the relationship between emotion dysregulation and borderline personality disorder among inner-city substance users in residential treatment. *Journal of Psychiatric Research*. 2008; 42:717–726. [PubMed: 17868698]
- Bornoalova MA, Gratz KL, Delany-Brumsey A, Paulson A, Lejuez CW. Temperamental and environmental risk factors for borderline personality disorder among inner-city substance users in residential treatment. *Journal of Personality Disorders*. 2006; 20:218–231. [PubMed: 16776552]
- Calkins SD, Johnson MC. Toddler regulation of distress to frustrating events: Temperamental and maternal correlates. *Infant Behavior and Development*. 1998; 21:379–395.
- Carlson SM, Wang TS. Inhibitory control and emotion regulation in preschool children. *Cognitive Development*. 2007; 22:489–510.
- Centers for Disease Control (CDC). *Youth risk behavior surveillance system*. Centers for Disease Control; Atlanta, GA: 2001.
- Chorpita BF, Moffitt CE, Gray J. Psychometric properties of the Revised Child Anxiety and Depression Scale in a clinical sample. *Behaviour Research and Therapy*. 2005; 43:309–322. [PubMed: 15680928]
- Chorpita BF, Yim L, Moffitt C, Umemoto LA, Francis SE. Assessment of symptoms of DSM-IV anxiety and depression in children: A revised child anxiety and depression scale. *Behaviour Research and Therapy*. 2000; 38:835–855. [PubMed: 10937431]
- Cicchetti D, Lynch M. Toward an ecological/transactional model of community violence and child maltreatment: Consequences for children's development. *Psychiatry*. 1993; 53:96–118. [PubMed: 8488217]
- Cicchetti D, Rogosch FA. Equifinality and multifinality in developmental psychopathology. *Development and Psychopathology*. 1996; 8:597–600.
- Cicchetti D, Rogosch FA. A developmental psychopathology perspective on adolescence. *Journal of Consulting and Clinical Psychology*. 2002; 70:6–20. [PubMed: 11860057]
- Cicchetti D, Toth SL. The past achievements and future promises of developmental psychopathology: The coming of age of a discipline. *The Journal of Child and Adolescent Psychiatry*. 2009; 50:16–25.
- Cohen, P.; Crawford, TN. Developmental issues. In: Oldham, JM.; Skodol, AE.; Bender, DS., editors. *The textbook of personality disorders*. American Psychiatric Publishing; Washington, DC: 2005. p. 171-185.
- Cole PM, Michel MK, Teti LO. The development of emotion regulation and dysregulation: A clinical perspective. *Monographs of the Society for Research in Child Development*. 1994; 59:73–100. [PubMed: 7984169]
- Collins RP, Litman JA, Spielberg CD. The measurement of perceptual curiosity. *Personality and Individual Differences*. 2004; 36:1127–1141.
- Coolidge, FL. *Coolidge Personality and Neuropsychological Inventory for Children Manual: CPNI*. Author; Colorado Springs, CO: 2005.
- Coolidge FL, Segal DL, Stewart SE, Ellett JA. Neuropsychological dysfunction in children with borderline personality disorder features: A preliminary investigation. *Journal of Research in Personality*. 2000; 34:554–561.
- Coolidge FL, Thede LL. Heritability and the comorbidity of ADHD with behavioral disorders and executive function deficits: A preliminary investigation. *Developmental Neuropsychology*. 2000; 17:273–287. [PubMed: 11056845]

- Coolidge FL, Thede LL, Jang KL. Heritability of childhood personality disorders: A preliminary study. *Journal of Personality Disorders*. 2001; 15:33–40. [PubMed: 11236813]
- Coolidge FL, Thede LL, Stewart SE, Segal DL. The Coolidge Personality and Neuropsychological Inventory for Children (CPNI): Preliminary psychometric characteristics. *Behavior Modification*. 2002; 26:550–566. [PubMed: 12205827]
- Crick NR, Murray-Close D, Woods K. Borderline personality features in childhood: A short-term longitudinal study. *Development and Psychopathology*. 2005; 17:1051–1070. [PubMed: 16613430]
- Depue, RA.; Lenzenweger, MF. A neurobehavioral dimensional model. In: Livesley, WJ., editor. *Handbook of personality disorders: Theory, research, and treatment*. Guilford Press; New York, NY: 2001. p. 136-176.
- Donegan NH, Sanislow CA, Blumberg HP, Fulbright RK, Lacadie C, Skudlarski P, et al. Amygdala hyperreactivity in borderline personality disorder: Implications for emotional dysregulation. *Biological Psychiatry*. 2003; 54:1284–1293. [PubMed: 14643096]
- Dowson J, Bazanis E, Rogers R, Prevost A, Taylor P, Meux C, et al. Impulsivity in patients with borderline personality disorder. *Comprehensive Psychiatry*. 2004; 45:29–36. [PubMed: 14671734]
- Duggal S, Carlson EA, Sroufe LA, Egeland B. Depressive symptomology in childhood and adolescence. *Development and Psychopathology*. 2001; 13:143–164. [PubMed: 11346049]
- Eisenberg N, Guthrie IK, Fabes RA, Reiser M, Murphy BC, Holgren R, et al. The relations of regulation and emotionality to resiliency and competent social functioning in elementary school children. *Child Development*. 1997; 68:295–311. [PubMed: 9180003]
- Eisenberg N, Valiente C, Fabes RA, Smith CL, Reiser M, Shepard SA, et al. The relations of effortful control and ego control to children's resiliency and social functioning. *Developmental Psychology*. 2003; 39:761–776. [PubMed: 12859128]
- Ekstein R, Wallerstein J. Observations on the psychology of borderline and psychotic children. *The Psychoanalytic Study of the Child*. 1954; 9:344–372.
- Feinberg ME, Greenberg MT, Osgood DW, Sartorius J, Bontempo D. Effects of the communities that care model in Pennsylvania on youth risk and problem behaviors. *Preventative Science*. 2007; 8:261–270.
- Flett GL, Blankstein KR, Obertynski M. Affect intensity, coping styles, mood regulation expectancies and depressive symptoms. *Personality and Individual Differences*. 1996; 20:221–228.
- Fox, NA. *Emotion regulation: Behavioral and biological considerations*. Monographs of the Society for Research in Child Development. University of Chicago Press; Chicago, IL: 1994.
- Frijling-Schreuder E. Borderline states in children. *The Psychoanalytic Study of the Child*. 1969; 24:307–327. [PubMed: 5353368]
- Freud, A. *The Writings of Anna Freud*. Vol. Vol. 5. International Universities Press; New York: 1969. The assessment of borderline cases; p. 301-314.
- Gibb BE, Wheeler R, Alloy LB, Abramson LY. Emotional, physical, and sexual maltreatment in childhood versus adolescence and personality dysfunction in young adulthood. *Journal of Personality Disorders*. 2001; 15:505–511. [PubMed: 11778392]
- Gjerde, PF.; Block, J. A developmental perspective on depressive symptoms in adolescence: Gender differences in autocentric-allocentric modes of impulse regulation. In: Cicchetti, D.; Toth, SL., editors. *Adolescence: Opportunities and challenges*. Rochester symposium on developmental psychopathology. Vol. Vol. 7. University of Rochester Press; Rochester, NY: 1996. p. 167-196.
- Goldman SJ, D'Angelo EJ, DeMaso DR, Mezzacappa E. Physical and sexual abuse histories among children with borderline personality disorder. *American Journal of Psychiatry*. 1992; 149:1723–1726. [PubMed: 1443251]
- Goldman SJ, D'Angelo EJ, DeMaso DR. Psychopathology in the families of children and adolescents with borderline personality disorder. *American Journal of Psychiatry*. 1993; 150:1832–1835. [PubMed: 8238638]
- Gratz KL. Measurement of deliberate self-harm: Preliminary data on the Deliberate Self-Harm Inventory. *Journal of Psychopathology and Behavioral Assessment*. 2001; 23:253–263.

- Gratz KL. Risk factors for deliberate self-harm among female college students: The role and interaction of childhood maltreatment, emotional inexpressivity, and affect intensity/reactivity. *American Journal of Orthopsychiatry*. 2006; 76:238–250. [PubMed: 16719643]
- Gratz KL, Chapman AC. The role of emotional responding and childhood maltreatment in the development and maintenance of deliberate self-harm among male undergraduates. *Psychology of Men and Masculinity*. 2007; 8:1–14.
- Gratz KL, Conrad SD, Roemer L. Risk factors for deliberate self-harm among college students. *American Journal of Orthopsychiatry*. 2002; 72:128–140. [PubMed: 14964602]
- Gratz KL, Rosenthal MZ, Tull MT, Lejuez CW, Gunderson JG. An experimental investigation of emotion dysregulation in borderline personality disorder. *Journal of Abnormal Psychology*. 2006; 115:850–855. [PubMed: 17100543]
- Gratz KL, Tull MT, Baruch DE, Bornoalova MA, Lejuez CW. Factors associated with co-occurring borderline personality disorder among inner-city substance users: The roles of childhood maltreatment, negative affect intensity/reactivity, and emotion dysregulation. *Comprehensive Psychiatry*. 2008; 49:603–615. [PubMed: 18970909]
- Greenman DA, Gunderson JG, Cane M, Saltzman PR. An examination of the borderline diagnosis in children. *American Journal of Psychiatry*. 1986; 143:998–1003. [PubMed: 3728745]
- Gualtieri CT, Koriath U, Van Bourgondien ME. “Borderline” children. *Journal of Autism and Developmental Disorders*. 1983; 13:67–72. [PubMed: 6853440]
- Gunderson JG, Englund DW. Characterizing the families of borderlines: A review of the literature. *Psychiatric Clinics of North America*. 1981; 4:159–168. [PubMed: 7232234]
- Guzder J, Paris J, Zerkowitz P, Feldman R. Psychological risk factors for borderline pathology in school-age children. *Journal of the American Academy of Child and Adolescent Psychiatry*. 1999; 38:206–212. [PubMed: 9951221]
- Guzder J, Paris J, Zerkowitz P, Marchessault K. Risk factors for borderline psychology in children. *Journal of the American Academy of Child and Adolescent Psychiatry*. 1996; 35:26–33. [PubMed: 8567606]
- Hart D, Keller M, Edelstein W, Hofmann V. Childhood personality influences on social-cognitive development: A longitudinal study. *Journal of Personality and Social Psychology*. 1998; 74:1278–1289. [PubMed: 9599443]
- Hartup WW, van Lieshout CF. Personality development in social context. *Annual Review of Psychology*. 1995; 46:655–687.
- Henry C, Mitropoulou V, New AS, Koenigsberg HW, Silverman J, Siever LJ. Affective instability and impulsivity in borderline personality and bipolar II disorders: Similarities and differences. *Journal of Psychiatric Research*. 2001; 35:307–312. [PubMed: 11684137]
- Herpertz SC, Dietrich TM, Wenning B, Krings T, Erberich SG, Willmes K, et al. Evidence of abnormal amygdala functioning in borderline personality disorder: A functional MRI study. *Biological Psychiatry*. 2001; 50:292–298. [PubMed: 11522264]
- Hochhausen NM, Lorenz AR, Newman JP. Specifying the impulsivity of female inmates with borderline personality disorder. *Journal of Abnormal Psychology*. 2002; 111:495–501. [PubMed: 12150425]
- Hoyle RH, Stephenson MT, Palmgreen P, Lorch EP, Donohew RL. Reliability and validity of a brief measure of sensation seeking. *Personality and Individual Differences*. 2002; 32:401–414.
- Johnson DM, Shea MT, Yen S, Battle CL, Zlotnick C, Sanislow CA, et al. Gender differences in borderline personality disorder: Findings from the Collaborative Longitudinal Personality Disorders Study. *Comprehensive Psychiatry*. 2003; 44:284–292. [PubMed: 12923706]
- Johnson JJ, Smailes EM, Cohen P, Brown J, Bernstein DP. Associations between four types of childhood neglect and personality disorder symptoms during adolescence and early adulthood: Findings of a community-based longitudinal study. *Journal of Personality Disorders*. 2000; 14:171–187. [PubMed: 10897467]
- Kernberg, PF. Borderline conditions: childhood and adolescent aspects. In: Robinson, KS., editor. *The Borderline Child*. McGraw-Hill; New York: 1983. p. 101-119.

- Kilpatrick T, Joyce E, Milton J, Duggan C, Tyrer P, Rogers RD. Altered emotional decision-making in prisoners with borderline personality disorder. *Journal of Personality Disorders*. 2007; 21:243–261. [PubMed: 17536938]
- Koenigsberg HW, Harvey PD, Mitropoulou V, Schmeidler J, New AS, Goodman M, et al. Characterizing affective instability in borderline personality disorder. *American Journal of Psychiatry*. 2002; 159:784–788. [PubMed: 11986132]
- Kristensen H, Torgersen S. The association between avoidant personality traits and motor impairment in a population-based sample of 11–12 year-old children. *Journal of Personality Disorders*. 2007; 21:87–97. [PubMed: 17373892]
- Kumra S, Jacobsen LK, Lenane M, Zahn TP, Wiggs E, Alaghband-Rad J, et al. “Multidimensionally Impaired Disorder”: Is it a variant of very early-onset Schizophrenia? *Journal of the American Academy of Child and Adolescent Psychiatry*. 1998; 37:91–99. [PubMed: 9444905]
- Lejuez CW, Aklin WM, Bornoalova MA, Moolchan ET. Differences in risk-taking propensity across inner-city adolescent ever-and never-smokers. *Nicotine and Tobacco Research*. 2005; 7:71–79. [PubMed: 15804679]
- Lejuez CW, Aklin W, Daughters S, Zvolensky MJ, Kahler CW, Gwadz M. Reliability and validity of the youth version of the Balloon Analogue Risk Task (BART-Y) in the assessment of risk-taking behavior among inner-city adolescents. *Journal of Clinical Child and Adolescent Psychology*. 2007; 36:106–111. [PubMed: 17206886]
- Lejuez CW, Aklin WM, Zvolensky MJ, Pedulla CM. Evaluation of the Balloon Analogue Risk Task (BART) as a predictor of adolescent real-world risk-taking behaviors. *Journal of Adolescence*. 2003; 26:475–479. [PubMed: 12887935]
- Lenzenweger MF, Lane MC, Loranger AW, Kessler RC. DSM-IV personality disorders in the National Comorbidity Survey Replication. *Biological Psychiatry*. 2007; 62:553–564. [PubMed: 17217923]
- Levine D, Marziali E, Hood J. Emotion processing in borderline personality disorders. *Journal of Nervous and Mental Disease*. 1997; 185:240–246. [PubMed: 9114809]
- Linehan, MM. *Cognitive-behavioral treatment of borderline personality disorder*. Guilford Press; New York, NY: 1993.
- Livesley WJ, Jang KL, Vernon PA. Phenotypic and genetic structure of traits delineating personality disorder. *Archives of General Psychiatry*. 1998; 55:941–948. [PubMed: 9783566]
- Muehlenkamp JJ, Gutierrez PM. An investigation of differences between self-injurious behavior and suicide attempts in a sample of adolescents. *Suicide and Life-Threatening Behavior*. 2004; 34:12–23. [PubMed: 15106884]
- Muller D, Judd CM, Yzerbyt VY. When moderation is mediated and mediation is moderated. *Journal of Personality and Social Psychology*. 2005; 89:852–863. [PubMed: 16393020]
- Nigg JT, Silk KR, Stavro G, Miller T. Disinhibition and borderline personality disorder. *Development and Psychopathology*. 2005; 17:1129–1149. [PubMed: 16613434]
- Palombo J. Critical review of the concept of the borderline child. *Clinical Social Work Journal*. 1982; 10:246–264.
- Paris J. Childhood trauma as an etiological factor in the personality disorders. *Journal of Personality Disorders*. 1997; 11:34–49. [PubMed: 9113821]
- Paris, J. *Personality disorders over time: Precursors, course and outcome*. American Psychiatric Publishing; Washington, DC: 2003.
- Paris J. The development of impulsivity and suicidality in borderline personality disorder. *Development and Psychopathology*. 2005; 17:1091–1104. [PubMed: 16613432]
- Paris J, Zekowitz P, Guzder J, Joseph S, Feldman R. Neuropsychological factors associated with borderline pathology in children. *Journal of the American Academy of Child and Adolescent Psychiatry*. 1999; 38:770–774. [PubMed: 10361797]
- Paris J, Zweig-Frank H, Guzder J. Psychological risk factors for borderline personality disorder in female patients. *Comprehensive Psychiatry*. 1994a; 35:301–5. [PubMed: 7956187]
- Paris J, Zweig-Frank H, Guzder J. Risk factors for borderline personality in male outpatients. *Journal of Nervous and Mental Disease*. 1994b; 182:375–80. [PubMed: 8021636]

- Petti TA, Vela RM. Borderline disorders of childhood: An overview. *Journal of the American Academy of Child and Adolescent Psychiatry*. 1990; 29:327–337. [PubMed: 2189868]
- Pine, F. Borderline syndromes in childhood: A working nosology and its therapeutic implications. In: Robson, KS., editor. *The Borderline Child*. McGraw-Hill; New York: 1983. p. 83-100.
- Pukrop R. Dimensional personality profiles of borderline personality disorder in comparison with other personality disorders and healthy controls. *Journal of Personality Disorders*. 2002; 16:135–147. [PubMed: 12004490]
- Putnam KM, Silk KR. Emotion dysregulation and the development of borderline personality disorder. *Development and Psychopathology*. 2005; 17:899–925. [PubMed: 16613424]
- Reich DB, Zananini MC. Developmental aspects of borderline personality disorder. *Harvard Review of Psychiatry*. 2001; 9:294–301. [PubMed: 11600488]
- Reist C, Haier RJ, DeMet E, Chicz-DeMet A. Platelet MAO activity in personality disorders and normal controls. *Psychiatry Research*. 1990; 33:221–227. [PubMed: 2243899]
- Rosenbloom T. Risk evaluation and risky behavior of high and low sensation seekers. *Social Behavior and Personality*. 2003; 31:375–386.
- Rosenfeld SK, Sprince MP. An attempt to formulate the meaning of the concept “borderline.”. *The Psychoanalytic Study of the Child*. 1963; 18:603–635. [PubMed: 14147294]
- Rutter M. Connections between child and adult psychopathology. *European Child and Adolescent Psychiatry*. 1996; 5(Suppl 1):4–7. [PubMed: 9010655]
- Ryan RM. The developmental line of autonomy in the etiology, dynamics, and treatment of borderline personality disorders. *Development and Psychopathology*. 2005; 17:987–1006. [PubMed: 16613427]
- Sack A, Sperling MB, Fagen G, Foelsch P. Attachment style, history, and behavioral contrasts for a borderline and normal sample. *Journal of Personality Disorders*. 1996; 10:88–102.
- Santucci AK, Silk JS, Shaw DS, Gentzler A, Fox NA, Kovacs M. Vagal tone and temperament as predictors of emotion regulation strategies in young children. *Developmental Psychobiology*. 2008; 50:205–216. [PubMed: 18335488]
- Shields AM, Cicchetti D. Emotion regulation in school-age children: The development of a new criterion Q-sort scale. *Developmental Psychology*. 1997; 33:906–916. [PubMed: 9383613]
- Shields AM, Cicchetti D. Reactive aggression among maltreated children: The contributions of attention and emotion dysregulation. *Journal of Clinical Child Psychology*. 1998; 27:381–395. [PubMed: 9866075]
- Siever LJ, Davis KL. A psychobiological perspective on the personality disorders. *American Journal of Psychiatry*. 1991; 148:1647–1658. [PubMed: 1957926]
- Silverman JM, Pinkham L, Horvath TB, Coccaro EF, Klar H, Schear S, et al. Affective and impulsive personality disorder traits in the relatives of patients with borderline personality disorder. *American Journal of Psychiatry*. 1991; 148:1378–1385. [PubMed: 1897620]
- Skodol AE, Gunderson JG, Pfohl B, Widiger TA, Livesley WJ, Siever LJ. The borderline diagnosis I: Psychopathology, comorbidity, and personality structure. *Biological Psychiatry*. 2002; 51:936–950. [PubMed: 12062877]
- Skodol AE, Gunderson JG, Shea MT, McGlashan TH, Morey LC, Sanislow CA, et al. The Collaborative Longitudinal Personality Disorders Study (CLPS): Overview and implications. *Journal of Personality Disorders*. 2005; 19:487–504. [PubMed: 16274278]
- Snyder S, Pitts WM. Characterizing anxiety in the DSM-III borderline personality disorder. *Journal of Personality Disorders*. 1988; 2:93–101.
- Sperling MB, Sharp JL, Fishler PH. On the nature of attachment in a borderline population: A preliminary investigation. *Psychological Reports*. 1991; 68:543–546. [PubMed: 1862187]
- Stephenson MT, Hoyle RH, Palmgreen P, Slater MD. Brief measures of sensation seeking for screening and large-scale surveys. *Drug and Alcohol Dependence*. 2003; 72:279–286. [PubMed: 14643945]
- Tabachnick, BG.; Fidell, LS. *Using multivariate statistics*. Harper Collins College Publishers; New York, NY: 1996.

- Tangney JP, Baumeister RF, Boone AL. High self-control predicts good adjustment, less pathology, better grades, and interpersonal success. *Journal of Personality*. 2004; 72:271–322. [PubMed: 15016066]
- Torgersen S, Kringlen E, Cramer V. The prevalence of personality disorders in a community sample. *Archives of General Psychiatry*. 2001; 58:590–596. [PubMed: 11386989]
- Torgersen S, Lygren S, Oien PA, Skre I, Onstad S, Edvardsen J, et al. A twin study of personality disorders. *Comprehensive Psychiatry*. 2000; 41:416–425. [PubMed: 11086146]
- Towbin KE, Dykens EM, Pearson, Cohen DJ. Conceptualizing “borderline syndrome of childhood” and “childhood schizophrenia” as a developmental disorder. *Journal of the American Academy of Child and Adolescent Psychiatry*. 1993; 32:775–782. [PubMed: 8340298]
- Trull TJ. Structural relations between borderline personality disorder features and putative etiological correlates. *Journal of Abnormal Psychology*. 2001; 110:471–481. [PubMed: 11502090]
- Vela, RM.; Gottlieb, EH.; Gottlieb, HP. Borderline syndromes in childhood: a critical review. In: Robson, KS., editor. *The Borderline Child*. McGraw-Hill; New York: 1983. p. 32-48.
- Verheul, R. Impulsivity in the diagnosis of borderline personality disorder. Phenotype Conference of the Borderline Personality Disorder Research Foundation; New York, NY. 2006.
- Widiger TA, Weissman MM. Epidemiology of borderline personality disorder. *Hospital and Community Psychiatry*. 1991; 42:1015–1021. [PubMed: 1959892]
- Work Group on Borderline Personality Disorder. Practice guideline for the treatment of patients with borderline personality disorder. *American Journal of Psychiatry*. 2001; 158:1–52.
- Zanarini MC, Frankenburg FR. Pathways to the development of borderline personality disorder. *Journal of Personality Disorders*. 1997; 11:93–104. [PubMed: 9113824]
- Zanarini MC, Frankenburg FR, Dubo ED, Sickel AE, Trikha A, Levin A, et al. Axis I comorbidity of borderline personality disorder. *American Journal of Psychiatry*. 1998a; 155:1733–1739. [PubMed: 9842784]
- Zanarini MC, Frankenburg FR, Dubo ED, Sickel AE, Trikha A, Levin A, et al. Axis II comorbidity of borderline personality disorder. *Comprehensive Psychiatry*. 1998b; 39:296–302. [PubMed: 9777282]
- Zanarini MC, Frankenburg FR, Khera GS, Bleichmar J. Treatment histories of borderline inpatients. *Comprehensive Psychiatry*. 2001; 42:144–150. [PubMed: 11244151]
- Zanarini MC, Frankenburg FR, Ridolfi ME, Jager-Hyman S, Hennen J, Gunderson JG. Reported childhood onset of self-mutilation among borderline patients. *Journal of Personality Disorders*. 2006; 20:9–15. [PubMed: 16563075]
- Zanarini MC, Williams AA, Lewis RE, Reich RB, Soledad C, Vera MA, et al. Reported pathological childhood experiences associated with the development of borderline personality disorder. *American Journal of Psychiatry*. 1997; 154:1101–1106. [PubMed: 9247396]
- Zelkowitz P, Paris J, Guzder J, Feldman R. Diathesis and stressors in borderline pathology of childhood: The role of neuropsychological risk and trauma. *Journal of the American Academy of Child and Adolescent Psychiatry*. 2001; 40:100–105. [PubMed: 11195550]
- Zweig-Frank H, Paris J. Parents' emotional neglect and overprotection according to the recollections of patients with borderline personality disorder. *American Journal of Psychiatry*. 1991; 148:648–651. [PubMed: 2018169]




Figure 1. Interactive effect of sensation seeking and affective dysfunction on childhood borderline personality symptoms among overall sample ($N = 263$).

Table 1

Descriptive Data for and Correlations between Primary Variables of Interest (N = 263)

	1	2	3	4	5	6	7	8	Mean	SD
1. BP symptoms	---	.72**	-.02	.22**	-.33**	-.52**	.14*	.11	15.06	4.24
2. Affective dysfunction		---	-.06	.13*	-.32**	-.34**	.09	.08	14.64	4.28
3. Risk-taking			---	-.02	-.00	.04	-.07	-.03	31.49	13.86
4. Sensation seeking				---	-.10	-.21**	.24**	.42**	12.79	5.38
5. Emotion regulation					---	.04	.10	.01	27.79	3.29
6. Ego-control						---	-.21**	-.17**	91.14	19.28
7. Depression/anxiety							---	.29**	29.42	17.43
8. Delinquent behaviors								---	2.76	3.19

Note. BP = borderline personality.

* $p < .05$.

** $p < .01$.

Table 2

Hierarchical Regression Analyses Exploring the Roles of Sensation Seeking, Affective Dysfunction, and their Interaction in Childhood Borderline Personality Symptoms, Emotion Regulation, and Ego-Control, Controlling for Depression and Anxiety Symptom Severity and Delinquent Behaviors (N = 263)

	Borderline Personality Symptoms					Emotion Regulation					Ego-Control				
	β	b (SE)	Adj R ²	R ² _A	F	β	b (SE)	Adj R ²	R ² _A	F	β	b (SE)	Adj R ²	R ² _A	F
<i>Step 1</i>			.02	.03*	3.28*			.00	.01	1.36			.05	.06**	7.81**
Depression/anxiety	.11	.03 (.02)				.11	.02 (.01)				-.17**	-.19 (.07)			
Delinquent behaviors	.08	.11 (.09)				-.02	-.02 (.07)				-.12	-.74 (.38)			
<i>Step 2</i>			.53	.52***	76.02***			.11	.12***	9.33***			.16	.11***	13.15***
Sensation Seeking (SS)	.12**	.52 (.20)				-.12	-.38 (.21)				-.11	-2.17 (1.22)			
Affective dysfunction (AD)	.70***	2.99 (.18)				-.32***	-1.04 (.19)				-.31***	-6.02 (1.11)			
<i>Step 3</i>			.54	.01*	62.95***			.11	.00	7.45***			.16	.01	11.17***
SS × AD	.10*	.38 (.16)				.01	.04 (.18)				-.10	-1.71 (1.01)			

Note. SS × AD = sensation seeking × affective dysfunction interaction.

* $p < .05$.

** $p < .01$.

*** $p < .001$.

Table 3
 Hierarchical Regression Analysis Examining the Mediating Role of Emotion Regulation and Ego-Control in the Relationship between Sensation Seeking and Affective Dysfunction (and their Interaction) and Childhood Borderline Personality Symptoms (N = 263)

	β	b (SE)	R ² (Adj)	R ² _A	F
<i>Step 1</i>			.03 (.02)	.03*	3.28*
Depression/anxiety	.11	.03 (.02)			
Delinquent behaviors	.08	.11 (.09)			
<i>Step 2</i>			.54 (.53)	.52***	76.02***
Depression/anxiety	.05	.01 (.01)			
Delinquent behaviors	-.01	-.01 (.06)			
Sensation seeking	.12**	.52 (.20)			
Affective dysfunction	.70***	2.99 (.18)			
<i>Step 3</i>			.55 (.54)	.01*	62.95***
Depression/anxiety	.06	.01 (.01)			
Delinquent behaviors	-.01	-.01 (.06)			
Sensation seeking	.12*	.51 (.20)			
Affective dysfunction	.70***	2.95 (.18)			
Sensation seeking × Affective dysfunction	.10*	.38 (.16)			
<i>Step 4</i>			.63 (.62)	.08***	62.87***
Depression/anxiety	.04	.01 (.01)			
Delinquent behaviors	-.02	-.03 (.06)			
Sensation seeking	.07	.31 (.18)			
Affective dysfunction	.56***	2.39 (.18)			
Sensation seeking × Affective dysfunction	.07	.28 (.15)			
Emotion regulation	-.13**	-.56 (.17)			
Ego-control	-.30***	-1.25 (.18)			

* $p < .05$.

** $p < .001$.

Table 4
Regression Analyses Examining Mediated Moderation among Overall Sample and Girls

Criterion	Equation 1: BPS		Equation 2: EGO		Equation 3: EMOT		Equation 4: BPS	
	b	t	b	t	b	t	b	t
Overall Sample (N = 263)								
Predictors								
IV: Sensation Seeking	0.51	2.59**	-0.11	-1.75	-0.12	-1.80	0.29	1.58
MO: Affective Dysfunction	2.95	16.38***	-0.30	-5.30***	-0.32	-5.37***	2.24	11.72***
IV × MO: Sensation seeking × Affective dysfunction	0.38	2.33**	-0.09	-1.69	0.01	0.23	0.21	1.34
MEDI: Ego-control							-1.24	-6.92***
MED2: Emotion regulation							-0.56	-3.28***
MED1 × MO: Ego-control × Affective dysfunction							-0.19	-1.15
MED2 × MO: Emotion regulation × Affective dysfunction							-0.28	-1.89
Girls (n = 118)								
Predictors								
IV: Sensation Seeking	0.72	2.66**	-0.03	-0.37	0.06	.61	0.66	2.68**
MO: Affective Dysfunction	2.39	9.30***	-0.32	-3.62***	-0.28	-2.99**	1.67	6.22***
IV × MO: Sensation seeking × Affective dysfunction	0.64	2.69**	-0.17	-2.08*	-0.06	-0.66	0.33	1.43
MEDI: Ego-control							-0.97	-3.87***
MED2: Emotion regulation							-0.89	-3.75***
MED1 × MO: Ego-control × Affective dysfunction							-0.31	-1.33
MED2 × MO: Emotion regulation × Affective dysfunction							-0.19	-.84

Note: IV = independent variable; MO = moderator; MEDI = first mediator; MED2 = second mediator; BPS = borderline personality symptoms; EGO = ego-control; EMOT = emotion regulation; b = unstandardized beta. Each regression analysis controlled for depression and anxiety symptoms and delinquent behaviors (parameters not shown).

* $p < .05$.

** $p < .01$.

*** $p < .001$.

Hierarchical Regression Analyses Examining the Mediating Role of Emotion Regulation and Ego-Control in the Relationship between Sensation Seeking and Affective Dysfunction (and their Interaction) and Childhood Borderline Personality Symptoms for Girls and Boys Separately.

Table 5

	Girls (n = 118)				Boys (n = 145)					
	β	b (SE)	R ² (Adj)	R ² _A	F	β	b (SE)	R ² (Adj)	R ² _A	F
<i>Step 1</i>			.06 (.05)	.06*	3.79*			.01 (.00)	.01	1.03
Depression/anxiety	.13	.03 (.02)				.12	.03 (.02)			
Delinquent behaviors	.17	.23 (.13)				-.01	-.02 (.12)			
<i>Step 2</i>			.50 (.48)	.44***	28.47***			.60 (.59)	.58***	51.83***
Depression/anxiety	.00	.00 (.02)				.10	.03 (.02)			
Delinquent behaviors	.01	.01 (.10)				-.01	-.02 (.09)			
Sensation seeking	.19**	.74 (.28)				.08	.37 (.28)			
Affective dysfunction	.64***	2.45 (.26)				.75***	3.43 (.25)			
<i>Step 3</i>			.53 (.51)	.03**	25.48***			.60 (.58)	.00	41.24***
Depression/anxiety	.04	.01 (.02)				.10	.03 (.02)			
Delinquent behaviors	-.01	-.02 (.09)				-.01	-.02 (.09)			
Sensation seeking	.19**	.72 (.27)				.08	.36 (.28)			
Affective dysfunction	.63***	2.39 (.26)				.75***	3.42 (.25)			
Sensation seeking × Affective dysfunction	.18**	.64 (.24)				.02	.09 (.22)			
<i>Step 4</i>			.63 (.61)	.10***	27.21***			.68 (.66)	.08***	40.69***
Depression/anxiety	.01	.00 (.01)				.06	.02 (.01)			
Delinquent behaviors	-.03	-.04 (.08)				.01	.01 (.08)			
Sensation seeking	.19**	.74 (.24)				.00	.01 (.27)			
Affective dysfunction	.48***	1.84 (.25)				.63***	2.89 (.26)			
Sensation seeking × Affective dysfunction	.12	.42 (.22)				.02	.08 (.20)			
Emotion regulation	-.21**	-.80 (.23)				-.09	-.39 (.25)			
Ego-control	-.27***	-1.03 (.25)				-.31***	-1.41 (.25)			

* $p < .05$.** $p < .01$.*** $p < .001$.

 $p < .001$

NIH-PA Author Manuscript

NIH-PA Author Manuscript

NIH-PA Author Manuscript