



Published in final edited form as:

J Nat Prod. 2010 November 29; 73(11): 1792–1795. doi:10.1021/np100411d.

Four Diphenylpropanes and a Cycloheptadibenzofuran from *Bussea sakalava* from the Madagascar Dry Forest¹

Ende Pan[†], Liva Harinantana[†], Peggy J. Brodie[†], James S. Miller[‡], Martin W. Callmänder[§], Stephan Rakotonandrasana[±], Etienne Rakotobe[±], Vincent E. Rasamison[±], and David G. I. Kingston^{*†}

[†]Department of Chemistry, M/C 0212, Virginia Polytechnic Institute and State University, Blacksburg, Virginia 24061

[‡]Missouri Botanical Garden, P. O. Box 299, St. Louis, Missouri 63166-0299

[§]Missouri Botanical Garden, B. P. 3391, Antananarivo 101, Madagascar

[±]Centre National d'Application et Recherches Pharmaceutiques, B. P. 702, Antananarivo 101, Madagascar

Abstract

Investigation of the endemic Malagasy plant *Bussea sakalava* Du Puy & R. Rabev. (Fabaceae) for antiproliferative activity against the A2780 ovarian cancer cell line led to the isolation of the four new diphenylpropanes **1–4** and the new cycloheptadibenzofuran **5**; compound **5** has a previously unreported natural product skeleton. The structure elucidation of these compounds was based on the analysis of their 1D and 2D NMR and mass spectroscopic data. Compounds **1–5** were tested for antiproliferative activity against the A2780 human ovarian cancer cell line.

In our continuing search for biologically active natural products from tropical rainforests as part of an International Cooperative Biodiversity Group (ICBG) program, we obtained an ethanol extract from the roots of a plant identified as *Bussea sakalava* Du Puy & R. Rabev. (Fabaceae) from Madagascar. This extract showed moderate antiproliferative activity against the A2780 human ovarian cancer cell line with an IC₅₀ value of 10 µg/mL. The extract was selected for examination on the basis of this activity and the absence of previous phytochemical studies of the species.

Previous studies on the genus *Bussea* indicated the presence of azetidine-2-carboxylic acid and 3-hydroxyproline in seeds of different *Bussea* species,^{2,3} and the cytotoxicity and high trypanocidal activity of a methanol extract of stem bark of *Bussea occidentalis* has been reported.⁴


Fractionation of a dichloromethane fraction of an ethanol extract of *B. sakalava* by C-18 open column and high performance liquid chromatography (HPLC) yielded four new diphenylpropanes named bussealins A–D (**1–4**) and a cycloheptadibenzofuran derivative named bussealin E (**5**). Herein we report the structural elucidation of these new compounds and their antiproliferative properties against the A2780 human ovarian cancer cell line.

^{*}To whom correspondence should be addressed. Tel: (540) 231-6570. Fax: (540) 231-3255. dkingston@vt.edu.
current address: The New York Botanical Garden, New York

Supporting Information Available: ¹H, ¹³C, COSY, HMBC, HMQC, and NOESY spectra of bussealins A–E (**1–5**). This information is available free of charge via the Internet at <http://pubs.acs.org>.

Results and Discussion

Bussealin A (**1**) was obtained as an off-white amorphous solid. Its positive ESI-MS revealed a pseudomolecular ion peak at m/z 321.1338 $[M + H]^+$ corresponding to the molecular formula $C_{17}H_{21}O_6$. The IR spectrum showed absorptions of OH (3367 cm^{-1}) and aromatic groups. The ^1H NMR spectrum (Table 1) exhibited a singlet at δ_{H} 6.18 (2H, s) corresponding to a pair of aromatic protons of an A_2 system, two aromatic doublets [δ_{H} 6.50 (d, $J = 8.4$) and 6.38 (d, $J = 8.4$)] of an AB system, two OCH_3 groups [δ_{H} 3.75 (s) and 3.78 (s)], and a multiplet and two triplet methylene groups at δ_{H} 1.79 (2H, m), 2.52 (2H, t, $J = 7.7$) and 2.41 (2H, t, $J = 7.7$) respectively. The ^{13}C NMR spectrum of **1** exhibited signals for 17 carbons, including three methylene carbons (δ_{C} 36.5, 33.0, and 30.6), two OCH_3 groups (δ_{C} 56.5 and 60.8), and twelve aromatic carbons assignable to two isolated aromatic rings. Six of the aromatic carbons were oxygenated, as shown by their deshielded carbon chemical shifts (Table 1) and were consistent with the molecular formula. The above data suggested that **1** had a diphenyl propane skeleton. The complete ^1H and ^{13}C NMR assignments and the connectivities were determined from analysis of a combination of COSY, HMQC, and HMBC data. Three mutually coupled methylene groups were revealed by the cross peaks observed in the COSY spectrum. In the HMBC spectrum, H-1 (δ_{H} 2.41) showed correlations with C-2 (δ_{C} 33.0), C-3 (δ_{C} 30.6), C-1' (δ_{C} 140.2), and with C-2' and C-6', both of which had the same chemical shifts (δ_{C} 108.7). The A_2 substitution pattern of the A ring of **1** was established by HMBC correlations from the signal at δ_{H} 6.18 (H-2' and H-6') to C-1 (δ_{C} 36.5), C-1' (δ_{C} 140.2), C-3' (δ_{C} 151.3), C-4' (δ_{C} 134.7) and C-6' and C-2' (δ_{C} 108.7), as well as the correlation from one OCH_3 group at δ_{H} 3.75 to C-4' (δ_{C} 134.7). The proton substitutions on the B ring were assigned based on the 3J HMBC correlations between H-3 (δ_{H} 2.52) and C-6'' (δ_{C} 120.5), and between H-5'' (δ_{H} 6.38) and C-1'' (δ_{C} 123.4). Moreover, the H-5'' proton showed HMBC correlations to C-6'' (δ_{C} 120.5), C-4'' (δ_{C} 147.8) and C-3'' (δ_{C} 134.9). The location of the remaining OCH_3 group was at C-4'', as deduced from the HMBC correlation between the signal at δ_{H} 3.78 and that of C-4''. On the basis of the molecular formula of **1**, the remaining four OH groups were located at C-2'' (δ_{C} 144.7), C-3'' (δ_{C} 134.9), C-3' (δ_{C} 151.3), and C-5' (δ_{C} 151.3). Bussealin A is thus assigned the structure 3',5',2'',3''-tetrahydroxy-4',4''-dimethoxy-1,3-diphenylpropane (**1**).



- 1** $R^1 = R^2 = \text{OH}$, $R^3 = \text{H}$
2 $R^1 = \text{OH}$, $R^2 = \text{H}$, $R^3 = \text{OMe}$
3 $R^1 = R^3 = \text{H}$, $R^2 = \text{OH}$
4 $R^1 = R^3 = \text{OMe}$, $R^2 = \text{H}$

Bussealin B (**2**) was obtained as an off-white amorphous solid. Its positive ESI-MS revealed a pseudomolecular ion peak at m/z 335.1512 $[M + H]^+$ corresponding to molecular formula $C_{18}H_{23}O_6$. The ^1H NMR spectrum (Table 1) showed two singlets of an AX system at δ_{H} 6.58 (s) and 6.60 (s), two aromatic doublets of an AB system at δ_{H} 6.51 (d, $J = 8.4$) and 6.39 (d, $J = 8.4$), three OCH_3 groups [δ_{H} 3.76 (s), 3.80 (s) and 3.83 (s)], and one multiplet and


two triplet methylene groups at δ_{H} 1.76 and 2.54 (t, $J = 7.8$) and 2.50 (t, $J = 7.8$). Inspection of the ^1H and ^{13}C NMR spectra of **2** revealed close similarities with those of **1**, except for the presence of an additional OCH_3 signal and the chemical shifts of the AX system of ring A. The fact that the chemical shifts of the carbons of ring B of compounds **1** and **2** were superimposable (Table 1) indicated the presence of a 2'',3''-dihydroxy-4''-methoxyphenyl group in **2**. Interpretation of HMBC and NOESY experiments allowed us to determine the location of the OCH_3 groups to be at 2', 4', and 4''. The two singlet aromatic protons on ring A were assigned according to the observation of 3J HMBC correlations from H-6' (δ_{H} 6.60) to C-1 (δ_{C} 30.4) and from H-3' (δ_{H} 6.58) to C-1' (δ_{C} 124.7). Moreover, the proton signal of H-1 (δ_{H} 2.50) showed HMBC correlations with C-1' (δ_{C} 124.7), C-6' (δ_{C} 117.9) and the methoxylated carbon at C-2' (δ_{C} 152.2). This indicated that the third OCH_3 group must be at C-4' or C-5'. NOESY correlations from H-3' (δ_{H} 6.58) to 2'-OMe (δ_{H} 3.76) and to 4'-OMe (δ_{H} 3.83) established the location of the methoxy group at C-4' and the hydroxy group at C-5'. The structure of bussealin B was thus assigned as 5',2'',3''-trihydroxy-2',4',4''-trimethoxy-1,3-diphenylpropane.

Bussealin C (**3**) was obtained as an off-white amorphous solid. Its positive ESI-MS revealed a pseudomolecular ion peak at m/z 305.1384 $[\text{M} + \text{H}]^+$ corresponding to molecular formula $\text{C}_{17}\text{H}_{21}\text{O}_5$. Its ^1H NMR and ^{13}C NMR spectra (Table 1) indicated that **3** is also a diphenylpropane with a 2'',3''-dihydroxy-4''-methoxyphenyl group substituted at C-3. The 1,3,4-trisubstituted A ring was determined by the proton coupling constants and HMBC correlations from H-2' (δ_{H} 6.64) and H-6' (δ_{H} 6.60) to C-1 (δ_{C} 36.1), and COSY correlations between H-5' (δ_{H} 6.79) and H-6' (δ_{H} 6.60). Furthermore, the HMBC spectrum showed a 3J correlation from H-6' to the methoxylated carbon at C-4' (δ_{C} 147.0), which was confirmed by NOESY correlations between H-5' (δ_{H} 6.79) and 4'-OMe (δ_{H} 3.80). The above data coupled with the molecular formula led to assignment of the structure of bussealin C as 3', 2'',3''-trihydroxy-4',4''-dimethoxy-1,3-diphenylpropane.

Bussealin D (**4**) was obtained as an off-white amorphous solid. The positive ESI-MS exhibited a pseudomolecular ion peak at m/z 349.1648 $[\text{M} + \text{H}]^+$ corresponding to the molecular formula $\text{C}_{19}\text{H}_{25}\text{O}_6$. The ^1H NMR and ^{13}C NMR spectra (Table 1) indicated that **4** had the same tetrasubstituted B ring with an OCH_3 group at C-4'' as in compounds **1–3**. In its ^1H NMR spectrum, the coupling patterns and the locations of the aromatic proton resonances of ring A were very similar to those of **2**. The presence of three OCH_3 groups and the substitution pattern of ring A of compound **4** were deduced by interpretation of the 1D and 2D NMR data. The HMBC spectrum of **4** showed correlations from H-1 (δ_{H} 6.79) to C-1' (δ_{C} 124.3), C-6' (δ_{C} 116.3) and to the methoxylated carbon at C-2' (δ_{C} 153.3). Furthermore, a clear 3J long-range correlation from the singlet proton H-3' (δ_{H} 6.61) to C-1' (δ_{C} 124.3) was also observed. Thus, the two remaining OCH_3 groups were determined to be at C-4' (δ_{C} 149.1) and C-5' (δ_{C} 144.1). The structure of bussealin D was thus determined to be 2'',3''-dihydroxy-2',4',5',4''-tetramethoxy-1,3-diphenylpropane.

The positive ESI-MS of bussealin E (**5**) displayed a pseudomolecular ion peak at m/z 331.1181 $[\text{M} + \text{H}]^+$ corresponding to the molecular formula $\text{C}_{18}\text{H}_{19}\text{O}_6$. The ^1H NMR spectrum in CDCl_3 showed signals for a singlet aromatic proton at δ_{H} 6.70, two OH groups (δ_{H} 5.75 and 5.69), three OCH_3 groups at δ_{H} 4.24, 4.24, and 4.01, and three methylene groups as multiplets at δ_{H} 3.13, 3.12 and 2.17. The ^{13}C NMR spectrum of **5** exhibited 18 signals, assigned to three methylene (δ_{C} 35.5, 28.7, 24.3), three OCH_3 (δ_{C} 60.8, 60.8 and 61.7), and twelve aromatic carbons of two isolated aromatic rings. Seven of the aromatic carbons were oxygenated, based on their deshielded chemical shifts (Table 2). The ten degrees of unsaturation implied by the molecular formula $\text{C}_{18}\text{H}_{18}\text{O}_6$ required two additional rings. Interpretation of ^1H - ^1H COSY, HMQC, HMBC and NOESY spectra allowed assignment of the locations of the functionalities present in **5**. In the COSY spectrum, the

three methylene groups were mutually coupled. The assignment of a singlet aromatic proton was substantiated by the observation of HMBC correlations from H-1 (δ_{H} 6.70) to C-10 (δ_{C} 35.5), C-3b (δ_{C} 118.2), and two oxygenated aromatic carbons at C-2 (δ_{C} 146.5) and C-3 (δ_{C} 129.7). HMBC correlations from the signal at δ_{H} 5.69 to C-1 (δ_{C} 110.1), C-2 (δ_{C} 146.5) and the methoxylated carbon at C-3 (δ_{C} 129.7) were observed, substantiating the location of a hydroxy group at C-2. The other hydroxy group was assigned to position 7 based on the observation of HMBC correlations from the signal at δ_{H} 5.75 to the carbon signals at C-6 (δ_{C} 136.5), C-7 (δ_{C} 142.3) and C-7a (δ_{C} 115.0). In addition, the signal at δ_{H} 5.75 showed NOESY correlations to H-8 (δ_{H} 3.13) and 6-OMe (δ_{H} 4.01). These observations required that the remaining OCH₃ group be placed at C-5. Furthermore, the HMBC correlations observed from H-10 (δ_{H} 3.12) to C-1 (δ_{C} 110.1), C-10a (δ_{C} 131.7), C-3b (δ_{C} 118.2), C-8 (δ_{C} 28.7) and C-9 (δ_{C} 24.3) confirmed the location of the cycloheptadiene ring. The above data confirmed the cycloheptadibenzofuran skeleton of **5**. Assignments of the ¹³C NMR signals of C-3a, C-4a and C-4b were made by comparing the measured data with those calculated by ACD/ChemSketch version 11.01. The calculated shifts were in excellent agreement with the observed values, and were all within the standard deviation of the software (5 ppm), except for C-7a. Therefore, the structure of **5** was assigned as 9,10-dihydro-2,7-dihydroxy-3,5,6-trimethoxy-8H-cyclohepta[klm]dibenzofuran.



It is noteworthy that bussealin E is the first cycloheptadibenzofuran isolated from natural sources, and the cycloheptadibenzofuran skeleton is rare among synthetic compounds. The only simple synthetic compound with this ring system is 9,10-dihydro-1-methyl-8H-cyclohepta[klm]dibenzofuran (**6**) and its 8-keto derivative.⁵

The presence of diphenylpropanes in *B. sakalava* suggests that bussealin E is biosynthesized by oxidative coupling of an appropriate precursor diphenylpropane. This could be followed by nucleophilic attack from a phenolate anion on a carbonyl group followed by dehydration to afford the new cycloheptadibenzofuran skeleton (**5**) as indicated in Scheme 1.

The bioactivity of diphenylpropanes has been widely studied. The diphenylpropane broussonin A inhibited respiratory syncytial-virus (RSV) more effectively than the standard antiviral drug ribavirin,⁶ and its anti-aromatase activity has also been evaluated.⁷ Broussonin B moderately inhibited a chymotrypsin-like activity of the proteasome.⁸ The anti-inflammatory,⁹ antifungal,¹⁰ antivascular,¹¹ adipogenic,¹² and anti-hCNT3 (human concentrative nucleoside transporter 3)¹⁴ activities of diphenylpropane analogues have also been reported. Since there have been no previous studies on the properties of diphenylpropanes on human ovarian cancer cells, we investigated the antiproliferative activity of diphenylpropanes **1–4** against the A2780 human ovarian cancer cell line. Bussealins A–D (**1–4**) showed only weak antiproliferative activities, with IC₅₀ values of 36, 24, 36, and 40 μM , respectively. Bussealin E (**5**), with a new chemical skeleton, was also tested against the A2780 cell line, but it also only exhibited weak activity with an IC₅₀ value of 45 μM . The new skeleton of bussealin E thus does not appear to confer any novel antiproliferative activity beyond that which is normal for diphenylpropanes.

Experimental Section

General Experimental Procedures

UV and IR spectra were measured on a Shimadzu UV-1201 spectrophotometer and a MIDAC M-series FTIR spectrophotometer, respectively. NMR spectra were recorded in CD₃OD or CDCl₃ on either JEOL Eclipse 500 or Bruker Avance 600 spectrometers. The chemical shifts are given in δ (ppm) and coupling constants (J) are reported in Hz. Mass spectra were obtained on an Agilent 6220 LC-TOF-MS. HPLC was performed on a Shimadzu LC-10AT instrument with a semi-preparative C18 Varian Dynamax column (5 μ m, 250 \times 10 mm).

Antiproliferative Bioassays

Antiproliferative activities were obtained at Virginia Polytechnic Institute and State University against the drug-sensitive A2780 human ovarian cancer cell line as previously described, except that the samples were added in 1 μ L 100% DMSO per well instead of 20 μ L of 1:1 DMSO:H₂O; paclitaxel (IC₅₀ 0.017 μ M) was used as a positive control.¹⁵ The A2780 cell line is a drug-sensitive ovarian cancer cell line.¹⁶

Plant Material

A sample of root of *Bussea sakalava* Du Puy & R. Rabev. (Fabaceae) was collected on January 25, 2007, near Ambolobozobe, Madagascar at coordinates 12°31'26"S 49°31'29"E, at an elevation of 20 m. Its assigned collection number is Rakotonandrasana et al. 1079. The genus *Bussea* Harms is a small genus including 7 species (5 from Tropical Africa and 2 from Madagascar). *B. sakalava* is endemic to deciduous forest from western to northern Madagascar. The hard wood of this species is used in construction and as firewood.¹⁷ Voucher specimens have been deposited at the Parc Botanique and Zoologique de Tsimbazaza (TAN) and at the Centre National d'Application des Recherches Pharmaceutiques (CNARP) in Antananarivo, Madagascar; the Missouri Botanical Garden in St. Louis, Missouri (MO); and the Muséum National d'Histoire Naturelle in Paris, France (P).

Extraction and Isolation

Dried roots of *B. sakalava* (275 g) were ground in a hammer mill, then extracted with ethanol by percolation for 24 hours at room temperature to give the crude extract MG 4273 (14.4 g), of which 3.0 g was shipped to Virginia Polytechnic Institute and State University (VPISU) for bioassay-guided isolation. Sample MG 4273 (IC₅₀ 9.6 μ g/mL, 2.1 g) was suspended in aqueous MeOH (MeOH-H₂O, 9:1, 100 mL) and extracted with hexane (3 \times 100 mL portions). The aqueous layer was then diluted to 60% MeOH (v/v) with H₂O and extracted with CH₂Cl₂ (3 \times 150 mL portions). The hexane extract was evaporated in vacuo to leave 227 mg with an IC₅₀ value of 19 μ g/mL. 102.9 mg of residue from the CH₂Cl₂ extract had an IC₅₀ of 10 μ g/mL. The aqueous MeOH extract (1.7 g) was inactive. The CH₂Cl₂ extract was selected for fractionation using an SPE cartridge over C-18, and two fractions were collected. Fractions I and II (70.2 mg and 26.8 mg) had IC₅₀ values of 8.6 and 15 μ g/mL, respectively. Fraction I was separated by C-18 HPLC (65% MeOH-H₂O), and compounds **1** (3.3 mg t_R 12.5 min), **2** (1.7 mg t_R 18.6 min), **3** (2.0 mg t_R 22.0 min), **4** (1.1 mg t_R 29.5 min) and **5** (1.1 mg t_R 26.5 min) were isolated.

3',5',2",3"-Tetrahydroxy-4',4"-dimethoxy-1,3-diphenyl-propane (**1**)

off-white amorphous solid; UV (MeOH) λ_{\max} nm (log ϵ) 218 (4.40), 267 (3.69), 294 (3.52); IR ν_{\max} cm⁻¹: 3367, 1648, 1450, 1115, 1024. ¹H NMR (500 MHz, CD₃OD) and ¹³C NMR

(125 MHz, CD₃OD), see Table 1; ESI-MS m/z 321.1338 [M + H]⁺ (calcd for C₁₇H₂₁O₆, 321.1338).

5',2",3"-Trihydroxy-2',4',4"-trimethoxy-1,3-diphenyl-propane (2)

off-white amorphous solid; UV (MeOH) λ_{\max} nm (log ϵ) 214 (4.25), 229 (sh) (4.10), 290 (3.59) nm; IR ν_{\max} cm⁻¹: 3332, 1599, 1444, 1095, 1032; ¹H NMR (500 MHz, CD₃OD) and ¹³C NMR (125 MHz, CD₃OD), see Table 1; ESI-MS m/z 335.1512 [M + H]⁺ (calcd for C₁₈H₂₃O₆, 335.1495).

3',2",3"-Trihydroxy-4',4"-dimethoxy-1,3-diphenyl-propane (3)

off-white amorphous solid; UV (MeOH) λ_{\max} nm (log ϵ) 208 (4.15), 267 (3.54), 289 (3.47) nm; IR ν_{\max} cm⁻¹: 3338, 1656, 1450, 1115, 1024; ¹H NMR (500 MHz, CD₃OD) and ¹³C NMR (125 MHz, CD₃OD), see Table 1; ESI-MS m/z 305.1384 [M + H]⁺ (calcd for C₁₇H₂₁O₅, 305.1389).

3',2",3"-Trihydroxy-4',4"-dimethoxy-1,3-diphenyl-propane (4)

off-white amorphous solid; UV (MeOH) λ_{\max} nm (log ϵ) 210 (4.21), 229 (sh) (4.01), 289 (3.48) nm; IR ν_{\max} cm⁻¹: 3350, 1602, 1450, 1115, 1026; ¹H NMR (500 MHz, CD₃OD) and ¹³C NMR (125 MHz, CD₃OD), see Table 1; ESI-MS m/z 349.1648 [M + H]⁺ (calcd for C₁₉H₂₅O₆, 349.1651).

9,10-Dihydro-2,7-dihydroxy-3,5,6-trimethoxy-8H-cyclohepta[klm]dibenzofuran (5)

off-white amorphous solid; UV (MeOH) λ_{\max} nm (log ϵ) 218 (4.25), 270 (3.78), 294 (3.73), 316 (3.47) nm; IR ν_{\max} cm⁻¹: 3332, 1567, 1449, 1115, 1024; ¹H NMR (600 MHz, CD₃OD and CDCl₃) and ¹³C NMR (150 MHz, CD₃OD), see Table 2; ESI-MS m/z 331.1181 [M + H]⁺ (calcd for C₁₈H₁₉O₆, 331.1182).

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

This project was supported by the Fogarty International Center, the National Cancer Institute, the National Science Foundation, the National Heart, Lung and Blood Institute, the National Institute of Mental Health, the Office of Dietary Supplements, and the Office of the Director of NIH, under Cooperative Agreement U01 TW000313 with the International Cooperative Biodiversity Groups. This project was also supported by the Agricultural Food Research Initiative of the National Institute of Food and Agriculture, USDA, Grant #2008-35621-04732. These supports are gratefully acknowledged. This work was also supported by the National Science Foundation under Grant No CHE-0619382 for purchase of the Bruker Avance 600 NMR spectrometer and Grant No. CHE-0722638 for the purchase of the Agilent 6220 mass spectrometer. We thank Mr. B. Bebout and Dr. Mehdi Ashraf-Khorassani for obtaining the mass spectra and Dr. Hugo Azurmendi for assistance with the NMR spectra. Fieldwork essential for this project was conducted under a collaborative agreement between the Missouri Botanical Garden and the Parc Botanique et Zoologique de Tsimbazaza and a multilateral agreement between the ICBG partners, including the Centre National d'Applications des Recherches Pharmaceutiques. We gratefully acknowledge courtesies extended by the Government of Madagascar (Ministère des Eaux et Forêts).

References and Notes

1. Biodiversity Conservation and Drug Discovery in Madagascar, Part 44. For Part 43, see Cao S, Hou Y, Brodie P, Miller JS, Randrianaivo R, Rakotobe E, Rasamison VE, Kingston DGI. Chem. Biodiversity. 2010 doi: 10.1002/cbdv.201000061.
2. Evans CS, Bell EA. Phytochemistry. 1978; 17:1127–1129.
3. Watson R, Fowden L. Phytochemistry. 1973; 12:617–622.

4. Freiburghaus F, Kaminsky R, Nkunya MHH, Brun R. J. Ethnopharm. 1996; 55:1–11.
5. Cagniant P, Bellinger N, Cagniant D. C. R. Acad. Sci., Série C. 1973; 277:383–385.
6. Bae G, Yu JR, Lee J, Chang J, Seo EK. Chem. Biodivers. 2007; 4:2231–2235. [PubMed: 17886842]
7. Lee D, Bhat KPL, Fong HHS, Farnsworth NR, Pezzuto JM, Kinghorn AD. J. Nat. Prod. 2001; 64:1286–1293. [PubMed: 11678652]
8. Tsukamoto S, Wakana T, Koimaru K, Yoshida T, Sato M, Ohta T. Biol. Pharm. Bull. 2005; 28:1798–1800. [PubMed: 16141565]
9. Leu YL, Hwang TL, Chung YM, Hong PY. Chem. Pharm. Bull. 2006; 54:1063–1066. [PubMed: 16819234]
10. Sawle P, Moulton BE, Jarzykowska M, Green CJ, Foresti R, Fairlamb IJS, Motterlini R. Chem. Res. Toxicol. 2008; 21:1484–1494. [PubMed: 18547064]
11. Lopes NP, Kato MJ, Yoshida M. Phytochemistry. 1999; 51:29–33.
12. Ducki S, Rennison D, Woo M, Kendall A, Chabert JFD, McGown AT, Lawrence NJ. Bioorg. Med. Chem. 2009; 17:7698–7710. [PubMed: 19837593]
13. Youn UJ, Lee YS, Jeong H, Lee J, Nam JW, Lee YJ, Hwang ES, Lee JH, Lee D, Kang SS, Seo EK. J. Nat. Prod. 2009; 72:1895–1898. [PubMed: 19757853]
14. Gupte A, Buolamwini JK. Bioorg. Med. Chem. Lett. 2009; 19:917–921. [PubMed: 19097778]
15. Cao S, Brodie PJ, Miller JS, Randrianaivo R, Ratovoson F, Birkinshaw C, Andriantsiferana R, Rasamison VE, Kingston DGI. J. Nat. Prod. 2007; 70:679–681. [PubMed: 17323994]
16. Louie KG, Behrens BC, Kinsella TJ, Hamilton TC, Grotzinger KR, McKoy WM, Winker MA, Ozols RF. Cancer Res. 1985; 45:2110–2115. [PubMed: 3986765]
17. Du Puy, DJ.; Labat, J-N.; Rabevohitra, R.; Villiers, J-F.; Bossier, J.; Moat, J. The Leguminosae of Madagascar. Kew: Royal Botanical Gardens; 2002. p. 22-24.





Figure 1.
COSY, HMBC and NOESY correlations of 5



Scheme 1.
Possible biosynthesis of cycloheptadibenzofuran **5** in *B. sakalava*

Table 1

 ^1H and ^{13}C NMR data for Bussealin A–D (1–4)^a

position	1		2		3		4	
	^1H (J, Hz)	^{13}C	^1H (J, Hz)	^{13}C	^1H (J, Hz)	^{13}C	^1H (J, Hz)	^{13}C
1	2.41 t (7.9)	36.5	2.50 t (7.8)	30.4	2.49 t (7.9)	36.1	2.55 t (7.7)	30.4
2	1.79 m	33.0	1.76 m	31.9	1.81 m	33.2	1.79 m	31.8
3	2.52 t (7.7)	30.6	2.54 t (7.8)	30.7	2.54 t (7.7)	30.6	2.55 t (7.7)	30.6
1'		140.2		124.7		137.2		124.3
2'	6.18 s	108.7		152.2	6.64 d (2.0)	116.5		153.3
3'		151.3	6.58 s	99.4		147.3	6.61 s	99.6
4'		134.7		147.2		147.0		149.1
5'		151.3		141.0	6.79 d (8.2)	112.9		144.1
6'	6.18 s	108.7	6.60 s	117.9	6.60 dd (8.2, 2.0)	120.6	6.75 s	116.3
1''		123.4		123.7		123.5		123.6
2''		144.7		144.7		144.7		144.7
3''		134.9		134.9		135.0		134.9
4''		147.8		147.8		147.8		147.8
5''	6.38 d (8.4)	103.8	6.39 d (8.4)	103.8	6.39 d (8.4)	103.9	6.39 d (8.4)	103.9
6''	6.50 d (8.4)	120.5	6.51 d (8.4)	120.4	6.50 d (8.5)	120.5	6.51 d (8.3)	120.4
2'-OMe			3.76 s	56.8			3.78 s	56.6
4'-OMe	3.75 s	60.8	3.83 s	57.0	3.80 s	56.6	3.82 s	56.8
5'-OMe							3.76 s	57.6
4''-OMe	3.78 s	56.5	3.80 s	56.6	3.80 s	56.6	3.80 s	56.9

^aIn CD₃OD; δ (ppm) 500 MHz for ^1H and 125 MHz for ^{13}C ; multiplicities; J values (Hz) in parentheses.

Table 2

 ^1H and ^{13}C NMR Data for Bussealin E (5)

position	$^1\text{H}^a$	$^{13}\text{C}^a$	$^{13}\text{C}^b$	$^1\text{H}^c$
1	6.70 s	110.1	110.0	6.59 s
2		146.5	149.1	
3		129.7	131.3	
3a		146.2	148.7	
3b		118.2	113.3	
4a		140.7	139.7	
4b		120.6	117.4	
5		135.6	137.0	
6		136.5	137.9	
7		142.3	145.1	
7a		115.0	109.7	
8	3.13 m	28.7	28.9	3.08 m
9	2.17 m	24.3	24.2	2.12 m
10	3.12 m	35.5	34.5	3.07 m
10a		131.7	128.6	
3-OCH ₃	4.24 s	60.8	61.5	4.18 s
2-OH	5.69 s			
5-OCH ₃	4.24 s	60.8	61.6	4.08 s
6-OCH ₃	4.01 s	61.7	61.0	3.90 s
7-OH	5.75 s			

^aIn CDCl₃; δ (ppm) 600 MHz for ^1H and 150 MHz for ^{13}C ; multiplicities; *J* values (Hz) in parentheses.

^bCalculated using ACD/ChemSketch version 11.01.

^cIn CD₃OD; δ (ppm) 600 MHz for ^1H ; multiplicities; *J* values (Hz) in parentheses.