

RESEARCH ARTICLE

Open Access

Boron neutron capture therapy induces apoptosis of glioma cells through Bcl-2/Bax

Peng Wang^{1†}, Haining Zhen^{1†}, Xinbiao Jiang^{2†}, Wei Zhang¹, Xin Cheng³, Geng Guo¹, Xinggang Mao¹, Xiang Zhang^{1*}

Abstract

Background: Boron neutron capture therapy (BNCT) is an alternative treatment modality for patients with glioma. The aim of this study was to determine whether induction of apoptosis contributes to the main therapeutic efficacy of BNCT and to compare the relative biological effect (RBE) of BNCT, γ -ray and reactor neutron irradiation.

Methods: The neutron beam was obtained from the Xi'an Pulsed Reactor (XAPR) and γ -rays were obtained from [⁶⁰Co] γ source of the Fourth Military Medical University (FMMU) in China. Human glioma cells (the U87, U251, and SHG44 cell lines) were irradiated by neutron beams at the XAPR or [⁶⁰Co] γ -rays at the FMMU with different protocols: Group A included control nonirradiated cells; Group B included cells treated with 4 Gy of [⁶⁰Co] γ -rays; Group C included cells treated with 8 Gy of [⁶⁰Co] γ -rays; Group D included cells treated with 4 Gy BPA (p-borono-phenylalanine)-BNCT; Group E included cells treated with 8 Gy BPA-BNCT; Group F included cells irradiated in the reactor for the same treatment period as used for Group D; Group G included cells irradiated in the reactor for the same treatment period as used for Group E; Group H included cells irradiated with 4 Gy in the reactor; and Group I included cells irradiated with 8 Gy in the reactor. Cell survival was determined using the 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium (MTT) cytotoxicity assay. The morphology of cells was detected by Hoechst33342 staining and transmission electron microscope (TEM). The apoptosis rate was detected by flow cytometer (FCM). The level of Bcl-2 and Bax protein was measured by western blot analysis.

Results: Proliferation of U87, U251, and SHG44 cells was much more strongly inhibited by BPA-BNCT than by irradiation with [⁶⁰Co] γ -rays ($P < 0.01$). Nuclear condensation was determined using both a fluorescence technique and electron microscopy in all cell lines treated with BPA-BNCT. Furthermore, the cellular apoptotic rates in Group D and Group E treated with BPA-BNCT were significantly higher than those in Group B and Group C irradiated by [⁶⁰Co] γ -rays ($P < 0.01$). The clonogenicity of glioma cells was reduced by BPA-BNCT compared with cells treated in the reactor (Group F, G, H, I), and with the control cells ($P < 0.01$). Upon BPA-BNCT treatment, the Bax level increased in glioma cells, whereas Bcl-2 expression decreased.

Conclusions: Compared with γ -ray and reactor neutron irradiation, a higher RBE can be achieved upon treatment of glioma cells with BNCT. Glioma cell apoptosis induced by BNCT may be related to activation of Bax and downregulation of Bcl-2.

Background

Glioma is one of the most aggressive human malignancies. Despite aggressive surgery combined with adjuvant radiotherapy and chemotherapy, patient prognosis remains poor. It is thus essential to develop novel

therapeutic strategies to treat glioma. Boron neutron capture therapy (BNCT) is a highly selective treatment modality that can target the tumor without causing excessive radiation damage to the normal tissues [1]. The ability of [¹⁰B] to capture thermal neutrons, and to then disintegrate immediately into a He nucleus (an α particle) and a Li nucleus, is utilized in BNCT [2]. Both the He and Li nuclei have high lineal energy transfer (LET) values, with path-lengths approximately the diameter of a single cell; the lethality of these nuclei is thus

* Correspondence: xzhang@fmmu.edu.cn

† Contributed equally

¹Department of Neurosurgery of Xijing Hospital, Fourth Military Medical University, Xi'an, Shaanxi 710032, P.R.China

Full list of author information is available at the end of the article

primarily limited to boron-containing cells [3]. BNCT has been carried out in numerous studies using syngeneic rat gliomas [3-6]. Approaches such as administering [¹⁰B] by intra-arterial injection with or without blood-brain barrier (BBB) disruption [7-9], or intracerebral delivery of high molecular weight agents by means of convection enhanced delivery following neutron beam irradiation [2,10], confirm that BNCT is highly effective for treatment glioma xenograft models. In clinic, BNCT has been used in various countries to treat patients with high-grade glioma, melanoma, liver metastasis of colon adenocarcinoma, malignant melanomas, oral cancer, and therapeutically refractory, recurrent tumors of the head and neck [3,6,11-15]; while the mechanism of BNCT-induced cell death remains unclear.

We treated U87, U251, and SHG44 glioma cells with BNCT *in vitro*, using L-p-borono-phenylalanine (L-BPA) as the boron carrier and the thermal neutron source of the Xi'an Pulsed Reactor (XAPR) in China as the irradiator. The relative biological effect (RBE) of BNCT on tested cells, especially with respect to apoptosis induction, was assessed, and a possible mechanism of BNCT action was presented.

Methods

Glioma cell culture

The human glioblastoma multiforme cell lines U87 and U251 were obtained from the American Type Culture Collection (ATCC; Manassas, VA). The human anaplastic astrocytoma cell line SHG44 was purchased from the Institute of Biochemistry and Cell Biology (IBCB; Shanghai, China). Cells were cultured in Dulbecco's modified Eagle's medium (DMEM) (Gibco, Grand Island, NY) with 10% (v/v) fetal bovine serum (Gibco) and 0.5% (w/v) penicillin-streptomycin solution (Gibco), at 37°C under 5% CO₂ and 95% air (both v/v).

Boron uptake by glioma cells *in vitro*

[¹⁰B]-enriched L-BPA (Ryscor Inc., Raleigh, NC) was used in the study. L-BPA was added to the culture media of glioma cells to 50 µg [¹⁰B]/ml. Cells in the exponential phase of growth were cultured in boron-containing medium for 3, 6, 12, or 24 h before harvesting, and boron concentrations were determined by inductively coupled plasma atomic emission spectroscopy (ICP-AES) (VISTA-MPX; Varian Co., Walnut Creek, CA) according to a previously described method [16].

In vitro irradiation

In vitro irradiation of the three cell lines was performed using the neutron beam at the XAPR whereas *in vitro* γ-ray irradiation by the [⁶⁰Co] γ source was performed in the Fourth Military Medical University (Xi'an, China).

Cells were divided into nine groups. Group A was control nonirradiated cells; Group B cells were irradiated with 4 Gy of [⁶⁰Co] γ-rays; Group C cells were irradiated with 8 Gy of [⁶⁰Co] γ-rays; Group D cells were irradiated with 4 Gy BPA-BNCT; Group E cells were irradiated with 8 Gy BPA-BNCT; Group F cells were irradiated in the XAPR for the same treatment period as used for Group D; Group G cells were irradiated in the XAPR for the same treatment period as used for Group E; Group H cells were irradiated with 4 Gy in the XAPR; and Group I cells were irradiated with 8 Gy in the XAPR. In Group D and Group E with BPA-BNCT, cells were incubated for 24 h in medium with BPA and washed three times with phosphate-buffered saline (PBS); the medium was next replaced with boron-free medium; cells were exposed to the XAPR with irradiation doses of 4 Gy and 8 Gy, respectively. In the γ-ray-treated Group B and Group C, cells incubated in growth medium without BPA were exposed to the γ-rays with irradiation doses of 4 Gy and 8 Gy, respectively. In Group F and Group G, cells incubated in medium without BPA were exposed to the XAPR with identical treatment durations of Group D and Group E, respectively. In Group H and Group I, cells incubated in growth medium without BPA were exposed to the XAPR with irradiation doses of 4 Gy and 8 Gy, respectively. Irradiated and nonirradiated cells were placed at room temperature and under a normal atmosphere during irradiation. At the beginning of the experiment, all cells were in the exponential phase of growth and at a density of 3×10⁵/ml. The components of the reactor neutron source and the [⁶⁰Co] γ source were shown in Table 1; the irradiation periods of various groups were listed in Table 2; the dose components of BNCT were listed in Table 3 and Table 4. The irradiation doses were calculated according to previous studies [17,18].

Proliferation assays

Cell survival was determined using the 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium (MTT) (Sigma, St. Louis, MO) assay. After irradiation, cells were trypsinized and washed, and cell concentrations were adjusted to 10⁴ cells/ml. Cells were seeded into the

Table 1 Dose rates for reactor neutron source and [⁶⁰Co] γ source

	Components	Dose rate (Gy/min)
Dose rates for the reactor neutron source	Thermal neutron	1.04×10 ⁻³
	Epithermal neutron	2.48×10 ⁻⁴
	Fast neutron	1.07×10 ⁻¹
	γ-ray	1.57×10 ⁻²
	¹⁰ B(n,α) ⁷ Li	5.09×10 ⁻⁴ /ppm ¹⁰ B
Dose rate for the [⁶⁰ Co] γ source		1.40×10 ¹

Table 2 The irradiation periods of various groups

	Group A (min)	Group B (min)	Group C (min)	Group D (min)	Group E (min)	Group F (min)	Group G (min)	Group H (min)	Group I (min)
U87	64	0.29	0.57	27	54	27	54	32	64
U251	64	0.29	0.57	25	50	25	50	32	64
SHG44	64	0.29	0.57	28	56	28	56	32	64

wells of 96-well microplates (Nunc, Roskilde, Denmark) and incubated for 2, 4, or 6 days. The culture medium was removed and replaced with 100 µl aliquots of fresh medium, without serum, containing 0.5 mg/ml MTT, followed by 4 h of incubation at 37°C. Next, medium was aspirated from the wells and 100 µl dimethyl sulfoxide (DMSO) was added to each well, to dissolve formazan crystals. Optical density (OD) was measured using an enzyme-linked immunosorbent method. Cell numbers were derived from OD values by reference to a standard cell number-OD curve.

Hoechst staining

At 12 h, 24 h, 36 h, and 48 h after treatment with BPA-BNCT, cells were fixed for 10 min in 4% (v/v) paraformaldehyde, and then incubated with Hoechst 33342 dye (Sigma) (10 µg/ml) for 10 min. After washed with PBS, cells were observed using an inverted fluorescence microscope (IX70; Olympus, Tokyo, Japan).

Electron microscopy

Forty-eight hours after irradiation, cells were detached from plates, washed, suspended in PBS, and concentrated by centrifugation. Cell samples were fixed in 2.5% (v/v) glutaraldehyde in 0.1 M phosphate buffer (pH 7.4), postfixed in 2% (w/v) buffered osmium tetroxide for 2 h, and dehydrated in ethanol. Specimens for transmission electron microscopy were embedded in Epon. Thin sections were cut using an ultramicrotome and double-stained with uranyl acetate and lead citrate. Electron micrography was performed (JEM-2000EX; JEOL, Tokyo, Japan) using an operating voltage of 80 kV.

AnnexinV/PI analysis

Twelve hours, 24 h, 36 h, and 48 h after irradiation, cells were trypsinized, counted, washed twice in ice-cold PBS solution, and resuspended in 1×binding buffer (10 mM HEPES/NaOH [pH 7.4], 140 mM NaCl, and 2.5 mM CaCl₂). Next, 5 µl of annexin V-FITC (AV) (R&D

Systems Europe Ltd., Abingdon, UK) and 10 µl of propidium iodide (PI) (Sigma) were added to 100 µl of cell suspension, followed by incubation for 15 min at room temperature in the dark. Finally, 400 µl of binding buffer was added to each sample, which was held on ice prior to analysis on a FACSCalibur (Becton Dickinson Labware, Franklin Lakes, NJ) flow cytometer. Ten thousand cells per sample were analyzed.

Clonogenic assay

After irradiation, 200 treated cells were plated in six-well flat-bottomed microplates (Nunc) and incubated at 37°C in a humidified incubator under 5% (v/v) CO₂ for 2 weeks. The cells were fixed in methanol and stained with Giemsa's solution. Colonies ≥ 50 µm in diameter were counted [19].

Western blot analysis

Twelve hours and 24 h after irradiation, cells were washed twice in ice-cold PBS and protein extracts of U87, U251, and SHG44 cells were prepared by lysis in RIPA buffer (150 mM NaCl, 1% [v/v] NP-40, 0.5% [w/v] sodium deoxycholate, 0.1% [w/v] sodium dodecyl sulfate (SDS), 50 mM Tris HCl [pH 8], 10 mM EDTA, and 1 mM PMSF [Sigma]) for 30 min at 4°C. Samples were next centrifuged for 15 min at 15,000 g. Protein concentrations of supernatants were determined. For each sample, 60 µg of protein was loaded on a 12.5% (w/v) SDS-polyacrylamide gel, electrophoresed, and transferred to a nitrocellulose membrane (Protran; Schleicher and Schuell, Florham Park, NJ). Each membrane was blocked for 1 h at room temperature with blocking buffer (TBS containing 0.1% [v/v] Tween 20 [Sigma] and 5% [w/v] milk powder). Primary antibodies (applied for 1 h at room temperature, or overnight at 4°C) were: anti-Bcl-2 (Santa Cruz Biotechnology, Inc., Heidelberg, Germany), anti-Bax (Santa Cruz), and anti-actin (mouse monoclonal C-2, Santa Cruz). Antibodies were diluted 1:1,000, except for the anti-actin antibody (1:500).

Table 3 The dose components of BNCT 4 Gy

	Thermal neutron (Gy)	Epithermal neutron (Gy)	Fast neutron (Gy)	γ-ray (Gy)	¹⁰ B(n,α) ⁷ Li (Gy)
U87	0.028	0.007	2.889	0.423	0.653
U251	0.026	0.007	2.675	0.392	0.900
SHG44	0.029	0.007	2.996	0.439	0.529

Table 4 The dose components of BNCT 8 Gy

	Thermal neutron (Gy)	Epithermal neutron (Gy)	Fast neutron (Gy)	γ -ray (Gy)	$^{10}\text{B}(n,\alpha)^7\text{Li}$ (Gy)
U87	0.056	0.013	5.778	0.846	1.306
U251	0.052	0.013	5.350	0.748	1.800
SHG44	0.058	0.013	5.992	0.878	1.058

Thereafter, membranes were incubated for 1 h with HRP-labeled secondary antibodies (Amersham Pharmacia Biotech, Uppsala, Sweden), either sheep anti-mouse (diluted 1:2,500) or donkey anti-rabbit (1:5,000), and the blots were finally developed using an ECL system, according to the manufacturer's instructions (Amersham Bioscience, Buckinghamshire, UK).

Statistical analysis

Outcome variables are expressed as means \pm standard deviations (SDs). Statistical analysis was performed using SPSS Version 13.0 (SPSS, Chicago, IL). ANOVA was used for data analysis. All statistical tests were two-sided, and P values < 0.05 were considered to be statistically significant.

Results

Boron uptake

As expected, [^{10}B] was taken up by U87, U251, and SHG44 cells during incubation for 24 hours in medium containing 50 μg [^{10}B]/ml. The boron concentrations in cells eventually reached $2.72 \pm 0.25 \mu\text{g}/10^7$ cells, $9.78 \pm 0.49 \mu\text{g}/10^7$ cells, and $2.48 \pm 0.34 \mu\text{g}/10^7$ cells for the U87, U251, and SHG44 cell lines, respectively, at 24 h (Figure 1).

Tumor cell numbers fall after BPA-BNCT treatment

The inhibitory effect of BPA-BNCT on proliferation of U87 cells was much more significant than that afforded by γ -rays, as confirmed by the MTT assay ($P < 0.01$, Figure 2). Any inhibitory effect was marginal in cells

receiving 4 Gy or 8 Gy of γ -rays, whereas significant inhibition was observed in cells treated with 4 Gy or 8 Gy of BPA-BNCT. The number of viable U87 cells decreased in 48 h after treatment in all of Groups D, E, F, G, H, and I, and the most potent inhibition was observed in the group receiving 8 Gy of BPA-BNCT. Similar effects were observed in SHG44 and U251 cells (data not shown).

BNCT induces apoptosis

Typical apoptotic morphological changes were found in all of the U87, U251, and SHG44 cell lines 12 h after treatment with BPA-BNCT; the changes included shrinkage, deformation, and detachment from culture dishes. Nuclear condensation and chromatin margination were evident by Hoechst 33342 staining (Figure 3A, 3B), and chromatin margination, nuclear condensation, and segmentation were noted on transmission electron microscopy (Figure 3C).

The apoptotic frequency of U87 cells treated with 4 Gy or 8 Gy of BPA-BNCT was significantly higher than that of cells treated with equivalent doses of γ -rays ($P < 0.01$), as shown by annexin V/PI staining (Figure 4, 5). Furthermore, the apoptotic frequency of cells treated with 8 Gy




Figure 1 Time course of boron accumulation. Data represent the mean \pm SD of triplicate experiments.







Figure 2 Growth curves of U87 cells in different groups. Group A: control nonirradiated cells; Group B: cells treated with 4 Gy of [^{60}Co] γ -rays; Group C: cells treated with 8 Gy of [^{60}Co] γ -rays; Group D: cells treated with 4 Gy BPA-BNCT; Group E: cells treated with 8 Gy BPA-BNCT; Group F: cells irradiated in the reactor for the same treatment period as used for Group D; Group G: cells irradiated in the reactor for the same treatment period as used for Group E; Group H: cells irradiated with 4 Gy in the reactor; Group I: cells irradiated with 8 Gy in the reactor. Data represent the mean \pm SD of triplicate experiments.



BPA-BNCT was higher than that seen after treatment with 4 Gy BPA-BNCT ($P < 0.01$). Also, the apoptotic frequencies of U87 cells after BPA-BNCT treatment were greater than those noted after delivery of equivalent reactor doses of irradiation ($P < 0.05$). Similar effects were observed in SHG44 and U251 cells (Figure 4, 5).



BNCT inhibits colony formation

The survival rates of nonirradiated control cells were set at 100%. Cell survival proportions were shown in Table 5. Survival analysis of U87, U251, and SHG44 cells showed that all three cell lines were inhibited more potently by BNCT than by irradiation with ⁶⁰Co γ-rays ($P < 0.01$) or by general reactor neutron irradiation ($P < 0.01$).

Altered expression of Bcl-2 and Bax after BNCT

In all of U87, U251, and SHG44 cells, Bcl-2 was initially abundantly expressed, whereas Bax expression was low. After treatment with 4 Gy or 8 Gy of BPA-BNCT, Bcl-2 expression was down-regulated, while Bax expression increased simultaneously. Furthermore, 24 h after treatment with 4 Gy BPA-BNCT, Bcl-2 expression was lower than that seen 12 h after treatment with 4 Gy BPA-BNCT, and the level of Bax expression was higher. The levels of Bcl-2 and Bax expression were similar 12 h and 24 h after treatment with 8 Gy BPA-BNCT (Figure 6).

Discussion

Two criteria must be fulfilled when BNCT is to be applied: an appropriate source of high-flux thermal neutrons is necessary, along with a boron carrier that is

Table 5 Survival fractions of glioma cells in different groups

Group	Survival fraction (%)		
	U87	U251	SHG44
Unirradiated control (A)	100	100	100
Irradiated in ⁶⁰ Co by 4 Gy (B)	68.1	63.7	64.5
Irradiated in ⁶⁰ Co by 8 Gy (C)	10.9	13.6	12.4
Irradiated in reactor for the same treatment period as used for Group D (F)	3.1	2.3	2.5
Irradiated in reactor for the same treatment period as used for Group E (G)	1.6	1.4	1.6
Irradiated in reactor by 4 Gy (H)	2.9	2.2	2.6
Irradiated in reactor by 8 Gy (I)	1.4	0.9	1.2
Irradiated in reactor with BNCT by 4 Gy (D)*	<0.01%	<0.01%	<0.01%
Irradiated in reactor with BNCT by 8 Gy (E)**	<0.01%	<0.01%	<0.01%

*P <0.01 vs A, B, F and H **P <0.01 vs A, C, G, I and D

able to concentrate in glioma tissues. Therefore, we initially verified uptake of BPA by glioma cells *in vitro*. We found that the boron concentration in the three glioma cell lines increased constantly over the first 24 hours of incubation. Irradiation was conducted at this time, and the [¹⁰B] level attained the theoretical requirement, in that at least 10⁹ [¹⁰B] atoms accumulated in each glioma cell and the thermal neutron fluence could attain 10¹²-10¹³ n·cm⁻², to produce a lethal discharge of high-LET particles [20].

BNCT can inhibit cell proliferation because the RBE is high, as has been shown in animal models. In the present study, although the three glioma cell lines differed in boron concentration after exposure to BPA, the therapeutic efficiency of BNCT was similar in all three lines because the RBE was high. Proliferation of glioma cells was significantly suppressed by BNCT, and the clonogenicity of glioma cells after BNCT treatment was less than 0.01% of control values.

Apoptosis, an active process of cellular suicide, is an important means of radiation-induced cell death [21], and may contribute to the therapeutic efficacy of BNCT. Many triggers of apoptosis are known, including cytogenetic alterations such as micronuclei (MN) formation and chromosomal aberrations (CA) within cellular systems [22]. Furthermore, several studies have sought to correlate the extent of apoptosis after irradiation with

cellular radiosensitivity [23]. Given that BNCT works mainly through the discharge effects of high-LET Li nuclei and α particles, we investigated whether BNCT could induce apoptosis of glioma cells. Typical apoptotic morphological changes were confirmed in U87, U251, and SHG44 cells after BNCT treatment, using transmission electron microscopy and Hoechst 33342 staining. Furthermore, previous studies have shown that BPA has no toxicity effect on cells at a dosage up to 110 μg [¹⁰B]/ml [24]. Few apoptotic changes were apparent in nonirradiated control cells, showing, for the first time, that the pro-apoptotic effect of BNCT plays an important role in the reduction of cell proliferation. Masunaga et al. described that BNCT induced apoptosis *in vivo*, the proportion of apoptotic cells increased slightly 6 h after BNCT, and the apoptotic rate was relatively low [17,18]. Fujita et al. stated that apoptosis was a form of cell death induced by BNCT [25]. While Aromando et al. reported that apoptosis did not have a significant role in BNCT-induced hamster cheek pouch tumor control [26], and Kamida et al. described that differences in apoptotic cells pre- and post-BNCT in human oral squamous cell carcinoma xenografts were minimal [27]. Whether BNCT can induce significant apoptosis rate in glioma cells *in vivo* and whether the pro-apoptotic effect is cell-type-dependant deserved further study.

It is well established in mammalian systems that high-LET irradiations have more pronounced effects on biological systems than that of low LET [22], owing to the characteristic mode of energy deposition along a track. Such energy generates complex forms of radiation damage to biological molecules. The transfer of the energy of high-LET radiation to such molecules can break chemical bonds and ionize various cellular components.

In the present study, AnnexinV/PI analysis not only confirmed the pro-apoptotic effect of BNCT, but also showed that apoptosis frequencies were higher in groups treated with BNCT compared with γ-rays. Moreover, the




Figure 6 The expression of Bcl-2 and Bax proteins in (A) U87 cells, (B) U251 cells, and (C) SHG44 cells, 12 h and 24 h after treatment with 4 Gy or 8 Gy BPA-BNCT.

apoptotic frequencies in BPA-BNCT groups were higher than those of the irradiated controls. Additionally, when the absorbed doses were identical, both the proliferation and the clonogenicity of glioma cells were suppressed by BNCT to an extent significantly greater than seen after exposure to a neutron beam. Thus, the RBE of BNCT is higher than that of either a neutron beam or γ -rays. In the present study, the contribution of a boron-dependent dose to the total BNCT dose was low. Thus, with optimization of the reactor and improvement in the boron-dependent dose, the RBE of BNCT can be further increased.

Previous studies have shown that the intrinsic apoptotic pathway (involving the mitochondria) plays a key role in the regulation of apoptosis in response to irradiation [28]. Bcl-2 family proteins interact to permeabilize the mitochondrial membrane, thus playing an important role in irradiation-induced apoptosis [29-31]. Bcl-2 and Bax, the two main members of the Bcl-2 protein family, function as tumor anti-apoptotic and pro-apoptotic factors, respectively [32-34]. Bax can cause release of cytochrome c after formation of the hetero-oligomer Bax/Bak, whereas Bcl-2 can engage with activator proteins or Bax/Bak, thereby sequestering these proteins [28]. We detected expression of Bcl-2 and Bax in all of U87, U251, and SHG44 cells after BNCT treatment in efforts to understand the impact of BNCT on the mitochondrial pathway of apoptosis. Down-regulation of Bcl-2 and up-regulation of Bax synthesis after BNCT showed that apoptosis induced by BNCT may be mediated through the mitochondrial pathway.

A Bcl-2 antisense has been developed as a gene therapeutic agent to treat cancer, such as: melanoma carcinoma, multiple myeloma, and small cell lung cancer [28]. Elimination of Bcl-2 induced by BNCT indicates that a directed Bcl-2-targeting strategy may be combined with BNCT, thus possibly leading to synergistic reduction of Bcl-2; this possibility deserves further investigation.

Our investigation of BPA-BNCT therapy was conducted *in vitro*, and many constraints are involved in attempts to extrapolate *in vitro* data to the *in vivo* context. During BNCT, the normal brain tissue and tumors are non-specifically irradiated by reactor-generated mixed irradiation beam components, and the patient tolerance to BNCT could be impaired. When it is considered that the RBE of BNCT is higher than that of γ -rays, neutron beam, and X-rays, all of which are widely used as treatment options for patients with brain tumors, the absorbed dose from BNCT can be much lower than those strategies. This will limit irradiation damage to normal tissue. The boron level in blood and brain parenchyma also contributes to the irradiation dose to normal tissue. With the development of boron containing agents and the optimization of strategies of

delivery, [^{10}B] can be highly accumulated in tumor cells and the level in normal cells becomes relatively lower [2,13,35,36]. Moreover, BNCT relies mainly on the use of boron compounds, and not on a neutron beam *per se*, to selectively irradiate a tumor [37]. Thus, the optimization of selective boron delivery to tumor cells is an important area of BNCT research. As known the BBB can impede the uptake of boron-containing compounds by tumor cells, with the invasive properties of glioma tumors, the BBB can be destroyed and the boron uptake by tumor cells could be increased. Drugs such as mannitol that transiently compromise BBB integrity have been used to improve boron absorbance by tumor cells. Overall, the use of BNCT *in vivo* will likely become effective, with a reduction in side effects and improvements in tumor-targeting boron compounds, following further development.

Conclusions

Altogether, induction of apoptosis contributes to the main therapeutic efficacy of BNCT to glioma cells *in vitro*. Activation of Bax and downregulation of Bcl-2 are involvement in the apoptosis triggered by BNCT. Moreover, the RBE of BPA-BNCT is higher than that of γ -ray and reactor neutron irradiation. Understanding the mechanisms involved in BNCT will ultimately contribute to the enhancement of the therapeutic effectiveness of this therapeutic modality.

Acknowledgements

This research was supported by the National Natural Science Foundation (No: 10375047), Beijing, P.R. China.

Author details

¹Department of Neurosurgery of Xijing Hospital, Fourth Military Medical University, Xi'an, Shaanxi 710032, P.R.China. ²Northwest Institute of Nuclear Technology, Xi'an, Shaanxi 710024, PR China. ³Department of Biochemistry and Molecular Biology, Fourth Military Medical University, Xi'an, Shaanxi 710032, PR China.

Authors' contributions

Peng Wang, Haining Zhen, and Xinbiao Jiang carried out the cell culture and experiment of irradiation. Peng Wang and Xin Cheng carried out Proliferation assays, Hoechst staining, Electron microscopy assay, AnnexinV/PI analysis, Colony formation assay, and drafted the manuscript. Wei Zhang, Geng Guo, and Xinggang Mao participated in the design of the study and performed the statistical analysis. Xiang Zhang conceived of the study, and participated in its design and coordination. All authors read and approved the final manuscript.

Competing interests

The authors declare that they have no competing interests.

Received: 8 February 2010 Accepted: 2 December 2010

Published: 2 December 2010

References

1. Barth RF, Joensuu H: Boron neutron capture therapy for the treatment of glioblastomas and extracranial tumours: as effective, more effective or less effective than photon irradiation? *Radiother Oncol* 2007, **82**:119-122.

2. Yang W, Barth RF, Wu G, Huo T, Tjarks W, Ciesielski M, Fenstermaker RA, Ross BD, Wikstrand CJ, Riley KJ, Binns PJ: **Convection enhanced delivery of boronated EGF as a molecular targeting agent for neutron capture therapy of brain tumors.** *J Neurooncol* 2009, **95**:355-365.
3. Barth RF, Coderre JA, Vicente MG, Blue TE: **Boron neutron capture therapy of cancer: current status and future prospects.** *Clin Cancer Res* 2005, **11**:3987-4002.
4. Yang W, Barth RF, Wu G, Bandyopadhyaya AK, Thirumamagal BT, Tjarks W, Binns PJ, Riley K, Patel H, Coderre JA, Ciesielski MJ, Fenstermaker RA: **Boronated epidermal growth factor as a delivery agent for neutron capture therapy of EGF receptor positive gliomas.** *Appl Radiat Isot* 2004, **61**:981-985.
5. Barth RF, Yang W, Coderre JA: **Rat brain tumor models to assess the efficacy of boron neutron capture therapy: a critical evaluation.** *J Neurooncol* 2003, **62**:61-74.
6. Barth RF: **Boron neutron capture therapy at the crossroads: challenges and opportunities.** *Appl Radiat Isot* 2009, **67**:S3-S6.
7. Yang W, Barth RF, Carpenter DE, Moeschberger ML, Goodman JH: **Enhanced delivery of boronophenylalanine for neutron capture therapy by means of intracarotid injection and blood-brain barrier disruption.** *Neurosurgery* 1996, **38**:985-992.
8. Barth RF, Yang W, Rotaru JH, Moeschberger ML, Boesel CP, Soloway AH, Joel DD, Nawrocky MM, Ono K, Goodman JH: **Boron neutron capture therapy of brain tumors: enhanced survival and cure following blood-brain barrier disruption and intracarotid injection of sodium borocaptate and boronophenylalanine.** *Int J Radiat Oncol Biol Phys* 2000, **47**:209-218.
9. Barth RF, Yang W, Rotaru JH, Moeschberger ML, Joel DD, Nawrocky MM, Goodman JH, Soloway AH: **Boron neutron capture therapy of brain tumors: enhanced survival following intracarotid injection of either sodium borocaptate or boronophenylalanine with or without blood-brain barrier disruption.** *Cancer Res* 1997, **57**:1129-1136.
10. Yang W, Barth RF, Adams DM, Ciesielski MJ, Fenstermaker RA, Shukla S, Tjarks W, Caligiuri MA: **Convection-enhanced delivery of boronated epidermal growth factor for molecular targeting of EGF receptor-positive gliomas.** *Cancer Res* 2002, **62**:6552-6558.
11. Kato I, Fujita Y, Maruhashi A, Kumada H, Ohmae M, Kirihata M, Imahori Y, Suzuki M, Sakurai Y, Sumi T, Iwai S, Nakazawa M, Murata I, Miyamaru H, Ono K: **Effectiveness of boron neutron capture therapy for recurrent head and neck malignancies.** *Appl Radiat Isot* 2009, **67**:S37-S42.
12. Barth RF, Kaur B: **Rat brain tumor models in experimental neuro-oncology: the C6, 9L, T9, RG2, F98, BT4C, RT-2 and CNS-1 gliomas.** *J Neurooncol* 2009, **94**:299-312.
13. Kawabata S, Miyatake S, Nonoguchi N, Hiramatsu R, Iida K, Miyata S, Yokoyama K, Doi A, Kuroda Y, Kuroiwa T, Michiue H, Kumada H, Kirihata M, Imahori Y, Maruhashi A, Sakurai Y, Suzuki M, Masunaga S, Ono K: **Survival benefit from boron neutron capture therapy for the newly diagnosed glioblastoma patients.** *Appl Radiat Isot* 2009, **67**:S15-S18.
14. Miyatake S, Kawabata S, Yokoyama K, Kuroiwa T, Michiue H, Sakurai Y, Kumada H, Suzuki M, Maruhashi A, Kirihata M, Ono K: **Survival benefit of Boron neutron capture therapy for recurrent malignant gliomas.** *J Neurooncol* 2009, **91**:199-206.
15. Gonzalez SJ, Bonomi MR, Santa CG, Blaumann HR, Calzetta LO, Menendez P, Jimenez RR, Longhino J, Feld DB, Dagrosa MA, Argerich C, Castiglia SG, Batistoni DA, Liberman SJ, Roth BM: **First BNCT treatment of a skin melanoma in Argentina: dosimetric analysis and clinical outcome.** *Appl Radiat Isot* 2004, **61**:1101-1105.
16. Yoshida F, Matsumura A, Shibata Y, Yamamoto T, Nakauchi H, Okumura M, Nose T: **Cell cycle dependence of boron uptake from two boron compounds used for clinical neutron capture therapy.** *Cancer Lett* 2002, **187**:135-141.
17. Masunaga S, Ono K, Takahashi A, Sakurai Y, Ohnishi K, Kobayashi T, Kinashi Y, Takagaki M, Ohnishi T: **Impact of the p53 status of the tumor cells on the effect of reactor neutron beam irradiation, with emphasis on the response of intratumor quiescent cells.** *Jpn J Cancer Res* 2002, **93**:1366-1377.
18. Masunaga S, Ono K, Sakurai Y, Takagaki M, Kobayashi T, Kinashi Y, Suzuki M: **Evaluation of apoptosis and micronucleation induced by reactor neutron beams with two different cadmium ratios in total and quiescent cell populations within solid tumors.** *Int J Radiat Oncol Biol Phys* 2001, **51**:828-839.
19. Busse D, Doughty RS, Ramsey TT, Russell WE, Price JO, Flanagan WM, Shawver LK, Arteaga CL: **Reversible G(1) arrest induced by inhibition of the epidermal growth factor receptor tyrosine kinase requires up-regulation of p27(KIP1) independent of MAPK activity.** *J Biol Chem* 2000, **275**:6987-6995.
20. Barth RF, Soloway AH, Goodman JH, Gahbauer RA, Gupta N, Blue TE, Yang W, Tjarks W: **Boron neutron capture therapy of brain tumors: an emerging therapeutic modality.** *Neurosurgery* 1999, **44**:433-450, 450-451.
21. Verheij M, Bartelink H: **Radiation-induced apoptosis.** *Cell Tissue Res* 2000, **301**:133-142.
22. Pathak R, Dey SK, Sarma A, Khuda-Bukhsh AR: **Genotoxic effects in M5 cells and Chinese hamster V79 cells after exposure to 7Li-beam (LET = 60 keV/microm) and correlation of their survival dynamics to nuclear damages and cell death.** *Mutat Res* 2007, **628**:56-66.
23. Lee HJ, Kim JS, Moon C, Kim JC, Jo SK, Kim SH: **Relative biological effectiveness of fast neutrons in a multiorgan assay for apoptosis in mouse.** *Environ Toxicol* 2008, **23**:233-239.
24. Dahlstrom M, Capala J, Lindstrom P, Wasteson A, Lindstrom A: **Accumulation of boron in human malignant glioma cells in vitro is cell type dependent.** *J Neurooncol* 2004, **68**:199-205.
25. Fujita Y, Kato I, Iwai S, Ono K, Suzuki M, Sakurai Y, Ohnishi K, Ohnishi T, Yura Y: **Role of p53 mutation in the effect of boron neutron capture therapy on oral squamous cell carcinoma.** *Radiat Oncol* 2009, **4**:63.
26. Aromando RF, Heber EM, Trivillin VA, Nigg DW, Schwint AE, Itoiz ME: **Insight into the mechanisms underlying tumor response to boron neutron capture therapy in the hamster cheek pouch oral cancer model.** *J Oral Pathol Med* 2009, **38**:448-454.
27. Kamida A, Obayashi S, Kato I, Ono K, Suzuki M, Nagata K, Sakurai Y, Yura Y: **Effects of boron neutron capture therapy on human oral squamous cell carcinoma in a nude mouse model.** *Int J Radiat Biol* 2006, **82**:21-29.
28. Kang MH, Reynolds CP: **Bcl-2 inhibitors: targeting mitochondrial apoptotic pathways in cancer therapy.** *Clin Cancer Res* 2009, **15**:1126-1132.
29. Oltersdorf T, Elmore SW, Shoemaker AR, Armstrong RC, Augeri DJ, Belli BA, Bruncko M, Deckwerth TL, Dinges J, Hajduk PJ, Joseph MK, Kitada S, Korsmeyer SJ, Kunzer AR, Letai A, Li C, Mitten MJ, Nettesheim DG, Ng S, Nimmer PM, O'Connor JM, Oleksijew A, Petros AM, Reed JC, Shen W, Tahir SK, Thompson CB, Tomaselli KJ, Wang B, Wendt MD, Zhang H, Fesik SW, Rosenberg SH: **An inhibitor of Bcl-2 family proteins induces regression of solid tumours.** *Nature* 2005, **435**:677-681.
30. Hotchkiss RS, Strasser A, McDunn JE, Swanson PE: **Cell death.** *N Engl J Med* 2009, **361**:1570-1583.
31. Zerp SF, Stoter R, Kuipers G, Yang D, Lippman ME, van Blitterswijk WJ, Bartelink H, Rooswinkel R, Lafleur V, Verheij M: **AT-101, a small molecule inhibitor of anti-apoptotic Bcl-2 family members, activates the SAPK/JNK pathway and enhances radiation-induced apoptosis.** *Radiat Oncol* 2009, **4**:47.
32. Ciardiello F, Tortora G: **Inhibition of bcl-2 as cancer therapy.** *Ann Oncol* 2002, **13**:501-502.
33. Kim KW, Moretti L, Mitchell LR, Jung DK, Lu B: **Combined Bcl-2/mammalian target of rapamycin inhibition leads to enhanced radiosensitization via induction of apoptosis and autophagy in non-small cell lung tumor xenograft model.** *Clin Cancer Res* 2009, **15**:6096-6105.
34. Snyder CM, Shroff EH, Liu J, Chandel NS: **Nitric oxide induces cell death by regulating anti-apoptotic BCL-2 family members.** *PLoS One* 2009, **4**:e7059.
35. Capuani S, Gili T, Bozzali M, Russo S, Porcari P, Cametti C, Muolo M, D'Amore E, Maraviglia B, Lazzarino G, Pastore FS: **Boronophenylalanine uptake in C6 glioma model is dramatically increased by L-DOPA preloading.** *Appl Radiat Isot* 2009, **67**:S34-S36.
36. Barth RF, Yang W, Wu G, Swindall M, Byun Y, Narayanasamy S, Tjarks W, Tordoff K, Moeschberger ML, Eriksson S, Binns PJ, Riley KJ: **Thymidine**

kinase 1 as a molecular target for boron neutron capture therapy of brain tumors. *Proc Natl Acad Sci USA* 2008, **105**:17493-17497.

37. Coderre JA, Gavin PR, Capala J, Ma R, Morris GM, Button TM, Aziz T, Peress NS: Tolerance of the normal canine brain to epithermal neutron irradiation in the presence of p-boronophenylalanine. *J Neurooncol* 2000, **48**:27-40.

Pre-publication history

The pre-publication history for this paper can be accessed here:
<http://www.biomedcentral.com/1471-2407/10/661/prepub>

doi:10.1186/1471-2407-10-661

Cite this article as: Wang *et al.*: Boron neutron capture therapy induces apoptosis of glioma cells through Bcl-2/Bax. *BMC Cancer* 2010 **10**:661.

**Submit your next manuscript to BioMed Central
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

