

4-[2-(2,2-Dimethyl-4,6-dioxo-1,3-dioxan-5-ylidene)hydrazin-1-yl]benzotrile

Naki Çolak,^a Yılmaz Yıldırım,^b Barış Tercan,^c Emel Ermiş^d and Tuncer Hökelek^{e*}

^aDepartment of Chemistry, Hitit University, 19030 Ulukavak, Çorum, Turkey,

^bDepartment of Chemistry, Gazi University, 06500 Teknikokullar, Ankara, Turkey,

^cDepartment of Physics, Karabük University, 78050 Karabük, Turkey, ^dDepartment of Chemistry, Faculty of Science, Anadolu University, 26470 Yenibağlar, Eskişehir, Turkey, and ^eDepartment of Physics, Hacettepe University, 06800 Beytepe, Ankara, Turkey

Correspondence e-mail: merzifon@hacettepe.edu.tr


Received 17 June 2010; accepted 21 June 2010

Key indicators: single-crystal X-ray study; $T = 100$ K; mean $\sigma(\text{C}-\text{C}) = 0.002$ Å; R factor = 0.038; wR factor = 0.108; data-to-parameter ratio = 13.8.

In the title compound, $\text{C}_{13}\text{H}_{11}\text{N}_3\text{O}_4$, the dioxane ring adopts an envelope conformation with the C atom bonded to the dimethyl group in the flap position [deviation = 0.613 (1) Å]. The nitrile group and the attached benzene ring are roughly coplanar [maximum deviation = 0.087 (1) Å]. An intramolecular $\text{N}-\text{H}\cdots\text{O}$ hydrogen bond involving the hydrazinyl group generates an $S(6)$ ring. The $\text{N}-\text{N}$ and $\text{C}-\text{N}$ bond lengths indicate that the compound may be a mixture of the azo and hydrazone tautomeric forms but the presence of the N-bound H atom supports the hydrazone form. The crystal structure is stabilized by weak intermolecular $\text{C}-\text{H}\cdots\text{O}$, $\text{C}-\text{H}\cdots\text{N}$ and $\text{C}-\text{H}\cdots\pi$ interactions.

Related literature

For the applications of related azo compounds, see: Branger *et al.* (1997); Buchel *et al.* (1995); Gale *et al.* (1998); Ikeda & Tsutsumi (1995); Kang *et al.* (2000); Karıcı *et al.* (2004); Kim *et al.* (1995); Kobrakov *et al.* (2004); Natansohn *et al.* (1992); Rochon *et al.* (1995). For related hydrazone structures, see: Çolak *et al.* (2010); Pavlovic *et al.* (2009); Seferoğlu *et al.* (2008); Seferoğlu *et al.* (2009); Wojciechowski & Szymezak (2007).



Experimental

Crystal data

$\text{C}_{13}\text{H}_{11}\text{N}_3\text{O}_4$
 $M_r = 273.25$
 Monoclinic, $P2_1/n$
 $a = 9.7617$ (2) Å
 $b = 11.0023$ (2) Å
 $c = 11.4753$ (3) Å
 $\beta = 93.796$ (1)°
 $V = 1229.76$ (5) Å³
 $Z = 4$
 Mo $K\alpha$ radiation
 $\mu = 0.11$ mm⁻¹
 $T = 100$ K
 $0.46 \times 0.43 \times 0.29$ mm

Data collection

Bruker Kappa APEXII CCD area-detector diffractometer
 Absorption correction: multi-scan (SADABS; Bruker, 2005)
 $T_{\min} = 0.950$, $T_{\max} = 0.968$
 11746 measured reflections
 3098 independent reflections
 2591 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.029$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.038$
 $wR(F^2) = 0.108$
 $S = 1.04$
 3098 reflections
 225 parameters
 All H-atom parameters refined
 $\Delta\rho_{\text{max}} = 0.38$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.24$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

Cg1 is the centroid of the C5–C10 ring.

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{N}2-\text{H}2\cdots\text{O}3$	0.92 (2)	1.958 (16)	2.6674 (13)	132 (1)
$\text{C}9-\text{H}9\cdots\text{N}1^i$	0.94 (2)	2.624 (15)	3.5320 (16)	164 (1)
$\text{C}10-\text{H}10\cdots\text{N}3^{ii}$	0.99 (2)	2.485 (15)	3.3876 (16)	152 (1)
$\text{C}12-\text{H}12\cdots\text{O}2^{iii}$	0.97 (2)	2.527 (17)	3.4454 (15)	159 (1)
$\text{C}12-\text{H}12\cdots\text{Cg}1^{iv}$	0.98 (2)	2.491 (15)	3.4575 (13)	171 (1)

Symmetry codes: (i) $-x + \frac{3}{2}, y - \frac{1}{2}, -z + \frac{3}{2}$; (ii) $x - \frac{1}{2}, -y + \frac{1}{2}, z + \frac{1}{2}$; (iii) $-x, -y + 1, -z + 2$; (iv) $-x + \frac{3}{2}, y - \frac{1}{2}, -z + \frac{1}{2}$.

Data collection: APEX2 (Bruker, 2007); cell refinement: SAINT (Bruker, 2007); data reduction: SAINT program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: ORTEP-3 for Windows (Farrugia, 1997); software used to prepare material for publication: WinGX (Farrugia, 1999) and PLATON (Spek, 2009).

The authors are indebted to Anadolu University and the Medicinal Plants and Medicine Research Centre of Anadolu University, Eskişehir, Turkey, for the use of X-ray diffractometer.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: CI5104).

References

- Branger, C., Lequan, M., Lequan, R. M., Large, M. & Kajzar, F. (1997). *Chem. Phys. Lett.* **272**, 265–270.
 Bruker (2005). SADABS. Bruker AXS Inc. Madison, Wisconsin, USA.
 Bruker (2007). APEX2 and SAINT. Bruker AXS Inc. Madison, Wisconsin, USA.
 Buchel, M., Sekkat, Z., Paul, S., Weichart, B., Menzel, H. & Knoll, W. (1995). *Langmuir*, **11**, 4460–4466.
 Çolak, N., Aksakal, D., Andaç, Ö. & Büyükgüngör, O. (2010). *Acta Cryst.* **E66**, o1165–o1166.
 Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.

- Farrugia, L. J. (1999). *J. Appl. Cryst.* **32**, 837–838.
- Gale, P. A., Chen, Z., Drew, M. G. B., Heath, J. A. & Beer, P. D. (1998). *Polyhedron*, **4**, 405–412.
- Ikeda, T. & Tsutsumi, O. (1995). *Science*, **268**, 1873–1875.
- Kang, H. C., Lee, B. M., Yoon, J. & Yoon, M. (2000). *J. Colloid Interface Sci.* **231**, 255–264.
- Karçı, F., Şener, İ. & Deligöz, H. (2004). *Dyes Pigments*, **62**, 131–140.
- Kim, D. Y., Tripathy, S. K., Li, L. & Kumar, J. (1995). *Appl. Phys. Lett.* **66**, 1166–1168.
- Kobrakov, K. I., Glyadyaeva, O. Yu., Stankevich, G. S. & Kovtun, L. G. (2004). *Fibre Chem.* **36**, 41–42.
- Natansohn, A., Rochon, P., Gosselin, J. & Xie, S. (1992). *Macromolecules*, **25**, 2268–2273.
- Pavlovic, G., Racane, L., Cicak, H. & Tralic-Kulenovic, V. (2009). *Dyes Pigments*, **83**, 354–362.
- Rochon, P., Battalla, E. & Natansohn, A. (1995). *Appl. Phys. Lett.* **66**, 136–138.
- Seferoğlu, Z., Ertan, N., Hökelek, T. & Şahin, E. (2008). *Dyes Pigments*, **77**, 614–625.
- Seferoğlu, Z., Ertan, N., Kickelbick, G. & Hökelek, T. (2009). *Dyes Pigments*, **82**, 20–25.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Spek, A. L. (2009). *Acta Cryst.* **D65**, 148–155.
- Wojciechowski, K. & Szymezak, A. (2007). *Dyes Pigments*, **75**, 45–51.

supplementary materials

Acta Cryst. (2010). E66, o1784-o1785 [doi:10.1107/S1600536810024025]

4-[2-(2,2-Dimethyl-4,6-dioxo-1,3-dioxan-5-ylidene)hydrazin-1-yl]benzotrile

N. Çolak, Y. Yildirim, B. Tercan, E. Ermis and T. Hökelek

Comment

It has been known for many years that the azo compounds are widely used class of dyes due to their applications in various fields such as the dyeing of textile fibers, the coloring of different materials, colored plastics and electrochemical sensors (Kobrakov *et al.*, 2004; Karcı *et al.*, 2004; Gale *et al.*, 1998). Azo dyes have been attracting intensive interest for their potential use in optical data storage (Natansohn *et al.*, 1992), optical switching (Ikeda & Tsutsumi, 1995), polarization holography (Kim *et al.*, 1995; Rochon *et al.*, 1995), optical modulation (Buchel *et al.*, 1995), nonlinear optics (Branger *et al.*, 1997) and photolabile surfactants (Kang *et al.*, 2000).

The dyes may exist in two possible tautomeric forms, namely azo form and hydrazone form. The azo-hydrazone tautomerism is quite interesting from a theoretical and practical point of view because the two tautomers have different properties. Azo dyes are known to exist in the azo-hydrazone tautomeric forms (Çolak *et al.*, 2010; Pavlovic *et al.*, 2009; Seferoğlu *et al.*, 2009; Seferoğlu *et al.*, 2008; Wojciechowski & Szymezak, 2007). We herein report the crystal structure of the title compound, (I).

The title compound, (I), contains benzonitrile and 2,2-dimethyl-1,3-dioxane-4,6-dione groups connected *via* a hydrazinyl group (Fig. 1). In (I), N1—N2 [1.3082 (14) Å] bond length is between the N=N double bond [1.20–1.28 Å for a real azo compound] and N—N single bond [longer than 1.4 Å] lengths. The N1—C3 and N2—C5 bond lengths are 1.3116 (14) and 1.4075 (14) Å, respectively. The carbonyl O atoms O3 and O4 slightly deviate from the N1/C2—C4 plane by 0.255 (2) and 0.259 (2) Å, respectively. So, the title compound may exist both in azo and hydrazone tautomeric forms, and is mainly in the hydrazone tautomeric form. The C8—C13 [1.4418 (15) Å] bond length is longer for a C(sp²)—C(sp¹) bond, but in agreement with the previously reported value [1.442 (3) Å; Çolak *et al.*, 2010].

An intramolecular N2—H2···O3 hydrogen bond (Table 1) results in the formation of a nearly planar [with a maximum deviation of 0.056 (1) Å for atom C4] six-membered ring C (O3/N1/N2/H2/C3/C4), which is oriented with respect to the benzonitrile ring B (C5—C10) at a dihedral angle of 7.8 (43)°. Atoms N1, N2, N3 and C13 are displaced by -0.162 (2), 0.038 (2), 0.049 (2) and 0.028 (2) Å from the plane of ring B, respectively. The benzonitrile and hydrazinyl groups (4-hydrazinylbenzonitrile) are essentially coplanar [with a maximum deviation of -0.087 (1) Å for atom N1]. The dioxane ring A (O1/O2/C1—C4) is not planar having envelope conformation with atom C1 displaced by 0.613 (1) Å from the plane of the other ring atoms.

In the crystal structure, weak C—H···O and C—H···N hydrogen bonds (Table 1) may be effective in the stabilization of the crystal packing. There also exists a weak C—H···π interaction (Table 1).

Experimental

A hydrochloric acid solution (2.5 ml) of 4-aminobenzonitrile (1.18 g, 10 mmol) and an aqueous solution (10 ml) of sodium nitrite (0.69 g, 10 mmol) were mixed and stirred at 273 K for 1 h. To this solution, an ethanol solution (10 ml) of the coupling

supplementary materials

component 2,2-dimethyl-1,3-dioxane-4,6-dione (1.44 g, 10 mmol) was added and the stirring was continued at 273 K for 4 h. The resulting product was filtered and washed with water, dried and crystallized from ethanol (yield 1.88 g, 69%; m.p. 440–442 K).

Refinement

H atoms were located in difference Fourier maps and refined isotropically.

Figures




Fig. 1. The molecular structure of the title compound, with the atom-numbering scheme. Displacement ellipsoids are drawn at the 50% probability level. The dashed line indicates a hydrogen bond.

4-[2-(2,2-Dimethyl-4,6-dioxo-1,3-dioxan-5-ylidene)hydrazin-1-yl]benzonitrile

Crystal data

$C_{13}H_{11}N_3O_4$

$M_r = 273.25$

Monoclinic, $P2_1/n$

Hall symbol: -P 2yn

$a = 9.7617(2) \text{ \AA}$

$b = 11.0023(2) \text{ \AA}$

$c = 11.4753(3) \text{ \AA}$

$\beta = 93.796(1)^\circ$

$V = 1229.76(5) \text{ \AA}^3$

$Z = 4$

$F(000) = 568$

$D_x = 1.476 \text{ Mg m}^{-3}$

Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$

Cell parameters from 4103 reflections

$\theta = 2.6\text{--}28.4^\circ$

$\mu = 0.11 \text{ mm}^{-1}$

$T = 100 \text{ K}$

Block, yellow

$0.46 \times 0.43 \times 0.29 \text{ mm}$

Data collection

Bruker Kappa APEXII CCD area-detector diffractometer

Radiation source: fine-focus sealed tube graphite

φ and ω scans

Absorption correction: multi-scan (*SADABS*; Bruker, 2005)

$T_{\min} = 0.950$, $T_{\max} = 0.968$

11746 measured reflections

3098 independent reflections

2591 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.029$

$\theta_{\max} = 28.5^\circ$, $\theta_{\min} = 2.6^\circ$

$h = -13 \rightarrow 13$

$k = -14 \rightarrow 14$

$l = -15 \rightarrow 8$

Refinement

Refinement on F^2	Primary atom site location: structure-invariant direct methods
Least-squares matrix: full	Secondary atom site location: difference Fourier map
$R[F^2 > 2\sigma(F^2)] = 0.038$	Hydrogen site location: inferred from neighbouring sites
$wR(F^2) = 0.108$	All H-atom parameters refined
$S = 1.04$	$w = 1/[\sigma^2(F_o^2) + (0.0626P)^2 + 0.2729P]$
3098 reflections	where $P = (F_o^2 + 2F_c^2)/3$
225 parameters	$(\Delta/\sigma)_{\max} = 0.001$
0 restraints	$\Delta\rho_{\max} = 0.38 \text{ e } \text{\AA}^{-3}$
	$\Delta\rho_{\min} = -0.24 \text{ e } \text{\AA}^{-3}$

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
O1	0.16363 (8)	0.74524 (7)	0.85585 (7)	0.0172 (2)
O2	0.17892 (8)	0.56739 (7)	0.97179 (7)	0.01470 (19)
O3	0.35656 (9)	0.44276 (8)	0.96914 (7)	0.0187 (2)
O4	0.32521 (9)	0.79574 (8)	0.73794 (8)	0.0249 (2)
N1	0.44403 (10)	0.56888 (9)	0.76510 (9)	0.0165 (2)
N2	0.51381 (10)	0.46953 (9)	0.78943 (9)	0.0159 (2)
H2	0.4942 (16)	0.4243 (15)	0.8538 (15)	0.028 (4)*
N3	1.03756 (11)	0.29578 (10)	0.47015 (10)	0.0232 (2)
C1	0.08704 (11)	0.64589 (10)	0.90176 (10)	0.0143 (2)
C2	0.27845 (12)	0.71950 (11)	0.79881 (10)	0.0172 (2)
C3	0.34022 (12)	0.59901 (10)	0.82608 (10)	0.0155 (2)
C4	0.29477 (11)	0.52812 (10)	0.92521 (10)	0.0144 (2)
C5	0.62374 (11)	0.43844 (10)	0.72180 (10)	0.0148 (2)
C6	0.64458 (13)	0.49922 (11)	0.61746 (10)	0.0173 (2)
H6	0.5852 (15)	0.5636 (14)	0.5904 (13)	0.020 (4)*
C7	0.75216 (13)	0.46401 (11)	0.55177 (10)	0.0175 (2)
H7	0.7668 (15)	0.5059 (14)	0.4788 (13)	0.024 (4)*
C8	0.83788 (12)	0.36875 (11)	0.59035 (10)	0.0155 (2)

supplementary materials

C9	0.81720 (12)	0.30898 (11)	0.69557 (10)	0.0161 (2)
H9	0.8752 (14)	0.2457 (14)	0.7224 (12)	0.020 (4)*
C10	0.70992 (12)	0.34407 (11)	0.76131 (10)	0.0159 (2)
H10	0.6926 (14)	0.3023 (14)	0.8350 (13)	0.021 (4)*
C11	0.01442 (13)	0.57371 (11)	0.80405 (10)	0.0178 (2)
H111	0.0795 (15)	0.5338 (14)	0.7556 (13)	0.022 (4)*
H112	-0.0444 (15)	0.5101 (14)	0.8393 (13)	0.023 (4)*
H113	-0.0456 (15)	0.6270 (15)	0.7544 (14)	0.026 (4)*
C12	-0.00717 (13)	0.70098 (11)	0.98606 (10)	0.0169 (2)
H121	0.0494 (16)	0.7364 (15)	1.0534 (13)	0.026 (4)*
H122	-0.0616 (15)	0.7636 (14)	0.9446 (13)	0.021 (4)*
H123	-0.0673 (16)	0.6388 (16)	1.0133 (15)	0.035 (4)*
C13	0.94875 (12)	0.32918 (11)	0.52248 (10)	0.0177 (2)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
O1	0.0168 (4)	0.0111 (4)	0.0244 (4)	0.0003 (3)	0.0057 (3)	0.0024 (3)
O2	0.0132 (4)	0.0139 (4)	0.0169 (4)	0.0027 (3)	0.0012 (3)	0.0015 (3)
O3	0.0164 (4)	0.0178 (4)	0.0220 (4)	0.0033 (3)	0.0011 (3)	0.0027 (3)
O4	0.0244 (5)	0.0191 (5)	0.0321 (5)	0.0004 (4)	0.0096 (4)	0.0083 (4)
N1	0.0144 (5)	0.0146 (5)	0.0204 (5)	-0.0002 (4)	0.0009 (4)	-0.0023 (4)
N2	0.0151 (5)	0.0149 (5)	0.0179 (5)	0.0004 (4)	0.0031 (4)	-0.0005 (4)
N3	0.0227 (6)	0.0233 (6)	0.0244 (5)	0.0003 (4)	0.0076 (4)	0.0001 (4)
C1	0.0135 (5)	0.0111 (5)	0.0182 (5)	0.0005 (4)	0.0010 (4)	0.0024 (4)
C2	0.0152 (6)	0.0159 (6)	0.0208 (5)	-0.0001 (4)	0.0025 (4)	-0.0003 (4)
C3	0.0143 (5)	0.0140 (5)	0.0184 (5)	-0.0005 (4)	0.0017 (4)	-0.0006 (4)
C4	0.0127 (5)	0.0135 (5)	0.0170 (5)	-0.0010 (4)	0.0008 (4)	-0.0027 (4)
C5	0.0119 (5)	0.0143 (5)	0.0182 (5)	-0.0024 (4)	0.0014 (4)	-0.0037 (4)
C6	0.0178 (6)	0.0137 (5)	0.0205 (5)	-0.0001 (4)	0.0014 (4)	0.0001 (4)
C7	0.0196 (6)	0.0155 (5)	0.0178 (5)	-0.0026 (4)	0.0039 (4)	0.0004 (4)
C8	0.0134 (5)	0.0155 (5)	0.0179 (5)	-0.0031 (4)	0.0028 (4)	-0.0022 (4)
C9	0.0137 (5)	0.0160 (5)	0.0185 (5)	0.0003 (4)	0.0001 (4)	-0.0002 (4)
C10	0.0144 (5)	0.0177 (5)	0.0155 (5)	-0.0026 (4)	0.0010 (4)	-0.0004 (4)
C11	0.0174 (6)	0.0170 (6)	0.0188 (5)	0.0019 (5)	-0.0009 (4)	-0.0015 (4)
C12	0.0162 (6)	0.0141 (5)	0.0207 (5)	0.0032 (4)	0.0037 (4)	0.0007 (4)
C13	0.0178 (6)	0.0167 (6)	0.0188 (5)	-0.0029 (4)	0.0024 (4)	0.0006 (4)

Geometric parameters (\AA , $^\circ$)

O1—C1	1.4435 (13)	C6—C7	1.3878 (16)
O1—C2	1.3645 (13)	C6—H6	0.954 (15)
O2—C1	1.4493 (13)	C7—C8	1.3949 (17)
O2—C4	1.3533 (13)	C7—H7	0.974 (15)
O3—C4	1.2081 (14)	C8—C9	1.4012 (16)
O4—C2	1.2009 (14)	C8—C13	1.4418 (15)
N1—N2	1.3082 (14)	C9—C10	1.3854 (15)
N1—C3	1.3116 (14)	C9—H9	0.936 (15)
N2—C5	1.4075 (14)	C10—H10	0.986 (15)

N2—H2	0.922 (17)	C11—C1	1.5117 (16)
N3—C13	1.1473 (15)	C11—H111	0.976 (15)
C1—C12	1.5055 (15)	C11—H113	0.984 (16)
C2—C3	1.4812 (16)	C11—H112	1.007 (16)
C3—C4	1.4718 (15)	C12—H122	0.975 (15)
C5—C6	1.3981 (16)	C12—H121	0.999 (16)
C5—C10	1.3932 (17)	C12—H123	0.967 (17)
C2—O1—C1	118.66 (9)	C7—C6—H6	119.7 (9)
C4—O2—C1	118.29 (8)	C6—C7—C8	119.80 (11)
N2—N1—C3	120.47 (10)	C6—C7—H7	119.6 (9)
N1—N2—C5	119.38 (10)	C8—C7—H7	120.6 (9)
N1—N2—H2	119.3 (10)	C7—C8—C9	120.56 (10)
C5—N2—H2	121.1 (10)	C7—C8—C13	120.76 (10)
O1—C1—O2	109.71 (9)	C9—C8—C13	118.68 (11)
O1—C1—C11	110.85 (9)	C8—C9—H9	120.9 (9)
O1—C1—C12	106.43 (9)	C10—C9—C8	119.72 (11)
O2—C1—C11	109.92 (9)	C10—C9—H9	119.4 (9)
O2—C1—C12	105.40 (9)	C5—C10—H10	119.5 (9)
C12—C1—C11	114.31 (10)	C9—C10—C5	119.50 (10)
O1—C2—C3	114.85 (10)	C9—C10—H10	121.0 (9)
O4—C2—O1	119.33 (11)	C1—C11—H111	111.6 (9)
O4—C2—C3	125.66 (11)	C1—C11—H113	110.4 (9)
N1—C3—C2	115.57 (10)	C1—C11—H112	108.6 (9)
N1—C3—C4	124.16 (11)	H111—C11—H112	109.0 (12)
C4—C3—C2	119.88 (10)	H111—C11—H113	108.9 (13)
O2—C4—C3	116.02 (10)	H113—C11—H112	108.3 (12)
O3—C4—O2	119.38 (10)	C1—C12—H121	108.9 (9)
O3—C4—C3	124.52 (10)	C1—C12—H122	107.9 (8)
C6—C5—N2	121.03 (11)	C1—C12—H123	109.4 (10)
C10—C5—N2	117.89 (10)	H121—C12—H123	110.0 (13)
C10—C5—C6	121.08 (10)	H122—C12—H121	110.9 (13)
C5—C6—H6	120.9 (9)	H122—C12—H123	109.6 (13)
C7—C6—C5	119.34 (11)	N3—C13—C8	178.56 (13)
C2—O1—C1—O2	51.23 (12)	O4—C2—C3—N1	-9.81 (19)
C2—O1—C1—C11	-70.35 (13)	O4—C2—C3—C4	163.35 (12)
C2—O1—C1—C12	164.79 (10)	N1—C3—C4—O2	-174.73 (10)
C1—O1—C2—O4	163.20 (11)	N1—C3—C4—O3	8.59 (19)
C1—O1—C2—C3	-21.07 (14)	C2—C3—C4—O2	12.73 (16)
C4—O2—C1—O1	-50.36 (12)	C2—C3—C4—O3	-163.95 (11)
C4—O2—C1—C12	-164.59 (9)	N2—C5—C6—C7	-178.47 (10)
C4—O2—C1—C11	71.77 (12)	C10—C5—C6—C7	0.62 (18)
C1—O2—C4—O3	-163.46 (10)	N2—C5—C10—C9	178.44 (10)
C1—O2—C4—C3	19.68 (14)	C6—C5—C10—C9	-0.67 (18)
C3—N1—N2—C5	179.24 (10)	C5—C6—C7—C8	0.05 (18)
N2—N1—C3—C2	173.80 (10)	C6—C7—C8—C9	-0.65 (18)
N2—N1—C3—C4	0.97 (18)	C6—C7—C8—C13	178.87 (11)
N1—N2—C5—C6	-11.37 (17)	C7—C8—C9—C10	0.60 (18)
N1—N2—C5—C10	169.52 (10)	C13—C8—C9—C10	-178.93 (11)

supplementary materials

O1—C2—C3—N1	174.77 (10)	C8—C9—C10—C5	0.06 (17)
O1—C2—C3—C4	-12.07 (16)		

Hydrogen-bond geometry (Å, °)

Cg1 is the centroid of the C5–C10 ring.

<i>D</i> —H \cdots <i>A</i>	<i>D</i> —H	H \cdots <i>A</i>	<i>D</i> \cdots <i>A</i>	<i>D</i> —H \cdots <i>A</i>
N2—H2 \cdots O3	0.92 (2)	1.958 (16)	2.6674 (13)	132 (1)
C9—H9 \cdots N1 ⁱ	0.94 (2)	2.624 (15)	3.5320 (16)	164 (1)
C10—H10 \cdots N3 ⁱⁱ	0.99 (2)	2.485 (15)	3.3876 (16)	152 (1)
C12—H123 \cdots O2 ⁱⁱⁱ	0.97 (2)	2.527 (17)	3.4454 (15)	159 (1)
C12—H122 \cdots Cg1 ^{iv}	0.98 (2)	2.491 (15)	3.4575 (13)	171 (1)

Symmetry codes: (i) $-x+3/2, y-1/2, -z+3/2$; (ii) $x-1/2, -y+1/2, z+1/2$; (iii) $-x, -y+1, -z+2$; (iv) $-x+3/2, y-1/2, -z+1/2$.

Fig. 1

