



## A survey of scale insects (Sternorrhyncha: Coccoidea) occurring on table grapes in South Africa

Vaughn M. Walton<sup>1,a</sup>, Kerstin Krüger<sup>2,b</sup>, Davina L. Saccaggi<sup>3,c,\*</sup> and Ian M. Millar<sup>3,d</sup>

<sup>1</sup> 4079 Ag and Life Sciences, Building Oregon State University, Corvallis, OR 97331-7304, U.S.A

<sup>2</sup> Department of Zoology & Entomology, University of Pretoria, Pretoria, 0002 South Africa

<sup>3</sup> ARC-Plant Protection Research Institute, Private Bag X134, Queenswood, Pretoria, 0121 South Africa

### Abstract

Increasing international trade and tourism have led to an increase in the introduction of exotic pests that pose a considerable economic threat to the agro-ecosystems of importing countries. Scale insects (Sternorrhyncha: Coccoidea) may be contaminants of export consignments from the South African deciduous fruit industry to the European Union, Israel, United Kingdom and the United States, for example. Infestations of immature scale insects found on South African fruit destined for export have resulted in increasing rates of rejection of such consignments. To identify the risk posed by scale insect species listed as phytosanitary pests on table grapes to the abovementioned importing countries, a field survey was undertaken in 2004–2005 in vineyards throughout all grape-producing regions in South Africa. Coccoidea species found during the current field survey were *Planococcus ficus* (Signoret), *Pseudococcus longispinus* (Targioni Tozzetti), *Coccus hesperidum* L. and *Nipaecoccus viridis* (Newstead). With the exception of *Pl. ficus*, which has only been collected from *Vitis vinifera* (Vitaceae) and *Ficus carica* (Moraceae) in South Africa, these species are polyphagous and have a wide host range. None of the scale insect species found to occur in vineyards in South Africa pose a phytosanitary risk to countries where fruit are exported except for *Ferrisia malvastra* (McDaniel) and *N. viridis* that have not been recorded in the USA. All scale insects previously found in vineyards in South Africa are listed and their phytosanitary status discussed. The results of the survey show that the risk of exporting scale insect pests of phytosanitary importance on table grapes from South Africa is limited.

**Keywords:** mealybugs, soft scale insects, Pseudococcidae, distribution, quarantine, post harvest pests

**Correspondence:** <sup>a</sup>waltonv@hort.oregonstate.edu, <sup>b</sup>kruger@zoology.up.ac.za, <sup>c</sup>\*DavinaM@nda.agric.za, <sup>d</sup>MillarI@arc.agric.za

**Associate Editor:** J.P. Michaud was editor of this paper

\*Current address: National Department of Agriculture, Plant Health Diagnostic Services, Private Bag X 5015, Stellenbosch, 7599 South Africa

**Received:** 28 November 2007 | **Accepted:** 23 May 2008 | **Published:** 26 June 2009

**Copyright:** This is an open access paper. We use the Creative Commons Attribution 3.0 license that permits unrestricted use, provided that the paper is properly attributed.

**ISSN:** 1536-2442 | Vol. 9, Number 47

#### Cite this paper as:

Walton VM, Krüger K, Saccaggi DL, Millar IM. 2009. A survey of scale insects (Sternorrhyncha: Coccoidea) occurring on table grapes in South Africa. 6pp. *Journal of Insect Science* 9:47, available online: [insectscience.org/9.47](http://insectscience.org/9.47)

## Introduction

The South African table grape export industry is situated in mild Mediterranean and arid subtropical climates. Several scale insect (Sternorrhyncha: Coccoidea) species have successfully colonized hosts, including grapevine (*Vitis vinifera* L. (Vitaceae)), in these climates (Ben-Dov 1994). More than 80% of table grape production in South Africa occurs in the Western Cape region. Other production areas include the Northern Cape, Eastern Cape, Limpopo, Free State and Mpumalanga. During 2003, South African table grape exports totaled 198,264 metric tons (Hugh Campbell, Deciduous Fruit Producers Trust (DFPT) research, personal communication). Most table grapes were exported to central Europe (40%), followed by Great Britain (35%), Asia and the Far East (11%), the USA (4%), the Middle East and Mediterranean countries (6%) and other African countries (4%) (Hugh Campbell, personal communication).

Many mealybug (Pseudococcidae) species, such as *Planococcus ficus* (Signoret), the key pest on grapevines in South Africa, are able to complete their entire lifecycle on grapes (Walton and Pringle 2004). The almost year-long warm climate in most grape-growing regions in South Africa leads to overlapping generations of *Pl. ficus*, resulting in multiple life stages being present at any one time. Adult female scales on export grapes can be readily identified using morphological characters and consignments are often released following species identification. However, many consignments are rejected prior to shipment due to infestation with immature scale insects that are difficult to identify using morphological characters. As a result, immature scale insect identification tools have been developed in export-based fruit producing countries. These include both molecular (e.g. Beuning et al. 1999; Demontis et al. 2007; Saccaggi et al. 2008) and morphological (e.g. Gullan 2000; Wakgari and Giliomee 2005) methods. The risk of a quarantine pest occurring in certain production areas can be quantified by undertaking extensive area-wide field surveys. This study reports the results of an industry-wide field survey of scale insects in South African vineyards. In addition, a literature survey of scale insects previously reported on grapevines is presented.

## Materials and Methods

To determine which scale insect species occur on table grapes, field surveys were conducted in different vine-growing regions in South Africa. Field surveys were undertaken from 1 November 2004 to 30 April 2005 as this is the period with the highest prevalence of scale insects in several grape-growing areas in the country (Whitehead 1957; Walton 2003). A portion (10%) of all producers in each grape-growing area in South Africa was randomly selected, and 10% of each production unit (farms with

average sizes of between 30 and 100 hectares) was surveyed. A central systematic sampling system was used in vineyard blocks that ranged between three and ten hectares in size. One sample was collected in each block as follows: field scouts thoroughly inspected leaves, bunches, areas beneath the bark and areas in the root region in each of twenty evenly spaced plots consisting of five vines each. Whenever possible, scouts collected adult females in order to facilitate the identification process. Samples of between three and 50 scale insects from each block were placed into single vials containing 97% ethanol. Each vial was marked with GPS coordinates, production unit name, contact details, vine cultivar and sampling date.

Samples collected during the survey (2004–2005) were identified using molecular and morphological techniques. V. Walton carried out morphological identification of specimens using the Systematic Entomology Laboratory ARS, USDA Scale Insects Identification Tools for Species of Quarantine Significance (Miller et al. 2007). To confirm morphological identifications, mealybugs were also identified using a multiplex polymerase chain reaction (PCR) technique (Saccaggi 2006; Saccaggi et al. 2008) developed to identify the mealybugs *Pl. ficus*, *Planococcus citri* (Risso) and *Pseudococcus longispinus* (Targioni Tozzetti). *Pl. ficus* and *Ps. longispinus* are the most common mealybug species occurring in South African vineyards. *Pl. citri* has been historically associated with *V. vinifera* in South Africa and is common in citrus groves in this country. Mealybugs that could not be identified morphologically, or whose identification was doubtful, were identified by I.M. Millar.

A literature search of scale insect species occurring on grapevines, was performed on the ScaleNet website, [www.sel.barc.usda.gov/SCALENET/scalenet.htm](http://www.sel.barc.usda.gov/SCALENET/scalenet.htm), and used to determine historical records of these insects on vines in South Africa. (Ben-Dov et al. 2006). In addition, a query of ScaleNet was done to determine all scale insects found on *V. vinifera* worldwide, followed by a cross-search in order to determine which of these had been recorded in South Africa.

## Results and Discussion

Scouts collected 249 samples from all major grape-growing regions and 29 municipal districts during the survey (Table 1). Two hundred and twenty nine samples were identified using the morphological method by V. Walton and sub-samples were confirmed using the molecular identification method (128 samples). Mealybugs and scale insects that could not be identified morphologically by V. Walton were submitted for identification to I. M. Millar (20 samples). Of these, eighteen samples were identified as *Pl. ficus* and the remaining two were *Nipaecoccus viridis* (Newstead), collected in the Grootdrink municipal district (28.40 S 21.43 E) on ‘Thompson Seedless’ grapes, and

**Table 1.** Grape-growing regions sampled for Coccidae in six Provinces in South Africa, their total area, approximate number of production units, number of units sampled, total number of samples collected and number of coccids found. Samples were collected between November and April.

| Province            | Grape growing region                   | Total area (ha) | No. of production units | No. units sampled | No. samples/unit | <i>Planococcus ficus</i> | <i>Pseudococcus longispinus</i> | <i>Nipaecoccus viridis</i> | <i>Coccus hesperidum</i> |
|---------------------|--|-----------------|-------------------------|-------------------|------------------|--------------------------|---------------------------------|----------------------------|--------------------------|
| Gauteng             | Gauteng                                | 55              | 5                       | 2                 | 2                | 2                        |                                 |                            |                          |
| Limpopo             | Marble Hall                            | 535             | 25                      | 14                | 16               | 16                       |                                 |                            |                          |
| Mpumalanga          | Mpumalanga                             | 40              | 4                       | 2                 | 3                | 2                        | 1                               |                            |                          |
| North West Province | North West                             | 90              | 5                       | 2                 | 2                | 2                        |                                 |                            |                          |
| Northern Cape       | Lower Orange River                     | 8,730           | 250                     | 87                | 93               | 89                       | 3                               | 1                          | 1                        |
| Northern Cape       | Upper Orange River                     | 180             | 6                       | 2                 | 2                | 2                        |                                 |                            |                          |
| Western Cape        | Berg River (Paarl, Agterpaarl, Hermon) | 3,210           | 100                     | 57                | 57               | 53                       | 4                               |                            |                          |
| Western Cape        | Hex River Valley                       | 5,130           | 200                     | 57                | 61               | 61                       |                                 |                            |                          |
| Western Cape        | Little Karoo                           | 435             | 10                      | 2                 | 2                | 2                        |                                 |                            |                          |
| Western Cape        | Namaqualand                            | 450             | 10                      | 2                 | 3                | 3                        |                                 |                            |                          |
| Western Cape        | Piketberg                              | 1,030           | 30                      | 3                 | 4                | 3                        | 1                               |                            |                          |
| Western Cape        | Stellenbosch                           | 44              | 5                       | 1                 | 2                | 1                        | 1                               |                            |                          |
| Western Cape        | Wolseley                               | 65              | 5                       | 2                 | 2                | 2                        |                                 |                            |                          |
| <b>Total</b>        |  | <b>19,994</b>   | <b>655</b>              | <b>233</b>        | <b>249</b>       | <b>238</b>               | <b>10</b>                       | <b>1</b>                   | <b>1</b>                 |

*Coccus hesperidum* Linnaeus, collected in Groblershoop municipal district (28.53 S 21.59 E) on ‘Thompson Seedless’ grapes. Both species had not been previously recorded on *V. vinifera*, and these are the first known records of these species from these areas.

The majority of species (95%) were *Pl. ficus*, a cosmopolitan mealybug reported from Europe, Israel and the USA among other regions. This species was found in the grape growing areas in South Africa (Table 1). Of the remaining samples, ten were *Ps. longispinus* (4%), one *N. viridis* and one *C. hesperidum* (1%). *Ps. longispinus* was found at six locations in the Western Cape Province: Paarl area (33.42 S 19.10 E, Berg River, 2 samples), Agterpaarl (33.40 S 18.54 E, Berg River, 1 sample), Hermon (33.24 S 18.58 E, Berg River, 1 sample) Stellenbosch (33.54 S, 18.50 E, 1 sample), and Piketberg (32.56 S 18.40 E, 1 sample). Further records are from the Northern Cape: Augrabies (28.40 S 20.26 E, Lower Orange River, 1 sample), Groblershoop (28.53 S 21.59 E, Lower Orange River, 1 sample) and Kanoneiland (33.24 S 18.58 E, Lower Orange River, 1 sample) and Mpumalanga: Groblersdal (25.10 S 29.25 E, 1 sample). These are the first records from these localities.

The online literature search listed 1389 records of scale insect species found in South Africa (Ben Dov et al. 2006), including 281 species from the Pseudococcidae. A total of 101 scale species have been recorded on *V. vinifera* worldwide (Ben-Dov et al. 2006). From this list, we have listed forty-six species recorded from a variety of host plants in South Africa (Table 2) as these species have the highest risk of occurring on grapevines in South Africa. Of these, 11 were recorded on *V. vinifera* in South Africa. The following seven species belonging to the families Coccidae and Pseudococcidae were recorded during the survey: *C. hesperidum*, *Cryptinglisia lounsburyi* Cockerell, *Trijuba oculata* (Brain), *Ferrisia malvastra* (McDaniel), *N. viridis*, *Pl. ficus* and *Ps. longispinus*.

In addition, four giant scale species (Margarodidae), all phytosanitary pests, have been recorded on *V. vinifera* in South Africa (Table 2). *Margarodes capensis* Giard, *Margarodes greeni* Brain, *Margarodes prieskaensis* (Jakubski) and *Margarodes vredendalensis* De Klerk, were studied and described by De Klerk et al. (1983). These species are subterranean and therefore not likely to be found on fruit destined for export. Both *T. oculata* and *C. lounsburyi* are of phytosanitary concern but are not found in any country to which South Africa exports table grapes (Ben-Dov et al. 2006).

**Table 2.** Records of scale insects found on *Vitis vinifera* worldwide, and that occur on grapevines and/or other hosts in South Africa. Currently, eleven species are found on *V. vinifera* in South Africa, and six of these (indicated with an asterisk) are of phytosanitary concern to other parts of the world. Source: Ben Dov et al. 2006.

| Family            | Genus                   | Species                          | Reference                           | Cited host plant in South Africa                |
|-------------------|-------------------------|----------------------------------|-------------------------------------|---|
| Coccidae          | <i>Ceroplastes</i>      | <i>rusci</i>                     | Brain 1920 (a)                      | <i>Cydonia oblonga</i>                          |
|                   | <i>Coccus</i>           | <i>hesperidum</i>                | Brain 1920 (a), This study          | <i>Citrus, Vitis vinifera</i>                   |
|                   |                         | <i>longulus</i>                  | Brain 1920 (a)                      | <i>Acacia melanoxylon</i>                       |
|                   | <i>Cryptinglisia</i>    | <i>lounsburyi</i> *              | Brain 1920 (a)                      | <i>Vitis vinifera</i>                           |
|                   | <i>Parasaissetia</i>    | <i>nigra</i>                     | Brain 1920 (a)                      | <i>Persea gratissima, Ficus</i>                 |
|                   | <i>Saissetia</i>        | <i>coffaeae</i>                  | Brain 1920 (a)                      | Pot plants                                      |
|                   |                         | <i>oleae</i>                     | Brain 1920 (a)                      | Not specified, Widely distributed               |
|                   | <i>Trijuba</i>          | <i>oculata</i> *                 | Brain 1920 (a)                      | <i>Vitis vinifera</i>                           |
| Diaspididae       | <i>Aonidiella</i>       | <i>aurantii</i>                  | Brain & Kelly 1917                  | Not specified, Widely distributed               |
|                   |                         | <i>orientalis</i>                | Borchenius 1966                     | Various   |
|                   | <i>Aspidiotus</i>       | <i>destructor</i>                | Brain & Kelly 1917, Newstead 1917   | <i>Hevea brasiliensis</i>                       |
|                   |                         | <i>nerii</i>                     | Brain & Kelly 1917                  | Various   |
|                   | <i>Chrysomphalus</i>    | <i>aonidium</i>                  | Brain & Kelly 1917                  | Not specified, Widely distributed               |
|                   |                         | <i>dictyospermi</i>              | Brain & Kelly 1917                  | <i>Cupressus macrocarpa, Chaetacme aristata</i> |
|                   |                         | <i>pinnulifer</i>                | Balachowsky 1956                    | <i>Citrus, Hedera helix</i>                     |
|                   | <i>Diaspidiotus</i>     | <i>osborni</i>                   | Munting 1971                        | <i>Quercus robur</i>                            |
|                   |                         | <i>perniciosus</i>               | Brain & Kelly 1917                  | Various   |
|                   | <i>Hemiberlesia</i>     | <i>lataniae</i>                  | Brain 1918                          | Various   |
|                   |                         | <i>rapax</i>                     | Brain 1918                          | <i>Euonymus, Acacia melanoxylon</i>             |
|                   | <i>Lepidosaphes</i>     | <i>gloverii</i>                  | Brain 1920 (b)                      | <i>Citrus</i>                                   |
|                   |                         | <i>ulmi</i>                      | Balachowsky 1954                    | Various   |
|                   | <i>Neoselenaspis</i>    | <i>silvaticus</i>                | Brain 1918                          | Various   |
|                   | <i>Oceanaspidotus</i>   | <i>spinosus</i>                  | Munting 1965                        | <i>Palmae</i>                                   |
|                   | <i>Parlatoria</i>       | <i>cinerea</i>                   | Morrison 1939                       | Not specified                                   |
|                   |                         | <i>proteus</i>                   | Brain & Kelly 1917                  | Greenhouses                                     |
|                   | <i>Pinnaspis</i>        | <i>strachani</i>                 | Brain 1920 (b)                      | Not specified                                   |
|                   | <i>Pseudaonidia</i>     | <i>trilobitiformis</i>           | Brain 1919                          | <i>Litchi sinensis</i>                          |
|                   | <i>Pseudaulacaspis</i>  | <i>pentagona</i>                 | Balachowsky 1954                    | Not specified                                   |
|                   | <i>Pseudotargionia</i>  | <i>glandulosa</i>                | Brain 1919                          | <i>Acacia horrida</i>                           |
| <i>Selenaspis</i> | <i>articulatus</i>      | Marlatt 1908, Brain & Kelly 1917 | <i>Celastus laurinus</i>            |   |
| Margarodidae      | <i>Icerya</i>           | <i>purchasi</i>                  |                                     | <i>Citrus, Acacia, Rosa</i>                     |
|                   |                         | <i>seychellarum</i>              |                                     | Roses, Ficus, Palms                             |
|                   | <i>Margarodes</i>       | <i>capensis</i> *                | Giard 1897, Brain 1915              | <i>Vitis vinifera</i>                           |
|                   |                         | <i>greeni</i> *                  | Brain 1915                          | <i>Vitis vinifera</i>                           |
|                   |                         | <i>prieskaensis</i> *            | Jakubski 1965                       | <i>Vitis vinifera</i>                           |
|                   | <i>vredendalensis</i> * | De Klerk 1983                    | <i>Vitis vinifera</i>               |   |
| Pseudococcidae    | <i>Dysmicoccus</i>      | <i>brevipes</i>                  | De Villiers et al. 1987             | Pineapple                                       |
|                   | <i>Ferrisia</i>         | <i>virgata</i>                   | De Lotto 1964                       | Agave, <i>Codiaeum</i>                          |
|                   |                         | <i>malvastra</i>                 | Ben-Dov 1994, Walton & Pringle 2004 | <i>Vitis vinifera</i>                           |
|                   | <i>Nipaecoccus</i>      | <i>viridis</i>                   | Brain 1915, This Study              | <i>Acacia, Vitis vinifera</i>                   |
|                   | <i>Planococcus</i>      | <i>citri</i>                     | Ben-Dov 1994                        | <i>Bauvardia, Citrus</i>                        |
|                   |                         | <i>ficus</i>                     | Joubert 1943, This study            | <i>Vitis vinifera</i>                           |
|                   | <i>Pseudococcus</i>     | <i>calceolariae</i>              | Ben-Dov 1994                        | <i>Citrus</i>                                   |
|                   | <i>longispinus</i>      | Walton & Pringle 2004            | <i>Vitis vinifera</i>               |   |

**Table 2** (cont.)

| Family | Genus            | Species         | Reference    | Cited host plant in South Africa |
|--------|------------------|-----------------|--------------|----------------------------------|
|        |                  | <i>viburni</i>  | Ben-Dov 1994 | Various                          |
|        | <i>Rhizoecus</i> | <i>falcifer</i> |              | Ornamental plants                |

Whereas *F. malvastrae* and *N. viridis* are found in all countries importing fruit from South Africa except the USA, both species have been reported from Mexico (Ben-Dov et al. 2006). Only single samples of *N. viridis* and *C. hesperidum* were collected in the survey, suggesting that they are incidental species in vineyards in South Africa. *C. hesperidum* is cosmopolitan and occurs in Australia, China, Egypt, Europe, Guam, Hawaii, India, Iran, Iraq, Israel, Jordan, Mexico, Pakistan, South Africa, and the USA (Ben-Dov 1994; Longo et al. 1995).

This survey confirms results of previous studies which found that *Pl. ficus* is dominant in South African vineyards (Kriegler 1954; De Lotto 1975), whereas *Ps. longispinus* is less abundant. These two cosmopolitan species are now resident in all countries importing grapes from South Africa and, therefore, do not pose a phytosanitary risk (Ben-Dov 1994; Walton and Pringle 2004). However, *F. malvastrae* and *N. viridis* were during the survey and have not been recorded in the USA. Thus, with this exception, we conclude that the risk of exporting exotic scale insect pests to novel geographic regions on table grapes is limited. We also conclude that the multiplex PCR method is an ideal tool for accurately identifying common mealybug species such as *Pl. ficus* so that bulk shipments of table grapes containing these species can be cleared for export, although samples containing other species will need to be identified using morphological techniques.

## Acknowledgments

The authors would like to thank the Deciduous Fruit Producers Trust and Winetech for funding. Thanks to P. Bredell, M. de Villiers, J. de Waal, C. Jelbert, R. Stotter, M. van Straaten, P. Visser, A Vlok, and M. Wolfarther for specimen collection. Special thanks to Drs. P. Gullan and R. Ripa for reviewing previous drafts of this manuscript.

## References

- Balachowsky AS. 1954. Les cochenilles Paléarctiques de la tribu des Diaspidini. Mémoires Scientifiques de l'Institut Pasteur, Paris.
- Balachowsky AS. 1956. Les cochenilles du continent Africain Noir. V. 1 - Aspidiotini (1ère partie). *Annales du Musée Royal du Congo Belge (Sciences Zoologiques), Tervuren* 3: 1-142.
- Ben-Dov Y. 1994. *A systematic catalogue of the mealybugs of the world (Insecta: Homoptera: Coccoidea: Pseudococcidae and Putoidae) with data on geographical distribution, host plants, biology and economic importance*. Intercept Publications LTD, Andover.
- Ben-Dov Y, Miller DR, Gibson GAP. 2006. *ScaleNet: A database of the scale insects of the world*. Available online: <http://www.scl.barc.usda.gov/scalenet/scalenet.htm>.
- Beuning LL, Murphy P, Wu E, Batchelor TA, Morris BA. 1999. Molecular-based approach to the differentiation of mealybug (Hemiptera: Pseudococcidae) species. *Journal of Economic Entomology* 92: 463-472.
- Borchsenius NS. 1966. [A catalogue of the armoured scale insects (Diaspidoidea) of the world] (In Russian). Nauka, Moscow & Leningrad.
- Brain CK. 1915. The Coccidae of South Africa. *Transactions of the Royal Society of South Africa, Cape Town*. 5: 65-194.
- Brain CK. 1918. The Coccidae of South Africa - II. *Bulletin of Entomological Research* 9: 107-139.
- Brain CK. 1919. The Coccidae of South Africa - III. *Bulletin of Entomological Research* 9: 197-239.
- Brain CK. 1920(a). The Coccidae of South Africa - IV. *Bulletin of Entomological Research* 10: 95-128.
- Brain CK. 1920(b). The Coccidae of South Africa - V. *Bulletin of Entomological Research* 11: 1-41.
- Brain CK, Kelly AE. 1917. The status of introduced coccids in South Africa in 1917. *Bulletin of Entomological Research* 8: 181-185.
- Cavaliere V, Mazzeo G, Garzia GT, Buonocore E, Russo A, . 2008. Identification of *Planococcus ficus* and *Planococcus citri* (Hemiptera: Pseudococcidae) by PCR-RFLP of COI gene. *Zootaxa* 1816: 65-68.
- De Klerk CA. 1983. Two new species of *Margarodes* Guilding (Homoptera: Coccoidea: Margarodidae) from South Africa. *Phytophylactica* 15: 85-93.
- De Klerk CA, Ben-Dov Y, Giliomee JH. 1983. General morphology of South African species of *Margarodes* (Homoptera, Coccoidea, Margarodidae) with keys to nymphs and adult females. *Phytophylactica* 15: 133-144.
- De Lotto G. 1964. Observations on African mealy bugs (Hemiptera: Coccoidea). *Bulletin of the British Museum (Natural History) Entomology Series* 14: 343-397.
- De Lotto G. 1975. Notes on the vine mealybug (Homoptera: Coccoidea: Pseudococcidae). *Journal of the Entomological Society of Southern Africa* 38: 125-130.

- Demontis MA, Ortu S, Cocco A, Lentini A, Migheli Q. 2007. Diagnostic markers for *Planococcus ficus* (Signoret) and *Planococcus citri* (Risso) by random amplification of polymorphic DNA-polymerase chain reaction and species specific mitochondrial DNA primers. *Journal of Applied Entomology* 131: 59-64.
- De Villiers EA, Bedford ECG, Petty GT. 1987. Mealybugs and Australian bug. In: Myburgh AC, editor. *Crop Pests in Southern Africa, Vol 2: Citrus and other Subtropicals*. Pretoria, South Africa: Plant Protection Research Institute, Department of Agriculture and Water Supply 115 pp.
- Giard A. 1897. Sur la distribution géographique de cochenilles du genre *Margarodes* et sur deux espèces nouvelles de ce genre. *Comptes Rendus des Séances de la Société de Biologie* 4: 683-685.
- Gullan PJ. 2000. Identification of the immature instars of mealybugs (Hemiptera: Pseudococcidae) found on citrus in Australia. *Australian Journal of Entomology* 39: 160-166.
- Jakubski AW. 1965. *A critical revision of the families Margarodidae and Termitococcidae (Hemiptera, Coccoidea)*. Trustees of the British Museum (Natural History) 187
- Joubert CJ. 1943. Mealybugs on vines. *Bulletin of the Department of Agriculture of South Africa* 24320
- Kriegler PJ. 1954. *Bydrae tot die kennis van Planococcus citri (Risso) (Homoptera: Pseudococcidae)* M.Sc. Thesis, University of Stellenbosch, Stellenbosch, South Africa.
- Longo S, Marotta S, Pellizzari G, Russo A, Tranfaglia A. 1995. An annotated list of the scale insects (Homoptera: Coccoidea) of Italy. *Israel Journal of Entomology* 29: 113-130.
- Marlatt CL. 1908. The national collection of Coccidae. *United States Department of Agriculture, Bureau of Entomology, Technical Series* 16: 1-10.
- Miller DR, Rung A, Venable GL, Gill RJ, Willimas DJ. 2007. *Systematic Entomology Laboratory ARS, USDA Scale Insects Identification Tools for Species of Quarantine Significance* updated 27 August 2007. Available online: <http://www.sel.barc.usda.gov/ScaleKeys/index.html>.
- Morrison H. 1939. Taxonomy of some scale insects of the genus *Parlatoria* encountered in plant quarantine inspection work. *United States Department of Agriculture, Miscellaneous Publications* 344: 1-34.
- Munting J. 1965. New and little known armoured scales (Homoptera: Diaspididae) from South Africa. *Journal of the Entomological Society of Southern Africa* 28 : 2179-216.
- Munting J. 1971. On the identity of the grey scale *Diaspidiotus africanus* Marlatt and some related species from southern Africa (Homoptera: Diaspididae). *Journal of the Entomological Society of Southern Africa* 34 : 1119-143.
- Newstead R. 1917. Observations on scale-insects (Coccidae) – V. *Bulletin of Entomological Research* 8: 125-134.
- Saccaggi DL. 2006. *Development of molecular techniques to identify mealybugs (Hemiptera: Pseudococcidae) of importance on grapevine in South Africa* M.Sc. Thesis, University of Pretoria, Pretoria, South Africa.
- Saccaggi DL, Krüger K, Pietersen G. 2008. A multiplex PCR assay for the simultaneous identification of three mealybug species (Hemiptera: Pseudococcidae). *Bulletin of Entomological Research* 98: 27-33.
- Wakgari WM, Giliomec JH. 2005. Description of adult and immature females of six mealybug species (Hemiptera: Pseudococcidae) found on citrus in South Africa. *African Entomology* 13: 281-332.
- Walton VM. 2003. *Development of an integrated pest management system for vine mealybug, Planococcus ficus (Signoret), in vineyards in the Western Cape Province, South Africa* Ph.D. Thesis, University of Stellenbosch, Stellenbosch, South Africa.
- Walton VM, Pringle KL. 2004. A survey of mealybugs and associated natural enemies in vineyards in the Western Cape Province, South Africa. *South African Journal of Entomology and Viticulture* 25: 23-25.
- Whitehead VB. 1957. *A study of the predators and parasites of Planococcus citri (Risso) (Homoptera) on vines in the Western Cape Province, South Africa* M.Sc. Thesis, Rhodes University, Grahamstown, South Africa.